



Rank-order conjoint experiments: efficiency and design

Bart Vermeulen, Peter Goos and Martina Vandebroek

DEPARTMENT OF DECISION SCIENCES AND INFORMATION MANAGEMENT (KBI)

Rank-order conjoint experiments: efficiency and design

Bart Vermeulen *
Peter Goos †
Martina Vandebroek ‡

Abstract

In a rank-order conjoint experiment, the respondent is asked to rank a number of alternatives instead of choosing the preferred one, as is the standard procedure in conjoint choice experiments. In this paper, we study the efficiency of those experiments and propose a D -optimality criterion for rank-order conjoint experiments to find designs yielding the most precise parameter estimators. For that purpose, an expression of the Fisher information matrix for the rank-ordered multinomial logit model is derived which clearly shows how much additional information is provided by each extra ranking step made by the respondent. A simulation study shows that Bayesian D -optimal ranking designs are slightly better than Bayesian D -optimal choice designs and (near-)orthogonal designs and perform considerably better than other commonly used designs in marketing in terms of estimation and prediction accuracy. Finally, it is shown that improvements of about 50% to 60% in estimation and prediction accuracy can be obtained by ranking a second alternative. If the respondent ranks a third alternative, a further improvement of 30% in estimation and prediction accuracy is obtained.

*Corresponding author. Bart Vermeulen is funded by project G.0611.05 of the Fund for Scientific Research Flanders. Department of Decision Sciences and Information Management, Faculty of Business and Economics, Katholieke Universiteit Leuven, Naamsestraat 69, 3000 Leuven, Belgium, e-mail: bart.vermeulen@econ.kuleuven.be, tel: +32 16 32 69 63, fax: +32 16 32 66 24

†Faculty of Applied Economics, Universiteit Antwerpen

‡Faculty of Business and Economics, University Center for Statistics, Katholieke Universiteit Leuven

1 Introduction

A conjoint choice experiment measures the importance of the features of a good or service in making a purchase decision. This is achieved by asking each respondent to choose his/her preferred alternative from a number of choice sets. A rank-order conjoint experiment measures the importance of the features of a good or service by asking the respondent to rank a certain number of alternatives within the choice sets. Data from a rank-order experiment can be analyzed by the rank-ordered multinomial logit model or exploded logit model (Beggs et al., 1981, Hausman and Ruud, 1987 and Allison and Christakis, 1994), which is an extension of the multinomial logit model, and provides a better view on the preferences of a consumer than data from a choice experiment.

The design of an experiment has a significant impact on the accuracy of the estimated parameters of the fitted model. Choosing the appropriate alternatives and grouping them in choice sets in the best possible way according to an optimality criterion yields an optimal design which guarantees precise parameter estimates and therefore an accurate view on the preferences of the customer. We propose to use the D -optimality criterion which focuses on the accuracy of the estimates of the rank-ordered multinomial logit model its parameters. The central question is then whether the corresponding Bayesian D -optimal ranking design results in significantly more precise estimates and predictions than commonly used design strategies in marketing. Additionally, we value the contribution of each extra ranking step to the estimation and prediction accuracy.

In the next section, we introduce the rank-ordered multinomial logit model. In section 3, we derive the Bayesian ranking D -optimality criterion and present the benchmark designs. In section 4, we compare the design generated using the new criterion with benchmark designs, which are commonly used in marketing, in terms of the newly developed design optimality criterion. Furthermore, in section 4, we use a simulation study to compare the precision of the estimators and the prediction accuracy of the designs. Finally, we examine

how much improvement is obtained by each extra ranking step in terms of estimation and prediction accuracy.

2 The rank-ordered multinomial logit model

Any rank order can be regarded as a sequence of choices made by the respondent. This was used as the starting point for the extension of the multinomial logit model to the rank-ordered multinomial logit model by Beggs et al. (1981). In their approach, each ranking of a choice set is converted into a number of independent pseudo-choices. In this way, each ranking of alternatives in a choice set is considered as a sequential and conditional choice task. The alternative ranked first is perceived as the preferred alternative of the entire choice set. Alternatives ranked next are viewed as the preferred alternatives of the choice sets consisting of all alternatives except the ones with a better ranking. In the resulting rank-ordered multinomial logit model, a ranking of a set of J alternatives is thus seen as a series of $J - 1$ choices.

The rank of an alternative is determined by its utility. The utility of alternative i in choice set k experienced by respondent n is modeled as

$$u_{kin} = \mathbf{x}'_{ki}\boldsymbol{\beta} + \varepsilon_{kin}. \quad (1)$$

The deterministic part, $\mathbf{x}'_{ki}\boldsymbol{\beta}$, of the utility consists of a p -dimensional vector $\boldsymbol{\beta}$, which is assumed to be common for all respondents and which represents the importance of the attributes for the consumer in determining the utility, and a p -dimensional vector \mathbf{x}_{ki} containing the levels of the attributes. The error term ε_{kin} , which forms the stochastic part of the utility, captures the unobserved influences on this utility. The error terms are assumed to be identically and independently distributed according to an extreme value distribution.

The probability of ranking an alternative first in a choice set k with J alternatives corresponds to the logit probability of choosing this alternative as the preferred one. The probability of ranking an alternative second corresponds to the logit probability of choosing this alternative from the profiles remaining after the removal of the first-ranked alternative. This can be said for the $J - 1$ rankings made by the respondent. Because of the assumed independence between these choice tasks, the likelihood of a certain ranking of the alternatives in the entire choice set k is thus the product of J logit probabilities. This likelihood can be written as

$$\prod_{j=1}^J \frac{\exp(\mathbf{x}'_{kj}\boldsymbol{\beta})}{\sum_{j'=1}^J \delta_{kjj'} \exp(\mathbf{x}'_{kj'}\boldsymbol{\beta})}, \quad (2)$$

where the dummy variable $\delta_{kjj'}$ equals zero when alternative j' in choice set k has a better ranking than alternative j , and one otherwise. For a respondent who ranks K choice sets, the likelihood function for the rank-ordered multinomial logit model is given by

$$L(\boldsymbol{\beta}) = \prod_{k=1}^K \prod_{j=1}^J \frac{\exp(\mathbf{x}'_{kj}\boldsymbol{\beta})}{\sum_{j'=1}^J \delta_{kjj'} \exp(\mathbf{x}'_{kj'}\boldsymbol{\beta})}. \quad (3)$$

The maximum likelihood estimate $\hat{\boldsymbol{\beta}}$ for the parameter vector $\boldsymbol{\beta}$ is obtained by maximizing the likelihood function (3).

As the rank-order conjoint experiment requires the respondents to provide more information about their preferences than the classical conjoint choice experiment, it results in more accurate parameter estimates when the same number of choice sets is used. Stated differently, to achieve a desired degree of precision of the estimates, less choice sets are required in a rank-order conjoint experiment (Chapman and Staelin, 1982). The major disadvantage of using a rank-order conjoint experiment is the weak link with reality: in real life, respondents choose the alternative they like most and hardly ever select a second best item. According to Bateman et al. (2002) and Chapman and Staelin (1982), lower

rankings are less reliable if the number of alternatives to rank is large. In this paper, we therefore limit ourselves to three or four alternatives in a choice set.

3 Designs for the rank-ordered multinomial logit model

This section is devoted to the derivation of the Bayesian D -optimality criterion for the rank-ordered multinomial logit model and the corresponding D -error. In addition, we discuss how to create efficient designs using this D -optimality criterion. Finally, benchmark designs, with which the Bayesian D -optimal ranking design will be compared, are introduced.

3.1 Bayesian D -optimal designs for rank-order conjoint experiments

The issue of finding optimal designs for rank-order experiments has not yet been explored in the literature on conjoint experiments. Till now, the applications measuring preferences by means of a rank-order conjoint experiment have used random designs (see, e.g., Boyle et al. (2001)), nearly orthogonal designs (see, e.g., Baarsma (2003)), and fractional factorial designs (Gustafsson et al. (1999)). Although several design selection criteria have been proposed in the literature on experimental designs, none of them has ever been applied to generate an optimal design for a rank-order conjoint experiment. In this article, we show how optimal designs for this type of experiment can be found and we use the D -optimality criterion to do so.

A D -optimal design minimizes the determinant of the variance-covariance matrix on the unknown parameters β and thereby minimizes the volume of the confidence ellipsoids around them. As the variance-covariance matrix is inversely proportional to the Fisher information matrix, a D -optimal design also maximizes the information content of the experiment. In order to construct a D -optimal design for a rank-order conjoint exper-

iment, an expression for the Fisher information matrix on the unknown parameters in the corresponding model is required. Starting from (3), it is not hard to show that the information matrix for an experiment with K choice sets of size three can be written as

$$\begin{aligned} \mathbf{I}_{\text{rank}}(\boldsymbol{\beta}) &= \sum_{k=1}^K \mathbf{X}'_k (\mathbf{P}_k - \mathbf{p}_k \mathbf{p}'_k) \mathbf{X}_k + \sum_{k=1}^K \sum_{i=1}^3 P_{ki1} \mathbf{X}'_{(i)k} (\mathbf{P}_{(i)k} - \mathbf{p}_{(i)k} \mathbf{p}'_{(i)k}) \mathbf{X}_{(i)k} \\ &= \mathbf{I}_{\text{choice}}(\boldsymbol{\beta}) + \sum_{k=1}^K \sum_{i=1}^3 P_{ki1} \mathbf{X}'_{(i)k} (\mathbf{P}_{(i)k} - \mathbf{p}_{(i)k} \mathbf{p}'_{(i)k}) \mathbf{X}_{(i)k}, \end{aligned} \quad (4)$$

where

$$\mathbf{I}_{\text{choice}}(\boldsymbol{\beta}) = \sum_{k=1}^K \mathbf{X}'_k (\mathbf{P}_k - \mathbf{p}_k \mathbf{p}'_k) \mathbf{X}_k \quad (5)$$

denotes the information matrix of a conjoint choice experiment with choice sets of size three (see, e.g., Sándor and Wedel, 2001), and,

- P_{ki1} represents the probability of ranking alternative i ($i = 1, 2, 3$) in choice set k first,
- \mathbf{p}_k is a 3-dimensional vector of which the i^{th} element is P_{ki1} ($i = 1, 2, 3$),
- \mathbf{P}_k is a diagonal matrix with the elements of \mathbf{p}_k on its diagonal,
- $\mathbf{p}_{(i)k}$ is a 2-dimensional vector containing the probabilities of ranking alternative i' second ($i' = 1, 2, 3$ and $i' \neq i$), given that i was ranked first,
- $\mathbf{P}_{(i)k}$ is a diagonal matrix with the elements of $\mathbf{p}_{(i)k}$ on its diagonal,
- \mathbf{X}_k is the $(3 \times p)$ design matrix containing all attribute levels of the profiles in choice set k , and
- $\mathbf{X}_{(i)k}$ is the $(2 \times p)$ design matrix containing all attribute levels of the profiles in choice set k , except those of the first-ranked profile i .

The information matrix for a rank-ordered multinomial logit model corresponding to a rank-order conjoint experiment with choice sets of size three is thus a sum of two

parts: (i) the information matrix corresponding to an ordinary conjoint choice experiment with choice sets of three alternatives, and (ii) a weighted sum of the information matrices corresponding to three ordinary conjoint choice experiments with choice sets of size two. Each of the latter information matrices gives the information content of the choice sets after removing one of the alternatives. The weight of the i^{th} such information matrix is the probability that alternative i is ranked first. The expression for the information matrix of a rank-order experiment proves that asking the respondents to rank the alternatives in a choice set provides extra information. This is because the difference $\mathbf{I}_{\text{rank}}(\boldsymbol{\beta}) - \mathbf{I}_{\text{choice}}(\boldsymbol{\beta}) = \sum_{k=1}^K \sum_{i=1}^3 P_{ki1} \mathbf{X}'_{(i)k} (\mathbf{P}_{(i)k} - \mathbf{p}_{(i)k} \mathbf{p}'_{(i)k}) \mathbf{X}_{(i)k}$ is a nonnegative definite matrix, which ensures that the amount of information in a ranking experiment is larger than in a choice experiment. In other words, for a given choice set, it is always better to rank. By means of equation (4), the extra amount of information can be quantified easily.

It is possible to extend equation (4) to the case of ranking more than three alternatives per choice set. If the respondent ranks all alternatives in K choice sets of size four, the expression for the information matrix is given by

$$\begin{aligned}
\mathbf{I}_{\text{rank}}(\boldsymbol{\beta}) &= \mathbf{I}_{\text{choice}}(\boldsymbol{\beta}) \\
&+ \sum_{k=1}^K \sum_{i=1}^4 P_{ki1} \mathbf{X}'_{(i)k} (\mathbf{P}_{(i)k} - \mathbf{p}_{(i)k} \mathbf{p}'_{(i)k}) \mathbf{X}_{(i)k} \\
&+ \sum_{k=1}^K \sum_{i=1}^4 \sum_{i'=1, i' \neq i}^4 (P_{ki1} P_{ki'2} + P_{ki'1} P_{ki2}) \mathbf{X}'_{(ii')k} (\mathbf{P}_{(ii')k} - \mathbf{p}_{(ii')k} \mathbf{p}'_{(ii')k}) \mathbf{X}_{(ii')k},
\end{aligned} \tag{6}$$

where

- $\mathbf{I}_{\text{choice}}(\boldsymbol{\beta})$ denotes the information matrix of a conjoint choice experiment with choice sets of size four,
- P_{ki1} represents the probability of ranking alternative i ($i = 1, 2, 3, 4$) in choice set k

first,

- \mathbf{p}_k is a 4-dimensional vector of which the i^{th} element is P_{ki1} ($i = 1, 2, 3, 4$),
- \mathbf{P}_k is a diagonal matrix with the elements of \mathbf{p}_k on its diagonal,
- $P_{ki'2}$ represents the probability of ranking alternative i' ($i' = 1, 2, 3, 4$ and $i' \neq i$) in choice set k second given that i was ranked first,
- $\mathbf{p}_{(i)k}$ is a 3-dimensional vector of which the i^{th} element is $P_{ki'2}$ ($i' = 1, 2, 3, 4$ and $i' \neq i$),
- $\mathbf{P}_{(i)k}$ is a diagonal matrix with the elements of $\mathbf{p}_{(i)k}$ on its diagonal,
- $\mathbf{p}_{(ii')k}$ is a 2-dimensional vector containing the probabilities of ranking alternative i'' third ($i'' = 1, 2, 3, 4$, $i'' \neq i$ and $i'' \neq i'$) given that i and i' have been ranked first and second or vice versa,
- $\mathbf{P}_{(ii')k}$ is a diagonal matrix with the elements of $\mathbf{p}_{(ii')k}$ on its diagonal,
- \mathbf{X}_k is the $(4 \times p)$ design matrix containing all attribute levels of the profiles in choice set k ,
- $\mathbf{X}_{(i)k}$ is the $(3 \times p)$ design matrix containing all attribute levels of the profiles in choice set k , except those of the first-ranked profile i , and
- $\mathbf{X}_{(ii')k}$ is the $(2 \times p)$ design matrix containing all attribute levels of the profiles in choice set k , except those of the first- and second-ranked profiles i and i' .

Expression (6) illustrates again that asking respondents to rank the alternatives of a choice set provides the researcher with more information than a choice experiment.

Ideally, the information content of the rank-order experiment is as large as possible. In the experimental design literature, the most commonly used measure for the information content of a design is the determinant of the information matrix. A design that maximizes

this quantity is called D -optimal. A D -optimal design can also be found by minimizing the p^{th} root of the determinant of the inverse of the information matrix. This criterion is usually referred to as the D -error and is defined as

$$D\text{-error} = \{\det(\mathbf{I}_{\text{rank}}(\mathbf{X}, \boldsymbol{\beta})^{-1})\}^{\frac{1}{p}}. \quad (7)$$

In this article, we will use the D -error to quantify the performance of a design in the D -optimality criterion. The design having the smallest D -error is then the D -optimal design.

As can be seen in (4), (6) and (7), the information matrix and the D -error depend on the parameters in $\boldsymbol{\beta}$ which are unknown at the moment of designing the experiment. To circumvent this problem, we use the Bayesian approach of Sándor and Wedel (2001, 2005) and Kessels et al. (2006a) assuming a prior distribution $f(\boldsymbol{\beta})$ on the unknown parameters for the purpose of designing a rank-order experiment. The Bayesian version of the D -error is given by

$$D_b\text{-error} = E_{\boldsymbol{\beta}}[\{\det(\mathbf{I}_{\text{rank}}(\mathbf{X}, \boldsymbol{\beta})^{-1})\}^{\frac{1}{p}}] = \int_{\mathfrak{R}^p} \{\det(\mathbf{I}_{\text{rank}}(\mathbf{X}, \boldsymbol{\beta})^{-1})\}^{\frac{1}{p}} f(\boldsymbol{\beta}) d\boldsymbol{\beta}. \quad (8)$$

Since there is no analytical way to compute the integral in (8), it has to be approximated by taking a number of draws from the prior distribution $f(\boldsymbol{\beta})$ and averaging the D -error over all draws. The design which minimizes the average D -error over all draws is called the Bayesian D -optimal ranking design.

The designs discussed in this paper involve three three-level and two two-level attributes which are effects-type coded. This implies that the number of parameters, p , contained within $\boldsymbol{\beta}$, equals 8. As a Bayesian approach is used to circumvent the dependence of the optimal design on the unknown parameter vector $\boldsymbol{\beta}$, a prior distribution on $\boldsymbol{\beta}$ has to be specified. We used as prior distribution an 8-dimensional normal distribution with mean $[-0.5; 0; -0.5; 0; -0.5; 0; -0.5; -0.5]$ and variance $0.5 * \mathbf{I}_8$ with \mathbf{I}_8 being the

8-dimensional unity matrix. This prior distribution, which follows the recommendations formulated by Kessels et al. (2006c), assumes the utility of a good or service increases with the attribute levels.

To construct a Bayesian D -optimal ranking design, we implemented the adaptive algorithm developed by Kessels et al. (2006b), using a systematic sample of 100 draws from the prior distribution $f(\beta)$. This systematic sample was generated using a Halton sequence (Train, 2003).

3.2 Benchmark designs

In order to evaluate the performance of the Bayesian D -optimal ranking designs, we compared them to four types of benchmark designs: Bayesian D -optimal designs for a conjoint choice experiment (Kessels et al. (2006a) describe extensively how to generate these designs) and three sorts of standard designs generated by Sawtooth Software. This software offers the user three options to construct designs: (near-)orthogonal designs (labelled 'complete enumeration' in Sawtooth), random designs and designs with a balanced attribute level overlap in one choice set (denoted as 'balanced overlap' in Sawtooth). To construct a (near-)orthogonal design, all candidate profiles obtained by all possible combinations of attribute levels are considered. The alternatives are selected and grouped in choice sets so that the attribute level overlap within one choice set is minimal and so that the design is as close as possible to an orthogonal design. The random method leads to alternatives with randomly drawn attribute levels and possibly results in a strong attribute level overlap within a choice set. Contrary to the (near-)orthogonal design which severely restricts the overlap of attribute levels within a choice set, the third option, balanced overlap, allows a moderate attribute level overlap. In this sense, this option is the middle course between the orthogonal design and the random design.

4 Evaluating the Bayesian D -optimal ranking design

In this section, we compare the different designs in terms of the D_b -error and describe the findings of a simulation study by which the performance of the designs in terms of estimation and prediction accuracy was examined. In addition, we measure the improvement achieved by including extra ranking steps in the experiment. In the simulation study, we considered two cases: ranking nine choice sets of size four and ranking twelve choice sets of size three. Each of the cases assumes 50 respondents participated in the experiment. As an illustration, the Bayesian D -optimal ranking design and the four benchmark designs for the case of nine choice sets of size four are included in appendix.

4.1 Measurement of the estimation and prediction accuracy

Based on simulated observations for all choice sets of the different designs discussed above, we estimated the parameters of the rank-ordered multinomial logit model and used the estimates to calculate the probabilities of choosing an alternative of a choice set. Comparing the estimated parameters with their true values, i.e. the ones used to generate the data, and comparing the estimated probabilities with the probabilities calculated using the true parameter values allows us to evaluate the estimation and prediction accuracy, respectively. The metrics we used for that purpose are the expected mean squared errors for estimation and prediction, denoted by $EMSE_{\hat{\beta}}(\beta)$ and $EMSE_{\hat{p}}(\beta)$, and are computed as

$$EMSE_{\hat{\beta}}(\beta) = \int_{\mathbb{R}^p} (\hat{\beta} - \beta)' (\hat{\beta} - \beta) f(\hat{\beta}) d\hat{\beta}, \quad (9)$$

and

$$EMSE_{\hat{p}}(\beta) = \int_{\mathbb{R}^p} (\hat{p}(\hat{\beta}) - p(\beta))' (\hat{p}(\hat{\beta}) - p(\beta)) f(\hat{\beta}) d\hat{\beta}, \quad (10)$$

where $f(\hat{\beta})$ is the distribution of the parameter estimates, β and $\hat{\beta}$ are the vectors containing the true and estimated parameter values and $p(\beta)$ and $\hat{p}(\hat{\beta})$ are vectors containing the true and the predicted probabilities of choosing an alternative of a choice set in a random design. For the results reported in this paper, (9) and (10) were approximated

using 1000 draws from $f(\hat{\boldsymbol{\beta}})$. This was done for 75 values of $\boldsymbol{\beta}$ drawn from the prior distribution used to generate the Bayesian D -optimal ranking design. The smaller the value of $EMSE_{\hat{\boldsymbol{\beta}}}(\boldsymbol{\beta})$ and $EMSE_{\hat{\mathbf{p}}}(\boldsymbol{\beta})$, the more accurate the parameter estimators and predictions respectively.

4.2 Performance of Bayesian D -optimal ranking designs

The next section considers the scenarios when the respondents do and do not rank all alternatives of a choice set. We refer to a rank-order experiment where all alternatives in a set are ranked as a full rank-order experiment. An experiment where only the subset of the better alternatives of each choice set is ranked, is called a partial rank-order experiment.

4.2.1 Full rank-order experiments

In this section, we examine whether a Bayesian D -optimal ranking design performs better in terms of estimation and prediction accuracy than the benchmark designs in case of a full rank-order experiment. Table 1 compares the five different designs in terms of the D_b -error calculated for 100 values of $\boldsymbol{\beta}$ drawn from the prior distribution. This table displays a slightly smaller D_b -error for the Bayesian D -optimal ranking design than for the Bayesian D -optimal choice design. The difference between the D_b -errors for the D -optimal ranking and the random design and the difference between the former design and the design with balanced attribute level overlap are considerably larger, which suggests a poor estimation performance of the random and balanced design.

The accuracies of the estimated parameters and the predicted probabilities, as measured by the $EMSE_{\hat{\boldsymbol{\beta}}}(\boldsymbol{\beta})$ and $EMSE_{\hat{\mathbf{p}}}(\boldsymbol{\beta})$, are shown in Figures 1 to 4 for both cases considered. As expected, the Bayesian D -optimal ranking design results in the most precise estimates compared to the four other designs, which can be seen in Figures 1 and 3. The difference with the design characterized by balanced attribute level overlap and with the random design is considerably larger than the difference with the Bayesian choice design and

	D_b -error when ranking the complete choice set	
	Choice set size = 4	Choice set size = 3
D -opt. rank.	0.1181	0.1458
D -opt. choice	0.1210	0.1467
Balanced overlap	0.1439	0.1755
(Near-)Orthogonal	0.1301	0.1629
Random	0.1421	0.2276

Table 1: D_b -errors when ranking nine choice sets of four alternatives and twelve choice sets of three alternatives for the Bayesian D -optimal ranking design, the Bayesian D -optimal choice design and three standard designs




Figure 1: Estimation accuracy, as measured by $EMSE_{\hat{\beta}}(\beta)$, for the five different designs if the respondents rank all four alternatives of nine choice sets

the (near-)orthogonal design. Figures 2 and 4 reveal that the smallest predictive error is achieved also by using the Bayesian ranking design closely followed by the Bayesian choice design and the (near-)orthogonal design.




Figure 2: Prediction accuracy, as measured by $EMSE_{\hat{\beta}}(\beta)$, for the five different designs if the respondents rank all four alternatives of nine choice sets




Figure 3: Estimation accuracy, as measured by $EMSE_{\hat{\beta}}(\beta)$, for the five different designs if the respondents rank all three alternatives of twelve choice sets




Figure 4: Prediction accuracy, as measured by $EMSE_{\hat{\beta}}(\beta)$, for the five different designs if the respondents rank all three alternatives of twelve choice sets

4.2.2 Partial rank-order experiments

So far, we have assumed that each respondent completes the ranking of all alternatives in a choice set. In practice, it can happen that respondents are asked to rank only the subset of the best alternatives in a choice set. That is why we discuss in this section the partial rank-order experiment. Remark that the partial rank-order experiment where the respondents rank only one alternative of each choice set corresponds to a conjoint choice experiment.

Table 2 illustrates that the Bayesian D -optimal choice design outperforms the Bayesian D -optimal ranking design and the three standard designs in terms of the D_b -error. However, the difference between the Bayesian ranking and choice design is rather small. Compared to the scenario where the respondents rank the two best alternatives of nine choice sets of size four, the choice of only the preferred alternative leads to a slightly larger difference in D_b -error between the ranking and choice design.

	Choice set size = 4		Choice set size = 3
	D_b -error for choice experiment	D_b -error for ranking the two best alternatives	D_b -error for choice experiment
D -opt. rank.	0.3446	0.1755	0.2830
D -opt. choice	0.3242	0.1718	0.2733
Balanced overlap	0.4188	0.2108	0.3417
(Near-)Orthogonal	0.3668	0.1966	0.345
Random	0.4159	0.2045	0.4142

Table 2: D_b -errors corresponding to a partial rank-order experiment and a choice experiment with nine choice sets of four alternatives and twelve choice sets of three alternatives for the Bayesian D -optimal ranking design, the Bayesian D -optimal choice design and three standard designs

Figures 5 to 8 display the accuracy of the parameter estimates and predictions respectively obtained by using the different designs in case of a partial rank-order experiment.




Figure 5: Estimation accuracy, as measured by $EMSE_{\hat{\beta}}(\beta)$, for the five different designs if the respondents rank two of four alternatives in nine choice sets




Figure 6: Prediction accuracy, as measured by $EMSE_{\hat{\beta}}(\beta)$, for the five different designs if the respondents rank two of four alternatives in nine choice sets




Figure 7: Estimation accuracy, as measured by $EMSE_{\hat{\beta}}(\beta)$, for the five different designs if the respondents choose their preferred alternative in twelve choice sets of size three




Figure 8: Prediction accuracy, as measured by $EMSE_{\hat{\beta}}(\beta)$, for the five different designs if the respondents choose their preferred alternative in twelve choice sets of size three

Figures 5 and 7 show that the Bayesian D -optimal choice design leads to the most precise parameter estimates, although it is closely followed by the Bayesian ranking design. Furthermore, Figures 5 and 7 visualize the poor performance of the random design and the design with balanced attribute level overlap. Figure 6 and 8 allow us to draw similar conclusions for the predictive accuracy.

The scenario where the respondents rank only one of the four alternatives in nine choice sets, which is not visualized in a figure, leads to a slightly larger difference between the Bayesian ranking and choice design in estimation and prediction accuracy, in terms of $EMSE_{\hat{\beta}}(\beta)$ and $EMSE_{\hat{p}}(\beta)$.

In conclusion, if the respondents rank all alternatives in each of the choice sets, the Bayesian D -optimal ranking design leads to only slightly better parameter estimates and predictions compared to the Bayesian D -optimal choice design and the orthogonal design

but does considerably better than the other standard designs used in marketing. Even when the respondents do not rank all alternatives, we notice that the Bayesian D -optimal ranking design results in more precise estimates and predictions than the random design and the design with balanced attribute level overlap and yields only slightly less precise estimates and predictions than the Bayesian D -optimal choice design. This leads to the conclusion that a D -optimal choice design is also an appropriate design for the rank-order conjoint experiment. A striking feature of the optimal designs compared to the standard designs is that they never lead to very imprecise estimates or predictions. This can be seen best from Figures 3, 5, 7 and 8 where the three standard designs all produce very large $EMSE_{\hat{\beta}}(\boldsymbol{\beta})$ and $EMSE_{\hat{p}}(\boldsymbol{\beta})$ values occasionally. This is unlike the optimal designs and confirms earlier results by Woods et al. (2006) in the context of industrial experiments involving non-normal responses.

We were able to draw similar conclusions for full and partial rank-order experiments with two three-level attributes and one two-level attribute with nine and twelve choice sets of size four and three, respectively.

4.2.3 The improvement in estimation and prediction accuracy by including extra ranking steps

Tables 3 and 4 exhibit the benefits in terms of precision of the parameter estimates and predicted probabilities by including an extra ranking step in case of nine choice sets of size four.

Table 3 shows that there is a 60% improvement in estimation accuracy if the respondents indicate which alternatives they like second best, in addition to the one they like best. This implies that ranking two of four alternatives in nine choice sets requires 38% less respondents than a choice experiment in order to have the same precision of parameter estimates. If the respondents rank also the third alternative of four in each choice set (and thus provide a full ranking of each choice set), a further improvement of some 30% is

obtained. This means that the number of respondents can be reduced further by another 23% to obtain the same estimation accuracy as the scenario where the respondents only rank two of four alternatives. The $EMSE_{\hat{\beta}}(\beta)$ values in Table 4 show that we can extend these conclusions to the prediction accuracy. An accuracy gain of about 60% is acquired by ranking a second alternative. If the respondents rank the full choice set, the predictions are about 25% more accurate compared to the situation where the respondents rank only two alternatives.

	Improvement in $EMSE_{\hat{\beta}}$	
	after assigning second rank	after assigning third rank
D -opt. rank.	62.29%	30.45%
D -opt. choice	58.07%	23.09%
Balanced overlap	63.07%	34.23%
(Near-)Orthogonal	64.78%	31.35%
Random	61.69%	30.70%

Table 3: Improvement in the accuracy of the estimators of the model parameters, as measured by $EMSE_{\hat{\beta}}(\beta)$, in case of a full and partial rank-order experiment with choice sets of size four for the five different designs

	Improvement in $EMSE_{\hat{p}}$	
	after assigning second rank	after assigning third rank
D -opt. rank.	57.96%	30.73%
D -opt. choice	54.75%	23.70%
Balanced overlap	58.41%	31.66%
(Near-)Orthogonal	58.57%	27.79%
Random	66.72%	27.80%

Table 4: Improvement in the accuracy of the predictions, as measured by $EMSE_{\hat{p}}(\beta)$, in case of a full and partial rank-order conjoint experiment with choice sets of size four for the five different designs

For the case of twelve choice sets with three alternatives each, represented in Table 5, we can draw similar conclusions: if the respondents rank the full choice set, an improvement

of about 50% is obtained in terms of estimation and prediction accuracy.

	Improvement in	
	$EMSE_{\hat{\beta}}$	$EMSE_{\hat{p}}$
D -opt. rank.	54.91%	52.45%
D -opt. choice	50.12%	48.02%
Balanced overlap	53.59%	49.26%
(Near-)Orthogonal	54.47%	52.71%
Random	49.06%	46.37%

Table 5: Improvement in the accuracy of the estimators of the model parameters and predictions, as measured by $EMSE_{\hat{\beta}}(\beta)$ and $EMSE_{\hat{p}}(\beta)$, in case of a full and rank-order experiment with choice sets of size three for the five different designs

Because of those considerable improvements in estimation and prediction error, it is worthwhile to use rank-order conjoint experiments instead of conjoint choice experiments.

5 Conclusion

In a rank-order conjoint experiment, the respondent is asked to rank a number of alternatives in each choice set. This way of performing a conjoint experiment offers the important advantage that extra information is extracted about the preferences of the respondent which results in better estimated part-worths and better predicted probabilities.

Although rank-ordered conjoint experiments are a popular instrument to measure consumer preferences, the literature on experimental design for these experiments is very scarce. In this paper, we try to fill this gap by proposing a D -optimality criterion focusing on the precision of the estimates for the parameters of the ranked-ordered multinomial logit model. To achieve this goal, the Fisher information matrix for this model was given. This matrix clearly shows the additional information yielded by each extra ranking step and is used to construct Bayesian D -optimal ranking designs. We compared these designs with four benchmarks and concluded that the Bayesian D -optimal ranking design

performs slightly better in terms of the D -optimality criterion than its competitors if all alternatives of a choice set are ranked. It is, however, closely followed by the Bayesian D -optimal choice design.

We also used a simulation study to compare the Bayesian D -optimal ranking designs with the four benchmark designs in terms of the estimation and predictive accuracy. We conclude that the Bayesian D -optimal ranking designs result in the most accurate estimates and predictions if the respondents rank all alternatives. It should be pointed out, however, that the Bayesian ranking and choice designs exhibit only minor differences. This allows us to conclude that the Bayesian D -optimal conjoint choice designs are also appropriate designs for rank-order conjoint experiments.

When examining the improvements in the precision of the estimates and predictions yielded by each extra ranking step, we observed that a major gain in estimation and predictive accuracy is achieved by including a second ranking step. Including a third ranking step still offers a considerable improvement in estimation and predictive accuracy. Based on these substantial improvements in efficiency, we strongly recommend the use of rank-order conjoint experiments instead of conjoint choice experiments.

6 Acknowledgments

We are grateful to Qi Ding for his contribution to the development of the adaptive algorithm. He was funded by project NB/06/05 of the National Bank of Belgium.

Appendix

I. Bayesian D -optimal ranking design

Choice set	Profile	Attr. 1	Attr. 2	Attr. 3	Attr. 4	Attr. 5
1	1	3	1	3	1	1
	2	3	3	2	1	2
	3	1	2	1	2	2
	4	2	1	2	2	1
2	1	3	2	1	2	2
	2	2	2	3	1	1
	3	2	3	2	1	2
	4	1	1	2	2	1
3	1	3	1	1	2	2
	2	1	3	1	1	1
	3	1	2	3	1	2
	4	2	2	2	2	1
4	1	1	3	2	1	2
	2	1	2	3	2	1
	3	3	2	1	1	1
	4	2	1	1	2	2
5	1	3	2	2	1	1
	2	1	2	2	2	2
	3	2	1	3	1	2
	4	3	3	1	2	1
6	1	2	3	1	1	2
	2	3	1	3	2	1
	3	1	1	2	1	2
	4	2	2	2	2	1
7	1	2	2	1	2	1
	2	1	1	3	1	2
	3	1	3	3	2	1
	4	3	2	2	1	2
8	1	2	2	3	1	2
	2	1	1	1	2	2
	3	3	3	2	2	1
	4	2	3	1	1	1
9	1	2	1	2	2	2
	2	1	2	1	1	2
	3	1	3	3	2	1
	4	3	1	2	1	1

II. Bayesian D -optimal choice design

Choice set	Profile	Attr. 1	Attr. 2	Attr. 3	Attr. 4	Attr. 5
1	1	3	2	1	2	1
	2	2	2	2	1	1
	3	1	3	3	1	2
	4	2	1	2	2	2
2	1	2	3	2	2	2
	2	3	2	1	1	2
	3	1	2	3	2	1
	4	3	1	3	1	1
3	1	1	1	3	2	2
	2	3	2	2	1	2
	3	2	1	3	1	1
	4	2	3	1	2	1
4	1	2	1	2	1	1
	2	1	2	2	2	2
	3	3	3	1	1	2
	4	2	1	3	2	1
5	1	1	1	3	1	2
	2	3	3	2	1	1
	3	1	1	2	2	1
	4	2	2	1	2	2
6	1	3	1	3	1	2
	2	2	2	1	2	1
	3	1	2	1	1	2
	4	1	3	2	2	1
7	1	2	2	2	1	1
	2	3	1	1	2	2
	3	2	3	3	1	1
	4	1	2	2	1	2
8	1	1	1	1	2	2
	2	2	1	2	1	2
	3	2	3	1	1	2
	4	3	2	3	2	1
9	1	2	2	3	1	2
	2	3	1	2	2	1
	3	1	1	1	1	2
	4	1	3	1	2	1

III. (Near-)Orthogonal design

Choice set	Profile	Attr. 1	Attr. 2	Attr. 3	Attr. 4	Attr. 5
1	1	2	1	3	2	1
	2	3	2	2	1	1
	3	3	3	2	2	2
	4	1	3	1	1	2
2	1	1	1	3	1	1
	2	2	2	1	2	2
	3	2	2	3	1	2
	4	3	3	2	2	1
3	1	1	1	1	2	1
	2	2	2	2	1	1
	3	1	1	1	2	2
	4	3	3	3	1	2
4	1	2	1	2	1	2
	2	3	3	1	1	1
	3	1	2	3	2	1
	4	3	1	3	2	2
5	1	1	2	2	2	2
	2	3	1	3	1	2
	3	2	3	1	1	1
	4	2	3	2	2	1
6	1	3	2	1	2	2
	2	2	3	3	2	2
	3	3	1	2	1	1
	4	1	2	1	1	1
7	1	2	1	1	2	1
	2	1	1	2	1	2
	3	3	2	3	2	1
	4	1	3	2	1	2
8	1	2	2	1	1	2
	2	1	3	3	2	1
	3	3	1	2	2	2
	4	1	2	3	1	1
9	1	3	1	3	1	1
	2	1	2	2	2	2
	3	2	3	1	2	2
	4	2	3	1	1	1

IV. Balanced overlap design

Choice set	Profile	Attr. 1	Attr. 2	Attr. 3	Attr. 4	Attr. 5
1	1	3	2	2	2	2
	2	2	1	1	1	1
	3	1	2	1	1	1
	4	1	3	3	2	2
2	1	3	2	3	1	2
	2	2	3	2	2	1
	3	2	1	3	2	1
	4	1	1	1	1	2
3	1	2	2	2	2	1
	2	3	3	2	1	2
	3	3	1	3	2	2
	4	3	3	1	1	1
4	1	3	3	1	2	2
	2	1	2	2	1	1
	3	2	3	1	2	2
	4	3	1	3	1	1
5	1	2	2	2	1	2
	2	1	3	1	2	2
	3	2	3	3	1	1
	4	1	1	2	2	1
6	1	1	1	3	1	2
	2	2	2	1	2	2
	3	1	1	1	1	1
	4	3	2	2	2	1
7	1	3	2	3	2	2
	2	2	3	3	1	1
	3	1	2	3	2	1
	4	1	1	2	1	2
8	1	2	1	3	1	2
	2	3	1	2	1	1
	3	2	3	1	2	2
	4	1	2	3	2	1
9	1	3	3	1	1	1
	2	1	3	1	1	2
	3	2	1	2	2	1
	4	3	2	2	2	2

V. Random design

Choice set	Profile	Attr. 1	Attr. 2	Attr. 3	Attr. 4	Attr. 5
1	1	2	2	3	2	2
	2	2	1	3	2	2
	3	2	2	1	1	2
	4	1	3	3	2	1
2	1	3	2	3	1	2
	2	1	3	3	2	2
	3	3	3	2	1	1
	4	1	3	1	2	1
3	1	2	3	2	1	2
	2	1	3	1	1	1
	3	1	2	2	1	1
	4	1	2	1	2	2
4	1	2	2	3	1	1
	2	1	1	3	1	1
	3	1	3	3	1	2
	4	2	2	2	2	1
5	1	1	3	3	1	2
	2	2	1	2	1	2
	3	3	3	3	2	2
	4	1	2	3	1	2
6	1	2	3	2	2	2
	2	2	1	1	1	1
	3	2	1	3	2	2
	4	2	2	3	1	2
7	1	2	1	2	2	2
	2	1	2	1	1	2
	3	2	1	3	1	1
	4	3	1	1	2	2
8	1	1	3	1	2	2
	2	1	1	3	2	1
	3	2	1	1	1	2
	4	2	3	2	2	1
9	1	2	3	2	1	1
	2	1	3	2	1	1
	3	1	1	3	2	1
	4	3	3	2	2	1

References

- Allison, P.D. and Christakis, N.A. (1994), Logit models for sets of ranked items, *Sociological Methodology*, 24, 199-228
- Baarsma, B.E. (2003), The valuation of the Ijmeer Nature Reserve using conjoint analysis, *Environmental and Resource Economics*, 25, 343-356
- Bateman, I.J., Carson, R.T., Day, B., Hanemann, M., Hanley, N., Hett, J., Jones-Lee, M., Loomes, G., Mourato, S., Özdemiroglu, E., Pearce, D.W., Sugden, R. and Swanson, J. (2002), *Economic valuation with stated preference techniques. A Manual*, Cheltenham, Edward Elgar
- Beggs, S., Cardell, S. and Hausman, J. (1981), Assessing the potential demand for electric cars, *Journal of Econometrics*, 16, 1-19
- Boyle, K.J., Holmes, T.P., Teisl, M.F. and Roe, B. (2001), A comparison of conjoint analysis response formats, *American Journal of Agricultural Economics*, 83, 441-454
- Chapman, R.G. and Staelin, R. (1982), Exploiting rank ordered choice set data within the stochastic utility model, *Journal of Marketing Research*, 19, 288-301
- Gustafsson, A., Ekdahl, F. and Bergman, B. (1999), Conjoint analysis: a useful tool in the design process, *Total Quality Management*, 10, 327-343
- Hausman, J.A. and Ruud, P.A. (1987), Specifying and testing econometric models for rank-ordered data, *Journal of Econometrics*, 34, 83-104
- Kessels, R., Goos, P. and Vandebroek, M. (2006a), A comparison of criteria to design efficient choice experiments, *Journal of Marketing Research*, 43, 409-419
- Kessels, R., Jones, B., Goos, P. and Vandebroek, M. (2006b), An efficient algorithm for constructing Bayesian optimal choice designs, Research Report KBI_0616, Department of Decision Sciences and Information Management, Katholieke Universiteit Leuven, 29pp.
- Kessels, R., Jones, B., Goos, P. and Vandebroek, M. (2006c), Recommendations on the use of Bayesian optimal designs for choice experiments, Research Report KBI_0617, Department of Decision Sciences and Information Management, Katholieke Universiteit Leuven, 18pp.
- Sándor, Z. and Wedel, M. (2001), Designing conjoint choice experiments using manager's prior beliefs, *Journal of Marketing Research*, 38, 430-444
- Sándor, Z. and Wedel, M. (2005), Heterogeneous conjoint choice designs, *Journal of Mar-*

keting Research, 42, 210-218

Train, K. (2003), *Discrete Choice Methods with Simulations*, Cambridge University Press

Woods, D.C., Lewis, S.M., Eccleston, J.A. and Russel, K.G. (2006), Designs for generalized linear models with several variables and model uncertainty, *Technometrics*, 48, 284-292