***Citation**

Achten WMJ, Verchot L, Franken YJ, Mathijs E, Singh VP, Aerts R, Muys B 2008. *Jatropha* bio-diesel production and use. Biomass and Bioenergy 32(12), 1063-1084

DOI: [10.1016/j.biombioe.2008.03.003](https://doi.org/10.1016/j.biombioe.2008.03.003)

Authors: W.M.J. Achten^a, L. Verchot^b, Y.J. Franken^c, E. Mathijs^d, V.P. Singh^e, R. Aerts^a and B. Muys^{a,*}

^a Katholieke Universiteit Leuven, Division Forest, Nature and Landscape, Celestijnenlaan 200 E Box 2411, BE-3001 Leuven, Belgium

^b World Agroforestry Centre (ICRAF) Head Quarters , United Nations Avenue, P.O. Box 30677, Nairobi, Kenya

^c FACT Foundation, Horsten 1, 5612 AX Eindhoven, The Netherlands

^d Katholieke Universiteit Leuven, Division Agricultural and Food Economics, Willem de Croylaan 42 Box 2424, BE- 3001 Leuven, Belgium

^e World Agroforestry Centre (ICRAF) Regional office for South Asia, CG Block, 1st Floor, National Agricultural Science Centre, Dev Prakash Shastri Marg, Pusa, New Delhi - 110 012, India

* Corresponding author. Tel.: +32 (0) 16 329726; Fax: +32 (0) 16 329760; E-mail address: bart.muys@biw.kuleuven.be (B. Muys)

*Elsevier authorized the authors to post this revised personal version of the text of the final journal article on the author's institutional server, incorporating the complete citation and with a link to the Digital Object Identifier (DOI) of the article.

Creative Commons Attribution-Noncommercial-Share Alike 2.0 Belgium License

Permissions beyond the scope of this license may be available at

<http://www.elsevier.com/wps/find/authorsview.authors/copyright#whatrights>

Title

Jatropha bio-diesel production and use

Authors

W.M.J. Achten^a, L. Verchot^b, Y.J. Franken^c, E. Mathijs^d, V.P. Singh^e, R. Aerts^a and B. Muys^{a,*}

Affiliations

^a Katholieke Universiteit Leuven, Division Forest, Nature and Landscape, Celestijnenlaan 200 E Box 2411, BE-3001 Leuven, Belgium

^b World Agroforestry Centre (ICRAF) Head Quarters, United Nations Avenue, P.O. Box 30677, Nairobi, Kenya

^c FACT Foundation, Horsten 1, 5612 AX Eindhoven, The Netherlands

^d Katholieke Universiteit Leuven, Division Agricultural and Food Economics, Willem de Croylaan 42 Box 2424, BE- 3001 Leuven, Belgium

^e World Agroforestry Centre (ICRAF) Regional office for South Asia, CG Block, 1st Floor, National Agricultural Science Centre, Dev Prakash Shastri Marg, Pusa, New Delhi - 110 012, India

* Corresponding author. Tel.: +32 (0) 16 329726; Fax: +32 (0) 16 329760;

E-mail address: bart.muys@biw.kuleuven.be (B. Muys)

Abstract

The interest in using *Jatropha curcas* L. (JCL) as a feed stock for the production of bio-diesel is rapidly growing. The properties of the crop and its oil have persuaded investors, policy makers and Clean Development Mechanism (CDM) project developers to consider JCL as a substitute for fossil fuels to reduce greenhouse gas emissions. However, JCL is still a wild plant of which basic agronomic properties are not thoroughly understood and the environmental effects have not been investigated yet. Gray literature reports are very optimistic on simultaneous wasteland reclamation capability and oil yields, further fueling the *Jatropha* bio-diesel hype. In this paper, we give an overview of the currently available information on the different process steps of the production process of bio-diesel from JCL, being cultivation and production of seeds, extraction of the oil, conversion to and the use of the bio-diesel and the by-products. Based on this collection of data and information the best available practice, the shortcomings and the potential environmental risks and benefits are discussed for each production step. The review concludes with a call for general precaution and for science to be applied.

Keywords

Jatropha curcas; physic nut; cultivation; bio-energy; energy conversion, bio-fuel; environmental impact; greenhouse gas balance; land use impact; human health

1. Introduction

In a context of growing interest for renewable energy sources liquid bioenergy production from vegetable oils is proposed as one of the possible options to reduce greenhouse gas (GHG) emissions. Against this background bio-diesel production from *Jatropha curcas* L. (JCL) has become a booming business. The oil produced by this crop can be easily converted to liquid bio-fuel which meets the American and European standards [1-2]. Additionally, the press cake can be used as a fertilizer and the organic waste products can be digested to produce biogas (CH₄) [3-7]. The plant itself is believed to prevent and control soil erosion or can be used as a living fence or to reclaim waste land [8-12]. JCL is still a wild plant, which can grow without irrigation in a broad spectrum of rainfall regimes, from 250 up to 3000 mm per annum [13]. Furthermore, JCL is reported to have few pests and diseases [5,10], but this may change when it is grown in commercial plantations with regular irrigation and fertilization [14]. Based on these interesting properties, potentials and hyped claims, a lot of investors, policy makers and Clean Development Mechanism project developers are interested in JCL to tackle the challenges of energy supply and GHG emission reduction [12,15].

The essential minimum requirements for bio-fuels to be a more sustainable alternative for fossil fuels is that they should be produced from renewable raw material and that their use has a lower negative environmental impact [16]. Closer investigation is needed in order to conclude if both minimum requirements are fulfilled. Different sustainability evaluation tools and environmental impact assessment tools are available to investigate if an agricultural production process meets these requirements [17]. Life cycle

assessment (LCA) is such an instrument and has already shown its utility to evaluate the environmental balance of bio-fuel from other vegetable oils [18-21]. Using LCA or any other sustainability evaluation tool will need input data as a start to perform the evaluation or the assessment.

In this paper we present a state-of-the-art literature review of the whole JCL bio-diesel production process and use. For each production step the available published information on inputs and outputs is compiled. This collection of data and information enables us to discuss (i) the actual best practice(s) for production of JCL bio-diesel, (ii) the most persistent shortcomings, and hints of remedies and (iii) on the most prominent potential environmental issues, using a limited LCA approach (energy balance, greenhouse gas balance, land use impact). Information was compiled not only from peer-reviewed literature, but also from reports and conference proceedings. Doing so we could cover a wider range of information, which allowed us to obtain quantitative data presented as ranges, averages and standard deviations.

2. Botanical description of *Jatropha curcas* L.

JCL or Physic nut is a small tree or large shrub, up to 5-7 m tall, belonging to the *Euphorbiaceae* family, with a soft wood and a life expectancy of up to 50 years. The plant has its native distributional range in Mexico, Central America, Brazil, Bolivia, Peru, Argentina and Paraguay [22], although nowadays it has a pantropical distribution [12] with distinct JCL seed provenances. The plant develops a deep taproot and initially four shallow lateral roots [9]. The taproot may stabilize the soil against landslides while the shallow

roots are alleged to prevent and control soil erosion caused by wind or water, but this potential has not been investigated scientifically. The leaves are smooth, 4-6 lobed and 10-15 cm in length and width. The plant is monoecious and the terminal inflorescences contain unisexual flowers. The ratio of male to female flowers ranges from 13:1 to 29:1 [6,23] and decreases with the age of the plant [24]. Normally JCL flowers only once a year during the rainy season [23]. In permanently humid regions or under irrigated conditions JCL flowers almost throughout the year [9]. After pollination, the inflorescences form a bunch of green ellipsoidal fruits [6]. The blackish seeds of most provenances contain toxins, such as phorbol esters, curcun, trypsin inhibitors, lectins and phytates, to such levels that the seeds, oil and seed cake are not edible without detoxification (see e.g. Becker and co-workers [25-33]).

3. *Jatropha* cultivation

The cultivation of JCL trees for the production of oil-bearing fruits is considered the first production step towards bio-diesel production (Fig. 1). The main inputs are land area including the prevalent site characteristics, plantation establishment practices and plantation management practices including the production and use of all machines, infrastructure and energy (transport, power, etc.) needed for those inputs. The outputs are the seeds, other biomass elements including the husks, and GHG emissions. In this analysis, we did not consider erosion or nutrient losses to surface water as these are highly dependent on site conditions and difficult to generalize.

(Insert Fig. 1)

3.1. Site requirements

JCL's high ecological adaptability [9] allows it to grow in a wide range of conditions. As a succulent that sheds its leaves during the dry season, JCL is well adapted to semi-arid conditions, although more humid environmental conditions show to result in better crop performance. The documented seed provenances show average temperatures between 20°C and 28°C [9,31], but its occurrence has been observed in a rainfall range between 250 mm and 3000 mm [13]. JCL can tolerate high temperature extremes, but generally fears frost, which causes immediate damage [9,34]. In Nicaragua, it has an altitude range from sea level up to 1800m [13]. The plant is not sensitive to day length [9].

JCL can grow in a wide range of soils. Well drained sandy or gravelly soils with good aeration are preferred [9,13]. In heavy soils, root formation will be hampered [9]. JCL should never be planted on soils with risk of even ephemeral water logging, such as Vertisols or other heavy clay soils [35,36]. Soil depth should be at least 45 cm [34] and surface slope should not exceed 30° [6]. JCL has low nutritional requirements but the soil pH should not exceed 9 [6,36] and on very acidic soils JCL might require some Ca and Mg fertilization. JCL is well adapted to marginal soils with low nutrient content [9], but in order to support a high biomass production the crop shows a high demand for nitrogen and phosphorus fertilization [13]. Mycorrhiza assisting with the uptake of phosphorus and micro-elements were found on the root system [8,10,34,37]. Mycorrhiza inoculated JCL

showed a 30% increase in both biomass and seed production seven months after plantation of one-year-old saplings [6].

3.2. Propagation and plantation establishment

JCL is easily propagated by generative (direct seeding or pre-cultivated seedlings) and vegetative (direct planting of cuttings) methods [9,10,38]. The crop shows high initial establishment success and survival [39]. For quick establishment of living fences and plantations for erosion control, direct planting of cuttings is considered easier [9], although JCL plants propagated from cuttings do not develop a taproot. The plants only develop thin roots unable to grow deep in the soil, which makes the plants more susceptible to uprooting by wind [40]. In agroforestry and intercropping systems direct seeding should be preferred over pre-cultivated JCL plants, as the taproot of directly seeded plants is believed to penetrate in deeper soil layers [41] where it can access extra nutrient resources and where it competes less with the roots of the other crops [9]. If early seed yields are to be achieved, direct planting of stakes can be used as well [9].

Recommendations on vegetative propagation vary. Cuttings of 25-30 cm length from one-year-old branches [34] or longer cuttings up to 120 cm [38] are among the options. Kaushik & Kumar [42] report that the survival percentage depends on the origin of the source material (top, middle or base of the branch) and the length and diameter combination of the cutting [42]. Their study showed a survival percentage of 42% when the top of the branches were used as cuttings, while cuttings from the middle (72%) and base (88%) showed significantly better survival results. The product of the length and

diameter dimensions of the used cuttings had a positive correlation on the survival percentage as well. The longer and larger a cutting, the higher its survival rate. Survival percentages higher than 80% were obtained for the length – diameter combinations from 105 – 2.5 cm, 45 – 3.5 cm, 45 – 4.5 cm and onwards [42]. Cuttings can be planted directly in the field or in nursery beds or polyethylene bags for first root development [9,34,38,42]. They have to be placed 10-20 cm into the soil depending on their length and diameter. Planting of cuttings is best done in the rainy season [43].

Using generative propagation, direct seed sowing is recommended at the beginning of the rainy season, after the first rains when soil is wet, because it helps to develop a healthy taproot system [34]. Seedlings can be pre-cultivated in polythene bags or tubes or in seed beds under nursery conditions. The use of plastic bags or tubes is observed to induce root node formation and spin growth [40]. In the nursery, seeds should be sown three months before the rainy season in a soil with a high concentration of organic material (sandy loam soil – compost ratio 1:1 [42]; in case of more heavy soils, sand is added: sand – soil – compost ratio 1:1:2 [34]; sand – soil – farm yard manure ratio 1:1:1 [35]) and should be well watered [38]. Pre-soaked seeds (24 hours in cold water) germinate in 7-8 days in hot humid environment whereas the process continues for 10-15 days [34]. A study on the germination enhancement of JCL seeds showed best results for pre-soaking in cow-dung slurry for 12 h (96% germination). The traditional 24 h cold water treatment showed 72% germination. Nicking yields similar germination rates (pers. obs.), while pretreatments using hot water or H_2SO_4 (0.5 M) do not enhance germination [44].

At the onset of the rains the seedlings can be planted in the field. Planting distances of 2×2 m (2500 plants ha^{-1}), 2.5×2.5 m (1600 plants ha^{-1}) or 3×3 m (1111 plants ha^{-1})

are common practice [9]. Kaushik and Kumar [42] propose to use wider spacing patterns (4×2 m and 4×3 m) and agroforestry systems (spacing 5×2 m and 6×6 m) to optimize the yield of individual JCL plants. In 2.5-year-old plantations it was observed that with increasing spacing, seed yield tree⁻¹ increased significantly, while the seed yield ha⁻¹ decreased [45]. The recommended spacing in hedgerows for soil conservation is 15-25 cm within and between (in case of double fence) rows (4000-6700 plants km⁻¹) [10].

Field preparation for oil production plantations mainly consists of land clearing and preparation of the planting pits for the pre-cultivated plants. Although planting can be done without any clearing, for oil production purpose it is advisable to clear the land at least partially [34]. Tall trees can be left, but shrubs and bushes that cover the soil should be cut. Ploughing the field belongs to the possibilities as well [34]. After clearing, planting pits of 30-45 × 30-45 × 30-45 cm³ should be dug prior to the rainy season [34,35]. For good establishment the pits are best refilled with a mixture of the local soil, sand, organic matter such as compost and/or artificial fertilizer.

The best moment for planting is the warm season — if watering can be provided — or at the onset of the rains [10]. Gour [34] poses that seedlings require irrigation, especially during the first 2-3 months after planting. Of course, the water demand depends on local soil and climatic conditions.

3.3. Tending practice of the plantation

Besides propagation and spacing some publications mention weeding and pruning [9,10,46,47]. Recent publications [34,35,42] sketch a more complete view on management

and cultivation activities. Regular weeding operations should free the field from competitive weeds. Uprooted weeds can be left on the field as mulch. Pruning and canopy management is presented as an important crop architectural intervention, which is believed to help the production of more branches and to stimulate abundant and healthy inflorescence, thus eventually enhancing good fruit setting and seed yield [34]. At the age of six months it is useful to pinch off the terminal shoots in order to induce lateral branching [34,42]. Experiments reveal that pruning the main branch at 30 - 45 cm height — depending on the growth rate — is ideal [34]. At the end of the first year, the secondary and tertiary branches should be pinched or pruned to induce more branches. During the second year each side branch should be pruned up to two-thirds of the top portion, retaining one-third of the branches on the plant [34,42]. Pruning should be done in the dry or winter period after the trees have shed their leaves. This will result in a lower and wider tree shape, induce earlier seed production and facilitate manual harvesting. Once every 10 years, the entire plant has to be cut low, leaving a stump of 45 cm. The re-growth will be quick and the trees will start yielding again within about one year. This intervention will induce new growth and help to stabilize the yield [34]. Beside trimming hedgerows and pruning plantations annually, periodic thinning of plantations is proposed as well. Starting from 1600 seedlings per hectare, stand density should be thinned to 400-500 trees per hectare in the final mature stand [10].

It is clear that optimal fertilization and irrigation application can increase the seed and oil yield. However, permanent humid situations and/or situations with high irrigation and fertilizer application can induce high biomass but low seed production. The input levels to optimize the harvest index in given conditions are yet to be quantified. No

quantitative data on water need, water productivity and water use efficiency of JCL are available at present. In general application of super phosphate or NPK fertilizer is reported to increase the yield. The optimum application levels of inorganic N and P fertilizers are observed to be variable according to the age of the plantation [48]. On degraded sites JCL plants are found to respond better to organic manure than to mineral fertilizers [8]. Based on the nutrient composition of JCL fruit (compiled by Jongschaap et al. [41]) it can be estimated that harvesting the equivalent amount of fruits for a yield of 1 ton of seeds ha⁻¹ results in a net removal of 14.3-34.3 kg N, 0.7-7.0 kg P and 14.3-31.6 kg K ha⁻¹. So fertilization (artificial or organic) at least has to compensate this.

The susceptibility of JCL for pest and diseases is a source of discussion and is believed to depend on the management intensity. Early publications [9,49] already listed numerous pests, diseases and damaging insects observed on JCL. Furthermore it is believed that JCL can transmit the cassava superelongation disease (*Sphaceloma manihoticola*) and is a possible host for African Cassava Mosaic Virus (until now only observed in *Jatropha multifida* L.) [9]. A popular belief is that JCL is not prone to pests and diseases in such extent to cause economic damage. However, in continuous JCL monocultures in India economic damage has already been observed [50]. The major problems in JCL cultivation are caused by the scutellarid bug *Scutellera nobilis* and the inflorescence and capsule-borer *Pempelia morosalis* [50]. Grimm and Maes [49] identified *Pachycoris klugii* (Scutelleridae) and *Leptoglossus zonatus* (Coreidae) as the key pests in Nicaragua. Other possible pests are the blister miner *Stomphastis (Acrocercops) thraustica*, the semi loopers *Achaea janata* and the flower beetle *Oxycetonia versicolor* [50]. Regular

irrigation and fertilizer application is expected to enhance these pest and disease infestations in commercial monocultures [14].

3.4. Seed yield

For best oil yields, the seeds should be harvested at maturity. Seeds are mature if the color of the fruits has changed from green to yellow-brown. Maturity is reached 90 days after flowering [9], but the fruits do not mature all at the same moment. As such the fruits have to be harvested manually at regular intervals [9,35], making this step very labour intensive. The moment and length of harvest period is likely to vary according to the seasonal conditions of the locality [51]. In semi-arid regions the harvest is spread over a period of two months which implies daily or weekly harvests. In permanent humid situations weekly harvest can be necessary all year through. Separation of the seeds and husks can be done manually or mechanically [34].

JCL seed yield is still a difficult issue. Actually the mature seed yield per ha per year is not known, since systematic yield monitoring only started recently. Earlier reported figures exhibit a very wide range ($0.4 - 12 \text{ t ha}^{-1} \text{ yr}^{-1}$ [10]) and are not coherent [9] (Fig. 2 and Appendix 1), mainly because of incorrect extrapolation of annual yields of individual trees to $\text{ha}^{-1} \text{ yr}^{-1}$ yields. At present the effect of spacing, canopy management and crown form and surface on the yield is not known, making it impossible to make such extrapolation. Fig. 2 indicates positive trends in the influence of both average annual rainfall and age on the seed yield. Mainly the upper boundary of the yield in function of

the rainfall is interesting and shows a clear difference between low rainfall and high rainfall regimes.

(insert Fig. 2)

Yield depends on site characteristics (rainfall, soil type and soil fertility) [8,10,54], genetics [55], plant age [9,56] and management (propagation method, spacing, pruning, fertilizing, irrigation, etc.) [9,34,35]. Information on these yield influencing variables was generally not reported alongside. JCL has not yet undergone a careful breeding programme with systematic selection and improvement of suitable germplasm, which is why it can still be considered a wild plant that exhibits great variability in productivity between individuals. Annual seed production can range from about 0.2 kg to more than 2 kg per plant [8]. For semi-arid areas and cultural wasteland Heller [9] and Tewari [6] propose an achievable dry seed production of 2-3 ton ha⁻¹ yr⁻¹, which are confirmed by field data of Francis et al. [8]. When good sites (good soil and average annual rainfall of 900-1200 mm) are claimed and/or optimal management practice is used, 5 ton dry seed ha⁻¹ yr⁻¹ can be achieved [6,8,13]. Jongschaap et al. [41] conclude to a potential yield range of 1.5-7.8 ton dry seed ha⁻¹ yr⁻¹.

As mentioned earlier JCL is a hardy and highly adaptable crop that can grow in marginal soils from an average annual rainfall of 250 mm. As such JCL is capable to reclaim wasteland [57]. But is it able to produce ecologically and socio-economically viable amounts of energy in these barren situations?

Average shell:kernel ratio on mass basis of the JCL seeds is 37:63 (Fig. 3). The kernel mainly contains crude fat and protein and has an average calorific value of 30.4 MJ kg⁻¹ (Fig. 4). The shell is mainly composed of fiber and has a calorific value of 19.4 MJ kg⁻¹. Based on these figures the average oil content of dry seed on mass basis is 34.4%.

To date no information on the total biomass production of JCL is at hand. For an irrigated JCL project in Egypt, Henning prospects the future total biomass production to be 80 ton dry matter ha⁻¹ (11 ton dry matter ha⁻¹ yr⁻¹ including seeds) (spacing 2.5 × 2.5 m) representing 5.5 ton CO₂ ha⁻¹ yr⁻¹ [58]. For Indian wastelands the average annual CO₂ sequestration rate in the standing biomass ha⁻¹ is estimated to be ±2.25 tons CO₂ ha⁻¹ yr⁻¹ (excluding the 2-2.5 tons dry seed yield ha⁻¹ yr⁻¹) [8]. The wood density of JCL is reported to range from 0.33 to 0.37 g cm⁻³ [58] although personal observations on young plants (116 days old) noted a wood density of 0.29 ±0.1 g cm⁻³ at the base of the stem.

(Insert Fig. 3)

(Insert Fig. 4)

Makkar et al. [31] reported on 18 different provenances of JCL from countries in West and East Africa, the Americas and Asia including climatic data of these places. Large variations were found in contents of crude protein, crude fat, neutral detergent fiber and ash, but no causal links were analyzed. Recently Kaushik et al. [61] recorded coefficients of variance between 24 provenances of Haryana state, India, which indicate a dominant role of environment over genetics in seed size, seed weight and oil content.

An important by-product of the JCL seed production is the husk which stands for 35-40 % by weight on whole fruit basis [62]. The husk can be used for direct combustion and biogas production [3], but recently it was shown that JCL seed husks are an excellent feedstock for gasification as well [62]. Using an open core down draft gasifier, maximum efficiency was found to be 68.3% at a gas flow rate of $5.5 \text{ m}^3 \text{ h}^{-1}$ and specific gasification rate of $270 \text{ kg h}^{-1} \text{ m}^{-2}$. The gas had an energy value of 4.6 MJ m^{-3} which is comparable to wood [62].

4. Oil extraction

In the second step of the production chain for bio-diesel, the oil contained in the seeds has to be expelled or extracted. The main inputs for this process, besides the seeds, are the production and use of machines, infrastructure and energy. On the output side, the main products are the JCL oil and the seed or kernel cake, which is an important by-product (Fig. 5). The emissions of GHGs and waste water have to be accounted for in the outputs of the process as well.

The process of gaining oil from oilseeds is as old as mankind, but the possibilities, procedures and means have evolved and are subject of many publications. In the following, only the different extraction options and their performances for JCL seeds are reviewed. For extraction of the JCL oil two main methods have been identified: (i) mechanical extraction and (ii) chemical extraction [28,63].

(Insert Fig. 5)

Prior to oil extraction the JCL seeds have to be dried [38,64]. Seed can be dried in the oven (105°C) or sun dried (3 weeks). Mechanical expellers or presses can be fed with either whole seeds or kernels or a mix of both, but common practice is to use whole seeds. For chemical extraction only ground JCL kernels are used as feed. The shells can be used directly as a combustible by-product or gasification feedstock.

4.1. Mechanical expellers

For mechanical extraction of the oil from the seed, either a manual ram press (e.g. Yenga or Bielenberg ram press) or an engine driven screw press (e.g. Sundhara press [65]) can be used [60,63,66]. Henning [38] stated that engine driven screw presses extract 75-80% of the available oil, while the manual ram presses only achieved 60-65%. Oil extraction efficiencies calculated from data reported in more recent studies [6,63,67,68] are found to generally correspond to these ranges, although the efficiency range of engine driven screw presses can be broadened to 70-80% (Table 1). This broader range corresponds to the fact that seeds can be subjected to a different number of extractions through the expeller. Up to three passes is common practice. Pretreatment of the seeds, like cooking, can increase the oil yield of screw pressing up to 89% after single pass and 91% after dual pass [68]

(Insert Table 1)

4.2. Chemical extraction

Table 2 summarizes the reaction temperature, reaction pH, time consumption and oil yield of different chemical extraction methods tested on JCL. The *n*-hexane method is the most common and results in the highest oil yield, but also takes most time. In aqueous enzymatic oil extraction the use of alkaline protease gave the best results for both available studies [69,70]. Furthermore, it is shown that ultrasonication pretreatment is a useful step in aqueous oil extraction [70].

(Insert Table 2)

Adriaans [73] concludes that solvent extraction is only economical at a large scale production of more than 50 ton bio-diesel per day. Furthermore he does not recommend the conventional *n*-hexane solvent extraction because of environmental impacts (generation of waste water, higher specific energy consumption and higher emissions of volatile organic compounds) and human health impacts (working with hazardous and inflammable chemicals). Using aqueous enzymatic oil extractions greatly reduces these problems [73] as do the use of supercritical solvents (mainly supercritical CO₂) or bio-renewable solvents as bio-ethanol and isopropyl alcohol. Although the new generation *n*-hexane extraction units are very efficient and produce far less environmental burdens than the older units, further research on these alternative solvents is recommended as on their commercial viability. Foidl and Mayorga [74] presented the use of supercritical isopropanol or CO₂ in a continuous mechanical oil extraction system only leaving 0.3% by weight of oil in the cake.

367

368 4.3. *Jatropha* oil

369 The composition and characteristics of the crude JCL oil are given in Table 3. The
370 JCL oil meets the quality standard of rapeseed as a fuel [60].

371

372 (Insert Table 3)

373

374 It is important to note that the values of the free fatty acids, unsaponifiables, acid
375 number and carbon residue show a very wide range, although it is a small data set. This
376 indicates that the oil quality is dependent on the interaction of environment and genetics.
377 As for the seed size, seed weight and oil content [61], it is believed that also for the oil
378 quality the environmental conditions have a larger impact than the genetics. More research
379 is necessary. Project developers and decision makers should be aware of these wide ranges,
380 because these above-mentioned characteristics are important properties for the further
381 processing of the oil into bio-diesel [86-88].

382 The JCL oil contains more than 75% unsaturated fatty acid, which is reflected in the
383 pour and cloud point of the oil. The fatty acid composition of JCL oil is dominated by oleic
384 acid (C18:1) and linoleic acid (C18:2) (Fig. 6). The maturity stage of the fruits at the
385 moment of collection is reported to influence the fatty acid composition of the oil [89].

386

387 (Insert Fig. 6)

388

4.4. *Jatropha* seed cake

The average crude protein content of the seed cake is 58.1% by weight and has an average gross energy content of 18.2 MJ kg⁻¹ (Fig. 7).

(Insert Fig. 7)

In case of mechanical oil extraction from whole seeds, the oil content of the seed cake will be higher, due to the lower efficiency of the expellers. Based on the extraction efficiencies discussed above and the average oil content of the whole seed (34.4% on a mass basis), the seed cake will contain 9 – 12 % oil by weight. This content will of course influence the gross energy value of this cake as well.

Next to the high quality proteins (Fig. 7) this cake contains various toxins and is therefore not usable as fodder [8]. However, the raw kernel or seed cake can be valuable as organic nutrient source, as it contains more nutrients than both chicken and cattle manure [8]. Table 4 gives an overview of experiments which show that JCL seed cake is useful as fertilizer. The presence of the aforementioned bio-degradable toxins, mainly phorbol esters, makes the fertilizing cake simultaneously serve as biopesticide/insecticide and molluscicide [8,91]. Although the phorbol esters decompose completely within 6 days [91] it is advisable to check the absence of phorbol esters in the crops grown on JCL cake fertilized land, certainly crops for human consumption. Heller [9] warns about phytotoxicity of over-application of JCL cake. One study showed phytotoxicity to tomatoes, expressed in reduced germination, when high rates of up to 5 ton ha⁻¹ are applied.

(Insert Table 4)

The cake can serve as feed for biogas production through anaerobic digestion before using it as a soil amendment as well. Staubman et al. [4] obtained 0.446 m³ of biogas, containing 70% CH₄, per kg of dry seed press cake using pig manure as inoculum. Radhakishna [7], using specific developed microbial consortia as inoculum, achieved 0.5 m³ biogas kg⁻¹ of solvent extracted kernel cake and 0.6 m³ biogas kg⁻¹ of mechanically de-oiled cake. Other organic wastes of JCL can be digested. For example, Lopez et al. [3] showed that biogas could be produced from JCL fruit husks.

The fact that JCL seed cake can be used for different purposes makes it an important by-product. Recycling of wastes as a fertilizer can help to reduce inputs needed for both JCL cultivation and other agricultural cultivation or it can produce extra energy in the form of biogas. Digesting the cake, and bringing the effluent back to the field is thought to be the best practice at present from an environmental point of view. A number of questions concerning the long-term and cumulative impacts of JCL seed cake on soils have not been addressed. In the event that detoxification becomes viable, the use as animal feed will be more beneficial.

5. Production of bio-diesel

Vegetable oil can be used as base for liquid engine fuel in various ways (straight vegetable oil, oil blends, pyrolysis, micro-emulsification, transesterification). All general problems and benefits and all these procedures of using vegetable oils as liquid engine fuel,

are the subject of numerous publications [86-88,95-98]. In this section only the published experiences with JCL oil are reviewed.

JCL oil is mainly transesterified to (m)ethyl esters (bio-diesel) and glycerol. Taking JCL bio-diesel as the end product transesterification should be considered the next step in the production process (Fig. 8). Glycerol is an important by-product. It can be burned for heat or be used as feedstock in the cosmetic industry. But using JCL oil as a bio-fuel does not always include an extra unit process. Several publications [63,67,76,80,81,99] report the use of the pure JCL oil or JCL oil blends (see further).

(Insert Fig. 8)

5.1. Transesterification

Although the transesterification process is quite straightforward, the genetic and environmental background of the produced oil might require modification of the input ratios of the alcohol reagent and reaction catalyst as well as alterations to reaction temperature and time, in order to reach optimal bio-diesel production results. The optimal inputs for the transesterification of JCL oil (3.1% free fatty acids and acid number 6.2 mg KOH g⁻¹) are identified to be 20% methanol (by mass on oil basis) (molar ratio methanol:oil $\approx$ 5.5:1), 1.0 % NaOH by mass on oil basis [83]. Maximum ester yield is achieved after 90 minutes reaction time at 60°C [83]. Optimal conversion of JCL oil with high free fatty acids (14%) and high acid number (28 mg KOH g⁻¹) needs pretreatment reaction with methanol (molar ratio methanol:oil $\approx$ 6.5:1) using H₂SO₄ as catalyst (1.43%)

during 88 minutes at 60°C. After pretreatment a maximal conversion rate of more than 99% was achieved by transesterification with methanol (molar ratio methanol:oil $\approx$ 4:1) and 0.6% KOH by weight during 24 minutes[2].

Given the variability the best practice to maximize ester yield, is to determine the optimal inputs of oil samples per oil batch that has to be transesterified. The characteristics of the resulting JCL (m)ethyl esters generally meet the American and European standards [1,2] (Table 5).

(Insert Table 5)

Recently more advanced transesterification procedures are tested on JCL oil as well. Table 6 summarizes the key inputs and outputs of these methods.

(Insert Table 6)

6. Use of the *Jatropha* oil

Jatropha oil has various uses. Apart from its use as a liquid fuel, the oil has been used to produce soap and biocides (insecticide, molluscicide, fungicide and nematocide) [50]. The oil can be directly used in older diesel engines or new big motors running at constant speed (e.g. pumps, generator). Blending with fossil diesel and/or other fossil fuels belongs to the options as well. The oil can also be transesterified into JCL (m)ethyl esters that can be used in conventional diesel engines or diesel engines with adapted parameters.

477

478 6.1. Direct use of the oil

479 Tests with a low heat rejection diesel (LHR) engine showed that the use of pure JCL
480 oil results in a higher brake specific energy consumption (BSEC), lower brake thermal
481 efficiency (BTE), higher exhaust gas temperature (EGT) and lower NO_x emissions in
482 comparison with fossil diesel [99]. Preheating and increasing the injection pressure
483 decreased BSEC, increased BTE, increased EGT and increased NO_x emissions only
484 marginally [99]. Kumar et al. [80] compared the use of JCL oil and fossil diesel in a single
485 cylinder 4 stroke water cooled diesel engine and concluded that the soot (hydrocarbon)
486 emission is higher with JCL oil as compared to fossil diesel. At maximum output an
487 increase from 100 ppm, for fossil diesel, to 130 ppm, for JCL oil, was measured and similar
488 trends were observed in the case of CO emissions. Smoke level was higher with JCL oil
489 (4.4 BSU) compared to fossil diesel (3.8 BSU) as well. Furthermore they observed an
490 increase in ignition delay and combustion duration with JCL oil in comparison to fossil
491 diesel [80].

492

493 6.2. Use of transesterified oil

494 The use of the methyl ester of JCL resulted in a soot emission of 110 ppm compared
495 to 100 ppm for fossil diesel and in increased CO emissions [80]. *Jatropha* methyl ester
496 (JME) approached the smoke levels (4.0 BSU) of fossil diesel level (3.8 BSU). JME was
497 also observed to decrease the BTE and volumetric efficiency, to increase the ignition delay

and combustion duration [80]. In addition, the study of Prasad et al. [99] showed higher BSEC for JME use than for conventional diesel use. Since the calorific value of JME is lower than conventional diesel, this observation implies that the JME fuel consumption will be higher than conventional diesel as well. Prasad et al. [99] conclude that the use of transesterified JCL oil achieves similar results as the use of fossil diesel, although it causes less NO_x emissions. Furthermore, transesterified JCL oil shows little or no additional engine corrosion as compared to fossil diesel [103].

6.3. Use of blends

The use of pure JCL oil – fossil diesel blends has been tested in a single cylinder, water cooled open combustion chamber diesel engine [81]. Brake specific fuel consumption (BSFC) and EGT of the blends were found to be higher compared to fossil diesel and tended to increase with increasing proportion of JCL oil in the blend. The opposite applies for the BTE. However, blends of 30% JCL oil and 70% fossil diesel by volume and of 40% JCL oil and 60% fossil diesel by volume showed BSFC and BTE close to the values of fossil diesel. Both BSFC and BTE were found to be acceptable up to 50% by volume of JCL oil [81]. Furthermore, the study concluded that the long-term durability of the engine using bio-diesel as fuel requires further study.

Kumar et al. [80] also tested JCL oil – methanol blends and dual fuel operation. Blending JCL oil with methanol increases the BTE, lowers the HC and CO emissions and reduces smoke levels, but the blend could not reach the results of JME. Furthermore, the blend increased ignition delay [80]. The dual fuel operation with methanol induction and

JCL oil as the pilot fuel resulted in a significant increase in BTE (although, not achieving the JME results) and showed lowest smoke and NO levels. The penalty is an increase in HC and CO emission [80].

In a Lister model engine (single cylinder, air cooled, direct injection, four stroke diesel engine) a blend of 2.6% JCL oil and 97.4% fossil diesel by volume showed lowest BSFC and highest BTE [63] in comparison to fossil diesel and blends with higher JCL oil portion. In the oil extraction procedure of the used JCL oil a small portion of water was added which emulsified the JCL oil. This resulted in a reduction of EGT and, as such, a reduction in NO_x emission with increasing portion of JCL oil in the blend [63]. Attention is to be paid to the water content of the used fuel as it causes oxidation inside the injection equipment [60].

6.4. Alternative engine technology

Retarding the injection timing with enhanced injection rate of a single cylinder, constant speed, direct injection diesel engine, operating on neat JCL oil, showed to improve the engine performance and emission level significantly [76]. The measured emissions were even lower than fossil diesel. At full output HC emission level was observed to be 532 ppm against 798 ppm for fossil diesel, NO level was 1163 ppm against 1760 ppm and smoke was reduced to 2.0 BSU against 2.7 BSU. However, the achieved BTE with JCL oil (28.9%) was lower than with fossil diesel (32.1%) [76].

From the above summarized studies it can be concluded that JME generally achieves the best results in comparison to the use of pure JCL oil, straight or in a blend, although the scope for the use of the pure JCL oil can not be underrated. Certainly in tropical developing countries the use of pure JCL oil, straight or as a blend, is believed to have great potential [60]. The diesel engines in those countries, including those of old four-wheel-drive vehicles, rely on older technology, are easier to adapt to the characteristics of the used fuel and the tropical temperatures lower the viscosity of the oil [60]. Stationary diesel engines at low speed, such as irrigation pumps and electricity generators, are believed to be suitable to pure JCL oil without a too high environmental burden. Pre-chamber diesel engines are more suitable for the use of pure oil than direct injection engines, but simple conversion to a direct injection two tank system can overcome their problems [60]. Despite the reported difficult use of pure vegetable oils [86,87,95,98], there is still scope for improvement. Further research is recommended.

7. Discussion

7.1. Best available practice for the production of JCL bio-diesel and shortcomings

In this discussion an intuitive interpretation is given to the collected data and information on *Jatropha* bio-diesel production and use. The collected data are evaluated with the objective of describing best available practice for a production system in which JCL oil and/or bio-diesel is the main product.

7.1.1 *Jatropha* cultivation

Aiming mainly at oil production, block plantations are probably the best option. How such plantation is best established, is subject to much discussion yet. According to Heller [9] plants propagated by seeds are preferred for establishment of long living plantations for oil production. This can be supported by the fact that vegetative propagated plants do not develop a taproot, but only a superficial root carpet [40] which leads to more superficial water and nutrient competition. The taproot of generatively propagated plants will have more access to nutrients from deeper soil layers [41] and can reach deeper water resources.

The selection of basic material is a critical step (in case of vegetative as well as generative propagation). Basing this selection on successes of controlled breeding programmes would be the best option, but present results are not yet sufficient. In JCL provenances available in India only modest levels of genetic variation were observed, while wide variation was found between the Indian and Mexican genotypes [104]. This shows the need to characterization of provenances with broader geographical background [104]. Best available practice at the moment is to use planting material obtained from the best performing trees of the best performing provenance available in the location of interest. Trees with an annual yield above two kg dry seeds and seed oil content higher than 30% by weight can be considered a good source [34]. In generative propagation the selection of the heaviest and largest seeds for sowing results in significant growth increase of JCL seedlings [105]. Although germination rates, certainly after easy applicable pretreatments of the seeds (nicking, cold water), are quite high and although nursery bags can hamper initial

584 root formation, we would intuitively recommend plantation establishment through planting
585 of seedlings. As such the plants can be sufficiently protected in their initial growth stage,
586 when they are still quite susceptible for weather extremes or other possible events. Using
587 seedlings one has more control on the uniformity of the plantation as well. Further the
588 planting pits will guarantee a good establishment in the soil. The main drawback of this
589 practice is the influence of the polythene bags and pots on the root structure [40].

590 Due to root competition for water the optimal spacing is believed to be a function of
591 rainfall, where wider spacing should be used in semi-arid environments and denser
592 plantations can be appropriate for sub-humid environments. It was noted that spacing of
593 plants is a trade off between biomass and fruit production. A narrow spacing will lead to
594 fast canopy closure which results in higher water and light competition and lower
595 fruit:biomass ratio in the mature stadium. When planting JCL for live-fencing or hedges
596 for soil conservation a dense biomass is needed and close spacing is appropriate. When the
597 aim of the plantation is oil production, seedlings should be planted wide enough to ensure
598 high seed yields in the mature stage, but close enough to avoid unacceptable loss of
599 photosynthetic capacity in the juvenile stage. Thus, optimum spacing can only be
600 recommended after at least 5 years consecutive growth and yield observations and this in
601 different environmental conditions and using different provenances. The authors feel that
602 the best available practice at this moment is to start with a densely spaced block plantation
603 and gradually remove rows or individuals (thinning) according to the plant performances.

604 Contrary to popular believe, it should be made clear that plantations aiming at oil
605 production will need fertilization (artificial or organic). Fertilizer at least needs to
606 compensate the nutrient removal due to harvest or management practice (pruning – if not

used as propagation material). Irrigation will depend on the climatic conditions of the location. The minimum annual average rainfall at which JCL is known to yield a harvestable amount of seeds is 500-600 mm yr⁻¹. So, simultaneous reclamation of barren lands and bio-diesel production will inevitably imply use of fertilizer and irrigation. Although there are already several fertilization trials available [48,94] there is still insufficient information to account the nutrient need for specific environmental and genetic setups. The same applies for irrigation.

Reliable yield prediction still forms the biggest problem. At present there are no reliable field data on the dry JCL seed yield ha⁻¹ yr⁻¹ in a given set of conditions and at a certain level of input. It is believed that for well managed plantations in good environmental conditions a yield expectation of 4-5 ton dry seed ha⁻¹ is reasonable. In order to tackle this knowledge gap, it is absolutely necessary to systematically monitor the year-to- year seed yield in operational plantation conditions along with the influencing factors. Furthermore, research is necessary to quantify the causal effects of each of the influencing factors on the yield. It is important to give special attention to the interactions between the environmental and management requirements and the influence of the different provenances. Issues to address at the crop level are biogeochemical cycling, water use efficiency, drought resistance, total biomass production, pest management (inclusive hosting and transmitting capacity of pest and diseases infesting other crops), issues on invasiveness and land suitability of JCL.

JCL is still a wild plant with a wide variation in growth, production and quality characteristics. In order to work towards high yielding bio-diesel plantations, the best suitable germplasm has to be identified for different cultivation situations. This implies

characterization of provenances with broader geographical background in order to widen the genetic base of JCL. An intensive inventory of the finalized and ongoing provenance trials will give an idea of the available material and will indicate where more provenance trials are needed (this is ongoing in the Global *Jatropha curcas* evaluation, breeding and propagation programme of Plant Research International, Wageningen with principal investigator R.E.E. Jongschaap). Based on such information, systematic and selective breeding should be carried out in order to develop high and early yielding hybrids with high oil yield in given site conditions. Recently a method has been developed for identification of superior lines by assessing the phenotypic traits of JCL plants recorded *in-situ* [106]. According to the authors this method facilitates the selection of promising accessions for multi-location evaluation and hastens the process of utilization of germplasm.

In short it can be stated that more systematic research and complete reporting is necessary on the input-responsiveness of the production at different levels of inputs, including environmental, as well as genetic, physical, chemical and management inputs (e.g. spacing, soil conditions, pruning, fertilizer, irrigation). Seed yield and biomass production in different environmental and abiotic setups, using different provenances or accessions, applying different levels of the different inputs should be monitored in order to discover the input-responsiveness of those different inputs as well as the interactions between the different inputs and the interaction between the environmental and genetic setups and the different inputs.

7.1.2 Oil Extraction

The choice of extraction method is clearly dependent on the intended scale of the activity. The two extraction procedures, mechanical and chemical, are quite well established, although there is still scope for further research. Both of them have their advantages and disadvantages with respect to scale suitability, centralization, extraction efficiency and environmental and health risks. Further research should investigate efficiency improvement of mechanical oil extraction, the applicability of alternative solvents as supercritical CO₂, bio-ethanol and isopropyl alcohol and their economical viability. Decentralized processing technology should be considered as well [8]. Such development should go in synergy with the transesterification setup.

The seed/kernel cake is a very important by-product, which we recommend to be brought back on the JCL field. Although the use of the cake as fertilizer is already common practice, there are still questions to be addressed. More trials are needed where the growth effect on different kinds of crops are monitored (including phytotoxicity and bio-safety effects). The impacts on the soil structure, water holding capacity, soil decomposition, organic matter content and soil biological activity should be brought under detailed investigation as well.

7.1.3 Production and use of JCL bio-fuel

The production of bio-diesel from vegetable oils in general is well documented. Crucial research and development options lay in the maximization of the transesterification efficiency at minimal cost. An important issue in this is the improvement in the catalytic

process, certainly the recovery and the reuse of the catalyst [8]. As part of the option of decentralized processing units, low-cost, robust and versatile small scale oil transesterification designs should be developed [8].

The choice of using JCL bio-diesel (i.e. methyl esters) or the JCL oil depends on the goal of the use (e.g. electricity or transport) and the available infrastructure. Studies show that transesterified JCL oil achieves better results than the use of pure JCL oil, straight or in a blend, in unadjusted diesel engines. Changing engine parameters shows considerable improvement of both the performance and the emission of diesel engines operating on neat JCL oil. More trials on the use of straight JCL oil in different diesel engine setups should be tested and investigated. Accurate measuring and reporting on emissions contributing to global warming, acidification, eutrophication, photochemical oxidant formation and stratospheric ozone depletion is very relevant. The long term durability of the engines using bio-diesel as fuel requires further study as well.

7.2. Environmental issues

As mentioned before there are different tools available for assessment of environmental impacts or evaluating environmental sustainability. In this section we will discuss the environmental issues of JCL production and use through a limited life-cycle approach.

In LCA all inputs and outputs of each step of the complete production cycle are inventoried and the calculated impacts are compared with a reference system. In this case we propose a reference system producing the same amount of energy based on fossil energy

sources [107-110]. Most LCA studies of bioenergy from agriculture and forestry are limited studies focusing on the energy balance and the global warming potential [107-114] while there are several other impact categories to address [115]. Land use impact is one of those that is rarely included, although “flows” of land area, water, vegetation and biodiversity are certainly as important for the viability and sustainability of production systems occupying substantial portions of land [116-118]. The land use impact assessment will give us an idea of the renewable character of vegetable oil or bio-fuel from the production process of interest.

Based on the unit processes described in the former sections we discuss the potential life cycle impacts of JCL biofuel production on (i) energy balance, (ii) global warming potential and (iii) land use.

7.2.1 Energy balance

The life-cycle energy balance of the bio-diesel production from JCL is reported to be positive [64,119] (Fig. 9). Note that Fig. 9 gives energy input values after allocating the total energy requirement of the whole process to the different products (end-product and by-products). This allocation is a pro-rata distribution of input energy among the products based on the energy content of those products. In Fig. 9 the energy input was distributed among the JME (end-product), glycerin (by-product of transesterification) and seedcake (by-product of oil extraction). The other by-products (wood, fruit husks) are not included in this allocation because the use of these by-products in an energy-efficient way is not common practice. The use of allocation is only justified if the by-products to which input

energy is allocated are effectively used in reality. In case none of the by-products are used, the energy balance will be only slightly positive (886 MJ input for 1000 MJ JME output [119]) or even negative. On the other hand, if all by-products (including wood and fruit husks) would be used efficiently this total input of 886 MJ results in a total output of 17235 MJ [119], resulting in a allocated energy input of 160 MJ per 1000 MJ JME (own calculations based on [119]).

(Insert Fig. 9)

Fig. 9 shows the energy balance of high and low JCL cultivation input. The difference between the two applied cultivation intensities is clear. In the low intensity system the JCL cultivation step stands for 17% of the total primary energy input, while the step accounts for 38% of the total energy input in the high intensity systems. Irrigation and fertilizer are the most energy intensive cultivation practices. Irrigation stands for 46% of the total energy input in the JCL cultivation; while fertilizer consumes 45% [119]. In both available studies the transesterification step is shown to be a big energy consumer (Fig. 9). The oil extraction step accounts for a similar share ($\pm 8\%$) of the total life cycle primary energy requirement in both studies (Fig. 9).

Based on the available results it can be expected that the life cycle energy balance of bio-diesel is generally positive. How positive the balance is in reality, will mainly depend on how efficient the by-products of the system are used. The available information furthermore shows that energy balance improvement options lay in the transesterification and cultivation step.

Transesterification shows to be the biggest contributor in the allocated energy requirement for the bio-diesel end product (Fig. 9). This would mean that the use of the crude oil as an end-product would improve the energy balance significantly. However, the engine combustion of pure JCL oil is less energy efficient [99] and still causes some engine problems [87]. In case of using old, stationary, low and constant speed engines, the lower energy efficiency of the pure oil compared with the transesterified oil, will probably be of no significance.

The energy requirement shows to vary a lot depending on the cultivation intensity (Fig. 9). The comparison between the two limited LCA case studies performed on JCL [64,119] shows that intensified cultivation does not always completely pay off in an extra energy production in the form of bio-diesel [120]. A significant energy balance improvement option lies in the optimization of the input-yield relationship, as discussed above.

7.2.2 Impact on global warming potential

Both available LCA exercises [64,119] showed positive results on the GHG requirement of the production of bio-diesel from JCL in comparison to fossil diesel. The largest GHG contributors of the production process are irrigation (if applied: 26% [119]), fertilizer (if applied: 30% [119]) and transesterification (24% [119] and 70% [64] depending on the applied cultivation intensity).

Prueksakorn and Gheewala [119] found that 90% of the total life cycle GHG emissions are caused by the end-use. They calculated that the global warming potential of

the production and use of JCL bio-diesel is 23% of the global warming potential of fossil diesel.

In general the impact on the global warming potential can be expected to be positive in comparison to the use of fossil diesel. It is clear that intensification of the cultivation step and transesterification will increase the GHG requirement of the production process. But, since Prueksakorn & Gheewala [119] consider the end use as the main contributor (90% of the total) this increase can only be thought to be marginal in the overall life cycle impact on the global warming potential. However, both limited LCA case studies do not account for N₂O emissions due to N fertilization. The global warming potential of N₂O is 296 times higher than an equal mass of CO₂ [121]. According to the IPCC the N₂O release is equal to about 1% of the nitrogen input from mineral fertilizer or biologically fixed N [122]. As the reduction of global warming potential is one of the main aims of the JCL bio-diesel system, this confirms the research need on input-responsiveness of the JCL cultivation step. For the use of the by-products the same applies for both GHG balance and energy balance.

The removal of (semi-) natural forest for the introduction of JCL, on the other hand, is expected to have a significant negative effect on the GHG balance of the whole life cycle. The caused emission due to removal of (semi-) natural forest is a heavy burden on the initial GHG investment which will take a significant time span before it is paid back with the GHG emission reduction of the use of the bio-diesel.

7.2.3 Land use impact

Land use impact assessment methodologies within LCA are still being discussed, but there is consensus about the fact that land use change and land use occupation impacts on soils and local biodiversity have to be assessed [117]. No assessment of these issues has been executed for JCL so far, but it is expected that land occupation impact of JCL on the soil will be positive. JCL is observed to improve soil structure [123], is strongly believed to control and prevent soil erosion and sequesters carbon. No information is available on nutrient cycles and the impact on soil biological life. The growing concern on these issues must be checked by focused research. It is important to note that this occupation impact will heavily depend on the applied cultivation system and intensity. Heavy machinery and high fertilizer application are expected to be the main drivers towards a negative impact.

Being an exotic species in most actual growing areas, the impact of land use change towards JCL on biodiversity is expected to be negative, although this will largely depend on the mix of land use which is replaced by JCL and on how JCL is cultivated. Impact on biodiversity will be especially negative when (semi-) natural systems such as dryland forests are cleared. Using JCL on barren wastelands has even potential to help restoring the local biodiversity. Using JCL as intensive monoculture will have severe impacts on the local biodiversity, while using JCL as live fence or in intercropping and agroforestry systems might not have a significant impact. However, there are some reports stating that JCL is an invasive species [124-126]. JCL is considered and treated as invasive in South Africa and as weedy in Australia [127]. However, no studies have been performed to quantify the allelopathic effects of JCL on native vegetation.

It is obvious that the main concern for this impact category is in the JCL cultivation unit process. Within this cultivation the mix of original land use, the used cultivation system and the applied cultivation intensity will be the most important influencing factors.

7.2.4 Issues in other impact categories

Due to the toxicity of the JCL seeds and oils, some attention should be paid to the human health and work environment impact categories. The fruits contain irritants affecting pickers and manual dehullers [128]. Although JCL has a very long history as medicinal plant, accidental intake of seeds and/or oil can cause severe digestion problems. For safety reasons, intercropping edible crops with JCL should only be recommended during the period before JCL starts bearing fruit. Also the use of the seed cake as fertilizer in edible crop production raises bio-safety questions. Several publications [129,130] suggest that the phorbol esters in the JCL oil would promote skin tumor. On the other hand Lin et al. [131] and Luo et al. [132] showed anti-tumor effects of the curcin from JCL oil. Furthermore, Gressel [128] warns for a serious lack of information about the effects of burning JCL oil in closed quarters, which is an important human health issue as the oil is proposed as a cooking fuel as well as a feedstock for bio-diesel production. He also calls for precaution in the use of accessions with high initial phorbol ester content since available extraction procedures for the removal of the phorbol esters are insufficient to bring those accessions to acceptable toxicological level [128].

Since very limited information is available regarding acidification, eutrophication and other unmentioned LCA impact categories of the JCL production cycle, no statements

or prognoses are made concerning these issues. Increased investigation of the cultivation step of the production of JCL bio-diesel will enable researchers to assess the specific contribution of JCL in these impacts as well.

8. Conclusions

In general we can state that the inclusion of data from reports and conference proceedings compensates the lack of peer-reviewed data. This state-of-the-art literature review on the production and use of JCL, however, detects several knowledge gaps which need to be bridged before more large scale cultivation can be undertaken.

The main knowledge gaps are situated in the cultivation of the crop, for both a description of best practice as for describing the potential environmental risks or benefits.. From selection of basic plant material up to yield there are many options, with a lot of variation in available data and not enough information for optimization. For project coordinators or investors it is almost impossible to lay out a coherent and realistic business plan because almost every step in the cultivation stage is uncertain, foremost the yield.

It is obvious that the JCL system cannot meet all expectations, which have been attributed to JCL since it has been hyped, at the same time or place. The plant cannot perform all its functions together at the best level. In popular press the JCL functions are presented separately and are, without evidence, attributed to all provenances. In an agricultural system there are trade-offs between the different functions the system or the crop can serve, this is not different for JCL.

Based on the available information it is still difficult to conclude if JCL bio-diesel will meet the two essential minimum requirements for bio-fuels to be a more sustainable alternative for fossil fuels (i.e. (i) produced from renewable raw material and (ii) their use has a lower negative environmental impact). JCL is expected to be renewable, but it is not clear at which cost. The impact on the soil seems to be positive, depending on used practice and used soil types, but this contribution to soil restoration might find trade-offs in biodiversity loss. The environmental impacts discussed are lower than the fossil alternative as long as no (semi-) natural ecosystems are removed in favor of JCL and as long as the by-products of the bio-diesel production system are efficiently used. The human health issue, on the other hand, is a persistent problem which can not be neglected. The authors call for precaution and science to be applied. The booming interest at the moment and the current lack of knowledge can not support the present popularity of JCL and as such can drive unsustainable practice and this could in its turn hamper the exploration of the true JCL potential risks and benefits.

Acknowledgements

This research is funded by the Flemish Interuniversity Council – University Development Co-operation (VLIR-UDC) and the K.U.Leuven Research Fund, and is a collaboration between K.U.Leuven and the World Agroforestry Centre (ICRAF). Special thanks go to the ICRAF team in India for hosting the research, to the FORECOMAN team (K.U.Leuven) and Dr. S. Dondeyne for their contributions during the writing process. The

869 comments and recommendations of two anonymous reviewers and editor Ralph P. Overend
870 are greatly acknowledged.

871 Appendix

872 Appendix 1 – Collection of published JCL dry seed yields

Reference	Location	AAR ^a	Age	kg tree ⁻¹ yr ⁻¹	kg ha ⁻¹ yr ⁻¹
[13]	Nicaragua, Managua	1200	2		2327
	Nicaragua, Managua	1200	3		2786
	Nicaragua	1200	4		3484
[9]	Cape Verde	600	-	0.80	
	India		3		1733
	Nicaragua, Managua	1200	-		5000
	Mali	1020	-		2640
	Thailand	1470	-		2146
	Mali	1020	-		8000
	Madagascar	1370	-	3.25	
	Paraguay	1370	3		100
	Paraguay	1370	4		700
	Paraguay	1370	5		1000
	Paraguay	1370	6		2000
	Paraguay	1370	7		3000
	Paraguay	1370	8		4000
	Paraguay	1370	9		4000
	Cape Verde	220	-		1750
	Cape Verde	220	-		500
	Thailand	1470	1	0.32	794
	Thailand	1470	1	0.06	
	Burkina faso	815	-	0.96	
	semi arid areas	-	-		2500
[52]	Nicaragua	1200	5	4.50	5000
[65]	Mali	1020	-		3000
[10]	-	-	-		400-12000
	Mali	1020	-		2500-3500
[8]					5000
[1]	Indian wasteland		-		2500
[60]	-		-		700-1000
[64]	Nicaragua	1200	-	4.50	5000
	India		-	3.05	6700
[28]	India	-	-		5000
[56]	India	450	1.25	1.73	1733
Pers. Comm. Kumar 2005	India, Rajasthan	610	2.5	0.47	1172
	India, Rajasthan	610	2.5	0.13	313
Pers. Comm. Buismans 2005	Mali, Digini	1020	2	0.30	337
[46]	Zimbabwe	725	-	0.4	

873 ^a: AAR: Average Annual Rainfall (if not reported in publication, obtained from
874 <http://www.worldclimate.com>)
875

876

877 **References**

- 878 [1] Azam MM, Waris A, Nahar NM. Prospects and potential of fatty acid methyl esters
879 of some non-traditional seed oils for use as biodiesel in India. *Biomass &*
880 *Bioenergy* 2005; 29: 293-302.
- 881 [2] Tiwari AK, Kumar A, Raheman H. Biodiesel production from *Jatropha* (*Jatropha*
882 *curcas*) with high free fatty acids: an optimized process. *Biomass & Bioenergy*
883 2007; 31: 569-75.
- 884 [3] Lopez O, Foidl G, Foidl N. Production of biogas from *J. curcas* fruitshells. In:
885 Gübitz GM, Mittelbach M, Trabi M, editors. *Biofuels and industrial products from*
886 *Jatropha curcas* - Proceedings from the Symposium "Jatropha 97", Managua,
887 Nicaragua, Feb. 23-27, 1997. Graz, Austria: Dbv-Verlag; p. 118-122.
- 888 [4] Staubmann R, Foidl G, Foidl N, Gübitz GM, Lafferty RM, Valencia Arbizu VM.
889 Production of biogas from *J. curcas* seeds press cake. In: Gübitz GM, Mittelbach
890 M, Trabi M, editors. *Biofuels and industrial products from Jatropha curcas* -
891 Proceedings from the Symposium "Jatropha 97", Managua, Nicaragua, Feb. 23-27,
892 1997. Graz, Austria: Dbv-Verlag; p. 123-131.
- 893 [5] Gubitiz GM, Mittelbach M, Trabi M. Exploitation of the tropical oil seed plant
894 *Jatropha curcas* L. *Bioresource Technology* 1999; 67: 73-82.
- 895 [6] Tewari DN. *Jatropha & Biodiesel*. 1st ed. New Delhi: Ocean Books Ltd.; 2007.

- 896 [7] Radhakrishna P. Contribution of de-oiled cakes in carbon sequestration and as a
897 source of energy, in Indian agriculture - need for a policy initiative. In:
898 Proceedings of the 4th international biofuels conference, New Delhi, India, Feb. 1-
899 2, 2007. New Delhi: Winrock International India; p. 65-70.
- 900 [8] Francis G, Edinger R, Becker K. A concept for simultaneous wasteland
901 reclamation, fuel production, and socio-economic development in degraded areas in
902 India: need, potential and perspectives of *Jatropha* plantations. Natural Resources
903 Forum 2005; 29: 12-24.
- 904 [9] Heller J. Physic nut. *Jatropha curcas* L. Promoting the conservation and use of
905 underutilized and neglected crops. 1. PhD dissertation, Institute of Plant Genetic
906 and Crop Plant Research, Gatersleben, Germany & International Plant Genetic
907 Resource Institute, Rome, Italy; 1996.
908 <http://www.ipgri.cgiar.org/Publications/pdf/161.pdf>
- 909 [10] Openshaw K. A review of *Jatropha curcas*: an oil plant of unfulfilled promise.
910 Biomass & Bioenergy 2000; 19: 1-15.
- 911 [11] Henning RK. Fuel production improves food production: the *Jatropha* project in
912 Mali. In: Gübitz GM, Mittelbach M, Trabi M, editors. Biofuels and industrial
913 products from *Jatropha curcas* - Proceedings from the Symposium "Jatropha 97",
914 Managua, Nicaragua, Feb. 23-27, 1997. Graz, Austria: Dbv-Verlag; p. 92-97.
- 915 [12] Henning RK. The *Jatropha* System. <http://www.jatropha.de/>. Feb. 12, 2006; last

- 916 modified on Jan. 30, 2006.
- 917 [13] Foidl N, Foidl G, Sanchez M, Mittelbach M, Hackel S. *Jatropha curcas* L. as a
918 source for the production of biofuel in Nicaragua. *Bioresource Technology* 1996;
919 58: 77-82.
- 920 [14] Sharma N. Pest disease management. In: FACT seminar on *Jatropha curcas* L.
921 agronomy and genetics, Wageningen, The Netherlands, Mar. 26-28, 2007.
922 Wageningen: FACT Foundation; p. Article number 26.
- 923 [15] Rao VR. The *Jatropha* hype: promise and performance. In: Singh B, Swaminathan
924 R, Ponraj V, editors. *Proceedings of the biodiesel conference toward energy*
925 *independance - Focus of Jatropha*, Hyderabad, India, June 9-10, 2006. New Delhi:
926 Rashtrapati Bhawan; p. 16-20.
- 927 [16] Janulis P. Reduction of energy consumption in biodiesel fuel life cycle. *Renewable*
928 *Energy* 2004; 29: 861-71.
- 929 [17] Baelemans A, Muys B. A critical evaluation of environmental assessment tools for
930 sustainable forest management. In: Ceuterick D, editors. *Proceedings of the*
931 *international conference on Life Cycle Assessment in agriculture, agro-industry and*
932 *forestry.*, Brussels, Belgium, Dec 3-4, 1998. Brussels: p. 65-75.
- 933 [18] Fredriksson H, Baky A, Bernesson S, Nordberg A, Noren O, Hansson P-A. Use of
934 on-farm produced biofuels on organic farms - Evaluation of energy balances and
935 environmental loads for three possible fuels. *Agricultural Systems* 2006; 89: 184-

936 203.

937 [19] Tan RR, Culaba AB, Purvis MRI. Polcage 1.0 - a possibilistic life-cycle assessment
938 model for evaluating alternative transportation fuels. Environmental Modelling &
939 Software 2004; 19: 907-18.

940 [20] Tan RR, Culaba AB, Purvis MRI. Application of possibility theory in the life-cycle
941 inventory assessment of biofuels. International Journal of Energy Research 2002;
942 26: 737-45.

943 [21] Zemanek G, Reinhardt.G.A. Notes on life-cycle assessments of vegetable oils.
944 Lipid - Fett 1999; 101: 321-7.

945 [22] USDA ANGRPNGRLBM. *Jatropha curcas* information from Germplasm
946 Resources Information Network - (GRIN) [Online Database]. [http://www.ars-](http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?20692)
947 [grin.gov/cgi-bin/npgs/html/taxon.pl?20692](http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?20692). Nov. 9, 2007; last updated on May 2nd,
948 2000.

949 [23] Raju AJS, Ezradanam V. Pollination ecology and fruiting behaviour in a
950 monoecious species, *Jatropha curcas* L. (*Euphorbiaceae*). Current Science 2002;
951 83: 1395-8.

952 [24] Prakash AR, Patolia JS, Chikara J, Boricha GN. Floral biology and flowering
953 behaviour of *Jatropha curcas*. In: FACT seminar on *Jatropha curcas* L. agronomy
954 and genetics, Wageningen, The Netherlands, Mar. 26-28, 2007. Wageningen:
955 FACT Foundation; p. Article number 2.

- 956 [25] Makkar HPS, Francis G, Becker K. Bioactivity of phytochemicals in some lesser-
957 known plants and their effects and potential applications in livestock and
958 aquaculture production systems. *Animal* 2007; 1: 1371-91.
- 959 [26] Martinez-Herrera J, Siddhuraju P, Francis G, Davila-Ortiz G, Becker K. Chemical
960 composition, toxic/antimetabolic constituents, and effects of different treatments on
961 their levels, in four provenances of *Jatropha curcas* L. from Mexico. *Food*
962 *Chemistry* 2006; 96: 80-9.
- 963 [27] Makkar HPS, Aderibigbe AO, Becker K. Comparative evaluation of non-toxic and
964 toxic varieties of *Jatropha curcas* for chemical composition, digestibility, protein
965 degradability and toxic factors. *Food Chemistry* 1998; 62: 207-15.
- 966 [28] Aderibigbe AO, Johnson COLE, Makkar HPS, Becker K, Foidl N. Chemical
967 composition and effect of heat on organic matter- and nitrogen-degradability and
968 some antinutritional components of *Jatropha* meal. *Animal Feed Science and*
969 *Technology* 1997; 67: 223-43.
- 970 [29] Aregheore EM , Becker K, Makkar HPS. Detoxification of a toxic variety of
971 *Jatropha curcas* using heat and chemical treatments, and preliminary nutritional
972 evaluation with rats. *South Pacific Journal of Natural Science* 2003; 21: 50-6.
- 973 [30] Makkar HPS, Becker K, Schmook B. Edible provenances of *Jatropha curcas* from
974 Quintana Roo state of Mexico and effect of roasting on antinutrient and toxic
975 factors in seeds. *Plant Foods for Human Nutrition* 1998; 52: 31-6.

- 976 [31] Makkar HPS, Becker K, Sporer F, Wink M. Studies on nutritive potential and toxic
977 constituents of different provenances of *Jatropha curcas*. Journal of Agricultural
978 and Food Chemistry 1997; 45: 3152-7.
- 979 [32] Aregheore EM , Makkar HPS, Becker K. Lectin activity in toxic and non-toxic
980 varieties of *J. curcas* using a latex agglutination test. In: Gübitz GM, Mittelbach
981 M, Trabi M, editors. Biofuels and industrial products from *Jatropha curcas* -
982 Proceedings from the Symposium "Jatropha 97", Managua, Nicaragua, Feb. 23-27,
983 1997. Graz, Austria: Dbv-Verlag; p. 65-69.
- 984 [33] Makkar HPS, Becker K. Potential of *J. curcas* seed meal as a protein supplement to
985 livestock feed, constraints to its utilisation and possible strategies to overcome
986 constraints. In: Gübitz GM, Mittelbach M, Trabi M, editors. Biofuels and
987 industrial products from *Jatropha curcas* - Proceedings from the Symposium
988 "Jatropha 97", Managua, Nicaragua, Feb. 23-27, 1997. Graz, Austria: Dbv-Verlag;
989 p. 190-205.
- 990 [34] Gour VK. Production practices including post-harvest management of *Jatropha*
991 *curcas*. In: Singh B, Swaminathan R, Ponraj V, editors. Proceedings of the
992 biodiesel conference toward energy independance - Focus of *Jatropha*, Hyderabad,
993 India, June 9-10, 2006. New Delhi: Rashtrapati Bhawan; p. 223-251.
- 994 [35] Singh L, Bargali SS, Swamy SL. Production practices and post-harvest
995 management. In: Singh B, Swaminathan R, Ponraj V, editors. Proceedings of the
996 biodiesel conference toward energy independance - Focus of *Jatropha*, Hyderabad,

- 997 India, June 9-10, 2006. New Delhi: Rashtrapati Bhawan; p. 252-267.
- 998 [36] Biswas S, Kaushik N, Srikanth G. Biodiesel: technology and business opportunities
 999 - an insight. In: Singh B, Swaminathan R, Ponraj V, editors. Proceedings of the
 1000 biodiesel conference toward energy independance - Focus of *Jatropha*, Hyderabad,
 1001 India, June 9-10, 2006. New Delhi: Rashtrapati Bhawan; p. 303-330.
- 1002 [37] TERI. Mycorrized *Jatropha*: seeds of sustainability.
 1003 <http://www.teriin.org/jatropha.php>. Dec. 19, 2006; Last modified Dec. 19, 2006.
- 1004 [38] Henning RK. The *Jatropha* Booklet - A guide to the *Jatropha* system and its
 1005 dissemination in Zambia. 1st ed. Weissensberg: bagani GbR; 2000.
- 1006 [39] Zahawi RA. Establishment and growth of living fence species: an overlooked tool
 1007 for the restoration of degraded areas in the tropics. *Restoration Ecology* 2005; 13:
 1008 92-102.
- 1009 [40] Soares Severino L, de Lourdes Silva de Lima R, Leão AB, de Macedo Deltrão NE.
 1010 Root system characteristics of *Jatropha curcas* plants propagated through five
 1011 methods. In: FACT seminar on *Jatropha curcas* L. agronomy and genetics,
 1012 Wageningen, The Netherlands, Mar. 26-28, 2007. Wageningen: FACT Foundation;
 1013 p. Article number 31.
- 1014 [41] Jongschaap REE, Corré WJ, Bindraban PS, Brandenburg WA. Claims and facts on
 1015 *Jatropha curcas* L. Wageningen, The Netherlands: Plant Research International B.V; 2007.
 1016 http://www.fact-fuels.org/media_en/Claims_and_Facts_on_Jatropha_-

- 1017 WUR?session=isgsklbna58j7grfst888n5r7
- 1018 [42] Kaushik N, Kumar S. *Jatropha curcas* L. silviculture and uses. 2nd revised ed.
 1019 Jodhpur: Agrobios; 2006.
- 1020 [43] Gosh L, Singh L. Variation in rooting response in cuttings of *Jatropha curcas* L.
 1021 In: National seminar on "*Jatropha* cultivation and Bio-diesel production",
 1022 Allahabad, India, Jan. 27-28, 2007. Allahabad: Utthan - Centre for Sustainable
 1023 Development & Poverty Alleviation;
- 1024 [44] Brahmam M. Enhancing germination percentage of *Jatropha curcas* L. by pre-
 1025 sowing. In: Proceedings of the 4th international biofuels conference, New Delhi,
 1026 India, Feb. 1-2, 2007. New Delhi: Winrock International India; p. 175-180.
- 1027 [45] Chikara J, Ghosh A, Patolia JS, Chaudhary DR, Zala A. Productivity of *Jatropha*
 1028 *curcas* under different spacing. In: FACT seminar on *Jatropha curcas* L.
 1029 agronomy and genetics, Wageningen, The Netherlands, Mar. 26-28, 2007.
 1030 Wageningen: FACT Foundation; p. Article number 9.
- 1031 [46] Tigere TA, Gatsi TC, Mudita II, Chikuvire TJ, Thamangani S, Mavunganidze Z.
 1032 Potential of *Jatropha curcas* in improving smallholder farmers' livelihoods in
 1033 Zimbabwe: *An exploratory study of Makosa Ward, Mutoko District*. Journal of
 1034 Sustainable Development in Africa 2006; 8: 1-9.
- 1035 [47] Euler H, Gorriz D. Case study "*Jatropha curcas*". Frankfurt: Global Facilitation
 1036 Unit for Underutilized Species and Deutsche Gesellschaft für Technische

- 1037 Zusammenarbeit; 2004.
- 1038 [48] Patolia JS, Ghosh A, CHikara J, Chaudhary DR, Parmar DR, Bhuva HM. Response
1039 of *Jatropha curcas* grown on wasteland to N and P fertilization. In: FACT seminar
1040 on *Jatropha curcas* L. agronomy and genetics, Wageningen, The Netherlands, Mar.
1041 26-28, 2007. Wageningen: FACT Foundation; p. Article number 34.
- 1042 [49] Grimm C, Maes J-M. Arthropod fauna associated with *Jatropha curcas* L. in
1043 Nicaragua: A synopsis of species, their biology and pest status. In: Gübitz GM,
1044 Mittelbach M, Trabi M, editors. Biofuels and industrial products from *Jatropha*
1045 *curcas* - Proceedings from the Symposium "Jatropha 97", Managua, Nicaragua,
1046 Feb. 23-27, 1997. Graz, Austria: Dbv-Verlag; p. 31-39.
- 1047 [50] Shanker C, Dhyani SK. Insect pests of *Jatropha curcas* L. and the potential for
1048 their management. Current Science 2006; 91: 162-3.
- 1049 [51] Kaushik N. Quality planting material and seed standards in *Jatropha curcas*. In:
1050 Singh B, Swaminathan R, Ponraj V, editors. Proceedings of the biodiesel
1051 conference toward energy independance - Focus of *Jatropha*, Hyderabad, India, 9-
1052 10 June, 2006. New Delhi: Rashtrapati Bhawan; p. 179-196.
- 1053 [52] Foidl N, Eder P. Agro-industrial exploitation of *J. curcas*. In: Gübitz GM,
1054 Mittelbach M, Trabi M, editors. Biofuels and industrial products from *Jatropha*
1055 *curcas* - Proceedings from the Symposium "Jatropha 97", Managua, Nicaragua,
1056 Feb. 23-27, 1997. Graz, Austria: Dbv-Verlag; p. 88-91.

- 1057 [53] Jones N, Miller JH. *Jatropha curcas*: a multipurpose species for problematic sites.
1058 Washington DC: The World Bank; 1992.
- 1059 [54] Aker CL. Growth and reproduction of *J. curcas*. In: Gübitz GM, Mittelbach M,
1060 Trabi M, editors. Biofuels and industrial products from *Jatropha curcas* -
1061 Proceedings from the Symposium "Jatropha 97", Managua, Nicaragua, Feb. 23-27,
1062 1997. Graz, Austria: Dbv-Verlag; p. 2-18.
- 1063 [55] Ginwal HS, Rawat PS, Srivastava RL. Seed source variation in growth
1064 performance and oil yield of *Jatropha curcas* Linn. in Central India. *Silvae*
1065 *Genetica* 2004; 53: 186-92.
- 1066 [56] Sharma GD, Gupta SN, Khabiruddin M. Cultivation of *Jatropha curcas* as future
1067 source of hydrocarbon and other industrial products. In: Gübitz GM, Mittelbach M,
1068 Trabi M, editors. Biofuels and industrial products from *Jatropha curcas* -
1069 Proceedings from the Symposium "Jatropha 97", Managua, Nicaragua, Feb. 23-27,
1070 1997. Graz, Austria: Dbv-Verlag; p. 19-21.
- 1071 [57] Spaan WP, Bodnar F, Idoe O, de Graaff J. Implementation of contour vegetation
1072 barriers under farmer conditions in Burkina Faso and Mali. *Quarterly Journal of*
1073 *International Agriculture* 2004; 43: 21-38.
- 1074 [58] Benge M. Assessment of the potential of *Jatropha curcas*, (Biodiesel tree,) for
1075 energy production and other uses in developing countries. ECHO's website: 2006.
1076 http://www.echotech.org/mambo/index.php?option=com_docman&task=doc_view

1077 &gid=179&Itemid=68

1078 [59] Rivera-Locra JA, KU-Vera JC. Chemical composition of three different varieties of
1079 *J. curcas* from Mexico. In: Gübitz GM, Mittelbach M, Trabi M, editors. Biofuels
1080 and industrial products from *Jatropha curcas* - Proceedings from the Symposium
1081 "Jatropha 97", Managua, Nicaragua, Feb. 23-27, 1997. Graz, Austria: Dbv-Verlag;
1082 p. 47-52.

1083 [60] FACT Foundation. *Jatropha* Handbook. Eindhoven: FACT Foundation; 2006.
1084 <http://www.fact-fuels.org/images/jatropha%20handbook%20march%202006.pdf>

1085 [61] Kaushik N, Kumar K, Kumar S, Kaushik N, Roy S. Genetic variability and
1086 divergence studies in seed traits and oil content of *Jatropha* (*Jatropha curcas* L.)
1087 accessions. *Biomass & Bioenergy* 2007; 31: 497-502.

1088 [62] Vyas DK, Singh RN. Feasibility study of *Jatropha* seed husk as an open core
1089 gasifier feedstock. *Renewable Energy* 2007; 32: 512-7.

1090 [63] Forson FK, Oduro EK, Hammond-Donkoh E. Performance of *Jatropha* oil blends
1091 in a diesel engine. *Renewable Energy* 2004; 29: 1135-45.

1092 [64] Tobin J, Fulford DJ. Life Cycle Assessment of the production of biodiesel from
1093 *Jatropha*. Msc dissertation , The University of Reading; 2005.

1094 [65] Henning RK. Combating desertification - Fuel from *Jatropha* plants. In: UNIDO
1095 Symposium on development and utilization of biomass energy resources in

- 1096 developing countries, Vienna, Austria, Dec. 1995. Vienna: UNIDO, Environment
1097 and Energy Branch, Industrial Sectors and Environment Division;
- 1098 [66] Henning RK. Use of *Jatropha curcas* L. (JCL): A household perspective and its
1099 contribution to rural employment creation - Experiences of the *Jatropha* project in
1100 Mali, West Africa, 1987 to 1997. Presentation: Regional Workshop on the
1101 Potential of *Jatropha curcas* in Rural Development & Environmental Protection,
1102 Harare, Zimbabwe, May 1998. <http://www.jatropha.de/harare98.htm>
- 1103 [67] Rabé ELM, Somers LMT, Goey LPH. *Jatropha* oil in copression ignition engines -
1104 Effects on the engine, environment and Tanzania as supplying country. Msc
1105 dissertation, Eindhoven University of Technology; 2005.
- 1106 [68] Beerens P. Screw-pressing of *Jatropha* seeds for fueling purposes in less
1107 developed countries. Msc dissertation, Eindhoven University of Technology; 2007.
- 1108 [69] Winkler E, Gübitz GM, Foidl N, Staubmann R, Steiner W. Use of enzymes for oil
1109 extraction from *J. curcas* seeds. In: Gübitz GM, Mittelbach M, Trabi M, editors.
1110 Biofuels and industrial products from *Jatropha curcas* - Proceedings from the
1111 Symposium "Jatropha 97", Managua, Nicaragua, Feb. 23-27, 1997. Graz, Austria:
1112 Dbv-Verlag; p. 184-189.
- 1113 [70] Shah S, Sharma A, Gupta MN. Extraction of oil from *Jatropha curcas* L. seed
1114 kernels by combination of ultrasonication and aqueous enzymatic oil extraction.
1115 Bioresource Technology 2005; 96: 121-3.

- 1116 [71] Augustus GDPS, Jayabalan M, Seiler GJ. Evaluation and bioinduction of energy
1117 components of *Jatropha curcas*. Biomass & Bioenergy 2002; 23: 161-4.
- 1118 [72] Shah S, Sharma A, Gupta MN. Extraction of oil from *Jatropha curcas* L. seed
1119 kernels by enzyme assisted three phase partitioning. Industrial Crops and Products
1120 2004; 20: 275-9.
- 1121 [73] Adriaans, T. Suitability of solvent extraction for *Jatropha curcas*. Eindhoven:
1122 FACT Foundation; 2006:9.
- 1123 [74] Foidl N, Mayorga L. Produccion de ester metilico del tempate (EMAT) como
1124 sustituto del combustible diesel. In: FACT seminar on *Jatropha curcas* L.
1125 agronomy and genetics, Wageningen, The Netherlands, Mar. 26-28, 2007.
1126 Wageningen: FACT Foundation;
- 1127 [75] Kandpal JB, Madan M. *Jatropha curcas* - a renewable source of energy for meeting
1128 future energy needs. Renewable Energy 1995; 6: 159-60.
- 1129 [76] Reddy JN, Ramesh A. Parametric studies for improving the performance of a
1130 *Jatropha* oil-fuelled compression ignition engine. Renewable Energy 2006; 31:
1131 1994-2016.
- 1132 [77] Adebawale KO , Adedire CO. Chemical composition and insecticidal properties of
1133 the underutilized *Jatropha curcas* seed oil. African Journal of Biotechnology 2006;
1134 5: 901-6.

- 1135 [78] Kpoviessi DSS, Accrombessi GC, Kossouh C, Soumanou MM, Moudachirou M.
 1136 Physicochemical properties and composition of pignut (*Jatropha curcas*) non-
 1137 conventional oil from different regions of Benin. *Comptes Rendus Chimie* 2004; 7:
 1138 1007-12.
- 1139 [79] Akintayo ET. Characteristics and composition of *Parkia biglobbosa* and *Jatropha*
 1140 *curcas* oils and cakes. *Bioresource Technology* 2004; 92: 307-10.
- 1141 [80] Kumar MS, Ramesh A, Nagalingam B. An experimental comparison of methods to
 1142 use methanol and *Jatropha* oil in a compression ignition engine. *Biomass &*
 1143 *Bioenergy* 2003; 25: 309-18.
- 1144 [81] Pramanik K. Properties and use of *Jatropha curcas* oil and diesel fuel blends in
 1145 compression ignition engine. *Renewable Energy* 2003; 28: 239-48.
- 1146 [82] Banerji R, Chowdhury AR, Misra G, Sudarsanam G, Verma SC, Srivastava GS.
 1147 *Jatropha* seed oils for energy. *Biomass* 1985; 8: 277-82.
- 1148 [83] Chitra P, Venkatachalam P, Sampathrajan A. Optimisation of experimental
 1149 conditions for biodiesel production from alkali-catalysed transesterification of
 1150 *Jatropha curcas* oil. *Energy for Sustainable Development* 2005; 9: 13-8.
- 1151 [84] Vaitillingom G, Liennard A. Various vegetable oils as fuel for diesel and burners:
 1152 *J. curcas* particularities. In: Gübitz GM, Mittelbach M, Trabi M, editors. *Biofuels*
 1153 *and industrial products from Jatropha curcas* - Proceedings from the Symposium
 1154 "Jatropha 97", Managua, Nicaragua, Feb. 23-27, 1997. Graz, Austria: Dbv-Verlag;

- 1155 p. 98-105.
- 1156 [85] Zamora E, Sobalvarro E, Norori R, Foidl G, Foidl N. Trans-esterification of *J.*
 1157 *curcas* oil. In: Gübitz GM, Mittelbach M, Trabi M, editors. Biofuels and industrial
 1158 products from *Jatropha curcas* - Proceedings from the Symposium "Jatropha 97",
 1159 Managua, Nicaragua, Feb. 23-27, 1997. Graz, Austria: Dbv-Verlag; p. 113-117.
- 1160 [86] Ma F, Hanna MA. Biodiesel production: a review. Bioresource Technology 1999 ;
 1161 70: 1-15.
- 1162 [87] Meher LC, Vidya Sagar D, Naik SN. Technical aspects of biodiesel production by
 1163 transesterification - a review. Renewable and Sustainable Energy Reviews 2006;
 1164 10: 248-68.
- 1165 [88] Van Gerpen J. Biodiesel processing and production: biodiesel processing and
 1166 production. Fuel Processing Technology 2005; 86: 1097-107.
- 1167 [89] Raina AK, Gaikwad BR. Chemobotany of *Jatropha* species in India and further
 1168 characterisation of curcas oil. Journal of oil technology of India 1987; 19: 81-5.
- 1169 [90] Wani SP, Sreedevi TK. Strategy for rehabilitation of degraded lands and improved
 1170 livelihoods through biodiesel plantations. In: Proceedings of the 4th international
 1171 biofuels conference, New Delhi, India, Feb. 1-2, 2007. New Delhi: Winrock
 1172 International India; p. 50-64.
- 1173 [91] Rug M, Sporer F, Wink M, Liu SY, Henning R, Ruppel A. Molluscicidal properties

- 1174 of *J. curcas* against vector snails of the human parasites *Schistosoma mansoni* and
 1175 *S. japonicum*. In: Gübitz GM, Mittelbach M, Trabi M, editors. Biofuels and
 1176 industrial products from *Jatropha curcas* - Proceedings from the Symposium
 1177 "Jatropha 97", Managua, Nicaragua, Graz, Austria: Dbv-Verlag; p. 227-232 .
- 1178 [92] Henning R, Samaké F, Thiéro I. La valeur fertilisante du tourteau du pourghère.
 1179 Bamako, Mali: Projet Pourghère DNHE - GTZ; 1995.
- 1180 [93] Ngoma T. Fertilizer, manure and *Jatropha* cake application trial.
 1181 <http://www.jatropha.de/Zimbabwe/zw-b-fertil.htm>. Feb. 12, 2007; last modified on
 1182 Feb. 20, 2003.
- 1183 [94] Ghosh A, Patolia JS, Chaudhary DR, Chikara J, Rao SN, Kumar D, Boricha GN,
 1184 Zala A. Respons of *Jatropha curcas* under different spacing to *Jatropha* de-oiled
 1185 cake. In: FACT seminar on *Jatropha curcas* L. agronomy and genetics,
 1186 Wageningen, The Netherlands, Mar. 26-28, 2007. Wageningen: FACT Foundation;
 1187 p. Article number 8.
- 1188 [95] Barnwal BK, Sharma MP. Prospects of biodiesel production from vegetables oils in
 1189 India. Renewable & Sustainable Energy Reviews 2005; 9: 363-78.
- 1190 [96] Ramadhas AS, Jayaraj S, Muraleedharan C. Use of vegetable oils as I.C. engine
 1191 fuels - a review. Renewable Energy 2004; 29: 727-42.
- 1192 [97] Canakci M, Erdil A, Arcaklioglu E. Performance and exhaust emissions of a
 1193 biodiesel engine. Applied Energy 2006; 83: 594-605.

- 1194 [98] Agarwal AK. Biofuels (alcohols and biodiesel) applications as fuels for internal
1195 combustion engines. *Progress in Energy and Combustion Science* 2007; 33: 233-
1196 71.
- 1197 [99] Prasad CMV, Krishna MVSM, Reddy CP, Mohan KR. Performance evaluation of
1198 non-edible vegetable oils as substitute fuels in low heat rejection diesel engines.
1199 *Proceedings of the Institution of Mechanical Engineers Part D - Journal of*
1200 *automobile engineering* 2000; 214: 181-7.
- 1201 [100] Zhu HP, Wu ZB, Chen YX, Zhang P, Duan SJ, Liu XH et al. Preparation of
1202 biodiesel catalyzed by solid super base of calcium oxide and its refining process.
1203 *Chinese Journal of Catalysis* 2006; 27: 391-6.
- 1204 [101] Harvey AP, Khurana R, Lee JGM. In-situ transesterification of *Jatropha curcas*
1205 seed to produce biodiesel, using a mixture of alcohols. In: *Proceedings of the 4th*
1206 *international biofuels conference, New Delhi, India, Feb. 1-2, 2007. New Delhi:*
1207 *Winrock International India; p. 187-193.*
- 1208 [102] Rathore V, Madras G. Synthesis of biodiesel from edible and non-edible oils in
1209 supercritical alcohols and enzymatic synthesis in supercritical carbon dioxide. *Fuel*
1210 2007; 86: 2650-9.
- 1211 [103] Kaul S, Saxena RC, Kumar A, Negi MS, Bhatnagar AK, Goyal HB et al.
1212 Corrosion behavior of biodiesel from seed oils of Indian origin on diesel engine
1213 parts. *Fuel Processing Technology* 2007; 88: 303-7.

- 1214 [104] Basha SD, Sujatha M. Inter and intra-population variability of *Jatropha curcas* (L.)
 1215 characterized by RAPD and ISSR markers and development of population-specific
 1216 SCAR markers. *Euphytica* 2007; 156: 375-86.
- 1217 [105] Kaushik N, Kaushik JC, Kumar S. Response of *Jatropha* seedlings to seed size and
 1218 growing medium. *Journal of Non-Timber Forest Products* 2003; 10: 40-2.
- 1219 [106] Sunil N, Varaprasad KS, Sivaraj N, Suresh Kumar T, Abraham B, Prasad RBN.
 1220 Assessing *Jatropha curcas* L. germplasm in-situ--A case study. *Biomass &*
 1221 *Bioenergy* In Press, Corrected Proof.
- 1222 [107] Bernesson S , Nilsson D, Hansson PA. A limited LCA comparing large- and small-
 1223 scale production of ethanol for heavy engines under Swedish conditions. *Biomass*
 1224 *& Bioenergy* 2006; 30: 46-57.
- 1225 [108] Hill J, Nelson E, Tilman D, Polasky S, Tiffany D. Environmental, economic, and
 1226 energetic costs and benefits of biodiesel and ethanol biofuels. *Proceedings of the*
 1227 *National Academy of Sciences of the United States of America* 2006; 103: 11206-
 1228 10.
- 1229 [109] Lettens S, Muys B, Ceulemans R, Moons E, Garcia J, Coppin P. Energy budget and
 1230 greenhouse gas balance evaluation of sustainable coppice systems for electricity
 1231 production. *Biomass & Bioenergy* 2003; 24: 179-97.
- 1232 [110] Bernesson S , Nilsson D, Hansson PA. A limited LCA comparing large- and small-
 1233 scale production of rape methyl ester (RME) under Swedish conditions. *Biomass &*

- 1234 Bioenergy 2004; 26: 545-59.
- 1235 [111] Elsayed MA, Matthews R, Mortimer ND. Carbon and energy balances for a range
1236 of biofuel options. Sheffield: Resources Research Unit Sheffield Hallam
1237 University; 2003.
1238 <http://www.dti.gov.uk/files/file14925.pdf?pubpdfdownload=03%2F836>
- 1239 [112] Ahlvik P, Brandberg A. Well-to-wheel efficiency for alternative fuels from natural
1240 gas or biomass. Borlänge: Swedish National Road Administration; 2001.
1241 <http://www.vv.se/filer/publikationer/2001-85.pdf>
- 1242 [113] Elsayed MA, Mortimer ND. Carbon and energy modelling of biomass systems:
1243 conversion plant and data updates. Sheffield: Resources Research Unit Sheffield
1244 Hallam University; 2001.
1245 <http://www.dti.gov.uk/files/file14926.pdf?pubpdfdownload=01%2F1342>
- 1246 [114] Kim S, Dale BE. Life Cycle Assessment of various cropping systems utilized for
1247 producing biofuels: bioethanol and biodiesel. Biomass & Bioenergy 2005; 29: 426-
1248 39.
- 1249 [115] ISO 14040 - 14043. Brussels: Comité Européen de Normalisation (CEN); 1997.
- 1250 [116] Mattsson B, Cederberg C, Blix L. Agricultural land use in life cycle assessment
1251 (LCA): case studies of three vegetable oil crops. Journal of Cleaner Production
1252 2000; 8: 283-92.

- 1253 [117] Lindeijer E , Müller-Wenk R, Steen B. Impact assessment of resources and land
1254 use. in: Udo de Haes HA, Finnveden G, Goedkoop M, Hauschild M, Hertwich EG,
1255 Hofstetter P et al. editors. Life-cycle Impact Assessment: Striving towards Best
1256 Practice, Pensacola, FL: Society of Environmental Toxicology and Chemistry;
1257 2002, p. 11-64.
- 1258 [118] Wagendorp T , Gulinck H, Coppin P, Muys B. Land use impact evaluation in life
1259 cycle assessment based on ecosystem thermodynamics. *Energy* 2006; 31: 112-25.
- 1260 [119] Prueksakorn K, Gheewala SH. Energy and greenhouse gas implications of
1261 biodiesel production from *Jatropha curcas* L. In: Proceedings of The 2nd Joint
1262 International Conference on "Sustainable Energy and Environments (SEE 2006)",
1263 Bangkok, Thailand, 2006 Nov 21-23 .
- 1264 [120] Achten WMJ, Mathijs E, Verchot L, Singh VP, Aerts R, Muys B. *Jatropha*
1265 biodiesel fueling sustainability? *Biofuels, Bioproducts and Biorefining* 2007 .
- 1266 [121] Crutzen PJ, Mosier AR, Smith KA, Winiwarter W. N₂ O release from agro-biofuel
1267 production negates global warming reduction by replacing fossil fuels.
1268 *Atmospheric Chemistry and Physics Discussions* 2007; 7: 11191-205.
- 1269 [122] IPCC. N₂O emissions from managed soils, and CO₂ emissions from lime and urea
1270 application. Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K., editors.
1271 Hayama, Japan: IGES; 2006,
- 1272 [123] Ogunwole JO , Patolia JS, Chaudhary DR, Ghosh A, Chikara J. Improvement of

- 1273 the quality of a degraded entisol with *Jatropha curcas* L. under an Indian semi arid
1274 condition. In: FACT seminar on *Jatropha curcas* L. agronomy and genetics,
1275 Wageningen, The Netherlands, Mar. 26-28, 2007. Wageningen: FACT Foundation;
1276 p. Article number 10.
- 1277 [124] WILMA. Land rehabilitation, soil conservation, and biofuels production around the
1278 Masinga reservoir in Kenya. WILMA Concept Brief 2005; September: 1-10.
- 1279 [125] Vos P. Case studies on the status of invasive woody plant species in the Western
1280 Indian Ocean: 2. The Comoros Archipelago (Union of the Comoros and Mayotte).
1281 Forestry Department, Food and Agriculture Organization of the United Nations.
1282 Rome, Italy: FAO; 2004.
1283 <http://www.fao.org/forestry/webview/media?mediaId=6840&langId=1>
- 1284 [126] von Oertzen I, Smith J. Focus on - invasive alien species in Cambodia. Biodiversity
1285 Threats. [http://www.biodiversity-](http://www.biodiversity-kh.org/Englishpage/File%20Download/NBSAPEng/Status%20Report/Biodiversity%20Threat%20(Eng).pdf)
1286 [kh.org/Englishpage/File%20Download/NBSAPEng/Status%20Report/Biodiversity%20Threat%20\(Eng\).pdf](http://www.biodiversity-kh.org/Englishpage/File%20Download/NBSAPEng/Status%20Report/Biodiversity%20Threat%20(Eng).pdf)
1287
- 1288 [127] Low T, Booth C. The weedy truth about biofuels. Melbourne, Australia: Invasive
1289 Species Council; 2007.
- 1290 [128] Gressel J. Transgenics are imperative for biofuel crops. Plant Science In Press,
1291 Corrected Proof.
- 1292 [129] Horiuchi T, Fujiki H, Hirota M, Suttajit M, Suganuma M, Yoshioka A et al.

1293 Presence of tumor promoters in the seed oil of *Jatropha curcas* L. from Thailand.
1294 Japanese Journal of Cancer Research 1987; 78: 223-6.

1295 [130] Hirota M, Suttajit M, Suguri H, Endo Y, Shudo K, Wongchai V et al. A new tumor
1296 promoter from the seed oil of *Jatropha curcas* L., an intramolecular diester of 12-
1297 deoxy-16-hydroxyphorbol. Cancer Research 1988; 48: 5800-4.

1298 [131] Lin J, Yan F, Tang L, Chen F. Antitumor effects of curcin from seeds of *Jatropha*
1299 *curcas*. Acta Pharmacologica Sinica 2003; 24: 241-6.

1300 [132] Luo MJ, Yang XY, Liu WX, Xu Y, Huang P, Yan F et al. Expression, purification
1301 and anti-tumor activity of curcin. Acta Biochimica Et Biophysica Sinica 2006; 38:
1302 663-8.

1303

Fig. captions

Fig. 1. Inputs and outputs of the *Jatropha* cultivation unit process.

Fig. 2. Dry seed yield in relation with average annual rainfall (mm) and age of the JCL crop. The plotted points represent a mix of provenances, site conditions and plant age or average annual rainfall respectively.

Sources: [9,11,13,52,53,56] and personal communication Kumar 2005 and Buisman 2005.

Fig. 3. Average shell:kernel ratio of JCL seed and standard deviation based on 21 reported data sets.

Sources: [9,10,13,26,28,30,31,59,60]

Fig. 4. Average kernel and shell composition and standard deviation based on *n* reported data sets. ♦ - minimum; ◇ - maximum.

* Calculated from values obtained for fat free samples since high fat content interferes with the analyses.

Sources: [9,10,13,27,30,31,55,59,60]

Fig. 5. Inputs and outputs of the *Jatropha* oil extraction unit process.

Fig. 6. Fatty Acid composition (%). C16:0 = Palmitic Acid; C18:0 = Stearic Acid; C18:1 Oleic Acid; C18:2 = Linoleic Acid. Other Acids containing Capric Acid, Myristic Acid (C14:0), Palmitoleic Acid (C16:1), Linolenic Acid (C18:3), Arachidic Acid (C20:0),

1326 Behenic Acid (C22:0), cis-11-Eicosenoic Acid (C20:1) and cis-11,14-Eicosadienoic Acid
1327 (C20:2). n = number of observations used.

1328 *Sources:* [1,13,26,56,71,75,77-79,82]

1329

1330 Fig. 7. Average kernel cake composition and standard deviation based on n reported data
1331 sets of solvent extracted JCL kernels. ♦ - minimum; ◇ - maximum. Inset table: ranges of
1332 reported chemical composition indicating the percentages of N, P, K, Ca and Mg (based on
1333 $n=5$ data sets).

1334 *Sources:* [5,6,8,10,26,28,31,59,71,90]

1335

1336 Fig. 8. Inputs and outputs of the transesterification unit process

1337

1338 Fig. 9. Primary energy input for the production of 1000 MJ JME after pro-rata allocation of
1339 the total energy requirement of the whole production process over the JME product and the
1340 by-products based on the energy content of the JME product and the by-products. *Sources:*
1341 [64] with low cultivation intensity and [119] with high cultivation intensity. Comparison
1342 with reference systems rapeseed methyl ester (RME) and ultra low sulphur diesel (ULSD)
1343 from crude oil [111].

Figures

Fig 1.

1347 Fig 2.

1348

1349 Fig 3.

1350

1351 Fig 4.

1352

1353 Fig 5.

1354

1355 Fig 6.

1356

1359 Fig 8

1360

1361 Fig 9.

1362

1363 **Tables**

Table 1
Calculated oil yields (% of contained oil) of mechanical
extraction methods

Press	Reference	Oil yield (%)
Engine driven screw press	[67]	68.0
	[6]	80.0
	[68]	79.0
Ram press	[63]	62.5
	[6]	62.5

1364

Table 2

Reported oil yields (% of contained oil) for different chemical extraction methods and different reaction parameters

Extraction Method	Reference	Reaction temperature (°C)	Reaction pH	Time consumption (hours)	Oil yield (%)
<i>n</i> -hexane oil extraction (Soxhelt apparatus)	[5,9,63]	-	-	24	95-99
1 st Acetone	[71]	-	-	48	
2 nd <i>n</i> -hexane					
Aqueous Oil Extraction (AOE)	[69]	-	-	2	38
	[70]	50	9	6	38
AOE with 10 minutes of ultrasonication as pretreatment	[70]	50	9	6	67
Aqueous Enzymatic Oil Extraction (AEOE) (hemicellulase or cellulase)	[69]	60	4.5	2	73
AEOE (alkaline protease)	[69]	60	7	2	86
	[70]	50	9	6	64
AEOE (alkaline protease) with 5 minutes of ultrasonication as pretreatment	[70]	50	9	6	74
Three phase partitioning	[72]	25	9	2	97

1365

1366

Table 3
JCL oil composition and characteristics.

	Range	Mean	SD	<i>n</i>
Specific gravity (g/cm ³)	0.860 – 0.933	0.914	0.018	13
Calorific value (MJ/kg)	37.83 – 42.05	39.63	1.52	9
Pour point (°C)	–3			2
Cloud point (°C)	2			1
Flash point (°C)	210 – 240	235	11	7
Cetane value	38.0 – 51.0	46.3	6.2	4
Saponification number (mg/g)	102.9 – 209.0	182.8	34.3	8
Viscosity at 30°C (cSt)	37.00 – 54.80	46.82	7.24	7
Free fatty acids % (kg/kg*100)	0.18 – 3.40	2.18	1.46	4
Unsaponifiable % (kg/kg*100)	0.79 – 3.80	2.03	1.57	5
Iodine number (mg iodine/g)	92 – 112	101	7	8
Acid number (mg KOH/g)	0.92 – 6.16	3.71	2.17	4
monoglycerides % (kg/kg*100)	nd – 1.7			1
Diglycerides % (kg/kg*100)	2.50 – 2.70			2
Triglycerides % (kg/kg*100)	88.20 – 97.30			2
Carbon residue % (kg/kg*100)	0.07 – 0.64	0.38	0.29	3
Sulfur content % (kg/kg*100)	0 – 0.13			2

SD = standard deviation; n = number of observations used; nd = no data.

Sources: [5,13,63,71,75-85]

Table 4
Summary of case studies using JCL seed cake as fertilizer.

Reference	Country	Crop	Dosage ^a	Comments
[92]	Mali	Pearl millet	5 t ha ⁻¹	46% yield increase in comparison to zero-input • 40 – 113 % yield increase in comparison to zero-input
[93]	Zimbabwe	Cabbage	2.5 – 10 t ha ⁻¹	• free from pest and disease, while cutworm infestation occurred with cow manure application
[9]	Nepal	Rice	10 t ha ⁻¹	11% yield increase in comparison to zero-input
[94]	India	JCL	0.75-3 t ha ⁻¹	13 – 120 % yield increase in comparison to zero-input

^a: 1 ton seed cake is produced on 0.27-0.54 ha JCL plantation. (own calculation based on an expected seed yield of 2.5 – 5 t dry seed/ha/year with an oil content of 34.4 wt% and a mechanical extraction efficiency of 75%).

Table 5

JCL (m)ethyl ester composition and characteristics with the corresponding values of the European (EN 14214:2003), German (DIN V 51606) and the USA Standards (ASTM D 6751).

	JME				JEE	EN	DIN V	ASTM
	Range	Mean	SD	n	n=1	14214:2003	51606	D6751
Density (g/cm ³)	0.864 – 0.880	0.875	0.007	6	0.89	0.86 - 0.90	0.87-0.90	
Calorific value (MJ/kg)	38.45 – 41.00	39.65	1.28	3				
Flash point (°C)	170 – 192	186	11	4	190	min 120	min 110	min 130
Cetane value	50.0 – 56.1	52.3	2.3	5	59	min 51	min 49	min 47
Saponification number (mg/g)	202.6			1				
Viscosity at 30°C (cSt)	4.84 – 5.65	5.11	0.47	3	5.54	3.5-5.0 ^a	3.5-5.0 ^a	1.9-6.0 ^a
Iodine number (mg iodine/g)	93 – 106			2		max 120	max 115	max 115
Acid number (mg KOH/g)	0.06 – 0.5	0.27	0.22	3	0.08	max 0.5	max 0.5	max 0.5
Monoglycerides % (kg/kg*100)	0.24			1	0.55	max 0.8	max 0.8	
Diglycerides % (kg/kg*100)	0.07			1	0.19	max 0.2	max 0.4	
Triglycerides % (kg/kg*100)	nd			0	nd	max 0.2	max 0.4	
Carbon residue % (kg/kg*100)	0.02 – 0.50	0.18	0.27	3		max 0.3	max 0.3	max 0.05
Sulfur content % (kg/kg*100)	0.0036			1		max 0.01	max 0.01	max 0.015 ^b
Sulfated ash % (kg/kg*100)	0.005 – 0.010	0.013	0.002	4		max 0.02	max 0.03	max 0.02
Methyl ester content % (kg/kg*100)	99.6			1	99.3	min 96.5		
Methanol % (kg/kg*100)	0.06 – 0.09			2	0.05	max 0.2	max 0.3	
Water % (kg/kg*100)	0.07 – 0.10			1	0.16	max 0.5	max 0.3	max 0.5
Free glycerol % (kg/kg*100)	0.015 – 0.030			2	nd	max 0.02	max 0.02	max 0.02
Total glycerol % (kg/kg*100)	0.088 – 0.100			2	0.17	max 0.25	max 0.25	max 0.24

SD = standard deviation; n = number of observations used; nd = no data. Sources: [5,13,63,71,75-85]

^a The standards include viscosity in mm²/s at 40°C. Francis et al. [8] reported a value of 4.2 mm²/s at 40°C for JME.

^b maximum 0.015% for S 15 Grade and maximum 0.05% for S 500 Grade

Table 6

Summary of reported alternative transesterification procedures

Conversion Method	Reference	Maximal conversion	Catalyst % (kg/kg*100)	Temp. (°C)	Time (min.)	Alcohol:oil molar ratio
Transesterification using a solid super base catalyst	[100]	93 %	1.5	70	150	methanol:oil 9:1
In-situ transesterification (skipping the oil extraction step)	[101]	87 %	1.0	60	60	100 ml (m)ethanol for 20 g whole seeds
Transesterification in supercritical alcohols	[102]	95-99 %	0	200-250	40	Supercritical (m)ethanol:oil 50:1
Bio-diesel synthesized enzymatically in presence of supercritical CO ₂	[102]	60-70 %	0	45	480	(m)ethanol:oil 5:1