S. Bouillon · N. Koedam · A. V. Raman · F. Dehairs

Primary producers sustaining macro-invertebrate communities in intertidal mangrove forests

Received: 19 March 2001 / Accepted: 27 August 2001 / Published online: 6 October 2001
© Springer-Verlag 2001

Abstract In contrast to the large number of studies on the trophic significance of mangrove primary production to the aquatic foodweb, there have been few attempts to provide an overview of the relative importance of different primary carbon sources to invertebrates in the intertidal mangrove habitats. We determined carbon and nitrogen stable isotope ratios ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) in sediments, primary producers, and 22 invertebrate species from an intertidal mangrove forest located along the southeast coast of India in order to determine the contribution of mangrove leaf litter and other carbon sources to the invertebrate community. Organic matter in sediments under the mangrove vegetation was characterized by relatively high $\delta^{13}\text{C}$ values and low C:N ratios, indicating that mangrove-derived organic matter was not the principal source and that imported phytodetritus from the mangrove creeks and adjacent bay significantly contributed to the sediment organic matter pool. Invertebrates were found to display a wide range of $\delta^{13}\text{C}$ values, most being 3–11‰ enriched relative to the average mangrove leaf signal. The pulmonate gastropod *Onchidium* sp. showed unusually low $\delta^{15}\text{N}$ values ($-5.6 \pm 0.9\text{‰}$), but further work is needed to adequately explain these data. A compilation of stable isotope data from various sources indicates that significant assimilation of mangrove-derived carbon is only detectable in a limited number of species, and suggests that local and imported algal sources are a major source of carbon for benthic invertebrate commu-

nities in intertidal mangrove forests. These results provide new insights into carbon utilization patterns in vegetated tropical intertidal habitats and show a striking similarity with results from temperate salt marsh ecosystems where local plant production has often been found to contribute little to intertidal foodwebs.

Keywords Carbon · Intertidal foodwebs · Sediment · Trophic relations

Introduction

Tropical mangrove forests can attain high net primary production rates (Clough 1992), and although there remains some uncertainty about the fate of the leaf litter and its role in sustaining adjacent aquatic secondary production, it has become clear that this role has been overestimated in the past (e.g. Lee 1995, 1999; Bouillon et al. 2000, 2001; Dehairs et al. 2000). For the macro-invertebrate fauna inhabiting the intertidal regions, however, most studies assume or conclude that mangroves are the dominant primary producers sustaining these communities (e.g. Camilleri 1992), which are usually dominated – both in terms of numbers and biomass – by brachyuran crabs and gastropods (e.g. Sasekumar 1974; Wells 1984). Few studies have attempted to provide an overall evaluation of different primary carbon sources to the intertidal mangrove invertebrate community, despite the potential importance of these faunal communities in ecosystem carbon dynamics (Robertson et al. 1992) and as food sources for foraging fish during high tide (Sasekumar et al. 1984; Wilson 1989; Sheaves and Molony 2000). Stable isotope analysis can offer valuable insights into the relative importance of different primary producers, but such studies have only rarely been done on mangrove-inhabiting invertebrates (Rodelli et al. 1984; Newell et al. 1995), or have been limited to a specific invertebrate species (Slim et al. 1997; France 1998). The stable isotope approach is based on the assumptions that (1) different primary producers (can)

S. Bouillon (✉) · F. Dehairs
Department of Analytical and Environmental Chemistry,
Mangrove Management Group, Free University of Brussels,
Pleinlaan 2, 1050 Brussels, Belgium
e-mail: sbouillo@vub.ac.be
Tel.: +32-2-6293970, Fax: +32-2-6293274

N. Koedam
Laboratory of General Botany and Nature Management,
Mangrove Management Group, Free University of Brussels,
Pleinlaan 2, 1050 Brussels, Belgium

A. V. Raman
Department of Zoology, Andhra University,
Visakhapatnam, Andhra Pradesh, India

have different $\delta^{13}\text{C}$ values because of different photosynthetic pathways or different inorganic carbon sources, and (2) a consistent degree of fractionation occurs between the isotopic signal of the diet and that of the consumer. For $\delta^{13}\text{C}$, a small or negligible enrichment of on average 0–1‰ has been found to occur (DeNiro and Epstein 1978). For $\delta^{15}\text{N}$, a higher fractionation of on average 2.6‰ (Owens 1987) to 3.4‰ (Minagawa and Wada 1984) is usually assumed, but the actual degree of fractionation may vary considerably, and several processes have been found to result in deviations from this general pattern (e.g. Scrimgeour et al. 1995).

Sesarmid crabs are known to have a high impact on leaf litter dynamics as they can remove a large amount of leaf litter from the sediment surface and carry it into their burrows (e.g. Twilley et al. 1997; Lee 1998). On the other hand, it appears that many sesarmids are more likely to be omnivores than strict herbivores (Dahdouh-Guebas et al. 1999). Ocypodid crabs from mangrove forests such as *Uca* spp. have been considered as either bacteria feeders (e.g. Dye and Lasiak 1986, 1987) or microalgal (including cyanobacteria) feeders (Rodelli et al. 1984; France 1998), and even less is known about the feeding habits of mangrove-dwelling gastropods which are often referred to as 'deposit-feeders' (Plaziat 1984), with little information on their selectivity for mangrove detritus or algal food sources (e.g. Yipp 1980; Rodelli et al. 1984).

Benthic microalgal production in mangrove forests is often low due to light limitation and/or inhibition by soluble tannins (e.g. Alongi 1994), but they have been found to be an important component of foodwebs in other intertidal ecosystems such as salt marshes (e.g. Sullivan and Moncreiff 1990; Currin et al. 1995; Page 1997), and several authors have suggested that their potential role in mangrove ecosystems deserves further

study (e.g. Micheli 1993; Newell et al. 1995). The potential trophic importance of imported organic matter such as phytoplankton from creeks (Bouillon et al., unpublished data) has also not been investigated. In an attempt to evaluate the importance of different primary producers to mangrove-inhabiting fauna, we analysed carbon and nitrogen stable isotope ratios of 23 species of invertebrates from a mangrove forest on the east coast of India, near the mouth of the Gautami Godavari River, along with sediments and primary producers.

Materials and methods

Study area

Samples were collected in the Coringa Wildlife Sanctuary (Fig. 1), which is part of the mangrove-covered area between Kakinada Bay and the Gautami branch of the Godavari (between 82°15' and 82°22' E, 16°43' and 17°00' N). The Godavari is India's second largest river and opens into the Bay of Bengal in the south-eastern state of Andhra Pradesh. The Gautami Godavari also has several branches into Kakinada Bay, the most important of these being Coringa and Gaderu. The sanctuary is dominated by mangrove forests and tidal mudflats, the most abundant mangrove species being *Avicennia marina*, *A. officinalis*, *Excoecaria agallocha*, *Sonneratia apetala*, *Rhizophora mucronata* and *R. apiculata*. Tides are semidiurnal and tidal amplitude in the bay is about 0.5–2 m. Samples were collected at three sites within the sanctuary, located along the Gaderu creek and one of its side creeks (Fig. 1) during a 2-week period in November and December 1999, but some mangrove leaf samples were collected in June 1999 at several other sites in the sanctuary. At all three sites, vegetation is dominated by *A. officinalis* and *E. agallocha*, but site 3 was near to a patch of non-mangrove species, *Suaeda maritima* and *S. monoica*.

Sample collection and preparation

All samples of vegetation, surface sediments and fauna were collected by hand, while benthic microalgae were obtained by gently

Fig. 1 Location of the sampling sites. Darkest areas in panel B indicate the most important mangrove-covered areas

scraping them off the sediment where they formed a conspicuous layer. All faunal samples were kept in a cool box, transported to the field laboratory, washed and dried at 60°C for at least 48 h. For the smaller *Uca* spp. and *Metaplex* spp., the gut and intestinal system were first removed and muscle tissue of the body was used; for larger crab species muscle tissue was taken from the chelae. For the small *Assiminea* sp. four individuals were pooled as one sample. These tissues were ground to a fine powder, and subsamples for $\delta^{13}\text{C}$ and elemental (C:N) analysis were treated with dilute HCl to remove possible carbonates and redried. As this treatment has been reported to affect $\delta^{15}\text{N}$ values (Bunn et al. 1995), subsamples for $\delta^{15}\text{N}$ analysis were not acidified.

Measurement of elemental and stable isotope ratios

Elemental ratios (C:N) of sediment were determined with a Carlo Erba NA-1500 Elemental Analyser. Sediment and biotic samples for stable isotope analysis were similarly combusted, and the resulting gases (CO_2 and N_2) were separated by cryopurification using a manual extraction line. Stable isotope ratios were then measured on a Delta E Finnigan Mat isotope ratio mass spectrometer, and are expressed relative to the conventional standards, i.e. PDB limestone for carbon and atmospheric air for nitrogen, as δ values, defined as:

$$\delta R = \left[\frac{X_{\text{sample}} - X_{\text{standard}}}{X_{\text{standard}}} \right] \times 10^3 [\text{permil}] \quad (1)$$

where $R = ^{13}\text{C}$ or ^{15}N , and $X = ^{13}\text{C}/^{12}\text{C}$ or $^{15}\text{N}/^{14}\text{N}$. Internal reference materials used were ammonium sulphate (IAEA-N1, IAEA-N2) and ammonium nitrate (IAEA-NO-3) for ^{15}N , and sucrose (IAEA-C6) and polyethylene (IAEA-CH-7) for ^{13}C . The standard deviation on 10 aliquots of the same sample was lower than 0.2‰ for both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$.

Results

Primary producers and sediments

Leaves of *Avicennia officinalis* and *Excoecaria agallocha* showed an average $\delta^{13}\text{C}$ of -27.9‰ ($n=5$), close to the overall value for mangrove leaves from this area ($-28.6 \pm 1.4\text{‰}$, $n=17$). Nitrogen stable isotope ratios for these leaves averaged $+4.1\text{‰}$ ($n=5$), close to the average of $+4.6 \pm 1.1\text{‰}$ ($n=12$) for the area (Fig. 2A). *Suaeda* sp. showed similar values of -27.5‰ and $+3.7\text{‰}$ for carbon and nitrogen, respectively. Carbon stable isotope ratios of benthic microalgae scraped off the sediment were much more enriched, averaging $-17.3 \pm 1.7\text{‰}$ ($n=5$), but $\delta^{15}\text{N}$ values were lower than those for mangroves ($\delta^{15}\text{N} = +1.7 \pm 1.7\text{‰}$, $n=5$) (Fig. 2A). Three different mac-

roalgae which were found only in very small quantities at site 1 had the highest $\delta^{15}\text{N}$ values of all primary producers sampled ($+7.5\text{‰}$ for red algae, $+9.1\text{‰}$ for green algae, and $+11.5\text{‰}$ for filamentous algae on *Avicennia* pneumatophores), but had intermediate $\delta^{13}\text{C}$ values of -26.0‰ , -20.0‰ , and -20.9‰ respectively. Sediments under the mangrove vegetation had a low organic carbon content (0.8–1.2%), low C:N ratios (7.0–8.5), and a carbon isotope composition ($\delta^{13}\text{C} = -22.8$ to -20.7‰ , Fig. 2A) enriched by 6–8‰ relative to the dominant vegetation.

Stable isotope composition of invertebrates

Overall, invertebrates exhibited a remarkably wide range of values, between -26.9 and -16.6‰ for $\delta^{13}\text{C}$ and between -6.6 and $+12.3\text{‰}$ for $\delta^{15}\text{N}$ (Fig. 2A, B), and all invertebrates showed average $\delta^{13}\text{C}$ values at least 3‰ enriched relative to the mangrove $\delta^{13}\text{C}$ signal. Most depleted $\delta^{13}\text{C}$ values were found in the gastropods *Melampus fasciatus* ($\delta^{13}\text{C} = -25.5 \pm 0.4$, $n=6$), *Cassidula mustelina* ($\delta^{13}\text{C} = -25.4$ and -25.8 , $n=2$), and *Pythia plicata* ($\delta^{13}\text{C} = -25.2$ and -26.9 , $n=2$) and in the sesarmid *Par-*

Fig. 2 a Plot of $\delta^{15}\text{N}$ versus $\delta^{13}\text{C}$ for different primary producers, sediments, and gastropods collected in the intertidal mangrove forest of the Coringa Wildlife Sanctuary (data available on request). (n) Number of individuals/samples, error bars indicate 1 SD. As *Assiminea* sp. (6), BA brown macroalgae (1), BMA benthic microalgae (5), Cm *Cassidula mustelina* (2), Co *Cerithidea obtusa* (6), FA filamentous algae (1), GA green macroalgae (1), La *Littoraria articulata* (5), Ld *Littoraria delicatula* (2), Lm *Littoraria melanostoma* (6), M mangrove leaves (17), Mf *Melampus fasciatus* (6), Nv *Neritina violacea* (6), On *Onchidium* sp. (7), Pp *Pythia plicata* (2), S sediment, SU *Suaeda* spp. (1), Tt *Telescopium telescopium*. Numbers refer to sampling sites, if not from sites 1 or 2. b Plot of $\delta^{15}\text{N}$ versus $\delta^{13}\text{C}$ for different primary producers, sediments, and crustaceans collected in the intertidal mangrove forest of the Coringa Wildlife Sanctuary (data available on request). (n) Number of individuals/samples, error bars indicate 1 SD. BA Brown macroalgae (1), BMA benthic microalgae (5), Cc *Cardisoma carnifex* (8), Cl *Clibanarius longitarsis* (6), Et *Episesarma tetragonum* (6), Ev *E. versicolor* (5 at site 1, 3 at site 3), FA filamentous algae (1), GA green macroalgae (1), M mangrove leaves (17), Md *Metaplex distinctus* (9), Me *M. elegans* (12), Met *Metopograpsus messor* (1), S sediment, Pa *Parasesarma asperum* (5), Ss *Scylla serrata* (7), Su *Suaeda* spp. (1), Ur *Uca rosea* (3 at each site), Ut *U. triangularis bengali* (7 at site 1, 1 at site 3). Numbers refer to sampling sites, if not from site 1 or 2

asesarma asperum ($\delta^{13}\text{C} = -25.5 \pm 0.6$, $n=5$) (Fig. 2A, B). The omnivorous sesarmids *Episesarma versicolor* and *E. tetragonum* were characterized by more enriched $\delta^{13}\text{C}$ values (-24.2 to -21.9‰ , Fig. 2B), the latter showing highly variable $\delta^{15}\text{N}$ values ($+4.9$ to $+10.1\text{‰}$). Two other large brachyurans, *Cardisoma carnifex* and *Scylla serrata*, had $\delta^{13}\text{C}$ values within the same range, but *C. carnifex* had markedly higher $\delta^{15}\text{N}$ values ($+9.3 \pm 2.2\text{‰}$, $n=8$). $\delta^{13}\text{C}$ values comparable to those of the sediment organic matter were found in the surface grazing gastropods *Telescopium telescopium* ($\delta^{13}\text{C} = -22.0 \pm 1.5$, $n=6$) and *Neritina violacea* ($\delta^{13}\text{C} = -22.6 \pm 0.7$, $n=6$). A large number of invertebrate species had $\delta^{13}\text{C}$ values in between those of the sediment organic matter and benthic microalgae, including the fiddler crabs *Uca rosea* ($-20.1 \pm 0.5\text{‰}$, $n=3$, at site 1, -17.3 ± 1.0 , $n=3$, at site 3) and *U. triangularis* (-20.9 ± 0.8 , $n=7$, at site 1 and -18.9‰ ($n=1$) at site 3), the grapsids *Metaplex distinctus* (-18.7 ± 1.7 , $n=9$) and *M. elegans* (-18.4 ± 1.1 , $n=12$), and the gastropods *Cerithidea obtusa* (-19.3 ± 0.7 , $n=6$) and *Assiminea* sp. (-18.7 ± 0.5 , $n=6$).

Unusually depleted $\delta^{15}\text{N}$ values ($-5.6 \pm 0.9\text{‰}$, $n=7$) were found in the pulmonate gastropod *Onchidium* sp., and three species of the genus *Littoraria* showed relatively low $\delta^{15}\text{N}$ values between -1.7‰ and $+2.6\text{‰}$. No differences were found between the isotopic signatures of specimens collected at site 1 or site 2, but some species which were collected at both site 1 and site 3 (*E. versicolor*, *U. rosea*, and *U. triangularis*) were enriched in ^{13}C at site 3 (Fig. 2B).

Discussion

Primary producers and sediments

The $\delta^{13}\text{C}$ values found for mangrove leaves are typical for terrestrial C_3 -plants and are within the range reported for leaves of various mangrove species by others (e.g. Rao et al. 1994; Newell et al. 1995; Loneragan et al. 1997; Marguillier et al. 1997). The $\delta^{13}\text{C}$ signature of the benthic microalgae ($-17.3 \pm 1.7\text{‰}$) was very different from the mangrove leaves and similar to that reported earlier for benthic algae from mangroves and other intertidal ecosystems (e.g. Newell et al. 1995 and references therein; Dittel et al. 1997; Page 1997; Lee 2000; Wainright et al. 2000). Although a more appropriate technique than the one used has been described to collect these algae (e.g. Couch 1989), this was found to result in insufficient material in our study. However, the close correspondence of our $\delta^{13}\text{C}$ data to values reported in the literature and with the most enriched $\delta^{13}\text{C}$ data for invertebrates allows us to conclude that our results for microalgae are likely to be valid. A large discrepancy was observed between the $\delta^{13}\text{C}$ values of mangrove leaves (average -28.6‰ , Fig. 2A) and the underlying sediment (-22.8‰ at site 1 to -20.7‰ at site 3, Fig. 2A). Such an enrichment in ^{13}C in mangrove sediments has been reported in earlier studies (e.g. Rodelli et al. 1984; Stoner

and Zimmerman 1988; Hemminga et al. 1994; Dittel et al. 1997; Lallier-Verges et al. 1998; Machiwa 1999), although the discrepancy was usually less pronounced than that found in this study. This enrichment, in combination with the low organic carbon content and low C/N ratios found in sediments in this area, indicates substantial inputs of suspended matter from the mangrove creeks and adjacent bay (Bouillon et al., unpublished data). Suspended particulate organic matter in these creeks and in the adjacent bay has been found to have a highly variable $\delta^{13}\text{C}$ (between -19.2 and -30.9‰ in the study area, see Bouillon et al. 2000) and was estimated to contain a large contribution by phytoplankton (Bouillon and Dehairs 2000; Dehairs et al. 2000). There are thus three major types of primary carbon sources available for invertebrates on the sediment surface: mangrove litter, imported plant detritus, and microphytobenthos.

Macro-invertebrates

Only a limited number of species showed evidence for significant assimilation of mangrove-derived carbon. The gastropods *Melampus fasciatus*, *Cassidula mustelina*, and *Pythia plicata* had some of the most depleted $\delta^{13}\text{C}$ values encountered in this study (averages between -26‰ and -25‰ , Fig. 2A), and these data suggest that mangrove-derived carbon and sediment organic matter could contribute in roughly equal proportions to their diet, although the large differences in their $\delta^{15}\text{N}$ signature suggest that their trophic position may be more complex. Sesarmid crabs are often a dominant feature of mangrove invertebrate communities, and although most experimental studies on their feeding habits only include mangrove leaf material (e.g. Steinke et al. 1993; Kwok and Lee 1995), several authors have suggested that they exploit a wider range of food sources under natural conditions (e.g. Micheli 1993; Lee 1997). The three sesarmids sampled in the Coringa area were the most ^{13}C -depleted brachyurans in this study (Fig. 2B). Of these, *Parasesarma asperum* had the lowest $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (Fig. 2A), but the discrepancy with the mangrove $\delta^{13}\text{C}$ signature (about 3.2‰) indicates that other sources also contributed to its diet. Feeding off the sediment surface has been noted frequently in several sesarmid species (e.g. Micheli 1993; M. Skov, personal communication), including *P. asperum* (personal observation), and we hypothesize that the $\delta^{13}\text{C}$ data for this species reflect extensive use of this source as a food source. The wide range of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values found in *Episesarma tetragonum* and *E. versicolor* and the large difference of their $\delta^{13}\text{C}$ values and the mangrove $\delta^{13}\text{C}$ signature (on average 4.6 – 6.8‰ , Fig. 2B) suggest a heterogeneous and mixed diet which included mangrove litter to some extent but not as the major food source. Their higher $\delta^{15}\text{N}$ values compared to *P. asperum* indicate that other invertebrates or carrion contributed to their diet. The low N content of mangrove leaves has led several authors to suggest that these can not be sufficient to meet the ses-

Fig. 3 Relationship between $\delta^{13}\text{C}$ values of sediment organic matter and those of different species of sesarmid crabs ($R^2=0.71$, $P<0.02$). Black symbols This study, grey symbols data from Rodelli et al. (1984). Sediment data for Galle (Sri Lanka) were taken from Bouillon et al. (unpublished data). Error bars indicate 1 SD. Data available on request

armids' N demand (e.g. Micheli 1993; Kwok and Lee 1995; Lee 1997), and our data confirm that other sources can constitute an important contribution to the diet of sesarmids. When combining our data with those from the literature (Rodelli et al. 1984) and some data on *Chiromanthes* sp. from a Sri Lankan mangrove (S. Bouillon, unpublished data), a significant correlation between the $\delta^{13}\text{C}$ of sediment organic matter and of sesarmids (Fig. 3) is apparent, confirming the contribution of sediment organic matter in the diet of some sesarmids. *Cardisoma carnifex* stable isotope ratios (Fig. 2B) suggest an even more omnivorous and heterogeneous diet with only a minor contribution of assimilated mangrove leaf litter, contrary to previous reports of a mainly herbivorous diet (Micheli et al. 1991; Dahdouh-Guebas et al. 1999).

Two surface-grazing gastropods, *Telescopium telescopium* and *Neritina violacea*, had $\delta^{13}\text{C}$ values slightly enriched relative to that of the sediment on which they were found to forage (Fig. 2A) and their $\delta^{15}\text{N}$ signatures ($+6.9\pm1.0\text{‰}$ and $+7.5\pm1.1\text{‰}$, respectively) were higher by about 3.2–3.7‰ relative to this substrate. Our data suggest that these species feed rather indiscriminately on sediment organic matter, which in turn comprised mainly deposited phytoplankton/detritus (Bouillon et al., unpublished data). The pulmonate gastropod *Onchidium* sp. also had $\delta^{13}\text{C}$ values ($-23.1\pm1.1\text{‰}$, $n=7$) close to that of the sediment it was usually found on (-22.8‰), but it exhibited remarkably low $\delta^{15}\text{N}$ values ($-5.6\pm0.9\text{‰}$). The fact that no other invertebrates in this survey displayed such low $\delta^{15}\text{N}$ values renders unlikely the possibility that we have overlooked a very $\delta^{15}\text{N}$ -depleted food source. Such low $\delta^{15}\text{N}$ values have, to our knowledge, so far only been reported for organisms with endosymbiotic chemoautotrophic bacteria from hydrothermal vent systems, 'cold seeps' and reducing sediments (e.g. Fisher 1990; Conway et al. 1994), and have been attributed to either the utilization of NH_4^+ by the endosymbionts, non-limitation of inorganic N allowing maximal fractionation, or nitrogen fixation (see Conway et al. 1994). Clearly, fur-

Fig. 4 Relationship between the carbon stable isotopic composition ($\delta^{13}\text{C}$, expressed in ‰) and carapace width (mm) for *Metaplex distinctus* (open circles, $R^2=0.70$, $P<0.01$) and *M. elegans* (black circles, $R^2=0.66$, $P<0.01$)

ther work is needed to clarify the mechanism leading to the very low $\delta^{15}\text{N}$ values in *Onchidium* sp.

A large number of surface grazers and deposit feeders had $\delta^{13}\text{C}$ signatures intermediate between those of sediment organic carbon and benthic microalgae, reflecting different degrees of selectivity for the latter and indicating little or no assimilation of mangrove carbon. These include the abundant gastropods *Assiminea* sp. and *Cerithidea obtusa*, for which $\delta^{13}\text{C}$ data indicate selective assimilation or ingestion of benthic microalgae. Consistent with this hypothesis, their $\delta^{15}\text{N}$ values ($+4.8\pm0.7\text{‰}$ and $+4.9\pm0.5\text{‰}$, respectively) were higher by about 3‰ than those of benthic microalgae (Fig. 2A), suggesting that bacteria were not important trophic intermediates. The two species of fiddler crabs (*Uca tringularis* and *U. rosea*) and the grapsids *Metaplex elegans* and *M. distinctus* collected in this study had significantly more enriched $\delta^{13}\text{C}$ values than the sesarmids, *S. serrata* or *C. carnifex* (Fig. 2B). Although there continues to be some ambiguity on the importance of different food sources for fiddler crabs, our data confirm results from previous stable isotope studies (e.g. Rodelli et al. 1984; France 1998) that at least some species select for microphytobenthos such as diatoms and cyanobacteria. For both *Metaplex* species, $\delta^{13}\text{C}$ values increased with increasing carapace width, indicating a higher selectivity for benthic microalgae in larger individuals (Fig. 4). Similar to the findings of France (1998), no such ontogenetic shift was found for fiddler crabs, or for any other of the invertebrates sampled. Only one specimen of *Metopograpsus messor* was sampled in this study, and although it is not possible to draw definite conclusions on its diet, it clearly does not rely heavily on mangrove carbon, and its isotopic composition is consistent with extensive feeding on littorinids in this genus (Reid 1986b), although others have found a large contribution of mangrove leaves and macroalgae in the diet of *Metopograpsus* sp. (Dahdouh-Guebas et al. 1999).

The three species of *Littoraria*, typically found on mangrove leaves and stems, exhibited a fairly wide

Fig. 5 Frequency distribution of $\delta^{13}\text{C}$ values of mangrove leaves (white columns, this study and literature data, $n=200$) and $\delta^{13}\text{C}$ values of different invertebrate groups from intertidal mangrove forests. Black bars correspond to data from this study, grey bars refer to published data. A Sesamid crabs, B portunid crabs, C *Cardisoma carnifex*, D *Uca* spp., E *Metaplex* spp., F *Clibanarius* spp., G bivalves, and H gastropods. n Number of data. Data available on request

range of $\delta^{13}\text{C}$ values (-24.7‰ to -20.5‰), but were all characterized by their low $\delta^{15}\text{N}$ values (-1.7‰ to $+2.6\text{‰}$, Fig. 2A). Although it has sometimes been suggested that *Littoraria* spp. feed on the hairs of *Avicennia* leaves, it is generally accepted that they graze the surface layers of trunks and roots where they feed on microepiphytes (Reid 1986a; Blanco and Cantera 1999). Our data support the latter hypothesis, as the $\delta^{13}\text{C}$ values are within the range of those reported for cyanobacterial crusts (Ziegler and Luttge 1998), and the low $\delta^{15}\text{N}$ values of *Littoraria* spp. suggest that the epiphytes are N_2 -fixing, as the process of nitrogen fixation has been reported to result in a fractionation of $0\text{--}4\text{‰}$ (Kohl and Shearer 1980) relative to atmospheric N_2 ($\delta^{15}\text{N}_{\text{air}}=0\text{‰}$, Mariotti 1983). Nitrogen fixation by such cyanobacterial crusts on mangrove stems has also been demonstrated in other studies (Sheridan 1991).

The predatory crab *Scylla serrata*, which feeds almost exclusively on animal matter (e.g. Williams 1978; Hill 1979; Jones 1984) is usually considered to be the top predator of the mangrove benthic community. Although its carbon isotope composition corresponds well with this hypothesis, its $\delta^{15}\text{N}$ values ($+7.7 \pm 1.3\text{‰}$, $n=7$) are lower than might be expected from the $\delta^{15}\text{N}$ signatures of po-

tential prey items (Fig. 2B). A possible explanation could be that *Onchidium* sp., which exhibits very low $\delta^{15}\text{N}$ values around -5.6‰ , contributes to the diet of *S. serrata*. The fact that local fishermen collect *Onchidium* sp. as bait for the capture of *S. serrata* in the mangrove creeks provides circumstantial evidence for this hypothesis.

Finally, the anomuran *Clibanarius longitarsis* had variable but depleted $\delta^{13}\text{C}$ values consistent with its filter-feeding habit (Manjulatha and Babu 1991), as suspended matter in the mangrove creeks during the sampling season is relatively depleted in ^{13}C (Bouillon et al. 2000).

A broader perspective

A striking feature of Fig. 2A, B is the diversity in stable isotope signatures found in co-occurring mangrove invertebrates, indicating a fairly limited overlap in resource utilization. Overall, the data show that although mangrove carbon was assimilated by some invertebrate species (e.g. some gastropods and sesamid crabs), other sources formed a major part of these species' diet. For the majority of the species studied, freshly deposited phyto-detritus, benthic microalgae, and N_2 -fixing microepiphytes were found to be dominant carbon sources. The distribution of $\delta^{13}\text{C}$ data of some important mangrove-inhabiting invertebrate groups from this study and several published studies have been compiled in Fig. 5, whereby data for *Neosarmatium meinerti* and *N. smithi* from Gazi Bay (Kenya) and on *Chiromanthes* sp. and *Terebralia palustris* from Galle (Sri Lanka) have also been included (S. Bouillon, unpublished data). The distribution patterns of

invertebrate $\delta^{13}\text{C}$ values confirm that substantial amounts of mangrove carbon are assimilated only by a limited number of invertebrate groups (such as sesamid crabs, and some of the gastropods), but even in these cases there is only a limited overlap in the $\delta^{13}\text{C}$ distributions. The fairly large discrepancy found between the distribution of mangrove $\delta^{13}\text{C}$ values and other invertebrate groups (*Uca* spp., *Metaplex* spp., bivalves, most of the gastropod species) clearly demonstrates the important role of imported organic matter and microphytobenthos for the macro-invertebrate community in intertidal mangrove habitats. The overall average $\delta^{13}\text{C}$ value of all invertebrates is around -21.9‰ , that is 6.8‰ enriched relative to the average mangrove litter signature. These results show a remarkable similarity with those obtained in temperate salt marsh ecosystems, where a number of recent studies (Sullivan and Moncreiff 1990; Currin et al. 1995; France 1995; Créach et al. 1997; Page 1997; Riera et al. 1999) have demonstrated the trophic importance of imported phytoplankton and local microalgal sources. Although standing stocks of salt marsh vegetation are about an order of a magnitude lower than those of mangroves, their above-ground productivity is comparable (e.g. see Twilley et al. 1992; Middelburg et al. 1997). Mangrove ecosystems and saltmarshes have also been found to show analogous patterns of variability in the sources of organic carbon in surface sediments (Middelburg et al. 1997; Bouillon et al., unpublished data), which suggests that carbon pathways and utilization patterns in these two types of vegetated intertidal ecosystems may be quite similar. However, many aspects of mangrove intertidal foodwebs remain virtually unknown, e.g., there is as yet very little information on the carbon sources for many infaunal organisms, notably meiofauna, or on the role of microheterotrophs as trophic intermediates. Until more studies are undertaken in a variety of mangrove ecosystems, regional differences and the influence of the availability of carbon sources in structuring mangrove foodwebs also remain to be determined.

Acknowledgements Financial support was granted by the EC project 'Assessment of mangrove degradation and resilience in the Indian sub-continent' (contract ERB IC18-CT98-0295). We are much indebted to S. Rajugopal and Krupa Ratnam for their invaluable help during the field collections, and to F. Dahdouh-Guebas and D. Gillikin for providing additional samples from Sri Lanka and Kenya, respectively. We are grateful to G. Poppe who identified many of the gastropods, and to P. Davie for the identification of crabs and some of the *Littoraria* species. We also thank L. De Brabandere, F. Dahdouh-Guebas, D. Gillikin, and two anonymous referees for useful and constructive comments on this manuscript.

References

- Alongi DM (1994) Zonation and seasonality of benthic primary production and community respiration in tropical mangrove forests. *Oecologia* 98:320–327
- Blanco JF, Cantera JR (1999) The vertical distribution of mangrove gastropods and environmental factors relative to tide level at Buenaventura Bay, Pacific coast of Colombia. *Bull Mar Sci* 65:617–630
- Bouillon S, Dehairs F (2000) Estimating spatial and seasonal phytoplankton ^{13}C variations in an estuarine mangrove ecosystem. *Isot Environ Health Stud* 36:273–284
- Bouillon S, Chandra Mohan P, Sreenivas N, Dehairs F (2000) Sources of suspended matter and selective feeding by zooplankton in an estuarine mangrove ecosystem, as traced by stable isotopes. *Mar Ecol Prog Ser* 208:79–92
- Bouillon S, Raman AV, Dauby P, Dehairs F (2001) Carbon and nitrogen stable isotope ratios of subtidal benthic invertebrates in an estuarine mangrove ecosystem (Andhra Pradesh, India). *Estuar Coast Shelf Sci* (in press)
- Bunn SE, Loneragan NR, Kempster MA (1995) Effects of acid washing samples on stable isotope ratios of C and N in penaeid shrimps and seagrass: implications for food web studies using stable isotopes. *Limnol Oceanogr* 40:622–625
- Camilleri JC (1992) Leaf-litter processing by invertebrates in a mangrove forest in Queensland. *Mar Biol* 114:139–145
- Clough BF (1992) Primary productivity and growth of mangrove forests. In: Robertson AI, Alongi DM (eds) *Tropical mangrove ecosystems. Coastal and estuarine studies* 41. American Geophysical Union, Washington, pp 225–249
- Conway NM, Kennicutt II MC, Van Dover CL (1994) Stable isotopes in the study of marine chemosynthetic-based ecosystems. In: Lathja K, Michener RH (eds) *Stable isotopes in ecological and environmental science*. Blackwell, New York, pp 158–186
- Couch CA (1989) Carbon and nitrogen stable isotopes of meio-benthos and their food resources. *Estuar Coast Shelf Sci* 28: 433–441
- Créach V, Schriker MT, Bertru G, Mariotti A (1997) Stable isotopes and gut content analyses to determine feeding relationships in saltmarsh macroconsumers. *Estuar Coast Shelf Sci* 44:599–611
- Currin CA, Newell SY, Paerl HW (1995) The role of standing dead *Spartina alterniflora* and benthic microalgae in salt marsh food webs: considerations based on multiple stable isotope analysis. *Mar Ecol Prog Ser* 121:99–116
- Dahdouh-Guebas F, Giuggioli M, Oluoch A, Vannini M, Cannicci S (1999) Feeding habits of non-ocypodid crabs from two mangrove forests in Kenya. *Bull Mar Sci* 64:291–297
- Dehairs F, Rao RG, Chandra Mohan P, Raman V, Marguillier S, Hellings L (2000) Tracing mangrove carbon in suspended matter and aquatic fauna of the Gautami-Godavari Delta, Bay of Bengal (India). *Hydrobiologia* 431:225–241
- DeNiro MJ, Epstein S (1978) Influence of diet on the distribution of carbon isotopes in animals. *Geochim Cosmochim Acta* 42: 495–506
- Dittel AI, Epifanio CE, Cifuentes LA, Kirchman DL (1997) Carbon and nitrogen sources for shrimp postlarvae fed natural diets from a tropical mangrove system. *Estuar Coast Shelf Sci* 45:629–637
- Dye AH, Lasiak TA (1986) Microbenthos, meiobenthos and fiddler crabs: trophic interactions in a tropical mangrove sediment. *Mar Ecol Prog Ser* 32:259–267
- Dye AH, Lasiak TA (1987) Assimilation efficiencies of fiddler crabs and deposit-feeding gastropods from tropical mangrove sediments. *Comp Biochem Physiol* 87A:341–344
- Fisher CR (1990) Chemoautotrophic and methanotrophic symbioses in marine invertebrates. *Rev Aquat Sci* 2:399–436
- France RL (1995) Stable isotopic survey of the role of macrophytes in the carbon flow of aquatic foodwebs. *Vegetatio* 124:67–72
- France R (1998) Estimation of the assimilation of mangrove detritus by fiddler crabs in Laguna Joyuda, Puerto Rico, using dual stable isotopes. *J Trop Ecol* 14:413–425
- Hemminga MA, Slim FJ, Kazungu J, Ganssen GM, Nieuwenhuize J, Krut NM (1994) Carbon outwelling from a mangrove forest with adjacent seagrass beds and coral reefs (Gazi Bay, Kenya). *Mar Ecol Prog Ser* 106:291–301
- Hill BJ (1979) Aspects of the feeding strategy of the predatory crab *Scylla serrata*. *Mar Biol* 55:209–214
- Jones DA (1984) Crabs of the mangal ecosystem. In: Por FD, Dor I (eds) *Hydrobiology of the mangal*. Junk, The Hague, pp 89–109

- Kohl PL, Shearer G (1980) Isotopic fractionation associated with symbiotic N_2 fixation and uptake of NO_3^- by plants. *Plant Physiol* 66:51–56
- Kwok PW, Lee SY (1995) The growth performance of two mangrove crabs, *Chiromantes bidens* and *Parasesarma plicata* under different leaf litter diets. *Hydrobiologia* 295:141–148
- Lallier-Verges E, Perrussel BP, Disnar J-R, Baltzer F (1998) Relationships between environmental conditions and the diagenetic evolution of organic matter derived from higher plants in a modern mangrove swamp system (Guadeloupe, French West Indies). *Org Geochem* 29:1663–1686
- Lee SY (1995) Mangrove outwelling: a review. *Hydrobiologia* 295:203–212
- Lee SY (1997) Potential trophic importance of the faecal material of the mangrove sesarminid crab *Sesarma messa*. *Mar Ecol Prog Ser* 159:275–284
- Lee SY (1998) Ecological role of grapsid crabs in mangrove ecosystems: a review. *Mar Freshw Res* 49:335–343
- Lee SY (1999) Tropical mangrove ecology: physical and biotic factors influencing ecosystem structure and function. *Aust J Ecol* 24:355–366
- Lee SY (2000) Carbon dynamics of Deep Bay, eastern Pearl River estuary, China. II. Trophic relationship based on carbon and nitrogen stable isotopes. *Mar Ecol Prog Ser* 205:1–10
- Loneragan NR, Bunn SE, Kellaway DM (1997) Are mangroves and seagrasses sources of organic carbon for penaeid prawns in a tropical Australian estuary? A multiple stable-isotope study. *Mar Biol* 130:289–300
- Machiwa J (1999) Fate of organic carbon in a tropical mangrove ecosystem. Doctoral thesis, Stockholm University
- Manjulatha C, Babu DE (1991) Functional organisation of mouth parts, and filter feeding, in *Clibanarius longitarsus* (Crustacea: Anomura). *Mar Biol* 109:121–127
- Marguillier S, van der Velde G, Dehairs F, Hemminga MA, Rajagopal S (1997) Trophic relationships in an interlinked mangrove-seagrass ecosystem as traced by ^{13}C and ^{15}N . *Mar Ecol Prog Ser* 151:115–121
- Mariotti A (1983) Atmospheric nitrogen is a reliable standard for natural ^{15}N abundance measurements. *Nature* 303:685–687
- Micheli F (1993) Feeding ecology of mangrove crabs in North Eastern Australia: mangrove litter consumption by *Sesarma messa* and *Sesarma smithii*. *J Exp Mar Biol Ecol* 171:165–186
- Micheli F, Gherardi F, Vannini M (1991) Feeding and burrowing ecology of two east African mangrove crabs. *Mar Biol* 111:247–254
- Middelburg JJ, Nieuwenhuize J, Lubberts RK, van de Plassche O (1997) Organic carbon isotope systematics of coastal marshes. *Estuar Coast Shelf Sci* 45:681–687
- Minagawa W, Wada E (1984) Stepwise enrichment of ^{15}N along food chains: further evidence and the relation between ^{15}N and animal age. *Geochim Cosmochim Acta* 48:1135–1140
- Newell RIE, Marshall N, Sasekumar A, Chong VC (1995) Relative importance of benthic microalgae, phytoplankton, and mangroves as sources of nutrition for penaeid prawns and other coastal invertebrates from Malaysia. *Mar Biol* 123:595–606
- Owens NJP (1987) Natural variations in ^{15}N in the marine environment. *Adv Mar Biol* 24:389–451
- Page HM (1997) Importance of vascular plant and algal production to macro-invertebrate consumers in a southern California salt marsh. *Estuar Coast Shelf Sci* 45:823–834
- Plaziat JC (1984) Mollusk distribution in the mangal. In: Por FD, Dor I (eds) *Hydrobiology of the mangal*. Junk, The Hague
- Rao RG, Woitichik AF, Goeyens L, van Riet A, Kazungu J, Dehairs F (1994) Carbon, nitrogen contents and stable isotope abundance in mangrove leaves from an east African coastal lagoon (Kenya). *Aquat Bot* 47:175–162
- Reid DG (1986a) The littorinid mollusks of mangrove forests in the Indo-Pacific region. The genus *Littoraria*. British Museum, London
- Reid DG (1986b) Intense predation by crabs on mangrove Littorinids. *Am Malac Bull* 4:112
- Riera P, Stal LJ, Nieuwenhuize J, Richard P, Blanchard G, Gentil F (1999) Determination of food sources for benthic invertebrates in a salt marsh (Aiguillon Bay, France) by carbon and nitrogen stable isotopes: importance of locally produced sources. *Mar Ecol Prog Ser* 187:301–307
- Robertson AI, Alongi DM, Boto KG (1992) Food chains and carbon fluxes. In: Robertson AI, Alongi DM (eds) *Tropical mangrove ecosystems. Coastal and estuarine studies* 41. American Geophysical Union, Washington, pp 293–326
- Rodelli MR, Gearing JN, Gearing PJ, Marshall N, Sasekumar A (1984) Stable isotope ratios as a tracer of mangrove carbon in Malaysian ecosystems. *Oecologia* 61:326–333
- Sasekumar A (1974) Distribution of macrofauna on a Malayan mangrove shore. *J Anim Ecol* 43:51–69
- Sasekumar A, Ong TL, Thong KL (1984) Predation of mangrove fauna by marine fishes. *Proceedings As Symposium on the Mangrove Environment – Reserve Management*. University of Malaya and UNESCO, Kuala Lumpur, pp 378–384
- Scrimgeour SM, Gordon SC, Handley LL, Woodford JAT (1995) Trophic levels and anomalous ^{15}N of insects on raspberry (*Rubus idaeus* L.). *Isot Environ Health Stud* 31:107–115
- Sheaves M, Molony B (2000) Short-circuit in the mangrove food chain. *Mar Ecol Prog Ser* 199:97–109
- Sheridan RP (1991) Epicaulous, nitrogen-fixing microepiphytes in a tropical mangal community, Guadeloupe, French West Indies. *Biotropica* 23:530–541
- Slim FJ, Hemminga MA, Ochieng C, Jannick NT, Cocheret de la Morinière E, van der Velde G (1997) Leaf litter removal by the snail *Terebralia palustris* (Linnaeus) and sesarminid crabs in an East African mangrove forest (Gazi Bay, Kenya). *J Exp Mar Biol Ecol* 215:35–48
- Steinke TD, Rajh A, Holland AJ (1993) The feeding behaviour of the red mangrove crab *Sesarma meinerti* de Man, 1887 (Crustacea: Decapoda: Grapsidae) and its effect on the degradation of mangrove leaf litter. *S Afr J Mar Sci* 13:151–160
- Stoner AW, Zimmerman RJ (1988) Food pathways associated with penaeid shrimps in a mangrove-fringed estuary. *Fish Bull* 86:543–551
- Sullivan MJ, Moncreiff CA (1990) Edaphic algae are an important component of salt marsh food webs: evidence from multiple stable isotope analysis. *Mar Ecol Prog Ser* 62:149–159
- Twilley RR, Chen RH, Hargis T (1992) Carbon sinks in mangrove forests and their implications to the carbon budget of tropical coastal ecosystems. *Water Air Soil Pollut* 64:265–288
- Twilley RR, Poro M, Garcia VH, Rivera-Monroy VH, Zambrano R, Boderio A (1997) Litter dynamics in riverine mangrove forests in the Guayas River Estuary, Ecuador. *Oecologia* 111:109–122
- Wainright SC, Weinstein MP, Able KW, Currin CA (2000) Relative importance of benthic microalgae, phytoplankton and the detritus of smooth cordgrass *Spartina alterniflora* and the common reed *Phragmites australis* to brackish-marsh food webs. *Mar Ecol Prog Ser* 200:77–91
- Wells FE (1984) Comparative distribution of macromolluscs and macrocrustaceans in a North-western Australian mangrove system. *Aust J Mar Freshw Res* 35:591–596
- Williams MJ (1978) Opening of bivalve shells by the mud crab *Scylla serrata* Forskål. *Aust J Mar Freshw Res* 29:699–702
- Wilson KA (1989) Ecology of mangrove crabs: predation, physical factors and refuges. *Bull Mar Sci* 44:263–273
- Yipp MW (1980) The distribution of ground-dwelling gastropods in a small mangrove stand in Hong Kong. In: Morton BS, Tseng CK (eds) *Proceedings of the first international marine biological workshop: the marine flora and fauna of Hong Kong and Southern China*. Hong Kong University Press, Hong Kong, pp 705–721
- Ziegler H, Luttge U (1998) Carbon isotope discrimination in cyanobacteria of rocks of inselbergs and soils of savannas in the neotropics. *Bot Acta* 111:212–215