Conclusions: the results highlight premorbid, aetiological, and phenomenological differences between patients with the onset of a schizophrenia-like illness at the extremes of life, and suggest it is premature to consider the two groups to be merely different manifestations of the same illness.

707

INTENSIVE USE OF COCAINE IN PATIENTS WITH SCHIZOPHRENIA: DEMOGRAPHIC AND OTHER FEATURES

J. Cho, George E. Jaskiw, Milton E. Strauss, Peter Goyer, P. Eric Konicki

Cleveland Department of Veterans Affairs Medical Center, 10000 Brecksville Road, Brecksville, OH 44141

The high incidence of cocaine abuse by patients with schizophrenia is surprising, in light of conventional theories of pathophysiology, and the presumed action of cocaine. Most patients with schizophrenia who use cocaine, do so sporadically, and these are often the subject of investigations. Less is known about patients with schizophrenia who use cocaine intensively and chronically. We identified by chart review, those patients who i) received DSM-III-R diagnoses of schizophrenia as well as cocaine dependence and who ii) either failed at least two courses of treatment for cocaine addiction or who turned down two separate recommendations for such treatment. In this way, 13 high intensity cocaine abusers were identified. They were compared to a sample of 63 patients with schizophrenia, on a number of demographic and other variables. Compared to other schizophrenic patients, the high intensity cocaine abusers were younger, were more likely to be black, never-married, legally competent, had a greater total number of psychiatric admissions and a longer cumulative hospital stay, and were more likely to abuse other substances as well. These high intensity cocaine abusers are being recruited for extensive neuropsychological and other investigations to define correlates.

768

PSYCHIATRIC MORBIDITY, SOCIAL DEVELOPMENT AND COGNITION, AND LATER SCHIZOPHRENIA: PREMORBID ADJUSTMENT OR PRODROME?

A.S. David *, A. Malmberg, G. Lewis, P. Allebeck

Dept of Psychological Medicine, Institute of Psychiatry & King's College Hospital, London SE5 8AZ

We have previously reported on psychosocial and cognitive risk factors for the later development of schizophrenia using data obtained from a Swedish conscript cohort of 50,000 males recruited at age 18, linked to a national case register. The strongest factors identified can be placed into 3 groups: (1) globally reduced IQ; (2) difficulties with personal and sexual relationships including the number and quality of friendships;

(3) non-psychotic psychiatric diagnoses at conscription. By analysing all of these factors using multiple logistic regression we show that they exert independent effects. The question remains as to how we can distinguish 'prodromal' symptoms from true 'premorbid' factors. We have addressed this by examining age of initial diagnosis with the assumption that if putative risk factors are in fact part of a prodrome, they will be more strongly related to early onset. The results show that low IQ and difficulties with relationships are not related to early onset so may be regarded as premorbid factors. In contrast, anxiety and depressive disorders noted at age 18 are related to earlier onset, but diagnoses of personality disorders are not. Understanding the nature of prodromal, premorbid and 'neurodevelopmental' factors in schizophrenia can be improved using such approaches.

709

HIGH PREVALENCE OF HELICOBACTER PYLORI IN SCHIZOPHRENIC PATIENTS

Marc De Hert *, Dimitri De Wilde **, Jim Reemans *, Mark Hautekeete **, Jozef Peuskens *

**UC St. Jozef, Leuvensesteenweg 517, B-3070 Kortenberg, Belgium*

***VU Brussel, Belgium*

Helicobacter Pylori is the most common infectious disease in man. There is considerable controversy concerning the possible modes of transmission. Person-to-person transmission has been documented in children but it is unknown whether it occurs in adults. In order to explore this question we assessed the prevalence of *Helicobacter Pylori* in a cohort of hospitalized schizophrenic patients with a hypothesis of person-to-person transmission correlated to length of stay in hospital.

We adopted a case-control design with matching for age and gender. Controls were recruited from a blood bank. Schizophrenic patients were diagnosed according to DSM-IV-criteria. The cohort consisted of 3 groups, according to length of stay: <2 years; between 2 and 5 years; >5 years. The presence of *Helicobacter Pylori* was assessed by means of an enzyme-linked immunosorbent assay (Malakit, Biolab).

At present we evaluated 89 patients and controls. 21% of the patients had a documented history of gastric pathology. We found significant differences between patients and controls on prevalence of positive serology for *Helicobacter Pylori* (34.8% vs 14.6%, $p < 0.01$) with an Odds Ratio of 3.0 (CI 1.4-7.3). The difference was most pronounced in younger patients. There was a strong correlation with length of stay in hospital, with a progressive increase up to 5 years of hospital stay. After 5 years seropositivity reached a plateau at 38%.

The study confirms the hypothesis for a person-to-person transmission of *Helicobacter Pylori* in adults. The high prevalence of *Helicobacter Pylori* in schizophrenic patients should receive attention by clinicians in the assessment of general physical health due to the relation between *Helicobacter Pylori* and serious gastric disease.