PUBLISHED ARTICLE

Citation*

Aerts R., Maes W., November E., Negussie A., Hermy M., Muys B. 2006. Restoring dry Afromontane forest using bird and nurse plant effects: direct sowing of *Olea europaea* ssp. *cuspidata* seeds. Forest Ecology and Management **230**, 23-31.

DOI: [10.1016/j.foreco.2006.04.001](https://doi.org/10.1016/j.foreco.2006.04.001)

Authors: **Raf Aerts,^{1,2,*} Wouter Maes,¹ Eva November,^{2,3} Aklilu Negussie,² Martin Hermy,¹ and Bart Muys¹**

¹ Division Forest, Nature and Landscape, K.U. Leuven, Belgium

² Land Resources Management and Environmental Protection Department, Mekelle University, Ethiopia

³ Division Soil and Water Management, K.U. Leuven, Belgium

* Corresponding author: Raf Aerts, Celestijnenlaan 200E-2411, BE-3001 Leuven (Belgium).

Tel. +32-16-329721; Fax: +32-16-329760

E-mail: raf.aerts@biw.kuleuven.be

*Elsevier authorized the authors to post this revised personal version of the text of the final journal article on the author's institutional server, incorporating the complete citation and with a link to the Digital Object Identifier (DOI) of the article.

Creative Commons Attribution-Noncommercial-Share Alike 2.0 Belgium License

Permissions beyond the scope of this license may be available at

<http://www.elsevier.com/wps/find/authorsview.authors/copyright#whatrights>

TITLE PAGE

a. TITLE: Restoring dry Afromontane forest using bird and nurse plant effects:
direct sowing of *Olea europaea* ssp. *cuspidata* seeds

b. AUTHORS: Raf Aerts,^{1,2} Wouter Maes,¹ Eva November,^{2,3} Aklilu Negussie,²
Martin Hermy,¹ and Bart Muys¹

c. AFFILIATIONS: ¹Division Forest, Nature and Landscape, Katholieke
Universiteit Leuven, Celestijnenlaan 200E, BE-3001 Leuven, Belgium; ²Land
Resources Management and Environmental Protection Department, Mekelle
University, P.O.Box 231, Mekelle, Ethiopia; ³Division Soil and Water
Management, Katholieke Universiteit Leuven, Celestijnenlaan 200E, BE-3001
Leuven, Belgium.

d. CORRESPONDING AUTHOR: Raf Aerts, Tel. +32 16329721, fax +32
16329760, e-mail raf.aerts@biw.kuleuven.be

e. PRESENT ADDRESSES: same

ABSTRACT

To identify the perspectives of seed sowing for reforestation of degraded dry Afromontane savanna in exclosures in northern Ethiopia, seeds of a fleshy-fruited, secondary climax tree, *Olea europaea* ssp. *cuspidata*, were placed under two pioneer shrub species (*Euclea racemosa* and *Acacia etbaica*) and in open microhabitats. Seed removal and germination rates were examined. The effects of bird ingestion, manual pulp removal and mechanical endocarp treatments on germination rates were also studied.

Pulp removal promoted seed germination, in particular in seeds without endocarp manipulation. Seed removal, presumably by rodents, was concentrated in narrow strips along linear soil conservation structures, and not different between shrubs and open microhabitats. In the absence of water stress, humus types did not have a significant effect on germination rates. Under field conditions however, germination was higher under *Euclea* canopies. The observed facilitative effect of *Euclea* shrubs on germination may be attributed to moisture retention effects, related to the deep humus layer and canopy architecture, preventing desiccation and promoting imbibition of seeds. When readily available *Olea* seeds are not used for seedling production, direct sowing of seeds in the humus under dense *Euclea* shrubs could offer a cheap complement to planting of nursery-raised seedlings for forest restoration projects in recovering Afromontane savanna.

Key words: direct seeding, direct sowing, facilitation, nurse plant, recruitment, restoration

1. Introduction

Changing land use and land cover, population growth, unsustainable use of resources and climate change are some of the most widely recognized proximate causes and driving forces of land degradation and desertification in many parts of the world (Geist and Lambin, 2004). This particularly holds for the arid and semiarid regions of Sub-Saharan Africa (Darkoh, 1998; Zeleke and Hurni, 2001). The consequences of desertification, especially in the least-developed countries, are large-scale loss of biodiversity but also poverty and food insecurity. Biological conservation and restoration policies therefore need to take account of the diverse relationships between conservation needs and the demands of poverty reduction (Adams *et al.*, 2004).

Despite their low productivity, dry tropical forests are often of vital importance to rural communities in Sub-Saharan Africa, in particular in the Horn and eastern Africa (Lamprecht, 1989). Conversion to agriculture (e.g. Lemenih *et al.*, 2005), accidental and intentional fires, wood and gum collection, grazing by livestock and selective logging of valuable species continue to transform dry tropical forests to such significant extent that they are considered by some to be the most endangered ecosystem in the tropics (Gillespie, 1999; Kalacska *et al.*, 2004).

In northern Ethiopia, the regional government and local NGOs have implemented a forest rehabilitation strategy built on the needs of local communities by formally declaring community-managed protected areas where removal of remnant vegetation and free grazing are currently suspended (Tekle, 2001; Asefa *et al.*, 2003; Mengistu *et al.*, 2005). Land rehabilitation efforts in these set-aside areas or exclosures aim at restoring the natural Afromontane

1 forest vegetation (Aerts *et al.*, 2004; Nyssen *et al.*, 2004). Natural regeneration
2 of tree species in exclosures depends on seed dispersal from nearby forest
3 patches (Turner and Corlett, 1996; Duncan and Duncan, 2000) or vegetative
4 resprouting of pioneers because most Afromontane forest trees lack persistent
5 soil seed reserves (Teketay and Granström, 1997; Tekle and Bekele, 2000;
6 Wassie and Teketay, 2005). But as most of the Afromontane tree species
7 exhibit limited long-distance seed dispersal as well (Teketay and Granström,
8 1995), exclosures face serious limits to natural forest regeneration, especially in
9 a landscape where forest relics are small, fragmented and highly isolated (Aerts
10 *et al.*, 2006c). In such cases, planting of nursery-raised seedlings may
11 accelerate regeneration (e.g. Aide *et al.*, 2000; Martínez-Garza and Howe,
12 2003; Yirdaw and Luukkanen, 2003). An alternative, easier and cheaper
13 technique which is slowly being adopted in restoration ecology of degraded
14 tropical lands, is direct sowing of seeds (see e.g. Sun *et al.*, 1995; Hardwick *et*
15 *al.*, 1997; Guariguata and Pinard, 1998; Chapman and Chapman, 1999; Engel
16 and Parrotta, 2001; Camargo *et al.*, 2002). Particularly when combined with
17 facilitative effects of pioneer shrubs, which may help overcome seed predation
18 and desiccation (Woods and Elliot, 2004), the use of direct sowing may be an
19 efficient and easy to implement forest restoration technique. Assisting
20 regeneration may expedite the productivity of the exclosures, hereby serving
21 both conservation and poverty-alleviation.

22 In our study area, naturally established seedlings of *Olea europaea* ssp.
23 *cuspidata* (Wall. ex G. Don) Cif. (Oleaceae, African wild olive), an important
24 shade-tolerant, fleshy-fruited component of dry Afromontane forest (Friis, 1992),
25 are restricted to microhabitats under shrubs, primarily under the fruit-bearing

pioneer *Euclea racemosa* ssp. *schimperi* (A. DC.) White (Ebenaceae), and to a lesser extent under the dominant thorn shrub *Acacia etbaica* Schweinf. (Fabaceae) (Aerts *et al.*, 2006b). The seeds are dispersed in the summer rainy season (Teketay and Granström, 1997), primarily by frugivorous birds (see Alcántara *et al.* (2000b) for *O. europaea* ssp. *sylvestris*). Seeds may be dropped during flight and may thus land anywhere, including on bare soil. Nevertheless it is more likely that seeds are dropped when the bird is perched (McClanahan and Wolfe, 1993; Stiles, 2000; Schupp *et al.*, 2002), resulting in a primary seed rain concentrated under shrubs and trees (e.g. Alcántara *et al.*, 2000a, 2000b; Stiles, 2000; Holl, 2002). As secondary dispersal by run-off does not alter this pattern (Aerts *et al.*, 2006a), the absence of seedlings in the patches between shrubs may be explained by lower recruitment, i.e. higher seed removal or predation, less seed germination and seedling emergence, higher seedling mortality, or a combination of these factors (see e.g. Herrera *et al.*, 1994; Rey and Alcántara, 2000; Holl, 2002).

In this study, we analyzed the post-dispersal fate of *O. europaea* seeds to evaluate the possibilities and constraints of direct sowing for reforestation of dry Afromontane savanna. We test whether different microhabitats in exclosures have an effect on seed removal patterns and germination rates. We also test the effects of bird ingestion, pulp removal and mechanical endocarp treatment on seed germination. We hypothesize that avian gut treatment will provide highest germination and that germination in microhabitats under pioneer shrubs will be higher compared to bare soil patches due to humus and moisture effects, despite the expected higher seed predation by rodents under shrub cover (Alcántara *et al.*, 2000a; Mohr *et al.* 2003).

2. Materials and methods

2.1. Study species

Olea europaea ssp. *cuspidata* (African wild olive) is a sclerophyllous evergreen tree with a heavy branched and rounded crown. It grows to 5–15 (max. 25) m tall and is widely distributed in Afromontane forests, particularly in drier highland forests in association with *Juniperus procera* Hochst. ex Endl. (Cupressaceae, East African pencil cedar), forest margins, riverine forests and evergreen montane scrub from 1,250 to 3,000 m above sea level in Sudan, Somalia and southward through East tropical Africa into South Africa (Green, 2003). The ovoid-ellipsoid drupes have a fleshy mesocarp (pulp) and are green and dark purple-black when ripe (Green 2003). The seeds are dispersed in the main rainy season (Teketay and Granström, 1997), primarily by frugivorous birds. The seeds have a width $W_s = 6$ mm, height $H_s = 6$ mm and length $L_s = 8$ mm with a volume $V_s = 151 \text{ mm}^3$ and mass $M_s = 225$ mg (Aerts *et al.*, 2006a). In undisturbed forest, germination results in populations of shade-tolerant and slowly growing seedlings (Teketay and Granström, 1997), and both in set-aside areas and grazing land, natural regeneration has been documented under early-successional shrubs (Aerts *et al.*, 2006b). Once established, the tree is drought resistant, but because of its multiple uses (e.g. durable timber, traditional ox-ploughs, firewood and charcoal) both young and mature trees have been over-harvested dramatically in Ethiopia. As a result this valuable tree is now under threat of local extinction (Negash, 2003).

2.2. Site description

The experiments were set-up during the 2002 and 2003 summer rainy seasons in the Enderta and Doga Tembien districts of Tigray, northern Ethiopia (Table 1). Two study sites, Sesemat and Mheni, were located 1,800–1,900 m above sea level in the Geba river catchment of central Tigray, northern Ethiopia, on slopes over calcareous rocks and lacustrine silified Mesozoic Antalo limestone layers 20 km NW of the regional capital Mekelle, and are set-aside areas since 1996. A third site, Bubu Hills, was located at the foot of a dolerite cliff at the edge of Mekelle town next to a peri-urban eucalypt plantation. The vegetation in the three field sites was characterized by a discontinuous cover of shrubs and small regenerating trees in a matrix of herbs and bare soil (see Aerts *et al.* (2006b) for details). It can thus be defined as semiarid degraded savanna (Vetaas, 1992; De Villiers *et al.*, 2001). In the three sites, soil erosion rates are partially controlled by stone bunds. These are stone walls back-filled with soil along the slope contours and are widely used in the Ethiopian highlands as a soil and water conservation measure. Sediment accumulates behind these structures, which results in the development of progressive terraces (Herweg and Ludi, 1999; Hengsdijk *et al.*, 2005; Vancampenhout *et al.*, 2005).

The climate is related to the mountainous facies of the Sudanese zone with hot and dry winters (October–January) and two rainy seasons: a short and moderate, highly variable spring rain (*belg*, coefficient of variation 31–55%) in (February)–March–April–(May–June) and a long, heavy and more reliable summer rain (*kremt*, coefficient of variation 19–31%) in (June)–July–August–September (Meze-Hausken, 2004). Mean annual precipitation is 625 ± 155 mm

(spring rain 125 ± 11 mm; summer rain 466 ± 22 mm; means $\pm$ SE for the years 1960–2003; Meze-Hausken, 2004) and mean annual temperature 18°C . Total spring and summer rains for 2003 were 149.3 and 357.3 mm (Mekelle Quiha station, National Meteorological Services Agency). In our study area, dry Afromontane forest relics and Afromontane savanna woodland are communities dominated by *Olea europaea*, *Acacia etbaica* and *Combretum collinum* Fresen (Combretaceae) (Aerts *et al.*, 2006c).

[Insert Table 1]

2.3. Post-dispersal seed removal experiments

To measure seed removal, seeds were placed in $8 \times 8 \text{ cm}^2$ mesh wire seed containers (depots), with 10 seeds and 3 stones as dummies per depot. The depots had a 2-cm high side that prevented seeds from washing away (abiotic secondary seed dispersal, see Aerts *et al.*, 2006a), but allowed for entry of seed predators (see Holl (2002) for a similar protocol). All seeds were collected under mature mother trees in a forest relic and were similar to naturally dispersed seeds (no fleshy mesocarp). Only viable seeds were used, determined by a floating test (Schatral and Fox, 1994), since rodents have the ability to detect and avoid sterile seeds (Kollman and Pirl, 1995). Depots were placed in three microhabitats: (1) Open, in areas of bare soil between shrubs; (2) *Euclea*, under the canopy of individuals of *E. racemosa*; and (3) *Acacia*, under the canopy of individuals of *A. etbaica*. On 12 August 2003, 60 depots were placed in the Mheni exclosure in two areas of approximately $100 \times 100 \text{ m}^2$ (twenty per treatment, one shrub or bare patch per depot). Remaining, intact

seeds were counted after 1, 3, 6, 9 and 12 days. Preliminary analyses showed that seed removal only occurred in depots near stone bunds. A second experiment was therefore carried out in the exclosure of Mheni to test the effect of these stone bunds on seed removal. On 2 September 2003, 52 depots were placed under seven treatments. Twenty-four depots were placed under *Acacia* and *Euclea* shrubs (2 × 8 depots) and in open areas (8 depots). The remaining depots were placed directly next to the base and on top of stone bunds (2 × 14 depots). Remaining, intact seeds were counted after 1, 3, 6, 9, 13, 15, 18, 21, 24 and 30 days. On 24 September 2003, a duplicate of this experiment was initiated in the Sesemat exclosure.

2.4. Direct sowing experiment

To measure *in-situ* germination, *Olea* seeds were sown in open areas, under *Euclea* and under *Acacia* shrubs in the exclosures of Mheni and Bubu Hills on 23–24 July 2003. In both sites, ten seeds were sown 1–2 cm deep in each of 30 randomly assigned patches (ten per treatment). Seed positions were marked with wooden tooth-picks to facilitate follow-up. Germination was monitored once a week for 14 weeks (well beyond the end of the rainy season, at 11 September 2003), and regularly followed up during the next spring and summer rains. Viable seeds without fleshy mesocarp, collected under mature trees in a nearby forest relic, were used.

2.5. Controlled germination experiments

The ectorganic layer under *Euclea* shrubs is deep with a very low bulk density and is generally characterized by a thick litter layer of slowly-

decomposing leathery leaves (oligomull). The ectorganic horizon under *Acacia* shrubs is thin, with few remains of leaves and a litter layer of small twig fragments and thorns (eumull) (K. Descheemaeker, unpub. data). There is little or no humus in open patches, only a cover of loose stones with very little organic objects. To measure the effects of these two humus types or the absence of humus on germination, independently from water stress, humus from under *Euclea* and *Acacia* shrubs (10-15 cm deep) and stony surface covers of open areas (5 cm deep) were collected from the enclosure of Mheni and carefully reconstructed in 15 1×0.5 m² plots in seedling beds of the Endayesus tree nursery (see Franco-Pizaña *et al.* (1996) for a similar protocol). On 24 July 2003, 20 *Olea* seeds were sown 1-2 cm deep in each treatment replication. Germination was monitored for 30 weeks. Viable seeds without fleshy mesocarp collected under mature trees in a nearby forest relic were used, and were soaked in water for 24 hours prior to sowing to promote imbibition and germination.

The effect of avian pulp removal (for instance, through gut passage) was tested using a second controlled germination experiment. The factors ‘pulp removal’ (removal of the fleshy mesocarp by birds, manual pulp removal and no pulp removal; see Samuels and Levey, 2005) and ‘mechanical treatment’ (scarification of the endocarp, splitting of the endocarp and no mechanical treatment) were combined to seven treatments (omitting the meaningless combinations no pulp removal × mechanical endocarp treatments). In each of 21 1×0.5 m² plots in seed beds of the Endayesus tree nursery (three per treatment), 50 seeds were sown 1–2 cm deep on 11 August 2002. Seeds of the ‘pulp removal by birds’ treatment were viable, regurgitated or defecated seeds.

Seeds for the manual pulp removal and control treatments were obtained from fresh, ripe drupes collected in a nearby forest relic. All treatments (including control) were soaked in water for 24 hours prior to sowing.

In both controlled germination experiments, germinants were recorded and then removed from the seed beds to avoid double counts. Grass mats 1 m above the beds provided shade; beds were watered twice daily.

2.6. *Data analysis*

For the first seed removal experiment, non-parametrical Kruskal-Wallis one-way ANOVA (KW) was used to compare seed removal in the three microhabitats. In the second seed removal experiment, seed survival was analyzed using a time-to-event approach, which measures the time to an event for each case (depot) (Altman and Bland, 1998). In this study the critical event is seed removal. Because seed removal is inherently linked to escape from removal or predation, this data is conventionally called survival data. Since rodents and other seed consumers often hoard food from a dense concentration of resources and frequently revisit such resources (Vander Wall, 1990), data of seed removal in a single depot on subsequent days does not reflect independent events. A depot was therefore treated as completely removed as soon as at least one seed had disappeared. The characteristic feature of survival data is that at the end of the follow up period, the critical event will probably not have occurred for all depots. For these depots, the survival time is said to be censored (Altman and Bland, 1998). Because of the censoring, we used Kaplan-Meier survival analysis, a model for censored survival data based on estimating conditional probabilities for each time interval in which an event

occurs (Bland and Altman, 1998). Equality of survival distributions for the seven different microhabitats was tested using the Breslow statistic (BS). After reclassification of the data in depots near stone bunds (<1 m) and those further away (>1 m), a second survival analysis was used to test for differences in seed removal between these two distance classes.

The first germination experiment (humus type) was analyzed using Kruskal-Wallis one-way ANOVA. The second germination experiment (pulp removal and mechanical treatment) was analyzed using two-way ANOVA on untransformed proportion data (Levene's test for homoscedacity indicated homogeneity of variance).

In the controlled germination experiment to test the effect of pulp removal by birds, one plot of the treatment pulp removal by birds × endocarp scarification was disturbed by nocturnal animals and omitted from analysis. The seed removal experiment in Sesemat was seriously disturbed between days four and six: 25 depots were destroyed. Analyses were performed using SPSS 11.0 and 12.0 for Windows (SPSS Inc., Chicago, IL).

3. Results

3.1. Seed removal

In the first experiment designed to capture seed removal differences between the three major microhabitats, seed removal was limited and highly localized in one site. Only 39 seeds were removed from the depots (6.5%): 19 (3.2%) from three open depots and 20 (3.3%) from two depots under *Acacia*. There was no significant effect of microhabitat on the number of remaining

seeds (KW $\chi^2 = 3.06$, $P = 0.216$), but interestingly, the five depots where seed removal occurred, were all positioned in the immediate vicinity of stone bunds.

For the second experiment, with depots explicitly placed near stone bunds, survival analysis showed that microhabitat had a significant effect on seed survival time (BS = 15.6, $P = 0.004$), but between treatments, only contrasts with depots near stone bunds were significant. Seed removal in depots near stone bunds (<1m) was significantly higher than in depots placed further away (>1m) (BS = 22.1, $P < 0.001$; Fig. 1). At the end of the experiment, 52±7% of all seeds deposited near stone bunds were removed, while this was only 3±2% for those deposited further away (Fig. 1). No seed removal was detected in Sesemat, possibly due to disturbance, but more probably because of factors related to movement of predators (see discussion).

[Insert Fig. 1]

3.2. Germination after direct sowing

In-situ germination was limited and site-dependent. In the Mheni enclosure, none of the seeds germinated. In the Bubu Hills enclosure, germination was significantly higher under *Euclea* cover (20±2%) compared to *Acacia* cover and open microhabitats (11±1 and 10±0%) (KW $\chi^2_{\text{week 8}} = 15.54$, $P < 0.001$; Fig. 2). Eight weeks after sowing, the summer rainy season stopped and no further germination occurred. No further germination could be detected during the following spring and summer rains.

[Insert Fig. 2; Insert Fig. 3]

3.3. *Controlled germination*

Humus type did not have a significant effect on controlled germination (KW $\chi^2_{\text{week 30}} = 0.005$, $P = 0.997$; Fig. 3). After 30 weeks, $34 \pm 4\%$ of all seeds pooled over treatments had germinated.

Germination in the pulp removal and endocarp treatment experiment was characterized by two distinct phases, in which both the pulp removal and the endocarp treatments had significant effects without interaction (Table 2, Fig. 4). Endocarp splitting resulted in rapid and uniform germination until the seventh week, while germination in the other treatments continued slowly and gradually. After 23 weeks, $32 \pm 4\%$ of the seeds with split endocarps had germinated compared to 14 ± 4 and $13 \pm 4\%$ for the scarified seeds and the controls (Table 2). At this stage, pulp removal was a crucial factor for germination since only marginal germination was observed in the overall control. In week 24, which coincided with the onset of the spring rains (27 january 2003), a second wave of germination occurred in the endocarp control treatments (Fig. 4). After 46 weeks, the highest germination was recorded in seeds with intact endocarp: first of all by those with pulp removed by birds ($63 \pm 10\%$), followed by those with manually removed pulp ($55 \pm 3\%$) and then the overall control ($37 \pm 7\%$). Endocarp splitting, combined with pulp removal, facilitated germination immediately after sowing, but ultimately, germination was more successful in seeds with intact endocarp.

[Insert Fig. 4; Insert Table 2]

4. Discussion

We examined removal and germination rates of seeds of an animal-dispersed secondary climax tree placed in three different microhabitats. We also studied the effects of bird ingestion, pulp removal by hand and mechanical endocarp treatments on germination rates. Our analysis confirms that full or partial passage through the digestive tract of frugivorous birds influences germination of *O. europaea* seeds (Fig. 4). In general, ingested seeds germinated in greater numbers and took less time to germinate than uningested seeds (c.f. Traveset and Verdú, 2002). Additional germination trials using seeds of three other fleshy-fruited Afromontane tree species, *Cordia africana* Lam. (Boraginaceae), *Juniperus procera* and *Afrocarpus falcatus* (Thunb.) C. N. Page (Podocarpaceae), also showed improved germination in seeds with naturally removed pulp, especially in those seeds not subjected to supplementary endocarp treatments (A. Negussie and R. Aerts, unpubl. data). This observation is in line with evidence provided by Murray *et al.* (1994) showing that germination success decreases as gut retention time, and thus mechanical and chemical endocarp scarification, increases (Stiles, 2000). In any case, removal of the lipid-rich pulp, which repels water and thus prevents imbibition of the embryo, is necessary for germination (Verdú and García-Fayos, 1998). Effective pulp removal may be achieved by birds or by hand (Wenny, 2000), but also by natural decay in the soil, all yielding different germination rates, but similar final germination proportions (Fig. 4). Many tropical species whose seeds disperse during the main rainy season, exhibit a strong innate dormancy, lasting 3–6 months, regardless of the watering regime

1 during this period (Garwood, 1989). But the germination in two distinct phases
2 (Fig. 4) suggests that seed passage through the gut of birds breaks this
3 physiological dormancy in a fraction of the seeds. Seeds germinating
4 immediately after dispersal have a lower predation and removal probability than
5 seeds germinating after the winter drought; but seedlings, on the other hand,
6 have a higher survival probability when emerging after the winter drought, since
7 only those seedlings can take advantage of a full rainy season before being
8 exposed to subsequent drought stress (see also Kitajima and Fenner, 2000).
9 The dual germination observed in *O. europaea* may thus be seen as a
10 mechanism to enhance offspring survival, by providing both dormant seeds and
11 overwintering seedlings.

12 *Olea* seeds may exceed the maximum size that can be removed and
13 predated by ants and other litter-dwelling invertebrates, but are in contrast very
14 attractive to vertebrate seed predators, such as rodents (Rey *et al.*, 2002;
15 Fornara and Dalling, 2005). Although we did not find direct evidence for
16 vertebrate seed predation, and cannot conclude that all seed removal resulted
17 in seed predation since secondary dispersal via scatter-hoarding is possible
18 (Vander Wall *et al.*, 2005), trails and droppings found along stone bunds
19 indicated considerable rodent activity, probably by species from the genera
20 *Mastomys* and *Arvicanthis*, the major rodent pests in the highlands of Ethiopia
21 (Bekele and Leirs, 1997; Gebreselassie *et al.*, 2004). The presence of refuges
22 for seed predators may explain high seed predation rates (Verdú and García-
23 Fayos, 1996). In our study area, nesting and burrowing activities of rodents are
24 often concentrated in stone bunds (R. Aerts, pers. obs.), hereby increasing the
25 probability of foraging animals to encounter seeds in the vicinity of these

1 structures (Fig. 1). But further away from stone bunds, post-dispersal seed
2 removal was fairly low and not higher under shrubs compared to open patches.
3 These results support the theory that predator population densities and within-
4 site differences can strongly influence patterns of seed removal and predation in
5 space and time (Holl and Lulow, 1997; Alcántara *et al.*, 2000a; Garcia, 2001;
6 Holl, 2002; Rey *et al.*, 2002; Jones *et al.*, 2003). Cessation of seed removal in
7 the exclosures approximately mid-September (17 September), the absence of
8 removal in the Sesemat site, and the overall low seed removal rates may be
9 related to the preference of rodents for ripened crops in fields near the
10 exclosures, as revealed by stomach content analysis of trapped animals in an
11 irrigation field near our study area (Gebresilassie *et al.*, 2004). Thus, in our
12 study and at least one other (Alcántara *et al.*, 2000a), the seed dispersal pattern
13 of *Olea europaea* is not reshaped by biotic post-dispersal seed removal,
14 because the initial seed rain is not altered significantly.

15 In Mheni seeds failed to germinate, probably due to the very poor
16 summer rains in that area during the study; in semiarid areas seeds may need
17 long and abundant rain events to germinate (Verdú and García-Fayos, 1998).
18 In Bubu Hills, which generally receives more rainfall (Table 1), seeds placed
19 under shrubs and in bare soil initially showed germination rates similar to those
20 in controlled germination experiments (Fig. 2), but germination stopped after the
21 summer rains and did not resume during the spring rains. In the absence of
22 water stress, microhabitat humus types did not have a significant effect on
23 germination rates (Fig. 3). Under field conditions, however, higher germination
24 rates were observed under *Euclea* canopies (Fig. 2). Since *Olea* removal rates
25 were not significantly different between *Acacia*, *Euclea* and bare soil treatments,

the observed facilitative effect of *Euclea* shrubs on germination may probably be attributed to moisture retention effects. This may be related to the humus structure, but also to canopy architecture. The sclerophyllous-evergreen *Euclea* shrubs provide more shade than the leguminous-deciduous *Acacia* shrubs (Aerts *et al.*, 2006b), reducing solar radiation, soil temperature and soil evaporation and thus preventing desiccation of seeds more effectively.

4. Conclusions

Overall, our results agree with previous studies that have shown that seed pre-treatment shortens seed dormancy and thus decreases the time available for seed removal or predation to occur. Duncan and Chapman (2002), commenting on previous seed sowing studies in degraded sites, found limited or no success in most cases. Nevertheless, the low germination and seedling survival typically found in direct sowing techniques (Engel and Parrotta, 2001) may be overcome by sowing more seeds, selecting larger, more vigorous seeds (Kidson and Westoby, 2000; Moles and Westoby, 2004; Rey *et al.*, 2004) or imbibing seeds prior to sowing (Camargo *et al.*, 2002). Seeds with naturally removed pulp, thus suitable for sowing, can usually be found in large quantities under fruiting *Olea* trees in forest relics, and under solitary trees in the vicinity of these relics (R. Aerts, pers. obs.). When these seeds are not used for seedling production (which is often the case), direct sowing of *Olea* seeds pre-treated by birds in the humus under dense *Euclea* shrubs could offer an effective, cheap complement to planting of nursery-raised seedlings for forest restoration projects in recovering Afromontane savanna. Sowing seeds 1–2 cm deep is preferable to broadcasting, since it may help to protect the seeds from removal

1 and predation and to prevent seed desiccation (Woods and Elliot, 2004). In
2 ecological terms, this technique would increase the seed density under safe
3 sites in degraded areas using seeds that would otherwise have been lost due to
4 dispersal limitation.

6 **Acknowledgements**

7 This research was funded by a Ph.D. grant of the Flemish Interuniversity
8 Council (VLIR) and the VLIR-OI project 'Forest Rehabilitation through Natural
9 Regeneration in Tigray, northern Ethiopia' of the Katholieke Universiteit Leuven
10 (Belgium) and Mekelle University (Ethiopia). We thank Yemane Belay and his
11 staff of the Endayesus tree nursery at Mekelle University for taking daily care of
12 the controlled germination experiments. Constructive comments by two
13 anonymous referees on an earlier draft of this paper are gratefully
14 acknowledged.

16 **References**

17 Adams, W.M., Aveling, R., Brockington, D., Dickson, B., Elliott, J., Hutton, J.,
18 Roe, D., Vira, B., Wolmer, W., 2004. Biodiversity conservation and the
19 eradication of poverty. *Science* 306, 1146-1149.

20 Aerts, R., Maes, W., November, E., Behailu, M., Poesen, J., Deckers, J.,
21 Hermy, M., Muys, B., 2006a. Surface runoff and seed trapping efficiency of
22 shrubs in a regenerating semiarid woodland in northern Ethiopia. *Catena* 65,
23 61-70.

24 Aerts, R., November, E., Van der Borgh, I., Behailu, M., Hermy, M., Muys, B.,

- 1 2006b. Effects of pioneer shrubs on the recruitment of the fleshy-fruited tree
- 2 *Olea europaea* ssp. *cuspidata* in Afromontane savanna. Applied Vegetation
- 3 Science 9. In press.
- 4 Aerts, R., Van Overtveld, K., Haile, M., Hermy, M., Deckers, J., Muys, B.,
- 5 2006c. Species composition and diversity of small Afromontane forest
- 6 fragments in northern Ethiopia. Plant Ecology. In press (doi:10.1007/s11258-
- 7 006-9137-0).
- 8 Aerts, R., Wagendorp, T., November, E., Behailu, M., Deckers, J., Muys, B.,
- 9 2004. Ecosystem thermal buffer capacity as an indicator of the restoration
- 10 status of protected areas in the northern Ethiopian highlands. Restoration
- 11 Ecology 12, 586-596.
- 12 Aide, T.M., Zimmerman, J.K., Pascarella, J.B., Rivera, L., Marciano-Vega, H.,
- 13 2000. Forest regeneration in a chronosequence of tropical abandoned
- 14 pastures: implications for restoration ecology. Restoration Ecology 8, 328-338.
- 15 Alcántara, J.M., Rey, P.J., Sánchez-Lafuente, A.M., Valera, F., 2000a. Early
- 16 effects of rodent post-dispersal seed predation on the outcome of the plant-
- 17 seed disperser interaction. Oikos 88, 362-370.
- 18 Alcántara, J.M., Rey, P.J., Valera, F., Sánchez-Lafuente, A.M., 2000b. Factors
- 19 shaping the seedfall pattern of a bird-dispersed plant. Ecology 81, 1937-1950.
- 20 Altman, D.G., Bland, J.M., 1998. Statistics notes - Time to event (survival) data.
- 21 British Medical Journal 317, 468-469.
- 22 Asefa, D.T., Oba, G., Weladji, R.B., Colman, J.E., 2003. An assessment of

- 1 restoration of biodiversity in degraded high mountain grazing lands in Northern
2 Ethiopia. *Land Degradation & Development* 14, 25-38.
- 3 Bekele, A., Leirs, H., 1997. Population ecology of rodents of maize fields and
4 grassland in central Ethiopia. *Belgian Journal of Zoology* 127, 39-48.
- 5 Bland, J.M., Altman, D.G., 1998. Statistics notes - Survival probabilities (the
6 Kaplan-Meier method). *British Medical Journal* 317, 1572.
- 7 Camargo, J.L.C., Ferraz, I.D.K., Imakawa, A.M., 2002. Rehabilitation of
8 degraded areas of Central Amazonia using direct sowing of forest tree seeds.
9 *Restoration Ecology* 10, 636-644.
- 10 Chapman, C.A., Chapman, L.J., 1999. Forest restoration in abandoned
11 agricultural land: a case study from East Africa. *Conservation Biology* 13, 1301-
12 1311.
- 13 Darkoh, M.B.K., 1998. The nature, causes and consequences of desertification
14 in the drylands of Africa. *Land Degradation & Development* 9, 1-20.
- 15 De Villiers, A.J., Van Rooyen, M.W., Theron, G.K., 2001. The role of facilitation
16 in seedling recruitment and survival patterns, in the Strandveld Succulent
17 Karoo, South Africa. *Journal of Arid Environments* 49, 809-821.
- 18 Duncan, R.S., Chapman, C.A., 2002. Limitations of animal seed dispersal for
19 enhancing forest succession on degraded lands. In: Levey, D.J., Silva, W.R.,
20 Galetti, M. (Eds.), *Seed dispersal and frugivory: ecology, evolution and*
21 *conservation*. CAB International, Oxon, UK, pp. 437-450.
- 22 Duncan, R.S., Duncan, V.E., 2000. Forest succession and distance from forest

- 1 edge in an Afro-tropical grassland. *Biotropica* 32, 33-41.
- 2 Engel, V.L., Parrotta, J.A., 2001. An evaluation of direct seeding for
3 reforestation of degraded lands in Central Sao Paulo State, Brazil. *Forest*
4 *Ecology and Management* 152, 169-181.
- 5 Fornara, D.A., Dalling, J.W., 2005. Post-dispersal removal of seeds of pioneer
6 species from five Panamanian forests. *Journal of Tropical Ecology* 21, 79-84.
- 7 Franco-Pizaña, J.G., Fulbright, T.E., Gardiner, D.T., Tipton, A.R., 1996. Shrub
8 emergence and seedling growth in microenvironments created by *Prosopis*
9 *glandulosa*. *Journal of Vegetation Science* 7, 257-264.
- 10 Friis, I., 1992. Forests and forest trees of Northeast Tropical Africa. Royal
11 Botanic Gardens, London.
- 12 Garcia, D., 2001. Effects of seed dispersal on *Juniperus communis* recruitment
13 on a Mediterranean mountain. *Journal of Vegetation Science* 12, 839-848.
- 14 García-Fayos, P., Verdú, M., 1998. Soil seed bank, factors controlling
15 germination and establishment of a Mediterranean shrub: *Pistacia lentiscus* L.
16 *Acta Oecologica-International Journal of Ecology* 19, 357-366.
- 17 Garwood, N.C., 1989. Tropical soil seed banks: a review. In: Leck, M.A.,
18 Simpson, R.L., Parker, V.T. (Eds.), *Ecology of Seed Banks*. Academic Press,
19 San Diego, pp. 149-209.
- 20 Gebresilassie, W., Bekele, A., Belay, G., Balakrishnan, M., 2004. Microhabitat
21 choice and diet of rodents in Maynugus irrigation field, northern Ethiopia.
22 *African Journal of Ecology* 42, 315-321.

- 1 Geist, H.J., Lambin, E.F., 2004. Dynamic causal patterns of desertification.
2 Bioscience 54, 817-829.
- 3 Gillespie, T.W., 1999. Life history characteristics and rarity of woody plants in
4 tropical dry forest fragments of Central America. Journal of Tropical Ecology 15,
5 637-649.
- 6 Green, P.S., 2003. 138. Oleaceae. In: Hedberg, I., Edwards, S., Nemomissa, S.
7 (Eds.), Flora of Ethiopia and Eritrea Volume 4, part 1, Apiaceae to
8 Dipsacaceae. The National Herbarium, Addis Ababa University, Ethiopia and
9 The Department of Systematic Botany, Uppsala University, Sweden, Addis
10 Ababa, Ethiopia & Uppsala, Sweden, pp. 79-86.
- 11 Guariguata, M.R., Pinard, M.A., 1998. Ecological knowledge of regeneration
12 from seed in Neotropical forest trees: implications for natural forest
13 management. Forest Ecology and Management 112, 87-99.
- 14 Hardwick, K., Healey, J., Elliott, S., Garwood, N., Anusarnsunthorn, V., 1997.
15 Understanding and assisting natural regeneration processes in degraded
16 seasonal evergreen forests in Northern Thailand. Forest Ecology and
17 Management 99, 203-214.
- 18 Hengsdijk, H., Meijerink, G.W., Mosugu, M.E., 2005. Modeling the effect of
19 three soil and water conservation practices in Tigray, Ethiopia. Agriculture,
20 Ecosystems & Environment 105, 29-40.
- 21 Herrera, C.M., Jordano, P., López-Soria, L., Amat, J.A., 1994. Recruitment of a
22 mast-fruited, bird-dispersed tree - bridging frugivore activity and seedling
23 establishment. Ecological Monographs 64, 315-344.

- 1 Herweg, K., Ludi, E., 1999. The performance of selected soil and water
2 conservation measures - case studies from Ethiopia and Eritrea . *Catena* 36,
3 99-114.
- 4 Holl, K.D., 2002. Effect of shrubs on tree seedling establishment in an
5 abandoned tropical pasture. *Journal of Ecology* 90, 179-187.
- 6 Holl, K.D., Lulow, M.E., 1997. Effects of species, habitat, and distance from
7 edge on post-dispersal seed predation in a tropical rainforest. *Biotropica* 29,
8 459-468.
- 9 Jones, F.A., Peterson, C.J., Haines, B.L., 2003. Seed predation in neotropical
10 pre-montane pastures: site, distance, and species effects. *Biotropica* 35, 219-
11 225.
- 12 Kalacska, M., Sanchez-Azofeifa, G.A., Calvo-Alvarado, J.C., Quesada, M.,
13 Rivard, B., Janzen, D.H., 2004. Species composition, similarity and diversity in
14 three successional stages of a seasonally dry tropical forest. *Forest Ecology*
15 *and Management* 200, 227-247.
- 16 Kidson, R., Westoby, M., 2000. Seed mass and seedling dimensions in relation
17 to seedling establishment. *Oecologia* 125, 11-17.
- 18 Kitajima, K., Fenner, M., 2000. Ecology of seedling regeneration. In: Fenner, M.
19 (Ed.), *Seeds. The ecology of regeneration in plant communities*. CAB
20 International, Oxon, UK, pp. 331-359.
- 21 Kollmann, J., Pirl, M., 1995. Spatial pattern of seed rain of fleshy-fruited plants
22 in a scrubland grassland transition. *Acta Oecologica-International Journal of*

- 1 Ecology 16, 313-329.
- 2 Lamprecht, H., 1989. Silviculture in the tropics. Tropical forest ecosystems and
3 their tree species - possibilities and methods for their long-term utilization.
4 Deutsche Gesellschaft für Technische Zusammenarbeit, Eschborn, Germany.
- 5 Lemenih, M., Karlun, E., Olsson, M., 2005. Assessing soil chemical and
6 physical property responses to deforestation and subsequent cultivation in
7 smallholders farming system in Ethiopia. Agriculture, Ecosystems &
8 Environment 105, 373-386.
- 9 Martínez-Garza, C., Howe, H.F., 2003. Restoring tropical diversity: beating the
10 time tax on species loss. Journal of Applied Ecology 40, 423-429.
- 11 Mcclanahan, T.R., Wolfe, R.W., 1993. Accelerating forest succession in a
12 fragmented landscape - the role of birds and perches. Conservation Biology 7,
13 279-288.
- 14 Mengistu, T., Teketay, D., Hulten, H., Yemshaw, Y., 2005. The role of
15 enclosures in the recovery of woody vegetation in degraded dryland hillsides of
16 Central and Northern Ethiopia. Journal of Arid Environments 60, 259-281.
- 17 Meze-Hausken, E., 2004. Contrasting climate variability and meteorological
18 drought with perceived drought and climate change in northern Ethiopia.
19 Climate Research 27, 19-31.
- 20 Mohr, K., Vibe-Petersen, S., Jeppesen, L.L., Bildsoe, M., Leirs, H., 2003.
21 Foraging of multimammate mice, *Mastomys natalensis*, under different
22 predation pressure: cover, patch-dependent decisions and density-dependent

1 GUDs. *Oikos* 100, 459-468.

2 Moles, A.T., Westoby, M., 2004. Seedling survival and seed size: a synthesis of
3 the literature. *Journal of Ecology* 92, 372-383.

4 Murray, K.G., Russell, S., Picone, C.M., Winnettmurray, K., Sherwood, W.,
5 Kuhlmann, M.L., 1994. Fruit laxatives and seed passage rates in frugivores -
6 consequences for plant reproductive success. *Ecology* 75, 989-994.

7 Negash, L., 2003. Vegetative propagation of the threatened African wild olive
8 [*Olea europaea* L. subsp. *cuspidata* (Wall. ex DC.) Ciffieri]. *New Forests* 26,
9 137-146.

10 Nyssen, J., Poesen, J., Moeyersons, J., Deckers, J., Haile, M., Lang, A., 2004.
11 Human impact on the environment in the Ethiopian and Eritrean highlands - a
12 state of the art. *Earth-Science Reviews* 64, 273-320.

13 Rey, P.J., Alcántara, J.M., 2000. Recruitment dynamics of a fleshy-fruited plant
14 (*Olea europaea*): connecting patterns of seed dispersal to seedling
15 establishment. *Journal of Ecology* 88, 622-633.

16 Rey, P.J., Alcántara, J.M., Valera, F., Sanchez-Lafuente, A.M., Garrido, J.L.,
17 Ramirez, J.M., Manzaneda, A.J., 2004. Seedling establishment in *Olea*
18 *europaea*: seed size and microhabitat affect growth and survival. *Ecoscience*
19 11, 310-320.

20 Rey, P.J., Garrido, J.L., Alcántara, J.M., Ramirez, J.M., Aguilera, A., Garcia, L.,
21 Manzaneda, A.J., Fernandez, R., 2002. Spatial variation in ant and rodent post-
22 dispersal predation of vertebrate-dispersed seeds. *Functional Ecology* 16, 773-

- 1 781.
- 2 Samuels, I.A., Levey, D.J., 2005. Effects of gut passage on seed germination:
3 do experiments answer the questions they ask? *Functional Ecology* 19, 365-
4 368.
- 5 Schatral, A., Fox, J.E.D., 1994. Quality and viability of seeds in the genus
6 *Hibbertia*. *Seed Science and Technology* 22, 273-284.
- 7 Schupp, E.W., Milleron, T., Russo, S.E., 2002. Dissemination limitation and the
8 origin and maintenance of species-rich tropical forests. In: Levey, D.J., Silva,
9 W.R., Galetti, M. (Eds.), *Seed dispersal and frugivory: ecology, evolution and*
10 *conservation*. CAB International, Oxon, UK, pp. 19-33.
- 11 Stiles, E.W., 2000. Animals as seed dispersers. In: Fenner, M. (Ed.), *Seeds.*
12 *The ecology of regeneration in plant communities*. CAB International,
13 Wallingford, UK, pp. 111-124.
- 14 Sun, D., Dickinson, G.R., Bragg, A.L., 1995. Direct seeding of *Alphitonia petriei*
15 (Rhamnaceae) for gully revegetation in tropical northern Australia. *Forest*
16 *Ecology and Management* 73, 249-257.
- 17 Teketay, D., Granström, A., 1997. Seed viability of Afromontane tree species in
18 forest soils. *Journal of Tropical Ecology* 13, 81-95.
- 19 Tekle, K., 2001. Natural regeneration of degraded hillslopes in Southern Wello,
20 Ethiopia: a study based on permanent plots. *Applied Geography* 21, 275-300.
- 21 Tekle, K., Bekele, T., 2000. The role of soil seed banks in the rehabilitation of
22 degraded hillslopes in Southern Wello, Ethiopia. *Biotropica* 32, 23-32.

- 1 Traveset, A., Verdú, M., 2002. A meta-analysis of the effect of gut treatment on
2 seed germination. In: Levey, D.J., Silva, W.R., Galetti, M. (Eds.), Seed dispersal
3 and frugivory: ecology, evolution and conservation. CAB International, Oxon,
4 UK, pp. 339-350.
- 5 Turner, I.M., Corlett, R.T., 1996. The conservation value of small, isolated
6 fragments of lowland tropical rain forest. *Trends in Ecology & Evolution* 11, 330-
7 333.
- 8 Vancampenhout, K., Nyssen, J., Gebremichael, D., Deckers, J., Poesen, J.,
9 Haile, M., Moeyersons, J., 2006. Stone bunds for soil conservation in the
10 northern Ethiopian highlands: Impacts on soil fertility and crop yield. *Soil and*
11 *Tillage Research*. In press (doi:10.1016/j.still.2005.08.004).
- 12 Vander Wall, S.B., 1990. Food hoarding in animals. University of Chicago
13 Press, Chicago.
- 14 Vander Wall, S.B., Kuhn, K.M., Beck, M.J., 2005. Seed removal, seed
15 predation, and secondary dispersal. *Ecology* 86, 801-806.
- 16 Verdú, M., García-Fayos, P., 1996. Postdispersal seed predation in a
17 Mediterranean patchy landscape. *Acta Oecologica-International Journal of*
18 *Ecology* 17, 379-391.
- 19 Vetaas, O.R., 1992. Micro-site effects of trees and shrubs in dry savannas.
20 *Journal of Vegetation Science* 3, 337-344.
- 21 Wassie, A., Teketay, D., 2006. Soil seed banks in church forests of northern
22 Ethiopia: Implications for the conservation of woody plants. *Flora* 201, 32-43.

- 1 Wenny, D.G., 2000. Seed dispersal, seed predation, and seedling recruitment
2 of a neotropical montane tree. *Ecological Monographs* 70, 331-351.
- 3 Woods, K., Elliott, S., 2004. Direct seeding for forest restoration on abandoned
4 agricultural land in northern Thailand. *Journal of Tropical Forest Science* 16,
5 248-259.
- 6 Yirdaw, E., Luukkanen, O., 2003. Indigenous woody species diversity in
7 *Eucalyptus globulus* Labill. ssp *globulus* plantations in the Ethiopian Highlands.
8 *Biodiversity and Conservation* 12, 567-582.
- 9 Zeleke, G., Hurni, H., 2001. Implications of land use and land cover dynamics
10 for mountain resource degradation in the Northwestern Ethiopian highlands.
11 *Mountain Research and Development* 21, 184-191.
- 12
- 13

1 Tables

2

Table 1. Study sites and seed removal and germination experiments carried out in northern Ethiopia

Site	Location	Elevation (m a.s.l.)	Rainfall (mm y ⁻¹)	Land use type	Experiments ^c			
					SR	DS	HG	BG
Endayesus	13°28'N, 39°30'E	2200	625 ± 155 ^a	Experimental tree nursery	–	–	×	×
Bubu Hills	13°30'N, 39°29'E	2150	≈ Endayesus ^b	Peri-urban exclosure	–	×	–	–
Mheni	13°36'N, 39°21'E	1860	< Endayesus ^b	Rural community exclosure	×	×	–	–
Sesemat	13°37'N, 39°19'E	1940	< Endayesus ^b	Rural community exclosure	×	–	–	–

^a Meze-Hausken (2004)

^b R. Aerts, pers. obs. (2001–2004)

^c SR: seed removal experiments, DS: direct sowing experiments, HG: controlled germination experiments in reconstructed humus, BG: controlled germination experiments after pulp and endocarp treatments.

3

4

Table 2. Two-way ANOVA significance and mean per cent seed germination $\pm$ SE of pulp removal and mechanical endocarp treatments in a controlled germination experiment of *Olea europaea* seeds

Pulp removal	Treatment means $\pm$ SE ^a			Factor significance ^{b,c}	
	Control	Manual	Bird	F _{2,13}	P
23 weeks after sowing	3 $\pm$ 1 ^a	17 $\pm$ 4 ^{ab}	27 $\pm$ 4 ^b	4.838	0.027*
46 weeks after sowing	37 $\pm$ 7 ^a	35 $\pm$ 6 ^a	44 $\pm$ 7 ^a	3.933	0.046*
Endocarp treatment	Control	Scarification	Splitting	F _{2,13}	P
23 weeks after sowing	13 $\pm$ 4 ^a	14 $\pm$ 4 ^a	32 $\pm$ 4 ^b	5.510	0.018*
46 weeks after sowing	52 $\pm$ 5 ^a	24 $\pm$ 5 ^b	32 $\pm$ 4 ^b	12.789	0.001***

^a Letters indicate significant differences between groups ($P < 0.05$; Tukey's HSD).

^b Significance: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

^c Interaction between pulp removal and endocarp treatment not significant: week 23 $P = 0.665$; week 46 $P = 0.920$.

1

2

Figure captions

Figure 1. Seed survival probability calculated as mean probability $\pm$ SE for depots with 10 *Olea* seeds to remain undetected i) near stone bunds (< 1 m; $N = 31$; $-\blacktriangle-$) and ii) further away from stone bunds (> 1 m; $N = 20$; $-\circ-$) in the Mheni exclosure, September 2003. Day 0 is 2 September 2003. After 20 September, no further seed removal was detected.

Figure 2. Germination probability of *Olea* seeds (mean $\pm$ SE; 10 seeds per replication) in three microhabitats of the Bubu Hills exclosure, 24 July – 17 September 2003: i) under *Euclea* cover ($N = 10$; $-\bullet-$), ii) under *Acacia* cover ($N = 10$; $-\Delta-$) and iii) in open patches between shrubs ($N = 10$; $-x-$).

Figure 3. Germination probability of *Olea* seeds (mean; 20 seeds per replication) in reconstructed humus of three microhabitats, 24 July 2003 – 20 February 2004: i) *Euclea* humus ($N = 5$; $-\bullet-$), ii) *Acacia* humus ($N = 5$; $-\Delta-$) and iii) stone cover of open patches between shrubs ($N = 5$; $-x-$). Standard errors of means were omitted for clarity.

Figure 4. Germination probability of *Olea* seeds (mean; 50 seeds per replication) after pulp removal and endocarp scarification treatments: pulp removed by birds, no endocarp treatment ($N = 3$; $-\blacksquare-$), pulp manually removed, no endocarp treatment ($N = 3$; $-\square-$), pulp removed by birds, endocarp split ($N = 3$; $-\blacktriangle-$), pulp manually removed, endocarp split ($N = 3$; $-\Delta-$), pulp removed by birds, endocarp scarification ($N = 2$; $-\bullet-$), pulp

- 1 manually removed, endocarp scarification (N = 3; —○—), intact drupe (overall
- 2 control) (N = 3; —x—). Standard errors of means were omitted for clarity.
- 3 Seeds were sown in the seed beds on 11 August 2002. The arrow indicates
- 4 the onset of the 2003 spring rains (27 January 2003).

1 **Figures**

2 **Fig. 1**

3

1 Fig. 2

2

1 Fig. 3

2

1 Fig. 4

2

3