0Ė ÁŮÇGDă*[!ã@ Á[!Á•ċ*Á@Á^}^!ăă^ăÁă{Á
[Á^ç^ă^ă!^~!^}&
0ă!ă^Áăă[ăă[}ĖŮăŮ{^|ă!•ÁăăăăŮ]ă\•{ă

DEPARTMENT OF DECISION SCIENCES AND INFORMATION MANAGEMENT (KBI)

An $O(n^2)$ Algorithm for Testing the Generalized Axiom of Revealed Preference

Fabrice Talla Nobibon^{*†}, Bart Smeulders[†], Frits C.R. Spieksma[†]

Abstract. This note presents an algorithm for testing the generalized axiom of revealed preference in time $O(n^2)$, where n is the number of observations in a given data set. Furthermore, we prove a lower bound of $\Omega(n \log n)$ on the running time of any algorithm for testing different axioms of revealed preference.

Keywords: generalized axiom of revealed preference; directed graph; strongly connected components; computational complexity; microeconomics.

1. Introduction

This note contains two contributions: (1) we propose an $O(n^2)$ algorithm for testing the so-called *generalized axiom of revealed preference*, and (2) we prove a lower bound of $\Omega(n \log n)$ on the running time of any algorithm for testing three well-known revealed preference axioms (here, n is the number of the observations in the data set).

Modeling and analyzing household consumption behavior is a central research topic in microeconomics since its introduction by Samuelson [8–10] for a unitary household. For a recent overview we refer to Varian [13] and Chapter 7 in Vohra [14]. Here, we restrict ourselves to describing three well-known axioms of revealed preference: the *weak axiom of revealed preference* (WARP), the *strong axiom of revealed preference* (SARP), and the *generalized axiom of revealed preference* (GARP). Consider a unitary household acting in an economy with m goods and suppose that we have observed n consumption quantity bundles $q^i = (q_1^i, \dots, q_m^i) \in \mathbb{R}_+^m$ (non-negative) with corresponding positive prices $p^i = (p_1^i, \dots, p_m^i) \in \mathbb{R}_{++}^m$, for $i = 1, \dots, n$. The component q_j^i (respectively p_j^i), for $j = 1, \dots, m$, corresponds with the quantity of good j bought by the household (respectively the unit price of good j) at the time of observation i . We denote the set of observations by $S = \{(p^i, q^i) : i \in N\}$ where $N = \{1, \dots, n\}$ and refer to S as the *data set*. For ease of exposition, we use $i \in N$ to refer to the observation (p^i, q^i) and denote the scalar product by $p^i q^j$ for $i, j \in N$.

^{*}PostDoc researcher for Research Foundation–Flanders. E-mail: Fabrice.TallaNobibon@econ.kuleuven.be

[†]ORSTAT, Faculty of Business and Economics, KULeuven, Belgium

In revealed preference theory, a unitary household is said to *directly prefer* bundle q^i over another bundle q^j if q^i was chosen while q^j was affordable (and could have been chosen); this translates into $p^i q^i \geq p^i q^j$. A household is said to *prefer* q^i over q^j if there exists a sequence (possibly empty) of observations $r, s, \dots, t$ such that $p^i q^i \geq p^i q^r$, $p^r q^r \geq p^r q^s$, $\dots$, and $p^t q^t \geq p^t q^j$. These notions of preference are used in the definition of WARP, SARP, and GARP [13].

Definition 1. A data set S satisfies WARP if and only if for any pair of distinct observations i and j satisfying $p^i q^i \geq p^i q^j$ we have $p^j q^j < p^j q^i$.

Definition 2. A data set S satisfies SARP if and only if for any sequence of distinct observations $i, r, s, \dots, t$, and j satisfying $p^i q^i \geq p^i q^r$, $p^r q^r \geq p^r q^s, \dots$, and $p^t q^t \geq p^t q^j$, we have $p^j q^j < p^j q^i$.

Definition 3. A data set S satisfies GARP if and only if for any sequence of distinct observations $i, r, s, \dots, t$, and j satisfying $p^i q^i \geq p^i q^r$, $p^r q^r \geq p^r q^s, \dots$, and $p^t q^t \geq p^t q^j$, we have $p^j q^j \leq p^j q^i$.

Varian [13] shows that the question of testing whether a given data set S satisfies WARP, SARP, or GARP can be answered in polynomial time. For testing WARP, we compare for each pair (i, j) of distinct observations $p^i q^i$ and $p^i q^j$, on the one hand, and $p^j q^j$ and $p^j q^i$, on the other hand. Because in total we perform $O(n^2)$ comparisons, we infer that testing WARP can be done in $O(n^2)$ time. For testing SARP, a directed graph G with n vertices is built where each vertex corresponds with an observation and there is an arc from vertex i to vertex j if and only if $p^i q^i \geq p^i q^j$. In term of this directed graph, the data set S satisfies SARP if and only if G is *acyclic*. (We recall that an acyclic graph is a graph that contains no cycle.) Observe that building the graph can be done in time $O(n^2)$ whereas checking whether G is acyclic is done in time $O(n + m) = O(n^2)$ using, for example, the topological ordering algorithm [1]. In total, testing SARP can be done in $O(n^2)$ time. The current algorithm for testing GARP is based on computing the transitive closure [12, 13] and proceeds as follows. A binary $n \times n$ matrix M is defined to summarize the direct preference relations. In the matrix M the (i, j) entry is given by $M_{ij} = 1$ if $p^i q^i \geq p^i q^j$ and $M_{ij} = 0$ otherwise. Next, the transitive closure MT of M is computed and based on MT , GARP is decided by checking whether there exists a pair (i, j) satisfying $MT_{ij} = 1$ and $p^j q^j > p^j q^i$. The data set satisfies GARP if and only if such pair (i, j) does not exist. The bottleneck of this procedure is the computation of the transitive closure of M . Varian [12] uses the $O(n^3)$ -algorithm proposed by Warshall [15] and mentions the possibility of computation the transitive

closure in time $O(n^{2.74})$ using the algorithm designed by Munro [7]. In the literature, the best-known algorithm for computation the transitive closure is based on matrix multiplication that runs in time $O(n^{2.376})$ [5]. Recently, several researchers have investigated a group-theoretic approach for computing matrix multiplication, with the hope of reducing the exponent (2.376) of this time complexity; we refer to [3,4,6] for more details. This approach, however, has led to smaller exponent only if some restrictive conditions are imposed on the structure of the matrices [3]. Even with these restrictions, the exponent is still greater than 2. Notice that the above complexity statements related to algorithms for computing WARP, SARP, and GARP assume that the scalar product $p^i q^j$ ($i, j \in N$) can be computed in constant time; we will follow this assumption throughout this note. In the next section, we present a simple algorithm for deciding GARP that runs in time $O(n^2)$ and we prove a lower bound of $\Omega(n \log n)$ on the running time of any algorithm devised for testing either WARP, SARP, or GARP.

2. Results

Given a data set $S = \{(p^i, q^i) : i \in N\}$, we build a directed graph $G = (V, A)$ with n vertices, where each vertex corresponds with an observation. Furthermore, the set of arcs is the union of two disjoint sets $A_>$ and $A_=-$; that is: $A = A_> \cup A_=-$. There is an arc from vertex i to vertex j (with $i \neq j$) in $A_>$ (respectively in $A_=-$) if and only if $p^i q^i > p^i q^j$ (respectively $p^i q^i = p^i q^j$). Notice that $A_>$ and $A_=-$ need not be non empty. Let us define the subgraphs $G_> = (V, A_>)$ and $G_-= (V, A_=-)$ of G . We recall that a *vertex-induced subgraph* of G (subsequently called induced subgraph) is a subset of vertices of G together with all arcs whose endpoints are both in that subset. Also, a *strongly connected component* (SCC) of G is a maximal induced subgraph $B = (V(B), A(B))$ of G , where for every pair of vertices $i, j \in V(B)$, there is a path (oriented) from i to j and a path from j to i in B . Each graph has a unique partition of its nodes into strongly connected components. We denote by $G_{scc} = (V, A_{scc})$ the subgraph of G containing only arcs that appear in a strongly connected component of G . We have the following result:

Lemma 1. *A data set S satisfies GARP if and only if $G_{scc} \subseteq G_=-$.*

Proof: $\Rightarrow$) Suppose that S satisfies GARP. We want to show that $G_{scc} \subseteq G_=-$. Clearly, if there is no arc in G_{scc} (each SCC is a singleton) then $G_{scc} \subseteq G_=-$. Let $i \rightarrow j$ be an arbitrary arc in G_{scc} .

Then, there exists a unique SCC of G , let us say $B = (V(B), A(B))$, such that $i \rightarrow j \in A(B)$. Because B is a SCC, there exists a sequence of arcs $j \rightarrow r \in A(B)$, $r \rightarrow s \in A(B)$, $\dots$, and $t \rightarrow i \in A(B)$ such that $i \rightarrow j \rightarrow r \rightarrow s \rightarrow \dots \rightarrow t \rightarrow i$ is a cycle in B . Given that the arc $j \rightarrow r$ (respectively $r \rightarrow s$, $\dots$, and $t \rightarrow i$) exists if and only if $p^j q^j \geq p^j q^r$ (respectively $p^r q^r \geq p^r q^s, \dots$, and $p^t q^t \geq p^t q^i$) we have, from the definition of GARP, that $p^i q^i \leq p^i q^j$. However, the existence of $i \rightarrow j$ implies that $p^i q^i \geq p^i q^j$. Because $p^i q^i \leq p^i q^j$ and $p^i q^i \geq p^i q^j$, we infer that $p^i q^i = p^i q^j$ and therefore $i \rightarrow j \in G_{=}$. This completes the proof that $G_{scc} \subseteq G_{=}$.

$\Leftarrow$) We now suppose that $G_{scc} \subseteq G_{=}$ and we want to show that S satisfies GARP. Consider an arbitrary sequence of observations $s, u, v, \dots, z$, and t satisfying $p^s q^s \geq p^s q^u$, $p^u q^u \geq p^u q^v$, $\dots$, and $p^z q^z \geq p^z q^t$. There are two options: either $p^t q^t > p^t q^s$ or $p^t q^t \leq p^t q^s$. We argue that the first option cannot occur. Indeed, if $p^t q^t > p^t q^s$ then together with the previous inequalities, this implies the existence of the cycle $\mathcal{C} : s \rightarrow u \rightarrow v \dots z \rightarrow t \rightarrow s$ in G and there exists a SCC of G , let us say $B = (V(B), A(B))$, such that $\mathcal{C} \subseteq B \subseteq G_{scc}$. However, the inequality $p^t q^t > p^t q^s$ implies that the arc $t \rightarrow s \in A_{>}$. We conclude that the cycle $\mathcal{C}$ is not included in $G_{=}$ and hence G_{scc} is not a subgraph of $G_{=}$. This would contradict the hypothesis that $G_{scc} \subseteq G_{=}$. Therefore, the only option left is $p^t q^t \leq p^t q^s$; which implies that S satisfies GARP. This completes the proof of Lemma 1. $\square$

We propose the following algorithm for testing whether a given data set S satisfies GARP. The

Algorithm 1 A graph algorithm for testing GARP

- 1: build the directed graph $G = (V, A)$
 - 2: compute G_{scc}
 - 3: **if** every arc present in G_{scc} is also in $G_{=}$ **then** return yes
 - 4: **else** return no
-

main difference between Algorithm 1 and the current testing procedure stems from the fact that the former algorithm computes the strongly connected components of a directed graph whereas the latter computes the transitive closure of a matrix that represents the same graph.

Theorem 1. *Algorithm 1 detects whether a given data set S satisfies GARP in time $O(n^2)$.*

Proof: The correctness of the algorithm results from Lemma 1. Let us now analyze its complexity. Observe that $|V| = n$, the number of observations, whereas $m = |A| = O(n^2)$. The first step

of the algorithm, building the directed graph, can be done in time $O(n^2)$ given the assumption that the scalar product is computed in constant time. Indeed, there are $O(n^2)$ comparisons to be made in order to build the graph G . The second step, computing G_{scc} , if implemented using Tarjan's algorithm [11], is completed in time $O(n + m) \equiv O(n^2)$. Finally, the last step (the **if** loop) has a running time of $O(m) \equiv O(n^2)$. Therefore, the overall running time of the algorithm is $3 \times O(n^2) = O(n^2)$. $\square$

We are now going to derive a lower bound of $\Omega(n \log n)$ on the running time of any algorithm devised for testing either WARP, SARP, or GARP on a data set S with n observations. This is achieved using a reduction from the following *decision problem*, also known as the *Element Distinctness* problem [2, 16].

Instance: A set $x_1, x_2, \dots, x_m$ of m integers.

Question: Is there a pair i, j with $i \neq j$ and $x_i = x_j$?

Using a topological method, Yao [16] proves that any algebraic computation tree that solves the m -Element Distinctness problem has a lower bound complexity of $\Omega(m \log m)$. We next show that an algorithm for testing WARP, SARP, or GARP can also be used for determining whether m integers are pairwise distinct.

Given an instance $x_1, x_2, \dots, x_m$ of the Element Distinctness problem, we build a data set S as follows. We assume that we are in an economy with m goods. To describe the prices and the quantities of all goods for each observation we make use of a 'default' price (quantity) for each good. The vector of default prices is $P = (x_1 - 0.1, x_2 - 0.1, \dots, x_m - 0.1)$ whereas that of default quantities is $Q = (0, 0, \dots, 0)$. Notice that to describe these default vectors, we need $O(m)$ operations. We consider a data set S with m observations where an observation is identified by the index of a good. This index means that for the considered observation that good, let us say j , has the price of $x_j + 0.1$ (instead of $x_j - 0.1$ as in the default vector) whereas the quantity of that good is now 1 (instead of 0 in the default quantity vector). The price (respectively the quantity) of each remaining good is exactly its default price (respectively quantity). Notice that this second part of our reduction required $O(m)$ operations because given the default price and quantity vectors, we need exactly $O(m)$ numbers to describe the data set S . We now prove that the data set S satisfies

WARP, SARP, or GARP if and only if the considered instance of the Element distinctness problem is a yes instance.

Consider the directed graph built from S and observe that there is an arc from i to j if and only if $p^i q^i \geq p^i q^j$; that is $x_i + 0.1 \geq x_j - 0.1$, which is equivalent to $x_i \geq x_j$ (because x_i and x_j are integers). Notice that in that graph, if there is a cycle $s \rightarrow u \rightarrow v \dots z \rightarrow t \rightarrow s$ then the two-cycles $s \rightarrow u \rightarrow s$, $u \rightarrow v \rightarrow u$, $\dots$, and $t \rightarrow s \rightarrow t$ are all present. This observation implies that if the graph does not contain any two-cycle then it does not contain any cycle and vice-versa. Stated in term of the data set, the observation implies that S satisfies WARP if and only if S satisfies SARP. Another remark is that for all pair of distinct observations i and j we have $p^i q^i = x_i + 0.1 \neq x_j - 0.1 = p^i q^j$, because x_i and x_j are integers. This means that for any pair of distinct observations i and j we have $p^i q^i \neq p^i q^j$. These inequalities imply that S satisfies GARP if and only if S satisfies SARP. Observe now that a two-cycle $i \rightarrow j \rightarrow i$ is present in our graph if and only if $x_i = x_j$; in other words, there is a two-cycle $i \rightarrow j \rightarrow i$ in our graph if and only if the two elements x_i and x_j of our set are identical. It immediately follows that the data set S satisfies WARP, SARP, or GARP if and only if the considered instance of the Element Distinctness problem is a yes instance. This proves that any algorithm for solving WARP, SARP, or GARP can be used to solve the Element Distinctness problem. Therefore, the lower bound of $\Omega(m \log m)$ inherent to any algorithm for solving the Element Distinctness problem is directly applicable to any algorithm for solving WARP, SARP, or GARP.

We formalize this result in the following theorem.

Theorem 2. *Any algorithm for testing either WARP, SARP, or GARP on a data set S with n observations has a running time bounded from below by $\Omega(n \log n)$.*

3. Conclusion

This note presents a new algorithm for testing GARP that runs in time $O(n^2)$, where n is the number of observations in the consumption data set. We also prove a lower bound of $\Omega(n \log n)$ on the running time of any algorithm for testing the weak axiom of revealed preference, the strong axiom of revealed preference, and the generalized axiom of revealed preference.

References

- [1] R.K. Ahuja, T.L. Magnanti, and J.B. Orlin. *Network Flows: Theory, Algorithms, and Applications*. Prentice Hall, 1993.
- [2] M. Ben-Or. Lower bounds for algebraic computation trees. In *Proceedings of 15th ACM Symposium on Theory of Computing*, pages 80–86, 1983.
- [3] H. Cohn, R. Kleinberg, B. Szegedy, and C. Umans. Group-theoretic algorithms for matrix multiplication. In *Foundations of Computer Science. 46th Annual IEEE Symposium*, pages 379–388, 2005.
- [4] H. Cohn and C. Umans. A group-theoretic approach to matrix multiplication. In *Foundations of Computer Science. 44th Annual IEEE Symposium*, pages 438–449, 2003.
- [5] D. Coppersmith and S. Winograd. Matrix multiplication via arithmetic progressions. *Journal of Symbolic Computation*, 9(3):251–280, 1990.
- [6] J. Demmel, I. Dumitriu, O. Holtz, and R. Kleinberg. Fast matrix multiplication is stable. *Numerische Mathematik*, 106:199–224, 2007.
- [7] I. Munro. Efficient determination of the transitive closure of a directed graph. *Information Processing Letters*, 1(2):56–58, 1971.
- [8] P.A. Samuelson. A note on the pure theory of consumer’s behaviour. *Economica*, 5(17):61–71, 1938.
- [9] P.A. Samuelson. The empirical implications of utility analysis. *Economica*, 6(4):344–356, 1938.
- [10] P.A. Samuelson. Consumption theory in terms of revealed preference. *Economica*, 15(60):243–253, 1948.
- [11] R.E. Tarjan. Depth-first search and linear graph algorithms. *SIAM Journal on Computing*, 1(2):146–160, 1972.
- [12] H.R. Varian. The nonparametric approach to demand analysis. *Economica*, 50(4):945–976, 1982.

- [13] H.R. Varian. Revealed preference. *Samuelsonian Economics and the 21st Century*, 2006.
- [14] R. Vohra. *Mechanism Design: A Linear Programming Approach*. Cambridge University Press, 2011.
- [15] S. Warshall. A theorem on boolean matrices. *Journal of the American Association of Computing Machinery*, 9(1):11–12, 1962.
- [16] A.C.-C. Yao. Lower bounds for algebraic computation trees with integer inputs. *SIAM Journal on Computing*, 20(4):655–668, 1991.