Dear Author

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For **fax** submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style.
- Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL:

<http://dx.doi.org/10.1007/s13105-012-0204-9>

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information, go to:

<http://www.springerlink.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us, if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

1	Article Title	Endurance training in mice increases the unfolded protein response induced by a high-fat diet	
2	Article Sub- Title		
3	Article Copyright - Year	University of Navarra 2012 (This will be the copyright line in the final PDF)	
4	Journal Name	Journal of Physiology and Biochemistry	
5	Corresponding Author	Family Name	Deldicque
6		Particle	
7		Given Name	Louise
8		Suffix	
9		Organization	K.U. Leuven
10		Division	Research Centre for Exercise and Health, Department of Biomedical Kinesiology
11		Address	Tervuursevest 101, Box 1500, Leuven 3001, Belgium
12		Organization	Université Catholique de Louvain
13	Author	Division	Institute of Neurosciences, Research Group in Muscle Exercise and Physiology
14		Address	Place Pierre de Coubertin, 1, Louvain-la-Neuve 1348, Belgium
15		e-mail	louise.deldicque@faber.kuleuven.be
16		Family Name	Cani
17	Author	Particle	
18		Given Name	Patrice D.
19		Suffix	
20		Organization	Université Catholique de Louvain
21		Division	Louvain Drug Research Institute, Metabolism and Nutrition research group
22		Address	Avenue E. Mounier, 73, PMNT 73/69, Brussels 1200, Belgium
23	Author	e-mail	
24		Family Name	Delzenne
25		Particle	
26		Given Name	Nathalie M.

27		Suffix	
28		Organization	Université Catholique de Louvain
29		Division	Louvain Drug Research Institute, Metabolism and Nutrition research group
30		Address	Avenue E. Mounier, 73, PMNT 73/69, Brussels 1200, Belgium
31		e-mail	
32	Author	Family Name	Baar
33		Particle	
34		Given Name	Keith
35		Suffix	
36		Organization	University of California Davis
37		Division	Department of Neurobiology, Physiology and Behavior
38		Address	One Shields Avenue, 181 Briggs Hall, Davis 95616, CA, USA
39		e-mail	
40	Author	Family Name	Francaux
41		Particle	
42		Given Name	Marc
43		Suffix	
44		Organization	Université Catholique de Louvain
45		Division	Institute of Neurosciences, Research Group in Muscle Exercise and Physiology
46		Address	Place Pierre de Coubertin, 1, Louvain-la-Neuve 1348, Belgium
47		e-mail	
48	Schedule	Received	19 September 2011
49		Revised	
50		Accepted	27 July 2012
51	Abstract	Certain conditions, such as several weeks of high-fat diet, disrupt endoplasmic reticulum (ER) homeostasis and activate an adaptive pathway referred as the unfolded protein response. When the unfolded protein response fails, the result is the development of inflammation and insulin resistance. These two pathological states are known to be improved by regular exercise training but the mechanisms remain largely undetermined. As it has recently been shown that the unfolded protein response is regulated by exercise, we hypothesised that concomitant treadmill exercise training (HFD+ex) prevents ER homeostasis disruption and its downstream consequences induced by a 6-week high-fat diet (HFD) in mice by	

activating the protective unfolded protein response. Several well-documented markers of the unfolded protein response were measured in the soleus and tibialis anterior muscles as well as in the liver and pancreas. In HFD mice, an increase in these markers was observed (from 2- to 15-fold, $P<0.05$) in all tissues studied. The combination of HFD+ex increased the expression of several markers further, up to 100 % compared to HFD alone ($P<0.05$). HFD increased inflammatory markers both in the plasma (IL-6 protein, 2.5 ± 0.52 -fold; MIP-1 α protein, 1.3 ± 0.13 -fold; $P<0.05$) and in the tissues studied, and treadmill exercise attenuated the inflammatory state induced by HFD ($P<0.05$). However, treadmill exercise could not reverse HFD-induced whole body glucose intolerance, assessed by OGTT (AUC, 1.8 ± 0.29 -fold, $P<0.05$). In conclusion, our results show that a HFD activated the unfolded protein response in mouse tissues in vivo, and that endurance training promoted this response. We speculate that the potentiation of the unfolded protein response by endurance training may represent a positive adaptation protecting against further cellular stress.

52	Keywords separated by ' - '	Skeletal muscle - ER stress - BiP/GRP78 - XBP1 - Caspase 12
----	-----------------------------	---

53	Foot note information	
----	-----------------------	--

Endurance training in mice increases the unfolded protein response induced by a high-fat diet

Louise Deldicque · Patrice D. Cani ·
Nathalie M. Delzenne · Keith Baar ·
Marc Francaux

Received: 19 September 2011 / Accepted: 27 July 2012
© University of Navarra 2012

Abstract Certain conditions, such as several weeks of high-fat diet, disrupt endoplasmic reticulum (ER) homeostasis and activate an adaptive pathway referred as the unfolded protein response. When the unfolded protein response fails, the result is the development of inflammation and insulin resistance. These two pathological states are known to be improved by regular exercise training but the mechanisms remain largely undetermined. As it has recently been shown

that the unfolded protein response is regulated by exercise, we hypothesised that concomitant treadmill exercise training (HFD+ex) prevents ER homeostasis disruption and its downstream consequences induced by a 6-week high-fat diet (HFD) in mice by activating the protective unfolded protein response. Several well-documented markers of the unfolded protein response were measured in the soleus and tibialis anterior muscles as well as in the liver and pancreas. In HFD mice, an increase in these markers was observed (from 2- to 15-fold, $P<0.05$) in all tissues studied. The combination of HFD+ex increased the expression of several markers further, up to 100 % compared to HFD alone ($P<0.05$). HFD increased inflammatory markers both in the plasma (IL-6 protein, 2.5 ± 0.52 -fold; MIP-1 α protein, 1.3 ± 0.13 -fold; $P<0.05$) and in the tissues studied, and treadmill exercise attenuated the inflammatory state induced by HFD ($P<0.05$). However, treadmill exercise could not reverse HFD-induced whole body glucose intolerance, assessed by OGTT (AUC, 1.8 ± 0.29 -fold, $P<0.05$). In conclusion, our results show that a HFD activated the unfolded protein response in mouse tissues in vivo, and that endurance training promoted this response. We speculate that the potentiation of the unfolded protein response by endurance training may represent a positive adaptation protecting against further cellular stress.

L. Deldicque (✉)
Research Centre for Exercise and Health,
Department of Biomedical Kinesiology, K.U. Leuven,
Tervuursevest 101, Box 1500, 3001 Leuven, Belgium
e-mail: louise.deldicque@faber.kuleuven.be

L. Deldicque · M. Francaux
Institute of Neurosciences, Research Group
in Muscle Exercise and Physiology,
Université Catholique de Louvain,
Place Pierre de Coubertin, 1,
1348 Louvain-la-Neuve, Belgium

P. D. Cani · N. M. Delzenne
Louvain Drug Research Institute, Metabolism and Nutrition
research group, Université Catholique de Louvain,
Avenue E. Mounier, 73, PMNT 73/69,
1200 Brussels, Belgium

K. Baar
Department of Neurobiology, Physiology and Behavior,
University of California Davis,
One Shields Avenue, 181 Briggs Hall,
Davis, CA 95616, USA

Keywords Skeletal muscle · ER stress · BiP/GRP78 ·
XBP1 · Caspase 12

Introduction

The endoplasmic reticulum (ER) is a key organelle, in which the folding and post-translational modifications of proteins occur. It is also crucial in the selection and transport of proteins to other compartments both within and outside of the cell. Certain stress conditions, such as high lipids, glucose deprivation and increased synthesis of secretory proteins, disrupt ER homeostasis and lead to the accumulation of unfolded or misfolded proteins in the ER lumen [36]. To cope with this stress, cells activate a signal transduction system that links the ER lumen to the cytoplasm and the nucleus; this process is referred to as the unfolded protein response [22, 27, 33]. The unfolded protein response is required for the restoration of normal ER function through three main unfolded protein response transducers: ATF6 (activating transcription factor 6); IRE1 α (inositol-requiring enzyme 1 alpha); and PERK (protein kinase R-like ER protein kinase). Each of these factors associates with the protein chaperone BiP/GRP78 (binding protein/glucose regulated-protein 78), a member of the Hsp70 (heat shock protein 70) family, in its inactive state. Upon accumulation of unfolded/misfolded proteins in the ER lumen, ATF6, IRE1 α , and PERK are released from BiP/GRP78 and become activated. The downstream effectors of these three pathways induce the expression of genes, such as XBP1 (X box binding protein 1), CHOP [C/EBP (CCAAT/enhancer binding protein) homologous protein] and ATF4 (activating transcription factor 4), which encode proteins that function to augment the ER protein-folding capacity.

When the unfolded protein response fails, the result is cell death, usually in the form of apoptosis triggered by the cleavage of pro-caspase 12 at the ER membrane [30, 35]. In addition to apoptosis, ER stress can increase inflammatory state [18] and decrease glucose tolerance by two distinct processes. First, ER stress in pancreatic islets results in dysfunctional beta-cells and in immature insulin release in the blood [21, 33]. Second, ER stress in peripheral insulin-sensitive tissues such as liver and adipose tissue leads to an increase in JNK (c-Jun N-terminal kinase)-mediated serine phosphorylation of IRS-1 (insulin receptor-substrate 1) and thereby to inhibition of insulin action via decreased signalling to Akt/PKB (protein kinase B) [31].

However, it is not clear whether similar effects occur in skeletal muscle. Using two different models

of high-fat-fed mice, we recently showed that the unfolded protein response was increased in skeletal muscles [6] although we could not confirm those results in human skeletal muscle [7]. Since exercise is known to prevent glucose intolerance and to reduce inflammatory state, endurance training has been hypothesised to reduce ER stress previously induced by high-fat feeding in rat adipose and hepatic tissues [5]. The conclusions of the previous study were that swimming reduced pro-inflammatory molecules and ER stress markers and increased Akt/PKB phosphorylation in adipose and hepatic tissues of diet-induced obese rats. While the results of the previous study are important for understanding the mechanisms of the beneficial effects of exercise in obesity and diabetes, one could ask if the same occurs in skeletal muscle, proportionally the most important tissue by weight in the body. The aim of the present study was thus to determine whether contractile activity alters the unfolded protein response in skeletal muscle and whole body glucose tolerance in 6-week high-fat-fed mice. Liver and pancreas were studied at the same time to detect whether the potential regulation of the unfolded protein response by exercise is tissue-specific.

Materials and methods

Animals and diets

Two-month-old female C57BL/6 J mice (Laboratory of Experimental Surgery) were housed in groups of six or seven per cage at 22 °C in a 12 h light/dark cycle and were given free access to diet and water. After 1 week acclimation, mice were randomly assigned to either a control group (Ctrl, $n=7$), a sedentary high-fat diet group (HFD, $n=7$) and a high-fat diet group that was simultaneously exercise trained (HFD+ex, $n=6$). The control group ate standard chow, while the two other groups received a diet containing 49.5 g fat (corn oil and lard)/100 g, 37 g protein (cow milk casein)/100 g and <1 g carbohydrate/100 g. This represents, in % total energy: 72 % fat, 28 % protein, and <1 % carbohydrate (UAR). This diet is known to induce a marked diabetic and metabolic stress state [2, 3]. Although a lack of carbohydrates per se has been shown to activate ER stress [13], we recently evidenced that a 45 % HFD containing 35 % carbohydrates induced the unfolded protein response in a way similar to the

144	present diet, indicating that a lack of carbohydrates is	vein blood samples were collected in EDTA tubes;	189
145	not the principal factor inducing the unfolded protein	after centrifugation (10 min at 1,500×g), plasma was	190
146	response in skeletal muscle [7]. All mice experiments	stored at −80 °C.	191
147	were approved by the local committee and the housing		
148	conditions were as specified by the Belgian Law of	Protein extraction, SDS/PAGE and immunoblotting	192
149	November 14, 1993 on the protection of laboratory		
150	animals (agreement no. LA 1220548).		
151	Exercise training protocol		
152	The results of the Ctrl and the HFD groups have been	Approximately 10–20 mg of frozen tissue were	193
153	partially published in a previous study [6], the purpose	ground in a mortar and homogenized in ice-cold buffer	194
154	of which was to evidence the presence of ER stress in	[20 mM Tris, pH 7.0, 270 mM sucrose, 5 mM EGTA,	195
155	skeletal muscle of sedentary high-fat-fed mice. The	1 mM EDTA, 1 % Triton X-100, 1 mM sodium	196
156	data of the HFD+ex group were acquired at the same	orthovanadate, 50 mM sodium β-glycerophosphate,	197
157	time as the Ctrl and HFD groups. In order to study the	5 mM sodium pyrophosphate, 50 mM sodium fluo-	198
158	effect of contractile activity, mice assigned to the HFD	ride, 1 mM DTT (1,4-dithiothreitol) and a protease	199
159	+ex group exercised on a treadmill 5 days per week	inhibitor cocktail containing 1 mM EDTA (Roche	200
160	for 6 weeks. Initially, mice were progressively accli-	Applied Science)] for 5 min on ice. The homogenates	201
161	ated to the treadmill during a period of 2 weeks from	were then centrifuged for 10 min at 10,000×g and the	202
162	7 m/min for 30–40 min to 12 m/min for 60 min. For	supernatants were immediately stored at −80 °C. Pro-	203
163	the remainder of the experiment, mice trained at a	tein concentration was determined using the DC pro-	204
164	treadmill speed of 12 m/min for 60 min. During the	tein assay kit (Bio-Rad Laboratories).	205
165	whole experiment, the slope of the treadmill was equal	Cell lysates (60 µg for skeletal muscle proteins and	206
166	to zero. According to pre-experiments on mice of	30 µg for pancreatic and hepatic proteins) were com-	207
167	exactly the same strain, same origin (Laboratory of	bined with Laemmli sample buffer and separated by	208
168	Experimental Surgery), same gender and same age,	SDS/PAGE. After electrophoretic separation at 40 mA	209
169	this speed corresponds to approximately 70 % of the	for 1 h, the proteins were transferred to a PVDF	210
170	maximal velocity of the mice. The maximal velocity	membrane at 80 V for 2 h for western blot analysis.	211
171	was determined thanks to an incremental exercise test	Membranes were then incubated in a 5 % Blotto	212
172	on a flat treadmill mill (no slope). The starting velocity	solution. Subsequently, membranes were incubated	213
173	was 8 m·min ^{−1} and was increased by 2 m·min ^{−1} every	with the following antibodies overnight at 4 °C: BiP,	214
174	2 min until exhaustion. The maximal velocity was	PDI (protein disulfide isomerase), MBTPS2 (mem-	215
175	defined as the velocity of the last stage completed by	brane-bound transcription factor protease site 2),	216
176	the animals.	IRE1α, PERK, phospho-PERK Thr 980, phospho-	217
177	Tissue and blood samples	p38 (p38 protein kinase) Thr 180/Tyr 182, phospho-	218
178	At the end of the 6 weeks and 48 h after the last	JNK Thr 183/Tyr 185, phospho-IKK (I-kappa-B ki-	219
179	exercise session, 6-h-fasted mice were terminally	nase) Ser 176/180, SDH (succinate dehydrogenase)	220
180	anaesthetized by intra-peritoneal injection of sodium	and GAPDH. All antibodies were from cell signalling	221
181	pentobarbital solution (using 60 mg/kg of body	except PERK and GAPDH from Abcam and SDH	222
182	weight, Nembutal®, Sanofi). Tibialis anterior and so-	from Santa Cruz.	223
183	leus muscles, liver, pancreas and visceral fat were	Membranes were washed in TBST and incubated	224
184	removed as fast as possible and immediately frozen	for 1 h at room temperature in a secondary antibody	225
185	in liquid nitrogen. The visceral fat corresponds to the	conjugated to horseradish peroxidase (1:10,000, Sig-	226
186	mesenteric fat present along the gastrointestinal tract.	ma). After an additional three washes, chemilumines-	227
187	This fat depot was separated from the pancreas and the	cence detection was carried out using an Enhanced	228
188	mesenteric ganglion was precisely dissected. Cava	Chemiluminescent Western blotting kit (ECL Plus,	229
		Amersham Biosciences). The films were then scanned	230
		on an ImageScanner using the Labscan software and	231
		quantified with the Image Master 1D Image Analysis	232
		Software (Amersham Biosciences). Results are	233
		reported relative to GAPDH. A value of 1 was arbi-	234
		trarily assigned to the control conditions which were	235

used as a reference for the high-fat and high-fat plus exercise values.

RNA extraction and quantitative real-time PCR

About 10–20 mg of frozen tissue samples were homogenized in TriPure reagent (Roche) using a Polytron. Total RNA was extracted according to the instructions provided by the manufacturer (Roche Diagnostics). RNA was quantified by spectrophotometry (260 nm) and its concentration adjusted to 1 µg/µl using RNase-free water. Since soleus muscles were very small (about 5–6 mg), the RNA concentrations obtained after extraction were too low to perform the reverse transcription reaction. For tibialis anterior and liver, cDNA was prepared by reverse transcription of 1 µg total RNA using the reverse transcription system (Promega). Real-time PCR was performed with a STEP one PLUS instrument and software (Applied Biosystems) using SYBR®Green PCR Master Mix (Applied Biosystems) for detection. Real-time PCR primers were designed (Table 1) for mouse CHOP, ATF4, spliced (s) XBP1, unspliced (u) XBP1, IL-1 (interleukin-1), IL-6 (interleukin-6), NADPHox (nicotinamide adenine dinucleotide phosphate oxidase) and RPL-19 (ribosomal protein L19). Specific primers were designed to recognize the spliced, or active, form of XBP1 (XBP1s) versus the unspliced form (XBP1u). RPL-19 was used as the “house keeping” gene. All tissues were run in duplicate in a single 96-well reaction plate (MicroAmp Optical, Applied Biosystems) and data were analysed according to the $2^{-\Delta CT}$ method. The identity and purity of the amplified product was checked through analysis of the melting curve carried out at the end of amplification. A value of 1 was arbitrarily assigned to the control condition to which the high-fat and the high-fat plus exercise values were reported.

Oral glucose tolerance test

An oral glucose tolerance test (OGTT) (gavage with 1 mg glucose/g body weight; 20 % glucose solution) was performed on 6-h-fasted mice at the end of the 6-week treatment and 24 h after the last training session. Blood glucose was determined with a glucose meter (Roche Diagnostics) on 3.5 µl of blood collected from the tip of the tail vein, 30 min before and 0, 15, 30, 60, 90 and 120 min following glucose injection.

Cytokines quantification

Cytokines were determined in 12 µl of plasma using a kit (Bio-Plex Multiplex; Bio-Rad) and measured using Luminex technology (Bio-Plex; Bio-Rad) as previously described [4]. Insulin was measured in 5 µl of plasma using an ELISA kit (Mercodia, Upssala, Sweden).

Statistical analysis

The difference between the three groups was tested for significance using a one-way analysis of variance. When significant, Student–Newman–Keuls post hoc tests were applied. The significance threshold was set to $P < 0.05$. Results are presented as means ± SEM.

Table 1 Primer sequences

	Forward	Reverse
CHOP	CCT AGC TTG GCT GAC AGA GG	CTG CTC CTT CTC CTT CAT GC
ATF4	GAG CTT CCT GAA CAG CGA AGT G	TGG CCA CCT CCA GAT AGT CAT C
XBP1u	TGA GAA CCA GGA GTT AAG AAC ACG C	TTC TGG GTA GAC CTC TGG GAG TTC C
XBP1s	GAG TCC GCA GCA GGT G	GTG TCA GAG TCC ATG GGA
IL-1	TCG CTC AGG GTC ACA AGA AA	CAT CAG AGG CAA GGA GGA AAA C
IL-6	ACA AGT CGG AGG CTT AAT TAC ACA T	TTG CCA TTG CAC AAC TCT TTT C
NADPHox	TTG GGT CAG CAC TGG CTC TG	TGG CGG TGT GCA GTG CTA TC
RPL-19	GAA GGT CAA AGG GAA TGT GTT CA	CCT TGT CTG CCT TCA GCT TGT

Sequences of primers used for mRNA quantification by real-time RT-PCR CHOP

C/EBP (CCAAT/enhancer binding protein) homologous protein, *ATF4* activating transcription factor 4, *XBP1u* unspliced X Box binding protein 1, *XBP1s* spliced X Box binding protein 1, *IL-1* interleukin-1, *IL-6* interleukin-6, *NADPHox* nicotinamide adenine dinucleotide phosphate oxidase, *RPL-19* ribosomal protein L19

Results

Endurance training reduces gain in body weight induced by high-fat diet

Throughout the 6 weeks of experiment, mice fed with a normal diet increased body weight by 20 % (Fig. 1a). At the end of the 6 weeks, HFD mice weighed 10 % more than control mice ($P<0.05$ at week 5 and $P=0.088$ at week 6). Running reduced the body weight gain induced by the HFD by about a half ($P<0.05$). In the same way, HFD increased visceral fat content by 50 % ($P<0.05$, Fig. 1b–c) compared to control mice whereas running while eating a HFD tended to reduce this gain.

Endurance training increases succinate dehydrogenase expression

The efficacy of the endurance training protocol was assessed by measuring SDH expression. The latter almost doubled in soleus ($P<0.001$) and tibialis

Fig. 2 Succinate dehydrogenase as a marker of endurance training. SDH expression in soleus, tibialis anterior and liver after 6 weeks HFD or HFD+ex. Results are expressed as the means±SEM. Ctrl, control; HFD, high-fat diet; HFD+ex, high-fat diet plus exercise. ** $P<0.01$, *** $P<0.001$ vs Ctrl; §§ $P<0.01$, §§§ $P<0.001$ vs HFD

anterior ($P<0.01$) muscles, as compared to untrained mice, whereas it was not modified in the liver (Fig. 2).

High-fat diet increases unfolded protein response in skeletal muscle

Several unfolded protein response markers were higher in skeletal muscles of HFD than in control mice (Fig. 3a and b and [6]). The changes were more pronounced in slow-type soleus muscle (Fig. 3a) than in fast-type tibialis anterior muscle (Fig. 3b). In the soleus, high-fat diet increased the protein expression of BiP ($P<0.05$) and MBTPS2 ($P<0.001$) eightfold as well as IRE1 α ($P<0.05$) and PERK ($P<0.05$) fivefold. The role of MBTPS2 is to cleave ATF6 into a functional transcription factor. In soleus, PERK was only detectable in its unactivated non-phosphorylated form. In tibialis anterior, HFD doubled BiP ($P<0.05$) and IRE1 α ($P<0.05$) levels and increased the expression of PERK 15-fold ($P<0.001$). Since PERK phosphorylation was unchanged by HFD, the ratio of phospho-PERK/PERK was decreased, suggesting that the relative activity of PERK was reduced in HFD (Fig. 3b). High-fat diet also induced the unfolded protein response in liver (Fig. 3c) and in pancreas (Fig. 3d). PDI, IRE1 α and MBTPS2 protein levels were more than doubled and PERK phosphorylation was increased fourfold in the liver ($P<0.05$). In the pancreas, IRE1 α expression was increased by threefold ($P<0.05$), MBTPS2 by twofold ($P<0.05$) and total PERK by sixfold ($P<0.01$). It is of note that the antibody against PDI used in this study resulted in a

Fig. 1 Body weight and visceral fat content. Mice were weighed each week (a). Visceral fat content was weighed at the end of the 6 weeks (b) and reported to body weight (c). Results are expressed as the means±SEM. Ctrl, control; HFD, high-fat diet; HFD+ex, high-fat diet plus exercise. * $P<0.05$ vs Ctrl; § $P<0.05$, §§ $P<0.01$ vs HFD

Fig. 3 Endoplasmic reticulum stress markers. Protein expression of BiP, PDI, IRE1 α , MBTPS2, PERK and phosphorylation state of PERK in soleus (a), tibialis anterior (b), liver (c) and pancreas (d) and mRNA level of CHOP, ATF4, XBP1s, XBP1u (e, f) and NADPHox (g, h) in tibialis anterior (e, g) and liver (f, h) after 6 weeks HFD or HFD+ex. Results are expressed as the means $\pm$ SEM. A value of 1 was arbitrarily assigned to the control conditions to which the HFD and HFD+ex values were reported and expressed as foldbasal. BiP binding protein, PDI protein disulfide isomerase, IRE1 inositol-requiring enzyme 1,

MBTPS2 membrane-bound transcription factor protease site 2, PERK protein kinase R-like ER protein kinase, CHOP C/EBP (CCAAT/enhancer binding protein) homologous protein, ATF4 activating transcription factor 4, XBP1u unspliced X Box binding protein 1, XBP1s, spliced X Box binding protein 1 NADPHox nicotinamide adenine dinucleotide phosphate oxidase. Ctrl control, HFD high-fat diet, HFD+ex high-fat diet plus exercise. * P <0.05, ** P <0.01, *** P <0.001 vs Ctrl; § P <0.05, §§ P <0.01, §§§ P <0.001 vs HFD

diffuse signal that was not quantifiable due to the two isoforms expressed only in the pancreas [8].

Since ER stress induces responses at both the translational and the transcriptional levels, we also analysed the mRNA levels of some well-documented ER stress markers: CHOP, ATF4, XBP1s and XBP1u (Fig. 3e and f) [11, 16]. Whereas the mRNA level of these markers was not affected by HFD in the tibialis anterior (Fig. 3e), CHOP was increased by 30 % (P <0.05) and XBP1u by 200 % (P <0.001) in the liver (Fig. 3f). NADPH oxidase is a major player in the production of reactive oxygen species which are known to trigger ER stress and to play a role in the pathogenesis of diabetes [14]. High-fat

diet doubled NADPHox mRNA level in both the tibialis anterior muscle (Fig. 3g) and the liver (Fig. 3h) (P <0.05).

Compared to the effects of HFD alone, endurance training did not induce many profound changes in ER stress markers. Endurance training increased the expression of a few markers more than HFD alone, as observed for PERK in the soleus (P <0.05, Fig. 3a) and in the liver (P <0.01, Fig. 3c), for BiP in the tibialis anterior (P <0.05, Fig. 3b) and for MBTPS2 in the pancreas (P <0.01, Fig. 3d). Interestingly, endurance training reversed the increase in NADPHox mRNA level induced by a HFD in the tibialis anterior (P <0.05, Fig. 3g).

Endurance exercising while fed with a high-fat diet increases caspase 12 expression and cleavage

High-fat diet alone had no effect on caspase 12 expression or cleavage in any tissue studied. The cleavage of caspase 12 however was increased in HFD+ex condition in the soleus ($P<0.05$, Fig. 4a), tibialis anterior ($P<0.001$, Fig. 4b), liver ($P<0.01$, Fig. 4c) and pancreas ($P<0.05$, Fig. 4d). It is of note that cleaved caspase 12 was not detectable in control conditions in the tibialis anterior (Fig. 4b) and in the liver (Fig. 4c) whereas total caspase 12 was observed. Endurance exercising while fed with a HFD also increased the total form of caspase 12 compared to control or HFD alone in the soleus ($P<0.05$, Fig. 4a) and liver ($P<0.01$, Fig. 4c).

Regulation of inflammation markers is tissue specific

JNK and IKK/NF κ B have been implicated in insulin resistance and inflammation induced by ER

stress through the IRE1 α pathway [12, 14]. In our hands, these two pathways seem to be regulated independently since phospho-IKK was increased by HFD in the soleus ($P<0.05$, Fig. 5a) whereas in the liver ($P<0.05$, Fig. 5c) and in the pancreas ($P<0.001$, Fig. 5d), HFD increased phospho-JNK without affecting phospho-IKK. Phospho-p38 was only enhanced in the tibialis anterior ($P<0.05$, Fig. 5b). All these results indicate that the pathways

Fig. 4 Apoptosis marker. Caspase 12 expression and cleavage in soleus (a), tibialis anterior (b), liver (c) and pancreas (d) after 6 weeks HFD or HFD+ex. Results are expressed as the means $\pm$ SEM. A value of 1 was arbitrarily assigned to the control conditions to which the HFD and HFD+ex values were reported and expressed as foldbasal. Ctrl control, HFD high-fat diet, HFD+ex high-fat diet plus exercise. * $P<0.05$, ** $P<0.01$, *** $P<0.001$ vs Ctrl; § $P<0.05$, §§ $P<0.01$, §§§ $P<0.001$ vs HFD

Fig. 5 Inflammation markers. Phosphorylation state of JNK, IKK and p38 in soleus (a), tibialis anterior (b), liver (c) and pancreas (d) and mRNA level of IL-1 and IL-6 in tibialis anterior (e) and liver (f) after 6 weeks HFD or HFD+ex. Results are expressed as the means $\pm$ SEM. A value of 1 was arbitrarily assigned to the control conditions to which the HFD and HFD+ex values were reported and expressed as foldbasal. JNK Jun N-terminal kinase, IKK I-kappa-B kinase, p38 p38 protein kinase, IL-1 interleukin-1, IL-6 interleukin-6. Ctrl control, HFD high-fat diet, HFD+ex high-fat diet plus exercise. * $P<0.05$, ** $P<0.01$, *** $P<0.001$ vs Ctrl; § $P<0.05$, §§ $P<0.01$, §§§ $P<0.001$ vs HFD

known to be switched on by inflammation were regulated in a tissue-specific manner by HFD feeding and that exercise did not significantly reverse this effect except in the pancreas, in which phospho-JNK mice reached a similar level in control and HFD+ex mice.

To evaluate the further implication of this inflammation-induced signalling by HFD, we measured the mRNA level of IL-1 and IL-6 in the tibialis anterior (Fig. 5e) and liver (Fig. 5f) as well as the plasma level of several pro- and anti-inflammatory cytokines (Table 2). IL-1 is a master mediator and initiator of inflammation in several tissues [24] and its mRNA expression has recently been shown to be also regulated in mouse skeletal muscle [20]. IL-1 mRNA level was increased ($P<0.001$, Fig. 5e) and IL-6 decreased ($P<0.05$, Fig. 5e) by HFD in the tibialis anterior and were unchanged in the liver (Fig. 5f). Exercise reversed the HFD-induced increase in IL-1 ($P<0.001$, Fig. 5e) but not the decrease in IL-6 mRNA in the tibialis anterior. The effects of exercise were more pronounced at the plasma level where it reduced the high-fat diet increase in IL-6 and MIP-1 α concentrations ($P<0.05$, Table 2). Endurance exercising while high-fat fed also decreased the levels of circulating IL-15 compared to sedentary normally fed mice and MCP-1 compared to sedentary

normally fed and high-fat-fed mice ($P<0.05$, Table 2).

High-fat diet-induced glucose intolerance is not reversed by endurance training

Mice fed a HFD were glucose intolerant as indicated by the OGTT performed at the end of the 6 weeks (Fig. 6a). For the same amount of glucose given orally, plasma glucose concentrations remained elevated up to 90 min after gavage ($P<0.05$) whereas plasma insulin concentrations were not different between groups (Fig. 6a). The area under the curve was increased by 1.8-fold by the HFD ($P<0.05$, Fig. 6b) and was not reduced by exercise. The OGTT indicates that low intensity running while eating a HFD did not reverse the glucose intolerant state induced by HFD alone.

Table 2 The cytokine concentrations are expressed in pg/ml

	Ctrl	HFD	HFD+ex
IL-1 α	34 $\pm$ 11.3	20 $\pm$ 3.5	17 $\pm$ 2.7
IL-1 β	26 $\pm$ 6.2	18 $\pm$ 2.5	32 $\pm$ 11.3
IL-6	22 $\pm$ 3.3	56 $\pm$ 11.3*	35 $\pm$ 6.1**
IL-10	43 $\pm$ 14.0	112 $\pm$ 24.3	232 $\pm$ 122.5
IL-15	306 $\pm$ 27.2	257 $\pm$ 17.8	210 $\pm$ 16.5*
IL-18	440 $\pm$ 107.7	593 $\pm$ 189.8	228 $\pm$ 52.0
MCP-1	107 $\pm$ 13.1	120 $\pm$ 12.4	72 $\pm$ 7.5***
MIP-1 α	139 $\pm$ 10.6	178 $\pm$ 18.7*	129 $\pm$ 12.6**
TNF- α	26 $\pm$ 6.4	14 $\pm$ 1.2	16 $\pm$ 3.6

Values are expressed as the means $\pm$ SEM

IL-1 α interleukin-1 alpha, IL-1 β interleukin-1 beta, IL-6 interleukin-6, IL-10 interleukin-10, IL-15 interleukin-15, IL-18 interleukin-18, MCP-1 monocyte chemoattractant protein-1, MIP-1 α macrophage inflammatory protein-1 alpha, TNF α tumor necrosis factor alpha, Ctrl control, HFD high-fat diet, HFD+ex high-fat diet plus exercise

* $P<0.05$ vs Ctrl; ** $P<0.05$ vs HFD

Fig. 6 OGTT test. An oral glucose tolerance test (OGTT) was performed on 6-h-fasted mice at the end of the 6-weeks treatment and 24 h after the last training session. (a) Plasma glucose and insulin concentrations after the oral glucose load. (b) Area under curve (AUC) of the glucose excursion after the oral glucose load. Results are expressed as the means $\pm$ SEM. Ctrl control, HFD high-fat diet, HFD+ex high-fat diet plus exercise. * $P<0.05$, ** $P<0.01$ vs Ctrl

Discussion

The main findings of the present study are that: (1) endurance training attenuated the inflammatory state induced by a HFD in mice without reversing whole body glucose intolerance; (2) endurance training increased the unfolded protein response induced by a HFD in mice.

The present results are different from those obtained in a recently published study showing that swimming reduced ER stress in adipose and hepatic tissues of diet-induced obese rats [5]. One major difference of the later study with the current study is that rats were high-fat-fed for 2 months before starting exercise training. One could postulate that rats were accustomed to the HFD and that a new homeostasis level was reached before the stress of physical activity was added, so that the beneficial effects of exercise could be exerted. This was not the case in the present study as both HFD and treadmill exercise began at the same time, each stress potentially exacerbating the other. It is also possible that the lipid percentage or nature of the diet and/or the intensity or nature of exercise could explain the opposite results. These hypotheses are further discussed below.

Our data show that several markers of the unfolded protein response were up-regulated following 6 weeks on a high-fat diet and that the effect was greatest in the postural muscles. In the tonically active soleus muscle, BiP, IRE1 α , MBTPS2 and PERK protein levels were increased more than fivefold. In the more phasically active tibialis anterior muscle, these changes were less pronounced. The difference could be due to greater susceptibility of oxidative fibers to a high-fat diet, an additive effect of contractile activity and high-fat diet on ER stress, or increased uptake of fatty acids in the soleus due to its higher metabolic requirements.

When the unfolded protein response is not sufficient to cope with ER stress, inflammation and insulin resistance may develop. The inflammation and insulin resistance may directly result from an activation of IRE1 α during the unfolded protein response, since IRE1 α , in turn, triggers the JNK and IKK/NF κ B pathways [12, 31]. Indeed, high-fat feeding activated IRE1 α as well as JNK and IKK, a kinase upstream of NF κ B. However, in no tissue was a simultaneous activation of JNK and IKK observed. High-fat diet also increased plasma levels of cytokines such as IL-6 and MIP-1 α and this was associated with whole

body glucose intolerance as demonstrated by the OGTT. Taken together, these data confirm that mild chronic inflammatory state and glucose intolerance were induced by our high-fat diet.

After having shown that a high-fat diet induced the unfolded protein response, a mild inflammatory state and glucose intolerance, we tested whether contractile activity in the form of endurance exercise could reduce these responses. The intensity of the exercise (12 m/min) corresponded to ~70 % of the average maximal running speed of untrained mice, as measured in preliminary experiments, or to ~75 % of VO₂max [28]. We chose this intensity as it has been shown to enhance lipid oxidation in mice skeletal muscle [9, 25]. Hence, we expected this form of exercise to reduce the deleterious effects of a high-fat diet by oxidizing more of the consumed fatty acids. The increase in muscle SDH expression observed after 6 weeks in the exercised mice indicates that the training was efficient in increasing aerobic metabolism. At the end of the training period, we observed an increase in markers of the unfolded protein response, some of them being even more elevated than in the HFD group without exercise. The increase in the unfolded protein response indicates a greater degree of ER stress with concomitant high-fat diet and exercise. Since exercise at 75 % of VO₂max increases the mobilization of fatty acids, exercising while consuming a high-fat diet could potentially result in a greater increase in fatty acid uptake [26] and therefore a lipid-dependent ER stress. Although not mentioned, it is likely that the intensity of swimming was less than running and that mobilization of fatty acids was less in the study of da Luz [5], potentially partially explaining the fact that exercise was able to reduce ER stress in obese rats.

Since exercise is known to improve glucose tolerance, we hypothesized that if ER stress is implicated in the development of glucose intolerance, exercise training would simultaneously reduce ER stress and glucose intolerance. Six weeks of treadmill running did reduce the plasma levels of several pro-inflammatory cytokines (IL-6, MCP-1 and MIP-1 α). However, the exercise protocol used in the current study was not effective in reducing either ER stress or whole body glucose intolerance induced by a high-fat diet. It is possible that, in the present study, the intensity and/or the duration of the exercise sessions were not sufficient to counteract glucose intolerance caused by a high-fat diet. It has been reported that treadmill

exercise had only a mild effect on high-fat diet-induced glucose intolerance when training sessions were performed for 1 h during 6–8 weeks [10, 32] or for 3 h during 3 weeks [29]. However, when animals have had free access to a wheel, and exercise for approximately five- to tenfold longer periods of time, insulin resistance induced by high-fat feeding had been shown to be improved [1, 19]. Another possibility is that the fat content of the diet was too high for allowing beneficial effect of exercise on whole body glucose tolerance to be observed. Yet, this percentage of fat is seen in diets used in human during carbohydrates free regimen. In the current study, 70 % of the energy in the high-fat diet was derived from fat, whereas studies showing a positive effect of exercise on glucose tolerance, either due to wheel running, treadmill running, or swimming, used diets containing about 40–45 % fat [1, 10] or used diets containing the same percentage of fat (60–70 %) for a shorter period of time (3 weeks) [15, 19]. It is also possible that the fatty acid composition of the diet, i.e. highly saturated, could have contributed to the lack of effect of the exercise training.

We recognize that a limitation of the present study is the lack of an exercised group not receiving a high-fat diet. This would have allow to discriminating the effect of exercise per se as high-fat feeding and a lack of carbohydrates are known to induce metabolic adaptations and to influence exercise performance [23]. However, the purpose was to determine molecular mechanisms by which exercise exerts beneficial effects in high-fat-fed and obese animals and our results allow to answer this question. Our results support the idea that exacerbation of the high-fat diet-induced unfolded protein response by endurance exercise might improve ER homeostasis and consequently might protect against inflammation and extend previous similar results in lean mice [34] and human [17]. A single moderate-intensity exercise bout activated the unfolded protein response in skeletal muscle of lean mice, whereas the activation was less or even repressed for some markers after several training sessions. The authors concluded that moderate exercise and the accompanying physiological ER stress in skeletal muscle may lead to adaptation and protect skeletal muscle against further stress [34]. The same potential mechanism of defence has been observed after a 200-km run in human skeletal muscle [17].

In conclusion, we have shown that endurance training attenuated the inflammatory state induced by a

HFD in mice without reversing whole body glucose intolerance and that endurance training increased the unfolded protein response induced by a HFD in mice. We speculate that the potentiation of unfolded protein response by endurance training may represent a positive adaptation protecting against further cellular stress. Different exercise protocols, in terms of frequency, intensity and duration should be tested to identify the effect of exercise per se on ER stress markers in skeletal muscle.

Acknowledgements The authors would like to thank Damien Naslain, Delphine Robin, Gang-Li An and Vincent d'Harveng for their technical assistance.

Grants This work was supported by the Fonds National de la Recherche Scientifique (Belgium), by the Université catholique de Louvain (FSR) and by the Chaire Delhaize (Belgium). At the time of data collection, L. Deldicque was a postdoctoral fellow from the FRS-FNRS (Fonds de la Recherche Scientifique), Belgium. P. Cani is Research Associate from the FRS-FNRS, Belgium.

Disclosure The authors have nothing to disclose.

References

- Bradley RL, Jeon JY, Liu FF, Maratos-Flier E (2008) Voluntary exercise improves insulin sensitivity and adipose tissue inflammation in diet-induced obese mice. *Am J Physiol Endocrinol Metab* 295:586–594
- Cani PD, Amar J, Iglesias MA, Poggi M, Knauf C, Bastelica D, Neyrinck AM, Fava F, Tuohy KM, Chabo C, Waget A, Delmee E, Cousin B, Sulpice T, Chamontin B, Ferrieres J, Tanti JF, Gibson GR, Casteilla L, Delzenne NM, Alessi MC, Burcelin R (2007) Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes* 56:1761–1772
- Cani PD, Knauf C, Iglesias MA, Drucker DJ, Delzenne NM, Burcelin R (2006) Improvement of glucose tolerance and hepatic insulin sensitivity by oligofructose requires a functional glucagon-like peptide 1 receptor. *Diabetes* 55:1484–1490
- Cani PD, Possemiers S, Van de Wiele T, Guiot Y, Everard A, Rottier O, Geurts L, Naslain D, Neyrinck A, Lambert DM, Muccioli GG, Delzenne NM (2009) Changes in gut microbiota control inflammation in obese mice through a mechanism involving GLP-2-driven improvement of gut permeability. *Gut* 58:1091–1103
- da Luz G, Frederico MJ, da SS, Vitto MF, Cesconetto PA, de Pinho RA, Pauli JR, Silva AS, Cintra DE, Ropelle ER, De Souza CT (2011) Endurance exercise training ameliorates insulin resistance and reticulum stress in adipose and hepatic tissue in obese rats. *Eur J Appl Physiol*
- Deldicque L, Cani PD, Philp A, Raymackers JM, Meakin PJ, Ashford ML, Delzenne NM, Francaux M, Baar K

- 634 (2010) The unfolded protein response is activated in skeletal
635 muscle by high-fat feeding: potential role in the downregu-
636 lation of protein synthesis. *Am J Physiol Endocrinol Metab*
637 299:E695–E705
- 638 7. Deldicque L, Van Proeyen K, Francaux M, Hespel P (2010)
639 The unfolded protein response in human skeletal muscle is
640 not involved in the onset of glucose tolerance impairment
641 induced by a fat-rich diet. *Eur J Appl Physiol*
- 642 8. Desilva MG, Notkins AL, Lan MS (1997) Molecular char-
643 acterization of a pancreas-specific protein disulfide isomerase,
644 PDip. *DNA Cell Biol* 16:269–274
- 645 9. Dzamko N, Schertzer JD, Ryall JG, Steel R, Macaulay SL,
646 Wee S, Chen ZP, Michell BJ, Oakhill JS, Watt MJ, Jorgensen
647 SB, Lynch GS, Kemp BE, Steinberg GR (2008) AMPK-
648 independent pathways regulate skeletal muscle fatty acid ox-
649 idation. *J Physiol* 586:5819–5831
- 650 10. Gauthier MS, Couturier K, Latour JG, Lavoie JM (2003)
651 Concurrent exercise prevents high-fat-diet-induced macro-
652 vesicular hepatic steatosis. *J Appl Physiol* 94:2127–2134
- 653 11. Gregor MF, Hotamisligil GS (2007) Thematic review series:
654 adipocyte biology. Adipocyte stress: the endoplasmic reticu-
655 lum and metabolic disease. *J Lipid Res* 48:1905–1914
- 656 12. Hu P, Han Z, Couvillon AD, Kaufman RJ, Exton JH (2006)
657 Autocrine tumor necrosis factor alpha links endoplasmic reticu-
658 lum stress to the membrane death receptor pathway through
659 IRE1alpha-mediated NF-kappaB activation and down-
660 regulation of TRAF2 expression. *Mol Cell Biol* 26:3071–3084
- 661 13. Ikesugi K, Mulhern ML, Madson CJ, Hosoya K, Terasaki T,
662 Kador PF, Shinohara T (2006) Induction of endoplasmic
663 reticulum stress in retinal pericytes by glucose deprivation.
664 *Curr Eye Res* 31:947–953
- 665 14. Kaneto H, Matsuoka TA, Nakatani Y, Kawamori D, Miyatsuka
666 T, Matsuhisa M, Yamasaki Y (2005) Oxidative stress, ER
667 stress, and the JNK pathway in type 2 diabetes. *J Mol Med*
668 83:429–439
- 669 15. Kim CH, Youn JH, Park JY, Hong SK, Park KS, Park SW,
670 Suh KI, Lee KU (2000) Effects of high-fat diet and exercise
671 training on intracellular glucose metabolism in rats. *Am J*
672 *Physiol Endocrinol Metab* 278:977–984
- 673 16. Kim DS, Jeong SK, Kim HR, Chae SW, Chae HJ (2007)
674 Effects of triglyceride on ER stress and insulin resistance.
675 *Biochem Biophys Res Commun* 363:140–145
- 676 17. Kim HJ, Jamart C, Deldicque L, An GL, Lee YH, Kim CK,
677 Raymackers JM, Francaux M (2011) Endoplasmic reticulum
678 stress markers and ubiquitin-proteasome pathway activity in
679 response to a 200-km run. *Med Sci Sports Exerc* 43:18–25
- 680 18. Kitamura M (2011) Control of NF-kappaB and inflammation
681 by the unfolded protein response. *Int Rev Immunol* 30:4–15
- 682 19. Kraegen EW, Storlien LH, Jenkins AB, James DE (1989)
683 Chronic exercise compensates for insulin resistance induced
684 by a high-fat diet in rats. *Am J Physiol* 256:242–249
- 685 20. Lang CH, Silvix C, Deshpande N, Nystrom G, Frost RA
686 (2003) Endotoxin stimulates in vivo expression of inflam-
687 matory cytokines tumor necrosis factor alpha, interleukin-
688 1beta, -6, and high-mobility-group protein-1 in skeletal
689 muscle. *Shock* 19:538–546
- 690 21. Lipson KL, Fonseca SG, Urano F (2006) Endoplasmic
691 reticulum stress-induced apoptosis and auto-immunity in
692 diabetes. *Curr Mol Med* 6:71–77
- 693 22. Malhotra JD, Kaufman RJ (2007) The endoplasmic reticu-
694 lum and the unfolded protein response. *Semin Cell Dev Biol*
695 18:716–731
- 696 23. Miller WC, Bryce GR, Conlee RK (1984) Adaptations to a
697 high-fat diet that increase exercise endurance in male rats. *J*
698 *Appl Physiol* 56:78–83
- 699 24. Mills KH, Dunne A (2009) Immune modulation: IL-1,
700 master mediator or initiator of inflammation. *Nat Med*
701 15:1363–1364
- 702 25. Miura S, Kai Y, Kamei Y, Bruce CR, Kubota N, Febbraio
703 MA, Kadowaki T, Ezaki O (2009) Alpha2-AMPK activity
704 is not essential for an increase in fatty acid oxidation during
705 low-intensity exercise. *Am J Physiol Endocrinol Metab*
706 296:E47–E55
- 707 26. Romijn JA, Coyle EF, Sidossis LS, Zhang XJ, Wolfe RR
708 (1995) Relationship between fatty acid delivery and fatty
709 acid oxidation during strenuous exercise. *J Appl Physiol*
710 79:1939–1945
- 711 27. Ron D, Walter P (2007) Signal integration in the endoplas-
712 mic reticulum unfolded protein response. *Nat Rev Mol Cell*
713 *Biol* 8:519–529
- 714 28. Schefer V, Talan MI (1996) Oxygen consumption in adult
715 and AGED C57BL/6J mice during acute treadmill exercise
716 of different intensity. *Exp Gerontol* 31:387–392
- 717 29. Straczkowski M, Kowalska I, Dzieńis-Straczkowska S,
718 Kinalski M, Gorski J, Kinalska I (2001) The effect of
719 exercise training on glucose tolerance and skeletal muscle
720 triacylglycerol content in rats fed with a high-fat diet. *Diab-*
721 *etes Metab* 27:19–23
- 722 30. Szegezdi E, Fitzgerald U, Samali A (2003) Caspase-12 and
723 ER-stress-mediated apoptosis: the story so far. *Ann N Y*
724 *Acad Sci* 1010:186–194
- 725 31. Urano F, Wang X, Bertolotti A, Zhang Y, Chung P, Harding
726 HP, Ron D (2000) Coupling of stress in the ER to activation
727 of JNK protein kinases by transmembrane protein kinase
728 IRE1. *Science* 287:664–666
- 729 32. Vieira VJ, Valentine RJ, Wilund KR, Antao N, Baynard T,
730 Woods JA (2009) Effects of exercise and low-fat diet on
731 adipose tissue inflammation and metabolic complications in
732 obese mice. *Am J Physiol Endocrinol Metab* 296:E1164–
733 E1171
- 734 33. Wu J, Kaufman RJ (2006) From acute ER stress to physi-
735 ological roles of the unfolded protein response. *Cell Death*
736 *Differ* 13:374–384
- 737 34. Wu J, Ruas JL, Estall JL, Rasbach KA, Choi JH, Ye L,
738 Bostrom P, Tyra HM, Crawford RW, Campbell KP, Rut-
739 kowski DT, Kaufman RJ, Spiegelman BM (2011) The un-
740 folded protein response mediates adaptation to exercise in
741 skeletal muscle through a PGC-1alpha/ATF6alpha complex.
742 *Cell Metab* 13:160–169
- 743 35. Yoneda T, Imaizumi K, Oono K, Yui D, Gomi F, Katayama
744 T, Tohyama M (2001) Activation of caspase-12, an endo-
745 plastic reticulum (ER) resident caspase, through tumor ne-
746 crosis factor receptor-associated factor 2-dependent
747 mechanism in response to the ER stress. *J Biol Chem*
748 276:13935–13940
- 749 36. Zhang K, Kaufman RJ (2006) Protein folding in the endo-
750 plasmic reticulum and the unfolded protein response. *Handb*
751 *Exp Pharmacol* 69:91

AUTHOR QUERY

AUTHOR PLEASE ANSWER QUERY.

No Query.

UNCORRECTED PROOF