

TRPV1 channel as new target for marine toxins: example of gigantoxin I, a sea anemone toxin acting via modulation of the PLA2 pathway

Eva Cuypers¹, Steve Peigneur¹, Sarah Debaveye¹, Kazuo Shiomi² and Jan Tytgat^{1*}

¹ Laboratory of toxicology, Campus Gasthuisberg, Onderwijs en navorsing 2, Herestraat 49, P.O. 922, B-3000 Leuven, Belgium

² Department of Food Science and Technology, Tokyo University of Marine Science and Technology, Konan-4, Minato-ku, Tokyo 108-8477, Japan

*** Corresponding author**

Jan Tytgat
Laboratory of toxicology
Campus Gasthuisberg
Onderwijs en navorsing 2
Herestraat 49, PO Box 922
B-3000 Leuven
Belgium
Phone: +32 16 32 34 03
Fax: +32 16 32 34 05
jan.tytgat@pharm.kuleuven.be
www.toxicology.be

Abstract

Gigantoxin I, isolated from sea anemone *Stichodactyla gigantea*, was previously described as the first epidermal growth factor (EGF)-like toxin from natural origin. In this study, we discovered the interaction between the transient receptor potential vanilloid subtype I (TRPV1) channels and gigantoxin I. The TRPV1 channel is a non-selective cation channel involved in pain sensation and is described as pharmacological target of cnidaria venom. Our results highlight the involvement of the epidermal growth factor receptor/phospholipaseA2/arachidonic acid/lipoxygenase (EGFR/PLA₂/AA/LOX) pathway in the indirect activation of TRPV1 channels by gigantoxin I. This is the first time that this pathway is described in the indirect activation of TRPV1 channels by toxins. This knowledge not only gives insights into the possible induced effects by this new group of toxins, but also leads to a better understanding of the regulatory mechanism of TRPV1 channels themselves.

Keywords

Capsaicin receptor, Phospholipase A2, Epidermal growth factor, Lipoxygenase, Arachidonic acid

Introduction

The phylum of Cnidaria represents exclusively aquatic animals such as sea anemones and jellyfish. With more than 7600 species, the Cnidaria is one of the largest phyla of marine animals. According to their morphological characters, the cnidarians are divided into 4 classes: the Anthozoa (meaning “flower animals”) including corals and sea anemones, the Hydrozoa (meaning “water animals”) which include for example the Portuguese man-of-war and fire corals, the Cubozoa (“cube animals”) which include the box jellyfish and the Scyphozoa (meaning “bowl animals”) which include the jellyfish. The name Cnidaria comes from the Greek word “cnidos” which means stinging nettle. Despite the fact that they are incredibly diverse in form, they are united because of their stinging cells called nematocysts. These stinging cells are used to capture and subdue the prey. The nematocyst is a coiled thread-like stinger. When the nematocyst is called upon to fire, the thread is uncoiled, and springs straight. The harpoon-like thread punctures through the cnidocyte wall into the prey. Most cnidarians have a toxin in their stinger that helps to immobilize the prey. The nematocyst is fired either by the tentacle touching prey or in some cases by a nerve impulse from the animal telling it to fire ¹. Most cnidarian nematocysts are not harmful to humans, as the stinger cannot penetrate sufficiently into human skin to inflict any harm. There are some jellyfish, however, which can deliver extremely painful and, in a few cases, even fatal stings to humans.

Sea anemones are a rich source of protein and peptide toxins which are generally divided into three classes: hemolysins (15-21 kDa) ²⁻⁵, sodium channel toxins (3-5 kDa) ⁶⁻⁹ and potassium channel toxins (3.5-6.5 kDa) ¹⁰⁻¹⁴. However, Shiomi and Honma isolated gigantoxin I, a peptide toxin from *Stichodactyla gigantea* that has 33% homologies with mammalian epidermal growth factor (EGF) ¹⁵ (Figure 1). In accordance with these sequence homologies, including the location of the six cysteine residues, this toxin exhibits EGF activity as rounding of human epidermoid carcinoma A431 cells. Herein, a redistribution of PLC- γ 1 to the plasma membrane area, in particular newly formed ruffles accompanied by increased actin polymerization, was observed ¹⁶. Also tyrosine phosphorylation of the EGF receptor was described ¹⁵. Interestingly, gigantoxin I is weakly lethal but potently paralytic to crabs. In spite of these known biological activities, the pharmacological target(s) of this toxin remain(s) unclear. In this study, we investigated the effect of gigantoxin I on the Transient Receptor Potential Vanilloid subtype 1 (TRPV1) channel that was previously described as pharmacological target of cnidaria (*e.g.* sea anemone) venoms and polycyclic ether toxins ¹⁷⁻

¹⁸. Only recently, the first sea anemone peptide, APHC1, isolated from *Heteractis crispata*, with analgesic and antagonistic activity on TRPV1 channels has been reported ¹⁹. Unfortunately, no molecular action working mechanism has been put forward and it is currently not known if APHC1 interacts directly or indirectly with TRPV1 channels. It is only very recently that the first action mechanism of a toxin activating TRPV1 channels was revealed, showing that the outer pore domain is the targeting mechanism of a bivalent tarantula toxin ²⁰.

Transient receptor potential (TRP) channels constitute a superfamily of cation permeable channels. The founding member of this family was identified as a *Drosophila* gene product required for visual transduction ²¹. Spontaneous mutations in the TRP-gene lead to TRP-deficient flies which are blinded by intense light ²². This is caused by a disruption of subsequent Ca²⁺-dependent adaptation and thus leads to a sustained Ca²⁺ entry via TRP ion channels. The name “transient receptor potential” is based on the transient (normal) rather than sustained (mutation) response to light of the flies carrying a mutant in the trp locus. Further studies on the *Drosophila melanogaster* phototransduction indicated that TRP is the target of a phosphoinositide cascade. This leads to the suggestion that phototransduction in *Drosophila* might be analogous to the general and widespread process of phosphoinositide-mediated Ca²⁺ influx in other cells ²³.

The TRPV1 channel, a ligand-gated and non-selective cation channel that is expressed in peripheral sensory neurons, is one of the key players in peripheral pain sensation (for an overview of its function and expression profile, refer to Table 1). TRPV1 is heat and acid sensitive and can be activated by pungent substances like capsaicin. Although capsaicin had been already known since ancient times to have an analgesic effect, the channel was only discovered in 1997 as the physiological target of capsaicin. Despite the fact that numerous analgesics are already on the market, pain treatment remains one of the major problems in the medical world. Therefore, the search for new and more effective analgesics is still going on. The discovery of TRPV1 channels started a new chapter in the research to novel generation analgesics. In order to prevent side effects, it is very important to know the mechanisms and pathways involved in TRPV1 channel functioning. Therefore, basic pharmacological research is indispensable.

Members of the TRP channel superfamily share the common structural features of six transmembrane segments (TM) which cross the cell membrane, a pore forming hydrophobic

loop between the fifth and the sixth segment and both cytosolic carboxy- and amino-termini (Figure 2). Although this basic topology is common to many ion channel families, such as voltage-gated potassium channels, TRP and voltage-gated potassium channels show relatively little sequence homology. A likely common feature between voltage-gated potassium channels and TRP channels is their tetrameric structure. Recently, the 19 Å structure of TRPV1 channels was partly revealed by using electrocryomicroscopy²⁴. It exhibits fourfold symmetry and comprises two distinct regions: a large open basket-like domain, likely corresponding to the cytoplasmic N- and C-terminal portions, and a more compact domain, corresponding to the transmembrane portion. Remarkably, a similar two-domain arrangement referred to as a “hanging gondola” has been observed in 3D structures of K_v1.2 channel. In general, this structure follows the basic structural organisation described above.

The TRPV1 channel is known as the heat transducer in sensory neurons because the channel is activated at a temperature above 42°C. Well-known direct activators are capsaicin, the pungent substance found in hot chilli peppers, and protons. *In vivo* and *in vitro* studies reveal that inflammatory mediators like bradykinin, prostaglandine E₂ (PGE₂), extracellular ATP, glutamate and nerve growth factor can indirectly sensitize TRPV1 channels²⁵⁻²⁷. This is possible via various mechanisms such as increasing TRPV1 channel expression levels in the membrane²⁸⁻²⁹, inducing TRPV1 channel phosphorylation by protein kinases³⁰⁻³¹ or releasing channel inhibition by phosphatidylinositol 4,5-biphosphate (PIP₂) and thus sensitizing the channel for agonist stimulation³⁰⁻³². In addition, these inflammatory mediators act on G-protein coupled receptors (GPCRs) or, in case of nerve growth factor, on tyrosine kinase pathways. These can induce on their turn, via activation of phospholipase C (PLC) and/or phospholipase A₂ (PLA₂), the release of arachidonic acid (AA) and lipoxygenase products of arachidonic acid (*e.g.* hydroperoxyeicosatetraenoic acid or HPETE), substances which are known as TRPV1 channel agonists³³⁻³⁴. Finally, the endogenous ligand anandamide has also been shown to activate TRPV1 channels³⁵.

In our work presented here, we have set out to elucidate the pharmacological mechanism of the molecular interaction between the sea anemone peptide gigantoxin I and TRPV1 channels, and have investigated several regulating and activating pathways of TRPV1 channels. It is concluded that the interaction between the toxin and TRPV1 channels is PLA₂ pathway dependent.

Materials and Methods

Isolation of gigantoxin I and determination of amino acid sequence

The isolation and amino acid determination of gigantoxin I was described previously by Shiomi and Honma¹⁵.

Molecular biology

cRNA transcripts were synthesized from XbaI-linearized VR1 cDNA templates using T7 RNA polymerase (Ambion).

Mutant S502A/S800A was made using Quick Change II site-directed mutagenesis kit (Stratagene). In a first step, mutant S502A was made using primers 5'-GGGGACTATTTCCGAGGCACCGGAGAGATCTTGTC-3' and 5'-GACAAGATCTCTCCGGTGCCTCGGAAATAGTCCCC-3'. Cycling parameters were set according to the manufacturer's guidelines. The mutated cDNA was transformed into XL-1 blue cells and cDNA was isolated using Wizard Miniprep kit (Promega). The construct was checked by cDNA sequencing. In a second step, mutant S800A was made starting from the S502A mutant cDNA. The following primers were used; 5'-CGTCAAGCGCACCCCTGGCCTTCTCCCTGAGGTCAG-3' and 5'-CTGACCTCAGGGAGAAGGCCAGGGTGCCTTGACG-3'. RNA was synthesized from the linearized plasmid using T7 mMessage mMachine transcription kit. A second cDNA sequencing confirmed the sequence of the full S502A/S800A construct.

Concentration of cRNA was determined by measuring the absorbance at 230, 260 and 280 nm.

Expression in Xenopus oocytes and electrophysiology

Oocytes were harvested from anaesthetized female *Xenopus laevis* frogs as previously described³⁶ and were injected with 0.5-5 ng TRPV1 cRNA. Oocytes were incubated in ND96 solution supplemented with 50 mg/L gentamycin. Two to seven days after injection, two-electrode voltage-clamp recording was performed. Currents were measured in ND96 solution at a holding potential of -90 mV during 400 s. The recording chamber was perfused at a rate of 2 mL min⁻¹ with a ND-96 solution containing (in mM) 96 NaCl, 2 KCl, 1.8 CaCl₂, 1 MgCl₂, 5 HEPES (pH 7.4). Temperature of the perfusate was kept at 22°C and controlled using a SC-20 dual in-line heater/cooler (Warner Instruments) and pH was kept at 7.4. As previously described³⁷, capsaicin (2 µM) was used as an agonist and capsazepine (10 µM) as an antagonist of TRPV1 channels. Capsaicin, capsazepine, Kt5720, U-73122, wortmannin and

5,8,11,14-eicosatetraynoic acid (ETYA) were all purchased from Sigma and dissolved in DMSO. The final dilution in ND96, used as test or incubation solution, contained maximal 0.5% DMSO. Oocytes were incubated for at least 30 minutes in Kt5720, U-73122, wortmannin or ETYA before measurements were done. All experiments were carried out on at least 3 different oocytes. Gigantoxin I was dissolved in ND96 and administered using the described perfusion system.

Results

Administration of 1.5 μ M and 3 μ M gigantoxin I on wild type TRPV1 channels induced no visible effect (n=3) (Figure 3A). In contrast, when the toxin (1 μ M) was administered in the presence of capsaicin (2 μ M), an activator of TRPV1 channels, an extra inward current was discernible (n=10) (Figure 3B). A similar phenomenon has been observed with cnidaria venom and polycyclic ether toxins and was defined as positive allosterism¹⁷⁻¹⁸.

In order to elucidate the mechanism behind this allosteric effect, regulatory factors of TRPV1 channels which might be involved in this effect were investigated step by step (Figure 4). The first factor we excluded was protein kinase C (PKC). This factor is known to sensitize the channel by phosphorylation on two serine residues (Ser-502 and Ser-800). A TRPV1 channel mutant was made replacing both serine residues into alanines (S502A/S800A mutant). This mutant has already been described and is generally accepted as research tool in the involvement of PKC in the effect of substances on TRPV1 channels³⁸. Tests of gigantoxin I on this mutated channel showed the same allosteric effect of the toxin when administered together with capsaicin (n=4), indicating that PKC induced phosphorylation is not the main working mechanism. The second intracellular regulator we checked was protein kinase A (PKA). Because four phosphorylation sites on the TRPV1 channel are known for PKA (serine-116, threonine-144, threonine-370 and serine-502), we have chosen to use the potent and selective PKA inhibitor, Kt5720. This pharmacological tool is widely used when investigating the role of PKA in the modulation of ion channels³⁹. Control tests were carried out in order to check the activity of Kt5720. Indeed, the capsaicin induced current after 30 min incubation in Kt5720 (3 μ M) was significantly decreased, indicating that the PKA dependent recovery from desensitization was prevented (data not shown)⁴⁰. Measurements on Kt5720 incubated oocytes showed the same allosteric effect of the toxin together with capsaicin (n=3), indicating that PKA is most probably not involved in the toxin effect. A third factor that might be involved in the described action of gigantoxin I is PIP₂. It has been shown that mutations that weaken the TRPV1-PIP₂ interaction induce a reduction of thresholds for

chemical and physical stimuli⁴¹. Therefore, it can be postulated that gigantoxin I reduces the threshold to capsaicin by promoting the hydrolysis of PIP₂ by phospholipase C (PLC) into diacylglycerol (DAG) and inositol (1,4,5)-triphosphate (IP₃). As a consequence, more TRPV1 channels will be activated by the same capsaicin concentration, leading to a greater inward current. In order to test this hypothesis, 3 μM U-73122, a PLC and PLA₂ inhibitor, was added to the incubation solution as described before⁴²⁻⁴³. Control experiments were carried out. Herein, desensitization of TRPV1 channels to capsaicin was clearly inhibited by U-73122 (data not shown). When gigantoxin I was tested on the incubated oocytes, no further effect was visible (n=5). This test leaves two open possibilities: the toxin decreases the PIP₂ concentration or PLA₂ is involved in the induction of the effect. Therefore, wortmannin (100 μM), an inhibitor of phosphoinositide 3-kinases (PI3K) was added to the incubation solution. PI3K phosphorylates PIP₂ into PIP₃ and thus decreases the PIP₂ concentration. Indeed, previous articles describe the depletion of PIP₂ in pre-incubated cells with high concentrations of wortmannin³⁴⁻⁴⁴. Interestingly, in our test the toxin still showed an allosteric effect with capsaicin (n=6). As a corollary, the only remaining possibility is a PLA₂ involvement. The involvement of this pathway was checked by adding ETYA, a non-specific inhibitor of cyclooxygenases (COX) and lipoxygenases (LOX). Both compounds are connected to the PLA₂ pathway (Figure 4). In this test, no effect was visible confirming the involvement of PLA₂ pathway in the effect of gigantoxin I on TRPV1 channels (n=3).

Since voltage-gated potassium and sodium channels are considered as the prime targets of sea anemone peptides, the effect of 1 μM gigantoxin I was also tested on a large collection of ion channels, including: K_v 1.1, K_v 1.2, K_v 1.3, K_v 1.4, K_v 1.5, K_v 1.6, *Shaker*IR, *hERG*, Na_v 1.1, Na_v 1.2, Na_v 1.4, Na_v 1.5, Na_v 1.6, Na_v 1.7, Na_v 1.8 and DmNa_v1. Here, no effect on these subtypes was found (data not shown), indicating a certain selectivity of gigantoxin I.

Discussion

Gigantoxin I, the first isolated EGF-like toxin of natural origin, was previously described to have crab toxicity with weakly lethal effect. Nevertheless, the physiological target of this toxin remained unclear. In this study, an allosteric effect on TRPV1 channels was shown. Remarkably, this effect corresponds with the previously described effect of cnidaria venom, indicating that EGF-like toxins might be the link between the cnidaria venom and TRPV1 channels. Because this new information may lead to the development of new analgesic treatments in cnidaria intoxication, it is important to find out the exact action mechanism of this TRPV1 channel modulating toxin. This study shows the involvement of the intracellular

PLA₂ dependent pathway. Keeping in mind that gigantoxin I possesses 33% homology with the mammalian EGF factor (see Figure 1), we hypothesize that gigantoxin I acts as an EGF factor by stimulating a tyrosine kinase activity of the EGF receptor and thus the calcium-dependent PLA₂ activity. As a consequence, PLA₂ releases AA which is oxidized by LOX generating metabolites such as HPETE which in turn act as TRPV1 channel agonist. Prostaglandine E2 (PGE₂), another AA metabolism product generated via COX, can act on G-protein coupled receptors (GPCR), inducing PKA and PKC activity and thus sensitizing TRPV1 channels. Since we ruled out the participation of PKA or PKC in this study, a role of the PLA₂ dependent pathway involving GPCRs is unlikely. Therefore, we can conclude that PLA₂/AA/LOX is most probably the main pathway that is involved in the gigantoxin I induced effect on TRPV1 channels. An important remaining question is why TRPV1 channels first need to be activated before visible effect is induced by gigantoxin I. It is known that EGF requires concomitant elevation of intracellular calcium for PLA₂ activity⁴⁵. The same conditions may be needed for the activation of PLA₂ when gigantoxin I is bound to EGF receptor. Therefore, the opening of TRPV1 channels by an agonist (*e.g.* capsaicin) inducing inward calcium current might be needed for the activation of the intracellular cascade leading to the described TRPV1 effect.

In conclusion, this study describes the physiological target of gigantoxin I and the involved intracellular EGFR/PLA₂/AA/LOX pathway in the action mechanism of the toxin. This is the first time that this pathway is described in the indirect activation of TRPV1 channels by toxins. This pathway and its coupling to TRPV1 channel activation can serve as a model in further research on toxins and ion channels including the APHC1 toxin mentioned before. This new knowledge also leads to a better understanding of the action mechanism of TRPV1 channels themselves. Since gigantoxin I is a link between the previously described painful cnidaria envenomation symptoms and the effect on TRPV1 channel, these new insights may lead to the development of new analgesic treatments in cnidaria intoxications.

Acknowledgements

We thank Dr. David Julius for providing the rat TRPV1 clone. This work was supported by the following grants: G.0330.06 and G.0257.08 (F.W.O.-Vlaanderen) and P6/31 (Interuniversity attraction Poles Programme - Belgian State - Belgian Science Policy).

Figures

	1	10	20	30	40
Gigantoxin I	DVG	VACTGQYASS	FCLN	GGTCRYIPEL	GEYYCICPGDYTGHRCEQMSV
Human EGF	NSDSE	CPLSHDG	YCLHD	GVCMYIEALDKYAC	NCVVG
Pig EGF	NSYSE	CPPSHDG	YCLHG	GVCMYIEAVDSYAC	NCVFGYVGERCQHRDLKWWELR
Mouse EGF	NSYPG	CPSSYDG	YCLN	GGVCMHIESLDSYTC	NCVIGYSGDRCQTRDLRWWELR

Figure 1. Sequence homology between gigantoxin I and EGF factors




Figure 2. Basic architecture of TRP (TRPV1) channels.

Tetrameric structure of the functional channel exists of four domains. One domain has six transmembrane segments and a pore forming hydrophobic loop between the fifth and the sixth segment. Ankyrin-binding domains (A) can be found in the N-terminus.




Figure 3A. Example of TRPV1 current induced by application of 2 μM capsaicin (CAP). No current is visible after application of 1.5 and 3 μM gigastrontoxin I (GIGANTx). Capsazepine is used as a TRPV1 channel blocker. **B.** Example of the extra inward current visible when 1 μM GIGANTx is added together with CAP (2 μM). In (A) and (B), capsazepine (CZP, 10 μM) was used as a TRPV1 channel blocker.





Figure 4. The involvement of different intracellular pathways in the gigantoxin I induced effect on TRPV1 channel is checked. Numbers indicated in the pathway figures refer to tests excluding specific pathways. The corresponding currents are shown.

1. Gigantoxin I effect is tested on S502A/S800A mutants in order to check the involvement of PLC. The allosteric effect is still visible, indicating no mayor involvement of this pathway. **2.** Kt5720 is used as PKA inhibitor. Current show visible effect of the toxin, indicating no mayor effect of PKA in the induction of the gigantoxin I effect. **3.** U-73122, used as non-specific PLC and PLA2 inhibitor, blocked the allosteric effect of gigantoxin I. **4.** Wortmannin incubation, leading to a higher PIP2 concentration via the block of PI3K, does not influence the

gigantoxin I effect on TRPV1 channels. 5. ETYA, a non-specific COX and LOX inhibitor, blocked the allosteric effect. The pathway that appears to be involved in the described effect is indicated with black arrows.

TRPV1, transient receptor vanilloid subtype 1; CAP, capsaicin; CZP, capsazepin; PIP3, phosphatidylinositol triphosphate; AC, adenylate cyclase; GIGANTx, gigantoxin I; EGFR, epidermal growth factor receptor; GPCR, G-protein coupled receptor; PLA2, phospholipase A2; AA, arachidonic acid; COX, cyclooxygenase; LOX, lipoxygenase; PIP2, phosphatidylinositol biphosphate; HPETE, hydroperoxyeicosatetraenoic acid; PKC, protein kinase C; PKA, protein kinase A; PLC, phospholipase C; DAG, diacylglycerol; IP₃, inositol (1,4,5)-triphosphate; PGE₂, Prostaglandine E2.

gene name	P_{Ca}/P_{Na}	proposed functions / possible disease connections	highest expression
TRPV subfamily			
TRPV1	3.8 (heat), 9.6 (vanilloids)	detection of mild noxious heat, taste/ functional bowel disease, inflammatory bowel disease, osteoarthritis, pancreatitis, impaired bladder contractions, impaired thermal hyperalgesia, asthma, cystitis, schizophrenia, gastroesophageal reflux	TG, DRG neurons, urinary bladder, testis, brain, respiratory tract, gastrointestinal tract
TRPV2	3	sensing thermal pain, mechanosensing/ role in cardiomyopathy, cardiac hypertrophy, interstitial fibrosis, muscular dystrophia, prostate cancer	DRG, spinal cord, brain, spleen, intestine
TRPV3	2.6	warm sensing, osmosensing/ defective environmental thermosensation, hairlessness, atopic dermatitis, breast cancer, non-insulin-dependent diabetes mellitus	TG, DRG, spinal cord, brain, keratinocytes, tongue
TRPV4	6	osmosensing, warm sensing, nociception, pressure sensing/ hypotonic hyperalgesia, impaired acidic nociception, impaired thermal hyperalgesia, asthma, neuropathic pain, hypertension, cardiopathy	DRG, kidney, lung, spleen, testis, heart, liver, keratinocytes, endothelia

Table 1. Overview of the function and expression profiles of TRPV channels (P_{Ca} , permeability for Ca ions; P_{Na} , permeability for Na ions; TG, trigeminal ganglia; DRG, dorsal root ganglia).

References

1. G. M. Watson and D. A. Hessinger, *Science* **1989**, 243,1589-91.
2. J. S. de Oliveira, A. J. Zaharenko, et al., *Biochim Biophys Acta* **2006**, 1760,453-61.
3. M. E. Lanio, V. Morera, et al., *Toxicon* **2001**, 39,187-94.
4. K. Shiomi, E. Tanaka, et al., *Toxicon* **1985**, 23,865-74.
5. G. Uechi, H. Toma, et al., *Protein Expr Purif* **2005**, 40,379-84.
6. G. J. Kelso and K. M. Blumenthal, *Toxicon* **1998**, 36,41-51.
7. L. Standker, L. Beress, et al., *Toxicon* **2006**, 48,211-20.
8. J. J. Smith and K. M. Blumenthal, *Toxicon* **2007**, 49,159-70.
9. F. Bosmans, A. Aneiros, et al., *FEBS Lett* **2002**, 532,131-4.
10. O. Castaneda, V. Sotolongo, et al., *Toxicon* **1995**, 33,603-13.
11. S. Diochot, E. Loret, et al., *Mol Pharmacol* **2003**, 64,59-69.
12. G. S. Gendeh, L. C. Young, et al., *Biochemistry* **1997**, 36,11461-71.
13. Y. Hasegawa, T. Honma, et al., *Toxicon* **2006**, 48,536-42.
14. R. S. Norton, M. W. Pennington, et al., *Curr Med Chem* **2004**, 11,3041-52.
15. T. Honma, H. Nagai, et al., *Biochim Biophys Acta* **2003**, 1652,103-6.
16. M. Diakonova, B. Payrastra, et al., *J Cell Sci* 108 (Pt 6),2499-509.
17. E. Cuypers, A. Yanagihara, et al., *FEBS Lett* **2006**, 580,5728-32.
18. E. Cuypers, A. Yanagihara, et al., *Biochem Biophys Res Commun* **1995**, 361,214-7.
19. K. Andreev, Koshelev, Ivanova, Monastyrnaya, Kozlovskaya, Grishin, *The Journal of Biological Chemistry* **2008**, 283,23914-23921.
20. P. Bohlen, Zhou, King, Siemens, Julius, *Cell* **2010**, 141,834-845.
21. C. Montell and G. M. Rubin, *Neuron* **1989**, 2,1313-23.
22. R. C. Hardie and B. Minke, *Neuron* **1992**, 8,643-51.
23. R. C. Hardie, *Annu Rev Physiol* **2003**, 65,735-59.
24. V. Y. Moiseenkova-Bell, L. A. Stanciu, et al., *Proc Natl Acad Sci U S A* **2008**, 105,7451-5.
25. J. B. Calixto, C. A. Kassuya, et al., *Pharmacol Ther* **2005**, 106,179-208.
26. M. Numazaki and M. Tominaga, *Curr Drug Targets CNS Neurol Disord* **2004**, 3,479-85.
27. M. Tominaga and T. Tominaga, *Pflugers Arch* **2005**, 451,143-50.
28. P. M. Bailey, M. Little, et al., *Med J Aust* **2003**, 178,34-7.
29. B. Liu, C. Zhang, et al., *J Neurosci* **2005**, 25,4835-43.
30. H. H. Chuang, E. D. Prescott, et al., *Nature* **2001**, 411,957-62.
31. D. P. Mohapatra and C. Nau, *J Biol Chem* **2003**, 278,50080-90.
32. S. V. Bhavé and P. L. Hoffman, *J Neurochem* **2004**, 88,359-69.
33. S. W. Hwang, H. Cho, et al., *Proc Natl Acad Sci U S A* **2000**, 97,6155-60.
34. H. Cho, J. Y. Hwang, et al., *J Biol Chem* **2002**, 277,27742-7.
35. M. Van Der Stelt and V. Di Marzo, *Eur J Biochem* **2004**, 271,1827-34.
36. E. R. Liman, J. Tytgat, et al., *Neuron* **1992**, 9,861-71.
37. M. J. Caterina, M. A. Schumacher, et al., *Nature* **1997**, 389,816-24.
38. G. P. Ahern, *J Biol Chem* **2003**, 278,30429-34.
39. A. Varga, K. Bolcskei, et al., *Neuroscience* **2006**, 140,645-57.
40. G. Bhavé, W. Zhu, et al., *Neuron* **2002**, 35,721-31.
41. E. D. Prescott and D. Julius, *Science* **2003**, 300,1284-8.
42. I. Vetter, B. D. Wyse, et al., *Eur J Pain* **2008**, 12,441-54.
43. V. Lukacs, B. Thyagarajan, et al., *J Neurosci* **2007**, 27,7070-80.
44. J. Benedikt, J. Teisinger, et al., *J Neurochem* **2007**, 100,211-24.
45. H. J. Goldberg, M. M. Viegas, et al., *Biochem J* **1990**, 267,461-5.

