CXCR3 ligands in disease and therapy

Katrien Van Raemdonck¹, Philippe E. Van den Steen², Sandra Liekens³, Jo Van Damme¹, Sofie Struyf¹

¹Laboratory of Molecular Immunology

²Laboratory of Immunobiology

³Laboratory of Virology and Chemotherapy

University of Leuven (KU Leuven), Department of Microbiology and Immunology,
Rega Institute for Medical Research, Leuven, Belgium

Corresponding author

Prof. Jo Van Damme

Laboratory of Molecular Immunology, Rega Institute for Medical Research,
University of Leuven (KU Leuven), Minderbroedersstraat 10, 3000 Leuven, Belgium

Tel: + 32 16 337342

Fax: + 32 16 337340

Jo.Vandamme@rega.kuleuven.be

ABSTRACT

Chemokines, binding their various G protein-coupled receptors, lead the way for leukocytes in health and inflammation. Yet chemokine receptor expression is not limited to leukocytes. Accordingly, chemokines are remarkably pleiotropic molecules involved in a range of physiological as well as pathological processes. For example, the CXCR3 chemokine receptor is expressed on activated T lymphocytes, dendritic cells and natural killer cells, but also fibroblasts and smooth muscle, epithelial and endothelial cells. In men, these cells express either CXCR3A, its splice variant CXCR3B or a balanced combination of both. The CXCR3 ligands, activating both receptor variants, include CXCL4, CXCL4L1, CXCL9, CXCL10 and CXCL11. Upon CXCR3A activation these ELR-negative CXC chemokines mediate chemotactic and proliferative responses, for example in leukocytes. In contrast, CXCR3B induces anti-proliferative and anti-migratory effects, as exemplified by angiostatic effects on endothelial cells. Taken together, the unusual and versatile characteristics of CXCR3 and its ligands form the basis for their pertinent involvement in a myriad of diseases. In this review, we discuss the presence and function of all CXCR3 ligands in various malignant, angiogenic, infectious, inflammatory and other disorders. By extension, we have also elaborated on the potential therapeutic applicability of CXCR3 ligand administration or blockade, as well as their additional value as predictive or prognostic biomarkers. This review illustrates the multifunctional, intriguing character of the various CXCR3-binding chemokines.

Keywords: CXCR3; chemokine; G protein-coupled receptor

1. AN INTRODUCTION TO CHEMOKINES

Originally discovered to be chemotactic cytokines leading the way for leukocytes in health and inflammation, the chemokines have now made a name for themselves as pleiotropic molecules involved in a range of physiological as well as pathological processes [1, 2]. Most of their effects are established by signaling through a set of chemokine receptors, typically G protein-coupled receptors (GPCR). Notably, chemokine ligand/receptor binding is both a promiscuous and a redundant interaction, meaning that a particular chemokine will show affinity for several chemokine receptors and a particular chemokine receptor can be activated by various chemokine ligands, respectively. Nevertheless, due to their unique temporal and spatial expression patterns, chemokines often do have non-redundant functions *in vivo* [3]. The biology of chemokines and their receptors is even further complicated by the identification of a set of atypical chemokine receptors (ACKR). Highly related to the classic G protein-coupled chemokine receptors, ACKR are a smaller subgroup of seven-transmembrane receptors, which bind chemokines with high affinity, but do not signal through G proteins [4]. They were originally considered silent scavenger receptors, however, the majority of ACKRs has since been shown to preserve β -arrestin-dependent signaling. Additionally, glycosaminoglycans (GAGs) also influence chemokine function, mostly by retaining chemokines at the endothelial cell surface and presenting them to passing leukocytes that express corresponding chemokine receptors.

The chemokines themselves are small proteins, not bigger than 8 to 10 kDa. They share a common three-dimensional structure defined primarily by four conserved cysteines, two near the amino-terminal tail of the protein, forming disulfide-bonds with two more internal cysteines. A functional distinction is sometimes made between inflammatory and homeostatic chemokines. However, more frequently, chemokine classes are subdivided on the basis of their amino acid sequence homology. Based on the exact configuration of the amino-terminal cysteine pair, a distinction can be made between different subfamilies. The CXC chemokines distinguish themselves from the CC chemokines by the inclusion of a single variable amino acid 'X' separating those amino-terminally located cysteine residues. Together, the CC and CXC subfamilies constitute the majority of chemokines, whereas the C

and CX₃C chemokines represent a limited number of rather unconventional chemokines. Though seemingly limited by the constraints of a highly conserved structure, the chemokines are without a doubt a diverse superfamily. This certainly also applies to the subfamily of CXC chemokines, which has gained extra attention because of its role in the regulation of angiogenesis, the formation of new blood vessels, branching of a pre-existing vascular network [5, 6]. Interestingly, a conserved ELR-motif, consisting of a glutamic acid, leucine and arginine, often precedes the most amino-terminal cysteine. The presence of this motif appears crucial in determining the chemokine's biological function as it defines receptor-binding [7]. All ELR-positive CXC chemokines bind the CXCR2 receptor, expressed on leukocytes, such as neutrophils, and on endothelial cells [8, 9]. Accordingly, the ELR-positive CXC chemokines attract neutrophils and promote angiogenesis, respectively. The ligands of CXCR1 also contain an ELR-motif and are particularly potent neutrophil chemoattractants. On the other hand, CXC chemokines lacking the ELR-motif are unable to bind CXCR2. For example, CXCL12/stromal cell-derived factor-1 (SDF-1) alternatively binds CXCR4 and ACKR3/CXCR7, receptors to which pro-angiogenic effects have been attributed as well [10]. However, most ELR-negative CXC chemokines bind the CXCR3 receptor, as do CXCL4/platelet factor-4 (PF-4), CXCL4L1/PF-4var, CXCL9/monokine induced by interferon- γ (Mig), CXCL10/interferon- γ -induced protein-10 (IP-10) and CXCL11/interferon-inducible T cell α chemoattractant (I-TAC) (Figure 1) [11-15]. As opposed to CXCR2, this CXCR3 receptor has been ascribed angiostatic properties, specifically its splice variant CXCR3B, a variant expressed on endothelial cells [11].

The CXCR3 ligands thus represent a distinct group of angiostatic chemokines, besides also retaining their classic chemotactic potential as they additionally direct migration of CXCR3A-expressing lymphocytes to sites of inflammation. Most CXCR3 ligands also bind several GAGs, despite considerable differences in affinity among these chemokines [15-17]. They thus compete with various angiogenic growth factors, such as vascular endothelial growth factor (VEGF) and basic fibroblast growth factor (FGF2), for binding to GAGs on the proteoglycans presented on the endothelial cell surface. Consequently, CXCR3 ligands hinder binding of these growth factors to their respective signaling receptors [18]. As indicated by recent studies, additionally, direct proteoglycan-mediated

signaling seems increasingly plausible. Indeed, GAG-binding has been held accountable for effects of CXCL4 on monocytes and neutrophils [19-22].

2. CXCR3 AND ITS LIGANDS

2.1. The CXCR3 receptor

The CXCR3 receptor is a seven-transmembrane GPCR, classified as a CXC-type receptor based on the structure of its chemokine ligands. This receptor was indeed originally cloned and characterized as an activated T lymphocyte-expressed GPCR selective for ELR-negative CXC chemokines CXCL9 and CXCL10 [13]. In contrast to other chemokine receptors, the human CXCR3 gene was allocated to the X chromosome, in the q13 region [23]. In culture, the percentage of CXCR3-positive T lymphocytes has been shown to increase upon interleukin (IL)-2 or IL-2/phytohemagglutinin (PHA) stimulation, resulting in up to 95% receptor-expressing cells [23]. T lymphocyte activation accordingly induces cell responsiveness to CXCR3 ligands. A heterogeneous group of effector T lymphocytes, including CD4⁺ T helper (Th) cells, CD8⁺ cytotoxic T lymphocytes (CTL), as well as memory T cells, either CD4⁺ or CD8⁺, show highly increased expression of CXCR3 in contrast to their naïve precursors [24, 25]. However, among the Th cell population, CXCR3 would particularly mark Th1 cells, rather than Th2 or Th17 cells [24, 26]. The transcription factor T-bet, a master switch for Th1 and CTL commitment, directly promotes CXCR3 expression [27, 28]. Interestingly, IFN- γ signaling, via T-bet, also promotes CXCR3 expression on a subset of Th1-specific regulatory T cells [29]. Additionally, natural killer (NK) cells were demonstrated to express CXCR3 [13]. Functional expression of CXCR3 was also demonstrated on dendritic cells [15, 30]. CXCR3 ligands have a broad spectrum of biological activities, acting on various CXCR3-expressing cell types. Though CXCR3 expression among immune cells is restricted, with no detectable expression on resting T lymphocytes, B lymphocytes, monocytes or granulocytes, a wide variety of non-immune cells have in fact been shown to express and display CXCR3 as well, including fibroblasts, endothelial, epithelial and smooth muscle cells [31]. These cells do display a variable, cell-dependent expression pattern of different human CXCR3 splice variants. Most cells, including leukocytes, predominantly express the originally identified CXCR3A. CXCR3A consists of 368 amino acids, encoded by just two exons, separated by a

single intron, and couples to a $G_{\alpha i}$ which mediates pro-migratory and proliferative signaling and increases intracellular calcium levels [11, 13, 23]. In contrast, endothelial cells exclusively express the splice variant CXCR3B, which instead couples to a $G_{\alpha s}$ subunit, initiating opposing signaling cascades [11]. CXCR3B is a 415 amino acid long receptor, generated by alternative splicing at the 5' end of the second exon. Additionally, the inserted sequence comprises an alternative start codon resulting in the loss of the four most amino-terminal residues encoded by the first exon, which are replaced by a new amino-terminal tail of 51 amino acids. Most CXCR3A-positive cells do exhibit low, parallel expression of CXCR3B. Existence of two presumably functionally divergent variants can explain the angiostatic effect attributed to CXCR3 ligands. No alternative splicing of the CXCR3 receptor was identified in mice, though murine CXCR3 on endothelial cells does exhibit CXCR3-mediated angiostatic effects [15]. A cell-dependent signaling mechanism could be involved, establishing differential G protein-coupling, independent of CXCR3 splicing. Different intracellular domains of the receptor have been shown to determine functional activity, in a ligand-dependent manner [32, 33]. The different domains are also of varying importance in establishing distinct, parallel downstream effects. However, the determinants of splice variant-dependent differential CXCR3 G protein-coupling remain thus far enigmatic. Furthermore, a third human splice variant generated by exon skipping, named CXCR3-alt, was later identified [14]. CXCR3-alt consists of only 267 amino acids. This variant shows considerable structural and functional dissimilarities attributed to a frame-shift. Only four transmembrane domains are conserved and a fifth domain within the alternative carboxy-terminal sequence has been postulated [14]. Despite drastically deviating from the classic seven transmembrane CXCR3A conformation, CXCR3-alt surprisingly retains affinity for CXCL11, whereas no other CXCR3 ligands are bound (Figure 1). Though affinity is lowered, CXCL11 can evoke a CXCR3-alt-mediated modest calcium increase and chemotactic response [14, 34].

Study on altered, chimeric CXCR3/CXCR1 receptors has taught that several distinct extracellular domains of human CXCR3A collectively ensure efficient ligand binding and following receptor activation, proposing a so-called multi-site model of ligand/receptor interaction [35]. CXCR3's amino-terminal tail and first extracellular loop, though dispensable for CXCL9-induced activation, both play

a more important role in CXCL10- and CXCL11-mediated activation. In fact, sulfation of two tyrosine residues in the amino-terminus of CXCR3A was later described to be a necessity for binding and activation by all three chemokines [36]. In line with earlier findings, this report did, however, also demonstrate CXCL10 and CXCL11 receptor binding to be more sensitive to amino-terminal truncation of the receptor compared to CXCL9 [36]. The second extracellular loop of CXCR3A, and in particular an arginine at position 216, proved crucial for receptor activation by all interferon (IFN)- γ -inducible chemokines [35, 36]. Surprisingly though, R216 is no absolute requirement for ligand/receptor binding nor CXCR3 internalization. Finally, CXCR3A's third extracellular loop appears necessary for CXCL9 and CXCL10 function [35]. In general, CXCL11 is suggested to be more tolerant to substantial changes within the extracellular loops. Accordingly, CXCL11 is indeed the only CXCR3 ligand known to induce CXCR3-alt-mediated signaling, as mentioned earlier [14].

Note that past research solely focused on the amino acid sequence and structure of splice variant CXCR3A, disregarding the binding properties of CXCR3B. The amino-terminal elongation of CXCR3B and the associated shift in affinity towards different CXCR3 ligands, can be reconciled though with the postulated multi-site model and the reported impact of amino-terminal changes on ligand binding. Beyond the altered 51 amino acid long amino-terminal tail, both splice variants are identical. Presumably, other findings on the structure and function of CXCR3A can thus be extended to CXCR3B.

Among the CXCR3-binding chemokines, two groups clearly can be distinguished based on their overall biology and biochemistry (Table 1). On one hand, the IFN- γ -inducible chemokines CXCL9, CXCL10 and CXCL11 constitute the traditional CXCR3 ligands. The platelet-derived chemokines CXCL4 and CXCL4L1, on the other hand, were only later recognized to bind both CXCR3A and CXCR3B as well and represent a distinct pair of CXCR3 ligands.

2.2. IFN- γ -inducible CXCR3 ligands

Three ELR-negative CXC chemokines, CXCL9, CXCL10 and CXCL11, can together be referred to as the IFN- γ -inducible CXCR3 ligands [12, 37, 38]. Though they share a common inflammatory

function, they do collectively gain fine control over leukocyte trafficking [3]. The mature IFN- γ -inducible chemokines display about 40% amino acid sequence identity between one another (Figure 2). However, the expression of CXCL9, CXCL10 and CXCL11 is regulated by unique promoters establishing distinct temporal and spatial expression patterns (Table 1). They are indeed differentially expressed by various cell types, induced by specific cytokines and Toll-like receptor (TLR) ligands or by combinations thereof [3, 39-41]. Furthermore, these chemokines all exhibit different receptor binding characteristics, different efficacy as well as potency. Generally, CXCL11 has the highest affinity for CXCR3A, whereas CXCL9 has the lowest. CXCL11 is also the most potent of the IFN- γ -inducible CXCR3 ligands, as measured by both mobilization of intracellular calcium and chemotactic response in CXCR3A-expressing cells [12]. However, CXCL10 expression appears more receptive to a range of different stimuli and has been associated accordingly with a multitude of physiological as well as pathological conditions [3]. CXCL10 is the most thoroughly investigated CXCR3 ligand and could be suggested a prototypical example of the IFN- γ -inducible chemokines.

2.3. Platelet-derived CXCR3 ligands

The mature chemokines CXCL4 and CXCL4L1 both show 37% sequence identity with the IFN- γ -inducible CXCR3 ligand CXCL10 (Figure 2). However, among themselves, these non-allelic variants show a striking 96% amino acid identity. CXCL4L1 is a non-allelic variant of CXCL4 which arose from a gene duplication unique to primates [42]. Only three amino acid substitutions near the carboxy-terminal end distinguish the mature chemokines. Yet this subtle difference suffices to alter the chemokine's three-dimensional structure drastically, reducing CXCL4L1's affinity for GAGs and resulting increased angiostatic potential of the variant CXCL4L1 [42, 43].

On the other hand, CXCL4 and CXCL4L1 clearly differ from the other CXCR3 ligands firstly based on their affinity for CXCR3A and CXCR3B, but also because their expression is not classically induced by IFN- γ . Even more, they are primarily released by activated platelets, therefore coined as the platelet-derived CXCR3 ligands [43, 44].

3. CXCR3 LIGANDS IN DISEASE

Given their pleiotropic nature, CXCR3 ligands play their part in a myriad of diseases. Involvement of all CXCR3 ligands has been observed in various angiogenesis-related pathologies as well as immunological disorders, the latter being mostly related to autoimmunity or infection [45, 46]. Here we discuss a varied and extensive, yet not exhaustive, repertoire of diseases having been associated with CXCL4, CXCL4L1, CXCL9, CXCL10 or CXCL11. The CXCR3 ligands are indeed characterized by their unique position at the crossroad of immunity and angiogenesis and accordingly play an exceptional role in health, disease and therapy.

3.1. Carcinoma and sarcoma

The CXCR3 ligands have been associated with many diseases, some new and still poorly understood, some already the subject of a long line of research. Their association with the pathogenesis and prognosis of cancer definitely belongs to the latter. The immunoangiostatic CXCR3 ligands are established players in numerous carcinomas and sarcomas [47]. Angiogenesis is indeed a critical process providing the expanding malignant tissue with oxygen and nutrients and clearing it of metabolic waste products [48, 49]. While these newly formed blood vessels allow successful tumor growth, the lymphatic vasculature will serve as an additional escape route for metastatic tumor cells, especially in later stages of tumor progression [50]. Recent reports reveal that CXCR3 ligands, specifically the platelet factors, are effective inhibitors of lymphangiogenesis as well [51] and hence support the belief that these chemokines are interesting, multifunctional anti-tumoral agents (Figure 3).

Furthermore, immunity also plays an ambiguous role. Generally, the tumor is believed to corrupt the immune system and suppress any anti-tumoral reactivity, possibly even attracting immune cells that will further promote tumor vascularization [52]. In contrast, immune polarization of the tumor microenvironment could potentially stimulate the anti-tumoral immune response, attracting CXCR3A-expressing, anti-tumoral CTLs, activated Th1 lymphocytes and NK cells [53]. It is no wonder that several chemokines, being immunological players attributed both angiogenic and angiostatic properties, also play their part in tumor development [45]. An abundance of studies have reported

overexpression or absence of CXCR3 ligands, especially CXCL10, in cancer patients [47, 54]. Despite a broad understanding of their potential anti-tumoral activity, the associated clinical significance often remains enigmatic. The impact of CXCR3 ligands on both tumor and tumor microenvironment is further complicated by some tumors' ability to express the CXCR3 receptor. CXCR3 expression was illustrated in many different cancers, formerly without distinguishing CXCR3A from CXCR3B [55-58]. Given tumoral G_{αi}-coupled CXCR3A surface expression, CXCR3 ligands will stimulate tumor cell migration and proliferation thereby promoting tumor invasiveness (Figure 4) [59]. CXCL4 and CXCL10 were accordingly reported to promote the metastatic potential of human, predominantly CXCR3A-expressing, prostate cancer cells *in vitro* [59]. In several other cancers, tumoral expression of CXCR3 was also associated with lymph node invasion or generally poor prognosis [60-62]. While mostly considered anti-tumoral agents, CXCR3 ligands can thus also have a detrimental, pro-tumoral effect under certain conditions. These findings require careful consideration to assure safe and efficient future therapeutic applicability. The impact of individual differences in tumoral CXCR3A and CXCR3B expression demands a personalized approach.

3.2. Leukemia, lymphoma and myeloma

Leukemia, lymphoma and myeloma, collectively referred to as hematopoietic malignancies, represent a rather unique set of cancers affecting leukocytes and their progenitors. All three diseases can be further divided into subclasses. As for leukemia, a distinction is made between acute or chronic, and lymphoid or myeloid disorders. Interestingly, the CXCR3 receptor and its ligands have been discussed in the context of all four major subtypes. In chronic lymphocytic leukemia (CLL), the CXCR3-signaling axis has mostly been linked to trafficking of CLL malignant B cells, which consistently express functionally active CXCR3, unlike B cells isolated from healthy subjects [63, 64]. CLL cells themselves were also frequently observed to co-express CXCL9 [64]. Furthermore, CXCL9, CXCL10 and CXCL11 were identified as part of an outcome-correlated cluster of cytokines, differentially expressed in CLL, with elevated serum levels serving as an indication of aggressive disease [65]. The cluster additionally consisted of CCL3, CCL4, CCL19, IL-5, IL-12 and IFN- γ . On the other hand, low levels of CXCR3 expression by CLL B cells have previously been reported to predict a poor outcome

and shorter survival as well [66]. In acute lymphoblastic leukemia (ALL), CXCL10 produced by monocytes in the vicinity of malignant blasts was hypothesized to have a direct pro-tumoral effect promoting tumor cell invasiveness [67]. CXCL10 serum levels in ALL patients were also proven to be increased compared to controls [68]. In contrast, decreased serum levels of another CXCR3 ligand, CXCL4, were discovered to characterize pediatric ALL [69]. Remarkably, involvement of CXCL4 in the oncogenesis of ALL was strongly supported by the identification of a translocation shared among a subset of ALL patients, with a breakpoint in 4q21, distal to the CXCL4 gene [70, 71]. Similarly, in acute myeloid leukemia (AML) CXCL4 serum levels were decreased as well, yet progressed in relation to therapy responsiveness, with increasing concentrations indicating AML remission [72, 73]. On the other hand, AML cells often constitutively produce CXCL10, with wide variations between patients potentially reflecting disease status [74]. Considering chronic myeloid leukemia (CML), a fourth leukemic subtype, CXCL10 mRNA levels were just recently reported to be upregulated in peripheral blood mononuclear cells isolated from CML patients, linking CXCR3 ligands to the pathogenesis of CML as well [75]. Dysregulation of CXCL10 gene expression has in fact been associated with several recurrent chromosomal translocations identified in different leukemic pathologies [76].

Hodgkin's lymphoma tissue also generally presents higher levels of CXCL9 and CXCL10 expression, though major differences are noted among different subtypes with predominant expression in Epstein-Barr virus (EBV)-positive cases and cases associated with mixed cellularity [77, 78]. Both chemokines were shown to be expressed by malignant cells as well as surrounding cells such as macrophages, lymphocytes and endothelial cells. This overexpression of both CXCL9 and CXCL10 has actually been associated with tumor tissue necrosis and vascular damage, consistent though with their general anti-tumoral activity. This duality was also illustrated in non-Hodgkin lymphoma, with on one hand reports of CXCL9 expression in B-cell lymphomas facilitating dissemination of CXCR3-positive malignant B cells [79-81] and on the other hand the suggestion that CXCL9 and CXCL10 are in fact associated with a protective immune response successfully targeting B cell cancer, comparable to their anti-tumoral activity reported in solid tumors [82].

Finally, for most multiple myeloma (MM) cases, freshly purified MM plasma cells have been shown to express CXCR3, making them responsive to chemokines like CXCL9, CXCL10 and CXCL11 [83]. Remarkably, myeloma cells express both CXCR3 splice variants, at varying ratios and in a cell cycle-dependent manner, potentially explaining the reported anti-proliferative effect of CXCL10 [84, 85]. As for CXCL4, shown to induce apoptosis of myeloma cells, this platelet factor was identified as tumor suppressor gene commonly silenced in MM as a consequence of promoter hypermethylation [86, 87].

In conclusion, most hematological cancers have, in one way or another, been linked to CXCR3 ligand expression. However, seldom their exact pathological or oncogenic function is really understood. This is a domain that requires further research, especially considering the prospects for leukemia patients and the shortcomings of current treatments. CXCL4 in particular appears to present promising new therapeutic opportunities as this chemokine may also serve to protect normal hematopoietic cells from the toxicity of chemotherapeutics, without affecting the viability of cancerous cells [88].

3.3. Inflammatory arthritis

Over the years, CXCR3 ligands and their function in autoimmune diseases such as rheumatoid arthritis (RA) have been extensively reported and reviewed [89]. In RA patients, recruitment of CXCR3-expressing T cells and mast cells to the inflamed synovium was indeed related to a preferential upregulation of CXCL9 and CXCL10 expression in the synovial fluid and tissue compared to chemokine levels in traumatic arthritis or osteoarthritis [90, 91]. This association was later also extended to enhanced CXCL4 expression concurrently with synovial inflammation in RA [92]. Elevated CXCL4 plasma levels have also been observed in a subset of RA patients, specifically concurrent with the formation of active vascular lesions [93]. Plasma CXCL4 concentrations might indeed prove to be a useful marker of endothelial injuries in rheumatic disorders [93]. Furthermore, synovial CXCL10 expression was also found to be increased in juvenile idiopathic arthritis [94]. Also in septic arthritis, inflammatory induction of CXCL9 and CXCL11 has been reported [41]. However, clearly distinguishing itself from other IFN- γ -inducible chemokines, evidence regarding increased secretion of CXCL11 in RA remains controversial [39, 95]. High synovial expression of CXCL11 in a

particular subpopulation of RA patients has, however, been identified as an indication of poor therapy responsiveness to adalimumab, a TNF-antagonist [96].

3.4. Inflammatory bowel disease

The CXCR3 axis has been implicated in the pathogenesis of both Crohn's disease and ulcerative colitis, two chronic gastrointestinal disorders collectively referred to as inflammatory bowel disease (IBD) [97, 98]. Raised plasma levels of CXCL4 were observed in Crohn's disease already in 1987 [99] and were furthermore correlated with IBD clinical activity [100]. Serum CXCL4 has more recently also been identified as a general biomarker for IBD in a proteomic serum profiling pilot study [101]. Though serum CXCL10 levels were also suggested to reflect disease activity in ulcerative colitis [102], expression of IFN- γ -inducible CXCR3 ligands has mostly been found to be increased locally, driving mucosal accumulation of inflammatory cells [97, 103, 104]. Accordingly, it has been suggested that these IFN- γ -inducible chemokines could contribute to the progression of IBD and worsen the associated inflamed state of the gastrointestinal tract [104, 105]. Involvement of the CXCR3 axis genes was supported by the report of a persistent overexpression of CXCL9, CXCL10, CXCL11 and CXCR3 in the inflamed colonic tissue of pediatric IBD patients [106]. A particular CXCL11 genotype, namely the rs6817952 A allele situated in intron 1, even proved to be a risk factor for both ulcerative colitis and pediatric Crohn's disease [106]. In a similar report, a CXCL9 polymorphism, 77147452G>A, showed an inverse association with pediatric Crohn's disease [107]. These genetic association studies underline the importance of the CXCR3 receptor and its ligands in the pathogenesis of IBD and could inspire new therapeutic strategies.

3.5. Diabetes

As an autoimmune disease, type 1 diabetes has repeatedly been associated with elevated levels of the typical Th1 chemokine CXCL10, a correlation unveiled by Shimada and colleagues and reassessed earlier this year by the group of Antonelli *et al.* [108, 109]. It is further hypothesized that CXCL10 serum levels would increase especially during the early Th1-driven stages of disease [110, 111]. However, there is also a minority of studies contesting this positive association between serum

CXCL10 and type 1 diabetes [112, 113]. Nevertheless, generally it is accepted that hyperglycemic and inflammatory CXCL10 production plays an important role in the pathogenesis of type 1 diabetes [114-116]. While monocytes are an important source of the chemokine, pancreatic beta cells in patient biopsies were actually shown to express and secrete CXCL10 themselves as well, contributing to autoreactive T cell islet infiltration and consequent beta cell destruction [117-119]. Potentially, beta cell-secreted CXCL9 may also attract CXCR3-expressing T cells [120]. Through TLR4 signaling, CXCL10 has even been described to interfere directly with pancreatic beta cell proliferation and viability, an effect originally assumed to be mediated by CXCR3 expressed on pancreatic beta cells [121-123]. Regardless of the controversy concerning involvement of CXCR3 itself, this direct effect on pancreatic beta cell numbers implicates CXCL10 not only in the progression of type 1 but also of type 2 diabetes, as a lack of proliferation prevents beta cells from meeting increased insulin demands, for example in obese or pregnant patients [122]. Accordingly, the latter study indeed showed that in type 2 diabetes patients, unlike healthy controls, pancreatic beta cells also express and secrete CXCL10.

Though not always clear-cut, several studies have investigated the association between platelet activation and coagulation factors on one hand and diabetes on the other. This also led to the observation that diabetic patients present increased CXCL4 plasma levels, presumably related to enhanced platelet activation [124, 125]. Like most CXCR3 ligands, CXCL4 has been studied regarding its involvement in complications of diabetes, such as proliferative diabetic retinopathy (PDR) [125, 126]. Characterized by distorted, pathological neovascularisation of the retina, several angiogenic factors have been shown to have a causative link to this pathology. But the distorted angiogenic balance in the diabetic eye also affects the angiostatic proteins involved, showing a counterintuitive upregulation of inhibitors of angiogenesis such as the CXCR3 ligands [127]. Chemokine concentrations measured in the vitreous fluid of patients correlate to clinical PDR manifestation. For example, vitreous CXCL9 levels have been shown to be elevated in patients with active PDR, but not in patients with inactive PDR [126]. A causal role for the CXCR3 ligands in this

severe complication in diabetes patients may remain enigmatic, but their potential as disease markers is apparent as is their potential therapeutic applicability.

3.6. Systemic sclerosis

Proteome-wide analysis recently revealed circulating and dermal plasmacytoid dendritic cells isolated from systemic sclerosis (SSc) patients to display markedly increased CXCL4 expression [128]. This study also confirmed plasma CXCL4 levels to be elevated, as had been previously described [129, 130]. More importantly, plasma levels of CXCL4 correlated with disease severity, specifically the presence and progression of complications such as pulmonary arterial hypertension and lung or skin fibrosis [128]. Increased CXCL4 levels had previously been measured in the bronchoalveolar lavage fluid of SSc patients [131]. Similarly, serum concentrations of the IFN- γ -inducible CXCR3 ligand CXCL10 appear indicative of disease activity [132]. Increased serum CXCL10 levels also correlate with the manifestation of pulmonary arterial hypertension [133]. In general, increased serum levels of CXCL9 and CXCL10 have been described consistently in SSc patients [134-136]. In contrast, low concentrations of CXCL11 in the lung have been associated with a greater likelihood of pulmonary function decline [137].

3.7. Transplant rejection

As potent T cell-chemoattractants, CXCL9 and CXCL10 have been demonstrated repeatedly to play a central role in the immune response against transplant tissue, leading to allograft rejection [138, 139], despite a few reports opposing involvement of the CXCR3 axis [140]. Regulating immune cell differentiation, migration and proliferation, local chemokine levels indeed closely relate to the host immune status, both pre- and post-transplantation [141-144]. Upregulation of CXCL10 expression, potentially serving as a biomarker predicting allograft failure, is indeed reflected by chemokine concentrations in serum and urine [139, 145], as is CXCL9 expression according to some studies [146]. These findings have been reported in various organs, including the transplanted kidney, lung, heart and liver [147-153]. Similar to CXCL9 and CXCL10, the third IFN- γ -inducible chemokine, CXCL11, has been observed to be upregulated in a number of transplant studies [151, 152, 154, 155].

As measured in endomyocardial biopsies, a persistent elevation in CXCL10 and CXCL11 mRNA expression was found to correlate with the occurrence of transplant vasculopathy, a severe complication among heart transplant recipients causing allograft ischemia [156].

A murine study also associated CXCL4 with allograft rejection in the context of cardiac transplantation. However, CXCL4 was actually considered a negative regulator of T cell reactivity to cardiac transplantation, by limiting Th17 differentiation [157].

CXCR3-binding chemokine levels have also been associated with patient prognosis after bone marrow transplantation, more specifically in the context of graft-versus-host-disease (GVHD), a complication in which transplanted donor cells regard the host tissue as foreign. The importance of CXCL10 in the pathogenesis of this disease was first recognized in acute GVHD, characterized by epidermal CXCL10 expression and recruitment of CXCR3-positive T cells to the skin [158]. Serum levels of this chemokine were significantly increased during acute GVHD. Furthermore, serum levels of all three IFN- γ -inducible CXCR3 ligands, CXCL9, CXCL10 and CXCL11, also correlated to episodes of chronic GVHD of the skin [159]. Correspondingly, during ocular manifestation of chronic GVHD, increased expression of CXCL9 and CXCL10 has been observed within the conjunctiva [160]. These observations thus suggest a central role for the CXCR3 ligands in the pathogenesis of both acute and chronic GVHD.

3.8. Malaria

The role of platelets in malaria, and with it a potential involvement of the platelet-specific protein CXCL4, has been recognized early on [161, 162]. In 1983, Essien and Ebhota described elevated CXCL4 plasma levels in patients upon acute *Plasmodium falciparum* infection [163]. On the other hand, the IFN- γ -inducible CXCR3 ligands were later implicated in the pathogenesis of malaria as well. Most importantly, the CXCR3 ligands have been associated with the occurrence of cerebral malaria (CM), a lethal complication of *Plasmodium falciparum* infection in humans [164, 165]. Aspiring to identify prognostic markers that may predict CM severity and hopefully improve intervention, CXCL10 and CXCL4, in contrast to CXCL9 or CXCL11, were found to be exceptional serum markers

associated with CM mortality [166, 167]. Following up on the potential value of CXCL10 in predicting CM associated mortality, a specific CXCL10 gene promoter polymorphism -1447A>G, associated with higher plasma CXCL10 levels, was identified as a genetic risk factor predicting clinical severity of malaria and susceptibility to CM [168]. The hypothesized key role for CXCL10 in cerebral malaria was further supported by preclinical data illustrating that this chemokine promotes accumulation of CXCR3-positive T lymphocytes in the brain, an essential part of the pathogenesis of CM [165, 169-171]. In parallel to CXCL10, a prominent role for CXCL4 in the pathogenesis of fatal CM was suggested as well [167]. Murine CXCL4 contributed to the vascular inflammation, driving both T cell and monocyte trafficking, which consequently leads to cerebral vascular injury typical of the development of experimental CM [172, 173]. Despite the lack of an identified correlation in patients, preclinical studies have also connected CXCL9 to this severe complication [174-177]. Moreover, in murine models of CM, CXCR3 expression or in contrast lack thereof has been associated with susceptibility to or protection against fatal murine CM, respectively [165, 169, 170, 176]. Protection goes hand in hand with a reduction of T lymphocyte recruitment in response to the IFN- γ -inducible CXCR3 ligands excessively expressed in the brain [170]. Neutralisation of CXCL10 in mice inhibits the recruitment of T cells from the spleen to the brain, thereby preventing cerebral inflammation and improving parasite clearance in the spleen [171]. In contrast, the protective, antimalarial effect of CXCL4 can be traced back to its remarkable quality to bind the Duffy-antigen receptor for chemokines (ACKR1/DARC) expressed on erythrocytes [178-180]. Activated platelets, bound to malaria-infected red blood cells, will locally release high concentrations of CXCL4, which in turn binds DARC leading to intraerythrocytic parasite-killing [179, 181, 182].

3.9. Acquired Immune Deficiency Syndrome

Serving as co-receptors for cell-specific human immunodeficiency virus (HIV) entry, CCR5 and CXCR4, as well as several other chemokine receptors, have been identified as critical pawns in HIV infection and pathogenesis, eventually leading to acquired immune deficiency syndrome (AIDS) [183, 184]. Though it was demonstrated that CXCR3 is also a potential co-receptor, given its marginal co-receptor efficiency, clinically CXCR3 is unlikely to take significant part in the process of HIV

infection [185]. On the other hand, the receptor and its ligands do play a role in the progression of this disease, even though it may not yet be entirely clear what that role entails.

Firstly, plasma levels of CXCL10 were recently suggested as a marker to predict viral load [186, 187]. Furthermore, CXCL10 has been detected in the cerebrospinal fluid (CSF) of HIV-1-infected patients, driving the accumulation of activated T cells and accordingly contributing to the occurrence of HIV-associated disorders of the central nervous system [188-190]. It was indeed repeatedly shown that a variety of cells, including dendritic cells, macrophages and lymphocytes but also astrocytes and microglia, can express significant amounts of CXCL9, CXCL10 and CXCL11 following HIV-infection [191-195]. Notably, CXCL10 was also identified as a direct neurotoxic factor [196, 197]. Most importantly, recruitment of susceptible CXCR3-expressing T cells to virus-infected macrophages, dendritic cells and lymph nodes is favored, ensuring ongoing propagation of the virus in newly supplied target cells and inflicting the general immunopathology of AIDS [192, 198]. Interestingly, the CXCR3 ligands also have a more direct impact on HIV infection as CXCL9, CXCL10 and CXCL11 were all shown to stimulate virus replication in a CXCR3-dependent manner [199].

Similar to the other CXCR3 ligands, CXCL4 also affects the pathogenesis of HIV. Remarkably, CXCL4 is believed to interfere specifically with the earliest events in the viral infectious cycle and would actually be a key player in the first line defense against HIV-1 [200, 201]. Activated platelets inhibit host cell entry by HIV through the release of CXCL4, which behaves as a broad-spectrum inhibitor, directly interacting with the major viral envelope glycoprotein, gp120, and thereby suppressing attachment and entry [200]. Unlike the other CXCR3 ligands, CXCL4 thus would be ascribed a protective role, supported by an inverse correlation between CXCL4 serum levels in HIV-1-infected patients and the clinical stage of their disease [200].

On a side note, CXCR3 ligands also seem to be involved in opportunistic infections manifesting in HIV-positive patients. In this context, the IFN- γ -inducible chemokines, especially CXCL10, have been linked to hepatitis C [202-205], tuberculosis [206, 207] and even malaria [208].

3.10. Hepatitis

In general, CXCR3-binding chemokines have been ascribed direct antimicrobial activity, mostly directed towards bacteria and malaria parasites [178, 182, 209, 210]. However their involvement in leukocyte recruitment and differentiation places the CXCR3 ligands at the front line of the host defense against different infectious diseases, including viral infections of the central nervous system [211, 212]. A link was also established between the IFN- γ -inducible chemokines and viral hepatitis, mostly hepatitis C [213]. However, the hepatitis C virus (HCV) can modulate the chemokine system, upregulating inflammatory chemokines yet subverting a specific immune response, postulated to result in the establishment of chronic low-grade inflammation and an impaired viral clearance [213, 214].

Both CXCL9 and CXCL10 have been reported to facilitate accumulation of T lymphocytes in the infected liver leading to patchy lobular inflammation, an important predictor of progressive liver injury [215-217]. Elevated CXCL10 plasma levels indeed mark hepatitis C patients with advanced fibrosis, whereas CXCL9 levels according to some studies mostly associate with advanced inflammation [218]. It has even been reported that CXCL9 may exert antifibrotic effects [219]. Additionally, Sahin and colleagues recently suggested CXCL10 has a direct pro-apoptotic effect on hepatocytes as well, potentially contributing to CXCL10-associated liver injury [220].

CXCL11 is also expressed by hepatocytes in chronic hepatitis C and promotes recruitment of pro-inflammatory T cells and subsequent portal and lobular inflammation [221]. Its intrahepatic expression may be mildly elevated in chronic hepatitis C patients with higher necroinflammation and fibrosis [222].

Finally, Zaldivar and colleagues revealed that patients with advanced HCV-induced fibrosis present increased CXCL4 serum concentrations and intrahepatic CXCL4 mRNA levels [223]. Also in non-viral, alcoholic hepatitis, increased expression of CXCL4 and CXCL10 in patient liver biopsies was reported [224]. Similarly, elevated serum concentrations and intrahepatic expression levels of CXCL9 and CXCL10 were shown to be positively associated with severity of liver fibrosis in chronic liver diseases of different etiologies, viral or otherwise [225].

3.11. Atherosclerosis

The initiation and progression of atherosclerosis, an arterial narrowing caused by the formation of progressive and degenerative plaques, is fundamentally intertwined with the chemoattraction and modulation of immune cells, most importantly monocytes and lymphocytes. Activated T lymphocytes accumulate in the atheroma early on, recruited at least in part by locally expressed CXCL9, CXCL10 and CXCL11 [226]. Different studies have indeed highlighted the effect of these chemokines on lesional T cell infiltration and as such on the consequent progression of atherosclerotic plaques [227-229]. On the other hand, CXCL4 differs from the others in that this chemokine mostly retains peripheral monocytes at the lesion site, where this proatherogenic chemokine is released at high concentrations following platelet activation [230]. Heterodimerization of CXCL4 with another platelet chemokine, CCL5/RANTES, even further enhances monocyte recruitment [231]. As reviewed by Aidoudi and Bikfalvi, many diverse regulatory mechanisms are involved in the achieved impact of CXCL4 on the progression of atherosclerosis [232]. CXCL4 instigates local monocyte differentiation to macrophages and controls their polarization [22, 233]. Remarkably, this platelet factor also modulates the low-density lipoprotein (LDL) cholesterol metabolism by inhibiting degradation of the LDL-receptor. CXCL4 simultaneously prevents native LDL to bind to its receptor, but selectively promotes the macrophage's uptake and following esterification of oxidized LDL, a crucial step in the development of foam cells which form the basis for future atherosclerotic plaques [232-234]. Accordingly, lesional CXCL4 deposition is shown to correlate with the clinical manifestation of atherosclerosis [235]. This enforces the hypothesis that CXCR3 ligands, though each via specific mechanisms, collectively contribute to the progression of atherosclerosis to a symptomatic disease [227, 235, 236].

3.12. Alzheimer's disease

Alzheimer's disease (AD) is a neurodegenerative disorder in which deposition of neurotoxic β -amyloid peptides, forming senile plaques, as well as formation of neurofibrillary tangles cause brain lesions leading to severe dementia [237]. Typically, AD patients also present with neuroinflammation. The physiological role of chemokines and their receptors in the brain has been established before, but

even more so are they considered to be associated with the pathogenesis of various neurological diseases, including AD [238]. Reactive astrocytes in AD were reported to display elevated CXCL10 expression and CXCL10- as well as CXCL9-induced signaling was confirmed in CXCR3-expressing neurons [239]. These findings were corroborated by reports of CXCL10-mediated calcium dysregulation and apoptosis in neurons, and thus neurotoxicity [240]. CXCL10 has also been reported to regulate migration of astrocytes themselves thereby promoting astrocyte aggregation around senile plaques [241]. Furthermore, CXCL10 concentrations in the CSF, as opposed to serum levels, were increased in AD patients, specifically during the early phase of disease, characterized by a mild cognitive decline and typically associated with more pronounced inflammatory events [242-244]. Increased CXCL10 expression was also observed in the brain of transgenic AD mice in which colocalization of β -amyloid and CXCL10 immunoreactivity was illustrated [245]. Taken together, evidence strongly suggests a causative role for CXCL10 in the pathogenesis of AD, possibly extendable to other neurodegenerative disorders.

On the other hand, involvement of CXCL9 remains controversial as opposing reports regarding CXCL9 plasma levels in AD patients have been published [246, 247]. Additionally, considering the role of platelet activation, CXCL4 has also been suggested to support chronic inflammation in AD patients, though this hypothesis remains to be further investigated [248, 249].

3.13. Thrombocytopenia

Most of the CXCR3 ligands display a remarkable affinity for glycosaminoglycans, though none as outspoken as CXCL4. Glycosaminoglycans are believed to act as co-receptors for CXCR3 ligands, particularly in the process of leukocyte rolling and efficient tissue intravasation [250]. Yet the formation of CXCL4/heparin complexes upon administration of heparin as an anti-coagulant can also lead to a rare and paradoxical complication as the odd patient will develop antibodies against a CXCL4/heparin-epitope in the vicinity of the platelet surface [251]. This complication is referred to as heparin-induced thrombocytopenia (HIT). Patients developing HIT are at risk of venous as well as arterial thrombosis, despite low platelet counts [252].

4. CXCR3 LIGANDS IN THERAPY

A great range of pathologies, mostly characterized by inflammation or vascular effects, have clearly been associated with changes in the expression levels of CXCR3 and CXCR3-binding chemokines (summarized in Table 2). More importantly, often the expression of these chemokine ligands and receptor correlates to pathogenesis, disease progression and prognosis or, under particular conditions, could reflect a failing compensation mechanism. Insight into CXCR3 biology has consequently inspired the development of new therapeutics and continues to highlight new avenues worth exploring. Generally speaking, most CXCR3 axis-based therapies can be assigned to either the anti-inflammatory approach, by preventing CXCR3 activation, or an angiostatic approach, through the administration of exogenous CXCR3 ligands or peptides derived thereof. Possibly, this CXCR3 activation may also promote effective immune responses.

4.1. CXCR3 ligands as angiostatic and anti-tumoral therapeutics

Cancer is a prototypical angiogenic disorder in which neovascularization plays a crucial role in tumor progression and thus offers a promising therapeutic target. The angiostatic chemokines may represent an opportunity in the optimization of current anti-angiogenic regimes. Earlier in this review, we have elaborated on the anti-tumoral characteristics of CXCR3-binding chemokines, granted the malignant cells themselves do not express the CXCR3A splice variant. Though currently clinical use has not yet been tested, a wide selection of publications has discussed the benefits of CXCR3-based anti-tumoral strategies in various preclinical mouse models (Table 3).

Numerous experimental approaches have been described: different cancer models, combinations with other therapeutics, and administration strategies ranging from intratumoral chemokine injections to treatment with viral vectors [253]. Moreover, use of the CXCR3 ligands is also being fine-tuned by testing the impact of different alterations on their anti-tumoral activity. Carboxy-terminal peptides derived from the platelet factors [254, 255], even a chimeric chemokine integrating aspects of both CXCL10 and CXCL11 [256] or synthetic peptides such as anginex, which contains structural elements of CXCL4, CXCL8 as well as bactericidal permeability-increasing protein [257], have been tested to

optimize therapeutic efficacy (not included in table). Commonly, despite some exceptions, administration of CXCR3 ligands or derived preparations has shown innovative therapeutic benefit, holding promise for the future of anti-angiogenic cancer treatments.

Besides their application in cancer, administration of angiostatic chemokines like the CXCR3 ligands may also benefit patients suffering from many other angiogenic disorders, for example pathological ocular neovascularization in diabetic retinopathy.

4.2. CXCR3 ligands as targets in anti-inflammatory therapy

As IFN- γ -inducible, T cell-attracting chemokines, CXCL9, CXCL10 and CXCL11 most definitely play an important role in host immunity, yet they can also instigate endless amplification loops leading the way for excessive, detrimental immune responses [89]. Inflammatory disorders could benefit from anti-inflammatory agents inhibiting CXCR3 ligands and thereby breaking the cycle. Human monoclonal antibodies against CXCL10 have been developed for clinical use (MDX-1100/BMS-936557 and NI-0801; ClinicalTrials.gov). Thus far, use of the anti-CXCL10 antibody BMS-936557 has been examined in the treatment of IBD and RA patients and phase II trials have been completed reporting clinical efficacy in patients who previously responded inadequately to methotrexate [258, 259]. Treatment was generally well tolerated in both patient populations. Studies in cirrhosis patients are also ongoing, with a phase II trial just recently having been finalized (Study ID NCT01430429; ClinicalTrials.gov). CXCL10 could be considered the more clinically prominent of inflammatory CXCR3 ligands and therefore an obvious therapeutic target, however other chemokines of this group represent valuable alternatives. For example, CXCL4 takes on an interesting role in atherosclerosis and has been proposed as a promising therapeutic target as well [260, 261]. Other inflammatory disorders that may benefit from CXCR3 ligand blockade include type 1 diabetes, multiple sclerosis, allograft rejection, *etc.* [262, 263].

4.3. CXCR3 ligands as prognostic and predictive biomarkers

Finally, in many diseases, CXCR3 ligands provide an accessible readout for patient diagnosis, prognosis or therapy responsiveness. Prognostic significance of CXCL4L1 serum concentrations, for

example, was evidenced in stable coronary artery disease, in that lower CXCL4L1 levels were associated with a higher event rate and overall worse outcome [264]. In both Crohn's disease and RA, high plasma levels of CXCL4 have been shown to predict non-responsiveness to anti-TNF α antibody (infliximab) therapy [265, 266]. CXCR3 ligands have also been identified as crucial instigators of transplant inflammation and following allograft rejection [138]. Their expression, both pre- and post-transplantation, is a reflection of the patient's immune status. Urinary CXCL9 and CXCL10 have accordingly been described as useful, non-invasive biomarkers to predict or diagnose allograft rejection before overt clinical manifestation or irreversible graft injury [139, 146]. In the context of HCV, an association was noted between, on one hand, either intrahepatic or plasma levels of CXCL10 and, on the other hand, the patient's overall response to pegylated IFN- α 2 and ribavirin combination therapy [267, 268]. Low baseline levels of CXCL10 proved to indicate a better outcome. These are just a few examples supporting the use of CXCR3 ligands as biomarkers to optimize individual treatment strategies.

Of note, interpretation of CXCL10 concentrations may be complicated considering recent reports on the detection of natural antagonistic forms of CXCL10, as observed in chronically HCV-infected patients [269]. Upon HCV infection, plasma CXCL10 is truncated by dipeptidyl peptidase IV (DPPIV)/CD26, generating its antagonistic form and thus interfering with its function as a pro-inflammatory leukocyte chemoattractant [270]. Amino-terminal truncation of the IFN- γ -inducible CXCR3 ligands by DPPIV was previously evidenced *in vitro* and indeed impaired their chemotactic activity, whilst retaining angiostatic potential [271]. Discovery of this functional difference between intact CXCL10 and its naturally occurring, amino-terminally processed form, challenges the significance of total, non-discriminatory CXCL10 concentrations detected in patients.

5. SUMMARY

CXCL4, CXCL4L1, CXCL9, CXCL10 and CXCL11 all bind, with varying affinity, to different splice variants of the CXCR3 receptor and are therefore collectively referred to as the CXCR3 ligands. Their pleiotropic character, regulating both leukocyte migration and angiogenesis, implicates these chemokines in a broad spectrum of diseases and their pathogenesis. Consequently, the CXCR3 ligands

have attracted attention as new therapeutic targets. However, the field is still lacking fundamental understanding of some of these chemokines. Ongoing research and clinical trials strive towards a better understanding of their basic biology, optimization of drug safety as well as efficacy. In conclusion, the CXCR3 ligands and their inhibitors represent a new generation of anti-angiogenic and anti-inflammatory strategies.

ACKNOWLEDGEMENTS

The authors would like to thank granted support by the Fund for Scientific Research of Flanders (FWO-Vlaanderen project G.0764.14 and G.0773.13), the Interuniversity Attraction Poles Programme initiated by the Belgian Science Policy Office (I.A.P. project P7/40) and the Concerted Research Actions of the Regional Government of Flanders (GOA13/014).

FIGURE LEGENDS

Fig. 1 Receptor binding by the IFN- γ -inducible chemokines, CXCL4 and CXCL4L1

CXCL4, CXCL4L1, CXCL9, CXCL10 and CXCL11, collectively acknowledged as the CXCR3 ligands, all bind both receptor splice variants CXCR3A and CXCR3B, whereas only CXCL11 binds the more distinctive CXCR3-alt [11-15]. CXCL11 also uniquely shows affinity for atypical chemokine receptor (ACKR) 3/CXCR7 [272]. On the other hand, both CXCL4 and CXCL11 bind to ACKR1/DARC [273, 274]. To some extent, all CXCR3-binding chemokines also display affinity for glycosaminoglycan branches on proteoglycans [16].

* Binding of CXCL4L1 to DARC has not yet been evaluated.

CXCL4	1	EAEEDGDLQCLCVKTTSQ	VRPRHITSLEVIKAGPHCPTAQLIATL	KNGRKICLDLQAPLY	60
CXCL4L1	1	EAEEDGDLQCLCVKTTSQ	VRPRHITSLEVIKAGPHCPTAQLIATL	KNGRKICLDLQALLY	60
CXCL9	1	-TPVVRKGRCS	CISTNQGTIHLQSLKDLKQFAPSPSCEKIEIIATL	KNGVQTCLNPDSADV	60
CXCL10	1	-VPLSRTVRCTCISISNQPVNPRSLEKLEIIPASQFCPRVEIIATM	KKKGEKRCLNPESKAI		61
CXCL11	1	-FPMFKRGRCLCIGPGVKAVKVADIEKASIMYPSNNCDKIEVIITL	KENKGQRCLNPKSKQA		61

CXCL4	61	KKIIK	KL	LES	70	37% identity	
CXCL4L1	61	KKIIK	EH	LES	70	37% identity	
CXCL9	61	KELIKKWEKQVSQ	KKKQKNGKKHQQKKV	LKVRSQ	RSRQKKTT	103	41% identity
CXCL10	62	KNLLKAVSKERSKRSP				77	100% identity
CXCL11	62	RLIIKK	VER	KNF		73	34% identity

Gap AA Substitution CXCL4 versus CXCL4L1
 % Sequence AA identity compared to CXCL10 (100%)

Fig. 2 Amino acid sequence alignment of the mature CXCR3-binding chemokines

The amino acid sequences of mature CXCL4, CXCL4L1, CXCL9 and CXCL11, without their respective signal peptides, were optimally aligned with respect to the mature CXCL10 chemokine. Their sequence identity compared to CXCL10 was calculated to give a general impression of the sequence homology among the CXCR3-binding chemokines. Mismatched amino acids (AA) are marked in grey. Gaps are marked in yellow. The differences between CXCL4 and CXCL4L1 are additionally marked in red.

Fig. 3 The anti-tumoral activity of CXCR3 ligands

The CXCR3 ligands affect the tumor environment in various ways. Their anti-tumoral activity is the net result of their angiostatic effect, preventing tumor neovascularization (A), of the recruitment of pro-inflammatory lymphocytes, promoting tumor immunosurveillance (B), and finally, of their lymphangiostatic effect, preventing metastatic escape of tumor cells via the lymphatic vasculature (C) [31, 51].

Fig. 4 The effect of CXCR3 ligands on tumor invasiveness

CXCR3 ligands may exert a direct effect on CXCR3-positive tumor cells, the nature of which depends on the specific splice variant expressed [59]. In case of CXCR3A expression, these chemokines will stimulate not only proliferation, but also migration of the CXCR3A-expressing tumor cells, especially to other peripheral sources of CXCR3 ligands, thereby promoting and directing tumor metastasis both to draining lymph nodes and distant organs, e.g. lung [275].

TABLES

Table 1. Fundamental characteristics of the human CXCR3 ligands

	CXCL4	CXCL4L1	CXCL9	CXCL10	CXCL11
Gene (HGNC)	NM_002619 4q13.3	NM_002620 4q13.3	NM_002416 4q21.1	NM_001565 4q21.1	NM_005409 4q21.1
Mature protein	70 AA	70 AA	103 AA	77 AA	73 AA
Producer cells	Platelets Monocytes T cells	Platelets Smooth muscle cells	HMVEC Fibroblasts Macrophages PBMC	HMVEC Fibroblasts Monocytes T cells Keratinocytes	HMVEC Fibroblasts Astrocytes PBMC
Inducers	Thrombin (platelets) Phorbol ester (T cells)	Thrombin (platelets)	IFN- γ	TNF- α (fibroblasts) IFN- α , IFN- β , IFN- γ	TNF- α (fibroblasts) IFN- γ , IFN- β
Processing					
- protease	N.D.	N.D.	DPPIV/CD26	DPPIV/CD26	DPPIV/CD26
- peptide product	CXCL4 ¹⁷⁻⁷⁰	N.D.	CXCL9 ³⁻¹⁰³	CXCL10 ³⁻⁷⁷	CXCL11 ³⁻⁷³
- effect on:					
chemotaxis	N.D.		↓	↓	↓
angiostasis	↑		=	=	=
Ref	[44, 276]	[43, 44]	[37, 39, 41, 271]	[38-40, 271, 277]	[12, 39, 41, 271]

HGNC = HUGO gene nomenclature committee; AA = amino acids; N.D. = not determined; HMVEC = Human microvascular endothelial cell; PBMC = Peripheral blood mononuclear cell; TNF = Tumor necrosis factor; DPPIV = Dipeptidyl peptidase IV

Table 2. CXCR3 ligands in disease

Disease	CXCR3 ligand expression ^a	Function of CXCR3 ligands ^b
CANCER		
Carcinoma and sarcoma	↑ / ↓	Recruitment of anti-tumoral lymphocytes via CXCR3A (CXCL9, CXCL10, CXCL11) Angiostatic (all ligands) and lymphangiostatic (CXCL4, CXCL4L1) via CXCR3B
	↑	Increased tumor invasiveness via CXCR3A (CXCL4, CXCL10)
Hematopoietic malignancies	↑	Increased tumor cell dissemination (CXCL9, CXCL10, CXCL11) Anti-tumoral activity: tumor tissue necrosis, vascular damage, protective immune response (CXCL9, CXCL10)
	↓	Tumor suppressor gene (CXCL4)
INFLAMMATORY DISORDERS		
Arthritis	↑	Synovial inflammation (CXCL4, CXCL9, CXCL10) and vascular lesions (CXCL4)
IBD	↑	Mucosal inflammation (CXCL9, CXCL10, CXCL11)
Diabetes	↑	T cell islet infiltration (CXCL9, CXCL10) Decreased pancreatic beta cell proliferation (CXCL10) Correlation to angiogenic complication PDR (CXCL4, CXCL9, CXCL10)
Systemic sclerosis	↑	Fibrotic complications in the skin or lung (CXCL4, CXCL10)
	↓	Pulmonary function decline (CXCL11)
Transplant rejection	↑	Biomarkers predicting allograft failure (CXCL9, CXCL10, CXCL11)
	N.A.	Negative regulation of T cell reactivity (CXCL4)
MICROBIAL PATHOLOGIES		
Malaria	↑	Positive association with CM occurrence and mortality (CXCL4, CXCL9, CXCL10) CD8+ recruitment to the brain in murine CM (CXCL9, CXCL10)
	↑	Intraerythrocytic parasite-killing (CXCL4)
AIDS	↑	T cell recruitment and virus propagation (CXCL9, CXCL10, CXCL11) Stimulated virus replication (CXCL9, CXCL10, CXCL11) Inhibited host cell attachment and entry (CXCL4)
Hepatitis	↑	Lobular inflammation and liver fibrosis (CXCL4, CXCL9, CXCL10, CXCL11)
OTHER		
Atherosclerosis	↑	Lesional T cell infiltration (CXCL9, CXCL10, CXCL11) Monocyte recruitment and macrophage polarization (CXCL4) Promoted uptake and esterification of oxidized LDL (CXCL4)
Alzheimer's disease	↑	Promoted astrocyte aggregation (CXCL10) Neurotoxicity (CXCL10)
Thrombocytopenia	N.A.	Generation of antibodies against CXCL4/heparin complexes

IBD = Inflammatory bowel disease; PDR = Proliferative diabetic retinopathy; CM = Cerebral malaria;
AIDS = Acquired immune deficiency syndrome; N.A. = Not applicable

^a ↑ / ↓ = Upregulated or downregulated CXCR3 ligand expression, respectively, depending on the study

^b For references *cfr.* citations in text

Table 3. Anti-tumoral effects of CXCR3 ligands: combination or monotherapies in preclinical models

Mouse model	CXCL4	CXCL4L1	CXCL9	CXCL10	CXCL11	Ref
B16 melanoma	✓	✓		✓		[278-282]
Lung adenocarcinoma (A11, A549, LLC)		✓	✓	✓		[278, 280, 283-286]
Colon adenocarcinoma (CT26, HCT116)	✓		✓	✓		[281, 284, 286, 287]
Capan-1 pancreatic adenocarcinoma		✓				[51]
Renal cell carcinoma (RENCA)			✓			[288]
Breast carcinoma (66.1, TA3)	✓		✓			[289, 290]
Ovarian carcinoma (SKOV3, ES2)	✓			✓		[290, 291]
Glioma (GL261, U87)	✓			✓		[292-294]
Malignant peripheral nerve sheath tumor	✓					[293, 295]
Meth A fibrosarcoma				✓		[296]
Angiosarcoma (H5V)				✓		[297]
Burkitt's lymphoma			✓			[298]
EL4 lymphoma					✓	[299]

Reference List

- [1] Rosenkilde MM, Schwartz TW. The chemokine system - a major regulator of angiogenesis in health and disease. *APMIS* 2004;112:481-95.
- [2] Zlotnik A, Yoshie O. Chemokines: a new classification system and their role in immunity. *Immunity* 2000;12:121-7.
- [3] Groom JR, Luster AD. CXCR3 ligands: redundant, collaborative and antagonistic functions. *Immunol Cell Biol* 2011;89:207-15.
- [4] Bachelier F, Graham GJ, Locati M, Mantovani A, Murphy PM, Nibbs R et al. New nomenclature for atypical chemokine receptors. *Nat Immunol* 2014;15:207-8.
- [5] Belperio JA, Keane MP, Arenberg DA, Addison CL, Ehlert JE, Burdick MD et al. CXC chemokines in angiogenesis. *J Leukoc Biol* 2000;68:1-8.
- [6] Bosisio D, Salvi V, Gagliostro V, Sozzani S. Angiogenic and antiangiogenic chemokines. *Chem Immunol Allergy* 2014;99:89-104.
- [7] Strieter RM, Polverini PJ, Kunkel SL, Arenberg DA, Burdick MD, Kasper J et al. The functional role of the ELR motif in CXC chemokine-mediated angiogenesis. *J Biol Chem* 1995;270:27348-57.
- [8] Addison CL, Daniel TO, Burdick MD, Liu H, Ehlert JE, Xue YY et al. The CXC chemokine receptor 2, CXCR2, is the putative receptor for ELR+ CXC chemokine-induced angiogenic activity. *J Immunol* 2000;165:5269-77.
- [9] Holmes WE, Lee J, Kuang WJ, Rice GC, Wood WI. Structure and functional expression of a human interleukin-8 receptor. *Science* 1991;253:1278-80.
- [10] Dai X, Tan Y, Cai S, Xiong X, Wang L, Ye Q et al. The role of CXCR7 on the adhesion, proliferation and angiogenesis of endothelial progenitor cells. *J Cell Mol Med* 2011;15:1299-309.
- [11] Lasagni L, Francalanci M, Annunziato F, Lazzeri E, Giannini S, Cosmi L et al. An alternatively spliced variant of CXCR3 mediates the inhibition of endothelial cell growth induced by IP-10, Mig, and I-TAC, and acts as functional receptor for platelet factor 4. *J Exp Med* 2003;197:1537-49.
- [12] Cole KE, Strick CA, Paradis TJ, Ogborne KT, Loetscher M, Gladue RP et al. Interferon-inducible T cell alpha chemoattractant (I-TAC): a novel non-ELR CXC chemokine with potent activity on activated T cells through selective high affinity binding to CXCR3. *J Exp Med* 1998;187:2009-21.
- [13] Loetscher M, Gerber B, Loetscher P, Jones SA, Piali L, Clark-Lewis I et al. Chemokine receptor specific for IP10 and mig: structure, function, and expression in activated T-lymphocytes. *J Exp Med* 1996;184:963-9.
- [14] Ehlert JE, Addison CA, Burdick MD, Kunkel SL, Strieter RM. Identification and partial characterization of a variant of human CXCR3 generated by posttranscriptional exon skipping. *J Immunol* 2004;173:6234-40.

- [15] Struyf S, Salogni L, Burdick MD, Vandercappellen J, Gouwy M, Noppen S et al. Angiostatic and chemotactic activities of the CXC chemokine CXCL4L1 (platelet factor-4 variant) are mediated by CXCR3. *Blood* 2011;117:480-8.
- [16] Handel TM, Johnson Z, Crown SE, Lau EK, Proudfoot AE. Regulation of protein function by glycosaminoglycans - as exemplified by chemokines. *Annu Rev Biochem* 2005;74:385-410.
- [17] Luster AD, Greenberg SM, Leder P. The IP-10 chemokine binds to a specific cell surface heparan sulfate site shared with platelet factor 4 and inhibits endothelial cell proliferation. *J Exp Med* 1995;182:219-31.
- [18] Shute J. Glycosaminoglycan and chemokine/growth factor interactions. *Handb Exp Pharmacol* 2012:307-24.
- [19] Kolset SO, Mann DM, Uhlin-Hansen L, Winberg JO, Ruoslahti E. Serglycin-binding proteins in activated macrophages and platelets. *J Leukoc Biol* 1996;59:545-54.
- [20] Pervushina O, Scheuerer B, Reiling N, Behnke L, Schroder JM, Kasper B et al. Platelet factor 4/CXCL4 induces phagocytosis and the generation of reactive oxygen metabolites in mononuclear phagocytes independently of Gi protein activation or intracellular calcium transients. *J Immunol* 2004;173:2060-7.
- [21] Petersen F, Bock L, Flad HD, Brandt E. A chondroitin sulfate proteoglycan on human neutrophils specifically binds platelet factor 4 and is involved in cell activation. *J Immunol* 1998;161:4347-55.
- [22] Scheuerer B, Ernst M, Durrbaum-Landmann I, Fleischer J, Grage-Griebenow E, Brandt E et al. The CXC-chemokine platelet factor 4 promotes monocyte survival and induces monocyte differentiation into macrophages. *Blood* 2000;95:1158-66.
- [23] Loetscher M, Loetscher P, Brass N, Meese E, Moser B. Lymphocyte-specific chemokine receptor CXCR3: regulation, chemokine binding and gene localization. *Eur J Immunol* 1998;28:3696-705.
- [24] Groom JR, Luster AD. CXCR3 in T cell function. *Exp Cell Res* 2011;317:620-31.
- [25] Rabin RL, Park MK, Liao F, Swofford R, Stephany D, Farber JM. Chemokine receptor responses on T cells are achieved through regulation of both receptor expression and signaling. *J Immunol* 1999;162:3840-50.
- [26] Bonecchi R, Bianchi G, Bordignon PP, D'Ambrosio D, Lang R, Borsatti A et al. Differential expression of chemokine receptors and chemotactic responsiveness of type 1 T helper cells (Th1s) and Th2s. *J Exp Med* 1998;187:129-34.
- [27] Szabo SJ, Kim ST, Costa GL, Zhang X, Fathman CG, Glimcher LH. A novel transcription factor, T-bet, directs Th1 lineage commitment. *Cell* 2000;100:655-69.
- [28] Taqueti VR, Grabie N, Colvin R, Pang H, Jarolim P, Luster AD et al. T-bet controls pathogenicity of CTLs in the heart by separable effects on migration and effector activity. *J Immunol* 2006;177:5890-901.

- [29] Koch MA, Tucker-Heard G, Perdue NR, Killebrew JR, Urdahl KB, Campbell DJ. The transcription factor T-bet controls regulatory T cell homeostasis and function during type 1 inflammation. *Nat Immunol* 2009;10:595-602.
- [30] Garcia-Lopez MA, Sanchez-Madrid F, Rodriguez-Frade JM, Mellado M, Acevedo A, Garcia MI et al. CXCR3 chemokine receptor distribution in normal and inflamed tissues: expression on activated lymphocytes, endothelial cells, and dendritic cells. *Lab Invest* 2001;81:409-18.
- [31] Billottet C, Quemener C, Bikfalvi A. CXCR3, a double-edged sword in tumor progression and angiogenesis. *Biochim Biophys Acta* 2013;1836:287-95.
- [32] Colvin RA, Campanella GS, Sun J, Luster AD. Intracellular domains of CXCR3 that mediate CXCL9, CXCL10, and CXCL11 function. *J Biol Chem* 2004;279:30219-27.
- [33] Dagan-Berger M, Feniger-Barish R, Avniel S, Wald H, Galun E, Grabovsky V et al. Role of CXCR3 carboxyl terminus and third intracellular loop in receptor-mediated migration, adhesion and internalization in response to CXCL11. *Blood* 2006;107:3821-31.
- [34] Korniejewska A, McKnight AJ, Johnson Z, Watson ML, Ward SG. Expression and agonist responsiveness of CXCR3 variants in human T lymphocytes. *Immunology* 2011;132:503-15.
- [35] Xanthou G, Williams TJ, Pease JE. Molecular characterization of the chemokine receptor CXCR3: evidence for the involvement of distinct extracellular domains in a multi-step model of ligand binding and receptor activation. *Eur J Immunol* 2003;33:2927-36.
- [36] Colvin RA, Campanella GS, Manice LA, Luster AD. CXCR3 requires tyrosine sulfation for ligand binding and a second extracellular loop arginine residue for ligand-induced chemotaxis. *Mol Cell Biol* 2006;26:5838-49.
- [37] Farber JM. A macrophage mRNA selectively induced by gamma-interferon encodes a member of the platelet factor 4 family of cytokines. *Proc Natl Acad Sci U S A* 1990;87:5238-42.
- [38] Luster AD, Ravetch JV. Biochemical characterization of a gamma interferon-inducible cytokine (IP-10). *J Exp Med* 1987;166:1084-97.
- [39] Loos T, Dekeyser L, Struyf S, Schutyser E, Gijsbers K, Gouwy M et al. TLR ligands and cytokines induce CXCR3 ligands in endothelial cells: enhanced CXCL9 in autoimmune arthritis. *Lab Invest* 2006;86:902-16.
- [40] Proost P, Vynckier AK, Mahieu F, Put W, Grillet B, Struyf S et al. Microbial Toll-like receptor ligands differentially regulate CXCL10/IP-10 expression in fibroblasts and mononuclear leukocytes in synergy with IFN-gamma and provide a mechanism for enhanced synovial chemokine levels in septic arthritis. *Eur J Immunol* 2003;33:3146-53.
- [41] Proost P, Verpoest S, Van de Borne K, Schutyser E, Struyf S, Put W et al. Synergistic induction of CXCL9 and CXCL11 by Toll-like receptor ligands and interferon-gamma in fibroblasts correlates with elevated levels of CXCR3 ligands in septic arthritis synovial fluids. *J Leukoc Biol* 2004;75:777-84.
- [42] Kuo JH, Chen YP, Liu JS, Dubrac A, Quemener C, Prats H et al. Alternative C-Terminal Helix Orientation Alters Chemokine Function: structure of the anti-angiogenic chemokine, CXCL4L1. *J Biol Chem* 2013;288:13522-33.

- [43] Struyf S, Burdick MD, Proost P, Van Damme J, Strieter RM. Platelets release CXCL4L1, a nonallelic variant of the chemokine platelet factor-4/CXCL4 and potent inhibitor of angiogenesis. *Circ Res* 2004;95:855-7.
- [44] Lasagni L, Grepin R, Mazzinghi B, Lazzeri E, Meini C, Sagrinati C et al. PF-4/CXCL4 and CXCL4L1 exhibit distinct subcellular localization and a differentially regulated mechanism of secretion. *Blood* 2007;109:4127-34.
- [45] Balestrieri ML, Balestrieri A, Mancini FP, Napoli C. Understanding the immunoangiostatic CXC chemokine network. *Cardiovasc Res* 2008;78:250-6.
- [46] Vandercappellen J, Van Damme J, Struyf S. The role of the CXC chemokines platelet factor-4 (CXCL4/PF-4) and its variant (CXCL4L1/PF-4var) in inflammation, angiogenesis and cancer. *Cytokine Growth Factor Rev* 2011;22:1-18.
- [47] Vandercappellen J, Van Damme J, Struyf S. The role of CXC chemokines and their receptors in cancer. *Cancer Lett* 2008;267:226-44.
- [48] Folkman J, Merler E, Abernathy C, Williams G. Isolation of a tumor factor responsible for angiogenesis. *J Exp Med* 1971;133:275-88.
- [49] Bergers G, Benjamin LE. Tumorigenesis and the angiogenic switch. *Nat Rev Cancer* 2003;3:401-10.
- [50] Cao Y. Opinion: emerging mechanisms of tumour lymphangiogenesis and lymphatic metastasis. *Nat Rev Cancer* 2005;5:735-43.
- [51] Prats AC, Van den Berghe L, Rayssac A, Ainaoui N, Morfoisse F, Pujol F et al. CXCL4L1-fibstatin cooperation inhibits tumor angiogenesis, lymphangiogenesis and metastasis. *Microvasc Res* 2013;89:25-33.
- [52] Sica A. Role of tumour-associated macrophages in cancer-related inflammation. *Exp Oncol* 2010;32:153-8.
- [53] Lakshmi Narendra B, Eshvendar RK, Shantikumar S, Ramakrishna S. Immune system: a double-edged sword in cancer. *Inflamm Res* 2013;62:823-34.
- [54] Verbeke H, Struyf S, Laureys G, Van Damme J. The expression and role of CXC chemokines in colorectal cancer. *Cytokine Growth Factor Rev* 2011;22:345-58.
- [55] Goldberg-Bittman L, Sagi-Assif O, Meshel T, Nevo I, Levy-Nissenbaum O, Yron I et al. Cellular characteristics of neuroblastoma cells: regulation by the ELR⁻-CXC chemokine CXCL10 and expression of a CXCR3-like receptor. *Cytokine* 2005;29:105-17.
- [56] Suyama T, Furuya M, Nishiyama M, Kasuya Y, Kimura S, Ichikawa T et al. Up-regulation of the interferon gamma (IFN-gamma)-inducible chemokines IFN-inducible T-cell alpha chemoattractant and monokine induced by IFN-gamma and of their receptor CXC receptor 3 in human renal cell carcinoma. *Cancer* 2005;103:258-67.
- [57] Goldberg-Bittman L, Neumark E, Sagi-Assif O, Azenshtein E, Meshel T, Witz IP et al. The expression of the chemokine receptor CXCR3 and its ligand, CXCL10, in human breast adenocarcinoma cell lines. *Immunol Lett* 2004;92:171-8.

- [58] Robledo MM, Bartolome RA, Longo N, Rodriguez-Frade JM, Mellado M, Longo I et al. Expression of functional chemokine receptors CXCR3 and CXCR4 on human melanoma cells. *J Biol Chem* 2001;276:45098-105.
- [59] Wu Q, Dhir R, Wells A. Altered CXCR3 isoform expression regulates prostate cancer cell migration and invasion. *Mol Cancer* 2012;11.
- [60] Ma X, Norsworthy K, Kundu N, Rodgers WH, Gimotty PA, Goloubeva O et al. CXCR3 expression is associated with poor survival in breast cancer and promotes metastasis in a murine model. *Mol Cancer Ther* 2009;8:490-8.
- [61] Kawada K, Hosogi H, Sonoshita M, Sakashita H, Manabe T, Shimahara Y et al. Chemokine receptor CXCR3 promotes colon cancer metastasis to lymph nodes. *Oncogene* 2007;26:4679-88.
- [62] Monteagudo C, Martin JM, Jorda E, Llombart-Bosch A. CXCR3 chemokine receptor immunoreactivity in primary cutaneous malignant melanoma: correlation with clinicopathological prognostic factors. *J Clin Pathol* 2007;60:596-9.
- [63] Trentin L, Agostini C, Facco M, Piazza F, Perin A, Siviero M et al. The chemokine receptor CXCR3 is expressed on malignant B cells and mediates chemotaxis. *J Clin Invest* 1999;104:115-21.
- [64] Jones D, Benjamin RJ, Shahsafaei A, Dorfman DM. The chemokine receptor CXCR3 is expressed in a subset of B-cell lymphomas and is a marker of B-cell chronic lymphocytic leukemia. *Blood* 2000;95:627-32.
- [65] Yan XJ, Dozmorov I, Li W, Yancopoulos S, Sison C, Centola M et al. Identification of outcome-correlated cytokine clusters in chronic lymphocytic leukemia. *Blood* 2011;118:5201-10.
- [66] Ocana E, Delgado-Perez L, Campos-Caro A, Munoz J, Paz A, Franco R et al. The prognostic role of CXCR3 expression by chronic lymphocytic leukemia B cells. *Haematologica* 2007;92:349-56.
- [67] Lee Y, Chittezhath M, Andre V, Zhao H, Poidinger M, Biondi A et al. Protumoral role of monocytes in human B-cell precursor acute lymphoblastic leukemia: involvement of the chemokine CXCL10. *Blood* 2012;119:227-37.
- [68] Khandany BK, Hassanshahi G, Khorramdelazad H, Balali Z, Shamsizadeh A, Arababadi MK et al. Evaluation of circulating concentrations of CXCL1 (Gro-alpha), CXCL10 (IP-10) and CXCL12 (SDF-1) in ALL patients prior and post bone marrow transplantation. *Pathol Res Pract* 2012;208:615-9.
- [69] Shi LN, Zhang J, Wu P, Feng K, Li J, Xie ZS et al. Discovery and identification of potential biomarkers of pediatric Acute Lymphoblastic Leukemia. *Proteome Sci* 2009;7.
- [70] Arthur DC, Bloomfield CD, Lindquist LL, Nesbit ME. Translocation 4-11 in Acute Lymphoblastic-Leukemia - Clinical Characteristics and Prognostic-Significance. *Blood* 1982;59:96-9.
- [71] Griffin CA, Emanuel BS, Larocco P, Schwartz E, Poncz M. Human platelet factor 4 gene is mapped to 4q12-q21. *Cytogenet Cell Genet* 1987;45:67-9.

- [72] Bai J, He AL, Zhang WG, Huang C, Yang J, Yang Y et al. Potential biomarkers for adult acute myeloid leukemia minimal residual disease assessment searched by serum peptidome profiling. *Proteome Sci* 2013;11.
- [73] Kim JY, Song HJ, Lim HJ, Shin MG, Kim JS, Kim HJ et al. Platelet factor-4 is an indicator of blood count recovery in acute myeloid leukemia patients in complete remission. *Mol Cell Proteomics* 2008;7:431-41.
- [74] Olsnes AM, Motorin D, Rynningen A, Zaritskey AY, Bruserud O. T lymphocyte chemotactic chemokines in acute myelogenous leukemia (AML): local release by native human AML blasts and systemic levels of CXCL10 (IP-10), CCL5 (RANTES) and CCL17 (TARC). *Cancer Immunol Immunother* 2006;55:830-40.
- [75] Wu B, Wang X, Ma W, Zheng W, Jiang L. Assay of OAZ1 mRNA levels in chronic myeloid leukemia combined with application of leukemia PCR array identified relevant gene changes affected by antizyme. *Acta Haematol* 2014;131:141-7.
- [76] Kim H, Gillis LC, Jarvis JD, Yang S, Huang K, Der S et al. Tyrosine kinase chromosomal translocations mediate distinct and overlapping gene regulation events. *BMC Cancer* 2011;11:528.
- [77] Teichmann M, Meyer B, Beck A, Niedobitek G. Expression of the interferon-inducible chemokine IP-10 (CXCL10), a chemokine with proposed anti-neoplastic functions, in Hodgkin lymphoma and nasopharyngeal carcinoma. *J Pathol* 2005;206:68-75.
- [78] Teruya-Feldstein J, Tosato G, Jaffe ES. The role of chemokines in Hodgkin's disease. *Leuk Lymphoma* 2000;38:363-71.
- [79] Ohshima K, Suefuji H, Karube K, Hamasaki M, Hatano B, Tutiya T et al. Expression of chemokine receptor CXCR3 and its ligand, mig, in gastric and thyroid marginal zone lymphomas. Possible migration and autocrine mechanism. *Leuk Lymphoma* 2003;44:329-36.
- [80] Suefuji H, Ohshima K, Karube K, Kawano R, Nabeshima K, Suzumiya J et al. CXCR3-positive B cells found at elevated frequency in the peripheral blood of patients with MALT lymphoma are attracted by MIG and belong to the lymphoma clone. *Int J Cancer* 2005;114:896-901.
- [81] Venetz D, Ponzoni M, Schiraldi M, Ferreri AJ, Bertoni F, Doglioni C et al. Perivascular expression of CXCL9 and CXCL12 in primary central nervous system lymphoma: T-cell infiltration and positioning of malignant B cells. *Int J Cancer* 2010;127:2300-12.
- [82] Haabeth OA, Lørvik KB, Hammarström C, Donaldson IM, Haraldsen G, Bogen B et al. Inflammation driven by tumour-specific Th1 cells protects against B-cell cancer. *Nat Commun* 2011;2:240.
- [83] Pellegrino A, Antonaci F, Russo F, Merchionne F, Ribatti D, Vacca A et al. CXCR3-binding chemokines in multiple myeloma. *Cancer Lett* 2004;207:221-7.
- [84] Giuliani N, Bonomini S, Romagnani P, Lazzaretti M, Morandi F, Colla S et al. CXCR3 and its binding chemokines in myeloma cells: expression of isoforms and potential relationships with myeloma cell proliferation and survival. *Haematologica* 2006;91:1489-97.
- [85] Barash U, Zohar Y, Wildbaum G, Beider K, Nagler A, Karin N et al. Heparanase enhances myeloma progression via CXCL10 downregulation. *Leukemia* 2014 (in press).

- [86] Cheng SH, Ng MHL, Lau KM, Liu HSY, Chan JCW, Hui ABY et al. 4q loss is potentially an important genetic event in MM tumorigenesis: identification of a tumor suppressor gene regulated by promoter methylation at 4q13.3, platelet factor 4. *Blood* 2007;109:2089-99.
- [87] Liang P, Cheng SH, Cheng CK, Lau KM, Lin SY, Chow EY et al. Platelet factor 4 induces cell apoptosis by inhibition of STAT3 via up-regulation of SOCS3 expression in multiple myeloma. *Haematologica* 2013;98:288-95.
- [88] Han ZC, Lu M, Li J, Defard M, Boval B, Schlegel N et al. Platelet factor 4 and other CXC chemokines support the survival of normal hematopoietic cells and reduce the chemosensitivity of cells to cytotoxic agents. *Blood* 1997;89:2328-35.
- [89] Antonelli A, Ferrari SM, Giuggioli D, Ferrannini E, Ferri C, Fallahi P. Chemokine (C-X-C motif) ligand (CXCL)10 in autoimmune diseases. *Autoimmun Rev* 2014;13:272-80.
- [90] Patel DD, Zachariah JP, Whichard LP. CXCR3 and CCR5 ligands in rheumatoid arthritis synovium. *Clin Immunol* 2001;98:39-45.
- [91] Ruschpler P, Lorenz P, Eichler W, Koczan D, Hanel C, Scholz R et al. High CXCR3 expression in synovial mast cells associated with CXCL9 and CXCL10 expression in inflammatory synovial tissues of patients with rheumatoid arthritis. *Arthritis Res Ther* 2003;5:R241-R252.
- [92] Erdem H, Pay S, Musabak U, Simsek I, Dinc A, Pekel A et al. Synovial angiostatic non-ELR CXC chemokines in inflammatory arthritides: does CXCL4 designate chronicity of synovitis? *Rheumatol Int* 2007;27:969-73.
- [93] Yamamoto T, Chikugo T, Tanaka Y. Elevated plasma levels of beta-thromboglobulin and platelet factor 4 in patients with rheumatic disorders and cutaneous vasculitis. *Clin Rheumatol* 2002;21:501-4.
- [94] Pharoah DS, Varsani H, Tatham RW, Newton KR, de Jager W, Prakken BJ et al. Expression of the inflammatory chemokines CCL5, CCL3 and CXCL10 in juvenile idiopathic arthritis, and demonstration of CCL5 production by an atypical subset of CD8+ T cells. *Arthritis Res Ther* 2006;8:R50.
- [95] Ueno A, Yamamura M, Iwahashi M, Okamoto A, Aita T, Ogawa N et al. The production of CXCR3-agonistic chemokines by synovial fibroblasts from patients with rheumatoid arthritis. *Rheumatol Int* 2005;25:361-7.
- [96] Badot V, Galant C, Nzeusseu Toukap A, Theate I, Maudoux AL, Van den Eynde BJ et al. Gene expression profiling in the synovium identifies a predictive signature of absence of response to adalimumab therapy in rheumatoid arthritis. *Arthritis Res Ther* 2009;11:R57.
- [97] Singh UP, Venkataraman C, Singh R, Lillard JW, Jr. CXCR3 axis: role in inflammatory bowel disease and its therapeutic implication. *Endocr Metab Immune Disord Drug Targets* 2007;7:111-23.
- [98] Yuan YH, ten HT, The FO, Slors JF, van Deventer SJ, te Velde AA. Chemokine receptor CXCR3 expression in inflammatory bowel disease. *Inflamm Bowel Dis* 2001;7:281-6.
- [99] Simi M, Leardi S, Tebano MT, Castelli M, Costantini FM, Speranza V. Raised plasma concentrations of platelet factor 4 (Pf4) in Crohn's disease. *Gut* 1987;28:336-8.

- [100] Vrij AA, Rijken J, van Wersch JWJ, Stockbrugger RW. Platelet factor 4 and beta-thromboglobulin in inflammatory bowel disease and giant cell arteritis. *Eur J Clin Invest* 2000;30:188-94.
- [101] Meuwis MA, Fillet M, Geurts P, de Seny D, Lutteri L, Chapelle JP et al. Biomarker discovery for inflammatory bowel disease, using proteomic serum profiling. *Biochem Pharmacol* 2007;73:1422-33.
- [102] Noguchi A, Watanabe K, Narumi S, Yamagami H, Fujiwara Y, Higuchi K et al. The production of interferon-gamma-inducible protein 10 by granulocytes and monocytes is associated with ulcerative colitis disease activity. *J Gastroenterol* 2007;42:947-56.
- [103] Uguccioni M, Gionchetti P, Robbiani DF, Rizzello F, Peruzzo S, Campieri M et al. Increased expression of IP-10, IL-8, MCP-1, and MCP-3 in ulcerative colitis. *Am J Pathol* 1999;155:331-6.
- [104] Singh UP, Singh S, Iqbal N, Weaver CT, McGhee JR, Lillard JW, Jr. IFN-gamma-inducible chemokines enhance adaptive immunity and colitis. *J Interferon Cytokine Res* 2003;23:591-600.
- [105] Papadakis KA, Prehn J, Zhu D, Landers C, Gaiennie J, Fleshner PR et al. Expression and regulation of the chemokine receptor CXCR3 on lymphocytes from normal and inflammatory bowel disease mucosa. *Inflamm Bowel Dis* 2004;10:778-88.
- [106] Schroepf S, Kappler R, Brand S, Prell C, Lohse P, Glas J et al. Strong overexpression of CXCR3 axis components in childhood inflammatory bowel disease. *Inflamm Bowel Dis* 2010;16:1882-90.
- [107] Lacher M, Kappler R, Berkholz S, Baurecht H, von Schweinitz D, Koletzko S. Association of a CXCL9 polymorphism with pediatric Crohn's disease. *Biochem Biophys Res Commun* 2007;363:701-7.
- [108] Antonelli A, Ferrari SM, Corrado A, Ferrannini E, Fallahi P. CXCR3, CXCL10 and type 1 diabetes. *Cytokine Growth Factor Rev* 2014;25:57-65.
- [109] Shimada A, Morimoto J, Kodama K, Suzuki R, Oikawa Y, Funae O et al. Elevated serum IP-10 levels observed in type 1 diabetes. *Diabetes Care* 2001;24:510-5.
- [110] Nicoletti F, Conget I, Di MM, Di MR, Mazzarino MC, Bendtzen K et al. Serum concentrations of the interferon-gamma-inducible chemokine IP-10/CXCL10 are augmented in both newly diagnosed Type I diabetes mellitus patients and subjects at risk of developing the disease. *Diabetologia* 2002;45:1107-10.
- [111] Antonelli A, Fallahi P, Ferrari SM, Pupilli C, d'Annunzio G, Lorini R et al. Serum Th1 (CXCL10) and Th2 (CCL2) chemokine levels in children with newly diagnosed Type 1 diabetes: a longitudinal study. *Diabet Med* 2008;25:1349-53.
- [112] Rotondi M, Romagnani P, Brozzetti A, Santeusano F, Serio M, Falorni A. Reply to Nicoletti F, Conget L, Di Mauro M et al. (2002) Serum concentrations of the interferon-alpha-inducible chemokine IP-10/CXCL10 are augmented in both newly-diagnosed Type I diabetes mellitus patients and subjects at risk of developing the disease. *Diabetologia* 45:1107-1110. *Diabetologia* 2003;46:1020-1.

- [113] Hanifi-Moghaddam P, Kappler S, Seissler J, Muller-Scholze S, Martin S, Roep BO et al. Altered chemokine levels in individuals at risk of Type 1 diabetes mellitus. *Diabet Med* 2006;23:156-63.
- [114] Rhode A, Pauza ME, Barral AM, Rodrigo E, Oldstone MB, von Herrath MG et al. Islet-specific expression of CXCL10 causes spontaneous islet infiltration and accelerates diabetes development. *J Immunol* 2005;175:3516-24.
- [115] Roep BO, Kleijwegt FS, van Halteren AG, Bonato V, Boggi U, Vendrame F et al. Islet inflammation and CXCL10 in recent-onset type 1 diabetes. *Clin Exp Immunol* 2010;159:338-43.
- [116] Devaraj S, Jialal I. Increased secretion of IP-10 from monocytes under hyperglycemia is via the TLR2 and TLR4 pathway. *Cytokine* 2009;47:6-10.
- [117] Tanaka S, Nishida Y, Aida K, Maruyama T, Shimada A, Suzuki M et al. Enterovirus infection, CXC chemokine ligand 10 (CXCL10), and CXCR3 circuit: a mechanism of accelerated beta-cell failure in fulminant type 1 diabetes. *Diabetes* 2009;58:2285-91.
- [118] Sarkar SA, Lee CE, Victorino F, Nguyen TT, Walters JA, Burrack A et al. Expression and regulation of chemokines in murine and human type 1 diabetes. *Diabetes* 2012;61:436-46.
- [119] Uno S, Imagawa A, Saisho K, Okita K, Iwahashi H, Hanafusa T et al. Expression of chemokines, CXC chemokine ligand 10 (CXCL10) and CXCR3 in the inflamed islets of patients with recent-onset autoimmune type 1 diabetes. *Endocr J* 2010;57:991-6.
- [120] Frigerio S, Junt T, Lu B, Gerard C, Zumsteg U, Hollander GA et al. Beta cells are responsible for CXCR3-mediated T-cell infiltration in insulinitis. *Nat Med* 2002;8:1414-20.
- [121] Morimoto J, Yoneyama H, Shimada A, Shigihara T, Yamada S, Oikawa Y et al. CXC chemokine ligand 10 neutralization suppresses the occurrence of diabetes in nonobese diabetic mice through enhanced beta cell proliferation without affecting insulinitis. *J Immunol* 2004;173:7017-24.
- [122] Schulthess FT, Paroni F, Sauter NS, Shu L, Ribaux P, Haataja L et al. CXCL10 impairs beta cell function and viability in diabetes through TLR4 signaling. *Cell Metab* 2009;9:125-39.
- [123] Shimada A, Oikawa Y, Yamada Y, Okubo Y, Narumi S. The role of the CXCL10/CXCR3 system in type 1 diabetes. *Rev Diabet Stud* 2009;6:81-4.
- [124] Ek I, Thunell S, Blomback M. Enhanced in vivo platelet activation in diabetes mellitus. *Scand J Haematol* 1982;29:185-91.
- [125] Roy MS, Podgor MJ, Rick ME. Plasma fibrinopeptide A, beta-thromboglobulin, and platelet factor 4 in diabetic retinopathy. *Invest Ophthalmol Vis Sci* 1988;29:856-60.
- [126] Nawaz MI, Van Raemdonck K, Mohammad G, Kangave D, Van Damme J, Abu El-Asrar AM et al. Autocrine CCL2, CXCL4, CXCL9 and CXCL10 signal in retinal endothelial cells and are enhanced in diabetic retinopathy. *Exp Eye Res* 2013;109:67-76.
- [127] Wakabayashi Y, Usui Y, Okunuki Y, Kezuka T, Takeuchi M, Goto H et al. Correlation of vascular endothelial growth factor with chemokines in the vitreous in diabetic retinopathy. *Retina* 2010;30:339-44.

- [128] van Bon L, Affandi AJ, Broen J, Christmann RB, Marijnissen RJ, Stawski L et al. Proteome-wide analysis and CXCL4 as a biomarker in systemic sclerosis. *N Engl J Med* 2014;370:433-43.
- [129] Lima J, Fonollosa V, Fernandez-Cortijo J, Ordi J, Cuenca R, Khamashta MA et al. Platelet activation, endothelial cell dysfunction in the absence of anticardiolipin antibodies in systemic sclerosis. *J Rheumatol* 1991;18:1833-6.
- [130] Maeda M, Kachi H, Mori S. Plasma levels of molecular markers of blood coagulation and fibrinolysis in progressive systemic sclerosis (PSS). *J Dermatol Sci* 1996;11:223-7.
- [131] Kowal-Bielecka O, Kowal K, Lewszuk A, Bodzenta-Lukaszyk A, Walecki J, Sierakowski S. Beta thromboglobulin and platelet factor 4 in bronchoalveolar lavage fluid of patients with systemic sclerosis. *Ann Rheum Dis* 2005;64:484-6.
- [132] Antonelli A, Ferri C, Fallahi P, Ferrari SM, Giuggioli D, Colaci M et al. CXCL10 (alpha) and CCL2 (beta) chemokines in systemic sclerosis - a longitudinal study. *Rheumatology* 2008;47:45-9.
- [133] George PM, Oliver E, Dorfmueller P, Dubois OD, Reed DM, Kirkby NS et al. Evidence for the involvement of type I interferon in pulmonary arterial hypertension. *Circ Res* 2014;114:677-88.
- [134] Hasegawa M, Fujimoto M, Matsushita T, Hamaguchi Y, Takehara K, Sato S. Serum chemokine and cytokine levels as indicators of disease activity in patients with systemic sclerosis. *Clin Rheumatol* 2011;30:231-7.
- [135] Rabquer BJ, Tsou PS, Hou Y, Thirunavukkarasu E, Haines GK, III, Impens AJ et al. Dysregulated expression of MIG/CXCL9, IP-10/CXCL10 and CXCL16 and their receptors in systemic sclerosis. *Arthritis Res Ther* 2011;13:R18.
- [136] Fujii H, Shimada Y, Hasegawa M, Takehara K, Sato S. Serum levels of a Th1 chemoattractant IP-10 and Th2 chemoattractants, TARC and MDC, are elevated in patients with systemic sclerosis. *J Dermatol Sci* 2004;35:43-51.
- [137] Sfriso P, Cozzi F, Oliviero F, Caso F, Cardarelli S, Facco M et al. CXCL11 in bronchoalveolar lavage fluid and pulmonary function decline in systemic sclerosis. *Clin Exp Rheumatol* 2012;30:S71-S75.
- [138] el-Sawy T, Fahmy NM, Fairchild RL. Chemokines: directing leukocyte infiltration into allografts. *Curr Opin Immunol* 2002;14:562-8.
- [139] Romagnani P, Crescioli C. CXCL10: a candidate biomarker in transplantation. *Clin Chim Acta* 2012;413:1364-73.
- [140] Halloran PF, Fairchild RL. The puzzling role of CXCR3 and its ligands in organ allograft rejection. *Am J Transplant* 2008;8:1578-9.
- [141] Heidt S, San SD, Shankar S, Mittal S, Muthusamy AS, Friend PJ et al. Peripheral blood sampling for the detection of allograft rejection: biomarker identification and validation. *Transplantation* 2011;92:1-9.
- [142] Lazzeri E, Rotondi M, Mazzinghi B, Lasagni L, Buonamano A, Rosati A et al. High CXCL10 expression in rejected kidneys and predictive role of pretransplant serum CXCL10 for acute rejection and chronic allograft nephropathy. *Transplantation* 2005;79:1215-20.

- [143] Rotondi M, Netti GS, Lazzeri E, Stallone G, Bertoni E, Chiovato L et al. High pretransplant serum levels of CXCL9 are associated with increased risk of acute rejection and graft failure in kidney graft recipients. *Transpl Int* 2010;23:465-75.
- [144] Crescioli C, Buonamano A, Scolletta S, Sottili M, Francalanci M, Giomarelli P et al. Predictive role of pretransplant serum CXCL10 for cardiac acute rejection. *Transplantation* 2009;87:249-55.
- [145] Karason K, Jernas M, Hagg DA, Svensson PA. Evaluation of CXCL9 and CXCL10 as circulating biomarkers of human cardiac allograft rejection. *BMC Cardiovasc Disord* 2006;6:29.
- [146] Kim SC, Page EK, Knechtle SJ. Urine proteomics in kidney transplantation. *Transplant Rev (Orlando)* 2014;28:15-20.
- [147] Agostini C, Calabrese F, Rea F, Facco M, Tosoni A, Loy M et al. Cxcr3 and its ligand CXCL10 are expressed by inflammatory cells infiltrating lung allografts and mediate chemotaxis of T cells at sites of rejection. *Am J Pathol* 2001;158:1703-11.
- [148] Belperio JA, Keane MP, Burdick MD, Lynch JP, III, Zisman DA, Xue YY et al. Role of CXCL9/CXCR3 chemokine biology during pathogenesis of acute lung allograft rejection. *J Immunol* 2003;171:4844-52.
- [149] Melter M, Exeni A, Reinders ME, Fang JC, McMahon G, Ganz P et al. Expression of the chemokine receptor CXCR3 and its ligand IP-10 during human cardiac allograft rejection. *Circulation* 2001;104:2558-64.
- [150] Miura M, Morita K, Kobayashi H, Hamilton TA, Burdick MD, Strieter RM et al. Monokine induced by IFN-gamma is a dominant factor directing T cells into murine cardiac allografts during acute rejection. *J Immunol* 2001;167:3494-504.
- [151] Lo DJ, Weaver TA, Kleiner DE, Mannon RB, Jacobson LM, Becker BN et al. Chemokines and their receptors in human renal allotransplantation. *Transplantation* 2011;91:70-7.
- [152] Goddard S, Williams A, Morland C, Qin S, Gladue R, Hubscher SG et al. Differential expression of chemokines and chemokine receptors shapes the inflammatory response in rejecting human liver transplants. *Transplantation* 2001;72:1957-67.
- [153] Asaoka T, Marubashi S, Kobayashi S, Hama N, Eguchi H, Takeda Y et al. Intragraft transcriptome level of CXCL9 as biomarker of acute cellular rejection after liver transplantation. *J Surg Res* 2012;178:1003-14.
- [154] Hu H, Aizenstein BD, Puchalski A, Burmania JA, Hamawy MM, Knechtle SJ. Elevation of CXCR3-binding chemokines in urine indicates acute renal-allograft dysfunction. *Am J Transplant* 2004;4:432-7.
- [155] Mitsuhashi N, Wu GD, Zhu H, Kearns-Jonker M, Cramer DV, Starnes VA et al. Rat chemokine CXCL11: structure, tissue distribution, function and expression in cardiac transplantation models. *Mol Cell Biochem* 2007;296:1-9.
- [156] Zhao DX, Hu Y, Miller GG, Luster AD, Mitchell RN, Libby P. Differential expression of the IFN-gamma-inducible CXCR3-binding chemokines, IFN-inducible protein 10, monokine induced by IFN, and IFN-inducible T cell alpha chemoattractant in human cardiac allografts: association with cardiac allograft vasculopathy and acute rejection. *J Immunol* 2002;169:1556-60.

- [157] Shi G, Field DJ, Ko KA, Ture S, Srivastava K, Levy S et al. Platelet factor 4 limits Th17 differentiation and cardiac allograft rejection. *J Clin Invest* 2014;124:543-52.
- [158] Piper KP, Horlock C, Curnow SJ, Arrazi J, Nicholls S, Mahendra P et al. CXCL10-CXCR3 interactions play an important role in the pathogenesis of acute graft-versus-host disease in the skin following allogeneic stem-cell transplantation. *Blood* 2007;110:3827-32.
- [159] Croudace JE, Inman CF, Abbotts BE, Nagra S, Nunnick J, Mahendra P et al. Chemokine-mediated tissue recruitment of CXCR3+ CD4+ T cells plays a major role in the pathogenesis of chronic GVHD. *Blood* 2012;120:4246-55.
- [160] Westekemper H, Meller S, Citak S, Schulte C, Steuhl KP, Homey B et al. Differential chemokine expression in chronic GVHD of the conjunctiva. *Bone Marrow Transplant* 2010;45:1340-6.
- [161] Kaplan KL, Owen J. Plasma levels of beta-thromboglobulin and platelet factor 4 as indices of platelet activation in vivo. *Blood* 1981;57:199-202.
- [162] Essien EM, Ebhota MI. Platelet hypersensitivity in acute malaria (*Plasmodium falciparum*) infection in man. *Thromb Haemost* 1981;46:547-9.
- [163] Essien EM, Ebhota MI. Platelet secretory activities in acute malaria (*Plasmodium falciparum*) infection. *Acta Haematol* 1983;70:183-8.
- [164] Lucchi NW, Jain V, Wilson NO, Singh N, Udhayakumar V, Stiles JK. Potential serological biomarkers of cerebral malaria. *Dis Markers* 2011;31:327-35.
- [165] Van den Steen PE, Deroost K, Van Aelst I, Geurts N, Martens E, Struyf S et al. CXCR3 determines strain susceptibility to murine cerebral malaria by mediating T lymphocyte migration toward IFN-gamma-induced chemokines. *Eur J Immunol* 2008;38:1082-95.
- [166] Armah HB, Wilson NO, Sarfo BY, Powell MD, Bond VC, Anderson W et al. Cerebrospinal fluid and serum biomarkers of cerebral malaria mortality in Ghanaian children. *Malar J* 2007;6:147.
- [167] Wilson NO, Jain V, Roberts CE, Lucchi N, Joel PK, Singh MP et al. CXCL4 and CXCL10 predict risk of fatal cerebral malaria. *Dis Markers* 2011;30:39-49.
- [168] Wilson N, Driss A, Solomon W, Dickinson-Copeland C, Salifu H, Jain V et al. CXCL10 gene promoter polymorphism -1447A>G correlates with plasma CXCL10 levels and is associated with male susceptibility to cerebral malaria. *Plos One* 2013;8:e81329.
- [169] Hansen DS, Bernard NJ, Nie CQ, Schofield L. NK cells stimulate recruitment of CXCR3+ T cells to the brain during *Plasmodium berghei*-mediated cerebral malaria. *J Immunol* 2007;178:5779-88.
- [170] Miu J, Mitchell AJ, Muller M, Carter SL, Manders PM, McQuillan JA et al. Chemokine gene expression during fatal murine cerebral malaria and protection due to CXCR3 deficiency. *J Immunol* 2008;180:1217-30.
- [171] Nie CQ, Bernard NJ, Norman MU, Amante FH, Lundie RJ, Crabb BS et al. IP-10-mediated T cell homing promotes cerebral inflammation over splenic immunity to malaria infection. *PLoS Pathog* 2009;5:e1000369.

- [172] Srivastava K, Cockburn IA, Swaim A, Thompson LE, Tripathi A, Fletcher CA et al. Platelet factor 4 mediates inflammation in experimental cerebral malaria. *Cell Host Microbe* 2008;4:179-87.
- [173] Srivastava K, Field DJ, Aggrey A, Yamakuchi M, Morrell CN. Platelet factor 4 regulation of monocyte KLF4 in experimental cerebral malaria. *Plos One* 2010;5:e10413.
- [174] Lacerda-Queiroz N, Lima OC, Carneiro CM, Vilela MC, Teixeira AL, Teixeira-Carvalho A et al. *Plasmodium berghei* NK65 induces cerebral leukocyte recruitment in vivo: an intravital microscopic study. *Acta Trop* 2011;120:31-9.
- [175] Palomo J, Fauconnier M, Coquard L, Gilles M, Meme S, Szeremeta F et al. Type I interferons contribute to experimental cerebral malaria development in response to sporozoite or blood-stage *Plasmodium berghei* ANKA. *Eur J Immunol* 2013;43:2683-95.
- [176] Campanella GS, Tager AM, El Khoury JK, Thomas SY, Abrazinski TA, Manice LA et al. Chemokine receptor CXCR3 and its ligands CXCL9 and CXCL10 are required for the development of murine cerebral malaria. *Proc Natl Acad Sci U S A* 2008;105:4814-9.
- [177] Morrell CN, Srivastava K, Swaim A, Lee MT, Chen J, Nagineni C et al. Beta interferon suppresses the development of experimental cerebral malaria. *Infect Immun* 2011;79:1750-8.
- [178] Love MS, Millholland MG, Mishra S, Kulkarni S, Freeman KB, Pan WX et al. Platelet Factor 4 activity against *P. falciparum* and its translation to nonpeptidic mimics as antimalarials. *Cell Host Microbe* 2012;12:815-23.
- [179] McMorran BJ, Burgio G, Foote SJ. New insights into the protective power of platelets in malaria infection. *Commun Integr Biol* 2013;6:e23653-1-4.
- [180] Szabo MC, Soo KS, Zlotnik A, Schall TJ. Chemokine class differences in binding to the Duffy antigen-erythrocyte chemokine receptor. *J Biol Chem* 1995;270:25348-51.
- [181] McMorran BJ, Marshall VM, de Graaf C, Drysdale KE, Shabbar M, Smyth GK et al. Platelets kill intraerythrocytic malarial parasites and mediate survival to infection. *Science* 2009;323:797-800.
- [182] McMorran BJ, Wieczorski L, Drysdale KE, Chan JA, Huang HM, Smith C et al. Platelet factor 4 and Duffy antigen required for platelet killing of *Plasmodium falciparum*. *Science* 2012;338:1348-51.
- [183] Murakami T, Yamamoto N. Roles of chemokines and chemokine receptors in HIV-1 infection. *Int J Hematol* 2000;72:412-7.
- [184] Wu Y, Yoder A. Chemokine coreceptor signaling in HIV-1 infection and pathogenesis. *PLoS Pathog* 2009;5:e1000520.
- [185] Hatse S, Huskens D, Princen K, Vermeire K, Bridger GJ, De Clercq E et al. Modest human immunodeficiency virus coreceptor function of CXCR3 is strongly enhanced by mimicking the CXCR4 ligand binding pocket in the CXCR3 receptor. *J Virol* 2007;81:3632-9.
- [186] Gray CM, Hong HA, Young K, Lewis DA, Fallows D, Manca C et al. Plasma interferon-gamma-inducible protein 10 can be used to predict viral load in HIV-1-infected individuals. *J Acquir Immune Defic Syndr* 2013;63:e115-e116.

- [187] Sarkar S, Kalia V, Murphey-Corb M, Montelaro RC, Reinhart TA. Expression of IFN-gamma induced CXCR3 agonist chemokines and compartmentalization of CXCR3+ cells in the periphery and lymph nodes of rhesus macaques during simian immunodeficiency virus infection and acquired immunodeficiency syndrome. *J Med Primatol* 2003;32:247-64.
- [188] Shacklett BL, Cox CA, Wilkens DT, Karl KR, Nilsson A, Nixon DF et al. Increased adhesion molecule and chemokine receptor expression on CD8+ T cells trafficking to cerebrospinal fluid in HIV-1 infection. *J Infect Dis* 2004;189:2202-12.
- [189] Cinque P, Bestetti A, Marenzi R, Sala S, Gisslen M, Hagberg L et al. Cerebrospinal fluid interferon-gamma-inducible protein 10 (IP-10, CXCL10) in HIV-1 infection. *J Neuroimmunol* 2005;168:154-63.
- [190] Kolb SA, Sporer B, Lahrtz F, Koedel U, Pfister HW, Fontana A. Identification of a T cell chemotactic factor in the cerebrospinal fluid of HIV-1-infected individuals as interferon-gamma inducible protein 10. *J Neuroimmunol* 1999;93:172-81.
- [191] Izmailova E, Bertley FM, Huang Q, Makori N, Miller CJ, Young RA et al. HIV-1 Tat reprograms immature dendritic cells to express chemoattractants for activated T cells and macrophages. *Nat Med* 2003;9:191-7.
- [192] Foley JF, Yu CR, Solow R, Yacobucci M, Peden KW, Farber JM. Roles for CXC chemokine ligands 10 and 11 in recruiting CD4+ T cells to HIV-1-infected monocyte-derived macrophages, dendritic cells, and lymph nodes. *J Immunol* 2005;174:4892-900.
- [193] Wetzel MA, Steele AD, Henderson EE, Rogers TJ. The effect of X4 and R5 HIV-1 on C, C-C, and C-X-C chemokines during the early stages of infection in human PBMCs. *Virology* 2002;292:6-15.
- [194] Asensio VC, Maier J, Milner R, Boztug K, Kincaid C, Moulard M et al. Interferon-independent, human immunodeficiency virus type 1 gp120-mediated induction of CXCL10/IP-10 gene expression by astrocytes in vivo and in vitro. *J Virol* 2001;75:7067-77.
- [195] Poluektova L, Moran T, Zelivyanskaya M, Swindells S, Gendelman HE, Persidsky Y. The regulation of alpha chemokines during HIV-1 infection and leukocyte activation: relevance for HIV-1-associated dementia. *J Neuroimmunol* 2001;120:112-28.
- [196] Mehla R, Bivalkar-Mehla S, Nagarkatti M, Chauhan A. Programming of neurotoxic cofactor CXCL-10 in HIV-1-associated dementia: abrogation of CXCL-10-induced neuro-glial toxicity in vitro by PKC activator. *J Neuroinflammation* 2012;9:239.
- [197] van Marle G, Henry S, Todoruk T, Sullivan A, Silva C, Rourke SB et al. Human immunodeficiency virus type 1 Nef protein mediates neural cell death: a neurotoxic role for IP-10. *Virology* 2004;329:302-18.
- [198] Reinhart TA. Chemokine induction by HIV-1: recruitment to the cause. *Trends Immunol* 2003;24:351-3.
- [199] Lane BR, King SR, Bock PJ, Strieter RM, Coffey MJ, Markovitz DM. The C-X-C chemokine IP-10 stimulates HIV-1 replication. *Virology* 2003;307:122-34.

- [200] Auerbach DJ, Lin Y, Miao HY, Cimbri R, DiFiore MJ, Gianolini ME et al. Identification of the platelet-derived chemokine CXCL4/PF-4 as a broad-spectrum HIV-1 inhibitor. *Proc Natl Acad Sci U S A* 2012;109:9569-74.
- [201] Tsegaye TS, Gnirss K, Rahe-Meyer N, Kiene M, Kramer-Kuhl A, Behrens G et al. Platelet activation suppresses HIV-1 infection of T cells. *Retrovirology* 2013;10:48.
- [202] Zeremski M, Markatou M, Brown QB, Dorante G, Cunningham-Rundles S, Talal AH. Interferon gamma-inducible protein 10: a predictive marker of successful treatment response in hepatitis C virus/HIV-coinfected patients. *J Acquir Immune Defic Syndr* 2007;45:262-8.
- [203] Qu J, Zhang Q, Li Y, Liu W, Chen L, Zhu Y et al. The Tat protein of human immunodeficiency virus-1 enhances hepatitis C virus replication through interferon gamma-inducible protein-10. *BMC Immunol* 2012;13:15.
- [204] Berenguer J, Fernandez-Rodriguez A, Jimenez-Sousa MA, Cosin J, Zarate P, Micheloud D et al. High plasma CXCL10 levels are associated with HCV-genotype 1, and higher insulin resistance, fibrosis, and HIV viral load in HIV/HCV coinfected patients. *Cytokine* 2012;57:25-9.
- [205] Roe B, Coughlan S, Hassan J, Grogan A, Farrell G, Norris S et al. Elevated serum levels of interferon- gamma -inducible protein-10 in patients coinfecting with hepatitis C virus and HIV. *J Infect Dis* 2007;196:1053-7.
- [206] Aabye MG, Ruhwald M, Praygod G, Jeremiah K, Faurholt-Jepsen M, Faurholt-Jepsen D et al. Potential of interferon-gamma-inducible protein 10 in improving tuberculosis diagnosis in HIV-infected patients. *Eur Respir J* 2010;36:1488-90.
- [207] Kabeer BS, Sikhamani R, Raja A. Comparison of interferon gamma and interferon gamma-inducible protein-10 secretion in HIV-tuberculosis patients. *AIDS* 2010;24:323-5.
- [208] Davenport GC, Hittner JB, Were T, Ong'echa JM, Perkins DJ. Relationship between inflammatory mediator patterns and anemia in HIV-1 positive and exposed children with *Plasmodium falciparum* malaria. *Am J Hematol* 2012;87:652-8.
- [209] Tang YQ, Yeaman MR, Selsted ME. Antimicrobial peptides from human platelets. *Infect Immun* 2002;70:6524-33.
- [210] Cole AM, Ganz T, Liese AM, Burdick MD, Liu L, Strieter RM. Cutting edge: IFN-inducible ELR-CXC chemokines display defensin-like antimicrobial activity. *J Immunol* 2001;167:623-7.
- [211] Liu MT, Armstrong D, Hamilton TA, Lane TE. Expression of Mig (monokine induced by interferon-gamma) is important in T lymphocyte recruitment and host defense following viral infection of the central nervous system. *J Immunol* 2001;166:1790-5.
- [212] Liu MT, Chen BP, Oertel P, Buchmeier MJ, Armstrong D, Hamilton TA et al. The T cell chemoattractant IFN-inducible protein 10 is essential in host defense against viral-induced neurologic disease. *J Immunol* 2000;165:2327-30.
- [213] Fahey S, Dempsey E, Long A. The role of chemokines in acute and chronic hepatitis C infection. *Cell Mol Immunol* 2014;11:25-40.

- [214] Larrubia JR, Benito-Martinez S, Calvino M, Sanz-de-Villalobos E, Parra-Cid T. Role of chemokines and their receptors in viral persistence and liver damage during chronic hepatitis C virus infection. *World J Gastroenterol* 2008;14:7149-59.
- [215] Harvey CE, Post JJ, Palladinetti P, Freeman AJ, Ffrench RA, Kumar RK et al. Expression of the chemokine IP-10 (CXCL10) by hepatocytes in chronic hepatitis C virus infection correlates with histological severity and lobular inflammation. *J Leukoc Biol* 2003;74:360-9.
- [216] Nguyen N, de Esch C, Cameron B, Kumar RK, Zekry A, Lloyd AR. Positioning of leukocyte subsets in the portal and lobular compartments of hepatitis C virus-infected liver correlates with local chemokine expression. *J Gastroenterol Hepatol* 2014;29:860-9.
- [217] Apolinario A, Majano PL, Alvarez-Perez E, Saez A, Lozano C, Vargas J et al. Increased expression of T cell chemokines and their receptors in chronic hepatitis C: relationship with the histological activity of liver disease. *Am J Gastroenterol* 2002;97:2861-70.
- [218] Zeremski M, Dimova R, Brown Q, Jacobson IM, Markatou M, Talal AH. Peripheral CXCR3-associated chemokines as biomarkers of fibrosis in chronic hepatitis C virus infection. *J Infect Dis* 2009;200:1774-80.
- [219] Wasmuth HE, Lammert F, Zaldivar MM, Weiskirchen R, Hellerbrand C, Scholten D et al. Antifibrotic effects of CXCL9 and its receptor CXCR3 in livers of mice and humans. *Gastroenterology* 2009;137:309-19, 319.
- [220] Sahin H, Borkham-Kamphorst E, do O NT, Berres ML, Kaldenbach M, Schmitz P et al. Proapoptotic effects of the chemokine, CXCL 10 are mediated by the noncognate receptor TLR4 in hepatocytes. *Hepatology* 2013;57:797-805.
- [221] Helbig KJ, Ruszkiewicz A, Semendric L, Harley HA, McColl SR, Beard MR. Expression of the CXCR3 ligand I-TAC by hepatocytes in chronic hepatitis C and its correlation with hepatic inflammation. *Hepatology* 2004;39:1220-9.
- [222] Zeremski M, Petrovic LM, Chiriboga L, Brown QB, Yee HT, Kinkhabwala M et al. Intrahepatic levels of CXCR3-associated chemokines correlate with liver inflammation and fibrosis in chronic hepatitis C. *Hepatology* 2008;48:1440-50.
- [223] Zaldivar MM, Pauels K, von Hundelshausen P, Berres ML, Schmitz P, Bornemann J et al. CXC chemokine ligand 4 (Cxcl4) is a platelet-derived mediator of experimental liver fibrosis. *Hepatology* 2010;51:1345-53.
- [224] Dominguez M, Miquel R, Colmenero J, Moreno M, Garcia-Pagan JC, Bosch J et al. Hepatic expression of CXC chemokines predicts portal hypertension and survival in patients with alcoholic hepatitis. *Gastroenterology* 2009;136:1639-50.
- [225] Tacke F, Zimmermann HW, Berres ML, Trautwein C, Wasmuth HE. Serum chemokine receptor CXCR3 ligands are associated with progression, organ dysfunction and complications of chronic liver diseases. *Liver Int* 2011;31:840-9.
- [226] Mach F, Sauty A, Iarossi AS, Sukhova GK, Neote K, Libby P et al. Differential expression of three T lymphocyte-activating CXC chemokines by human atheroma-associated cells. *J Clin Invest* 1999;104:1041-50.

- [227] Anger T, Pohle FK, Kandler L, Barthel T, Ensminger SM, Fischlein T et al. VAP-1, Eotaxin3 and MIG as potential atherosclerotic triggers of severe calcified and stenotic human aortic valves: effects of statins. *Exp Mol Pathol* 2007;83:435-42.
- [228] Ranjbaran H, Wang Y, Manes TD, Yakimov AO, Akhtar S, Kluger MS et al. Heparin displaces interferon-gamma-inducible chemokines (IP-10, I-TAC, and Mig) sequestered in the vasculature and inhibits the transendothelial migration and arterial recruitment of T cells. *Circulation* 2006;114:1293-300.
- [229] Heller EA, Liu E, Tager AM, Yuan Q, Lin AY, Ahluwalia N et al. Chemokine CXCL10 promotes atherogenesis by modulating the local balance of effector and regulatory T cells. *Circulation* 2006;113:2301-12.
- [230] Sachais BS, Turrentine T, Dawicki McKenna JM, Rux AH, Rader D, Kowalska MA. Elimination of platelet factor 4 (PF4) from platelets reduces atherosclerosis in C57Bl/6 and apoE^{-/-} mice. *Thromb Haemost* 2007;98:1108-13.
- [231] Koenen RR, von Hundelshausen P, Nesmelova IV, Zerneck A, Liehn EA, Sarabi A et al. Disrupting functional interactions between platelet chemokines inhibits atherosclerosis in hyperlipidemic mice. *Nat Med* 2009;15:97-103.
- [232] Aidoudi S, Bikfalvi A. Interaction of PF4 (CXCL4) with the vasculature: a role in atherosclerosis and angiogenesis. *Thromb Haemost* 2010;104:941-8.
- [233] Gleissner CA, Shaked I, Little KM, Ley K. CXC chemokine ligand 4 induces a unique transcriptome in monocyte-derived macrophages. *J Immunol* 2010;184:4810-8.
- [234] Nassar T, Sachais BS, Akkawi S, Kowalska MA, Bdeir K, Leitersdorf E et al. Platelet factor 4 enhances the binding of oxidized low-density lipoprotein to vascular wall cells. *J Biol Chem* 2003;278:6187-93.
- [235] Pitsilos S, Hunt J, Mohler ER, Prabhakar AM, Poncz M, Dawicki J et al. Platelet factor 4 localization in carotid atherosclerotic plaques: correlation with clinical parameters. *Thromb Haemostasis* 2003;90:1112-20.
- [236] Ardigo D, Assimes TL, Fortmann SP, Go AS, Hlatky M, Hytopoulos E et al. Circulating chemokines accurately identify individuals with clinically significant atherosclerotic heart disease. *Physiol Genomics* 2007;31:402-9.
- [237] Heneka MT, O'Banion MK, Terwel D, Kummer MP. Neuroinflammatory processes in Alzheimer's disease. *J Neural Transm* 2010;117:919-47.
- [238] Xia MQ, Hyman BT. Chemokines/chemokine receptors in the central nervous system and Alzheimer's disease. *J Neurovirol* 1999;5:32-41.
- [239] Xia MQ, Bacskai BJ, Knowles RB, Qin SX, Hyman BT. Expression of the chemokine receptor CXCR3 on neurons and the elevated expression of its ligand IP-10 in reactive astrocytes: in vitro ERK1/2 activation and role in Alzheimer's disease. *J Neuroimmunol* 2000;108:227-35.
- [240] Sui Y, Stehno-Bittel L, Li S, Loganathan R, Dhillon NK, Pinson D et al. CXCL10-induced cell death in neurons: role of calcium dysregulation. *Eur J Neurosci* 2006;23:957-64.

- [241] Lai W, Wu J, Zou X, Xie J, Zhang L, Zhao X et al. Secretome analyses of Abeta(1-42) stimulated hippocampal astrocytes reveal that CXCL10 is involved in astrocyte migration. *J Proteome Res* 2013;12:832-43.
- [242] Galimberti D, Schoonenboom N, Scheltens P, Fenoglio C, Venturelli E, Pijnenburg YA et al. Intrathecal chemokine levels in Alzheimer disease and frontotemporal lobar degeneration. *Neurology* 2006;66:146-7.
- [243] Galimberti D, Schoonenboom N, Scheltens P, Fenoglio C, Bouwman F, Venturelli E et al. Intrathecal chemokine synthesis in mild cognitive impairment and Alzheimer disease. *Arch Neurol* 2006;63:538-43.
- [244] Galimberti D, Venturelli E, Fenoglio C, Lovati C, Guidi I, Scalabrini D et al. IP-10 serum levels are not increased in mild cognitive impairment and Alzheimer's disease. *Eur J Neurol* 2007;14:e3-e4.
- [245] Duan RS, Yang X, Chen ZG, Lu MO, Morris C, Winblad B et al. Decreased fractalkine and increased IP-10 expression in aged brain of APP(swe) transgenic mice. *Neurochem Res* 2008;33:1085-9.
- [246] Hochstrasser T, Marksteiner J, DeFrancesco M, Deisenhammer EA, Kemmler G, Humpel C. Two Blood Monocytic Biomarkers (CCL15 and p21) Combined with the Mini-Mental State Examination Discriminate Alzheimer's Disease Patients from Healthy Subjects. *Dement Geriatr Cogn Dis Extra* 2011;1:297-309.
- [247] Lee KS, Chung JH, Lee KH, Shin MJ, Oh BH, Hong CH. Bioplex analysis of plasma cytokines in Alzheimer's disease and mild cognitive impairment. *Immunol Lett* 2008;121:105-9.
- [248] Skovronsky DM, Lee VM, Pratico D. Amyloid precursor protein and amyloid beta peptide in human platelets. Role of cyclooxygenase and protein kinase C. *J Biol Chem* 2001;276:17036-43.
- [249] Zhang W, Huang W, Jing F. Contribution of blood platelets to vascular pathology in Alzheimer's disease. *J Blood Med* 2013;4:141-7.
- [250] Proudfoot AE, Handel TM, Johnson Z, Lau EK, LiWang P, Clark-Lewis I et al. Glycosaminoglycan binding and oligomerization are essential for the in vivo activity of certain chemokines. *Proc Natl Acad Sci U S A* 2003;100:1885-90.
- [251] Prechel M, Walenga JM. Emphasis on the role of PF4 in the incidence, pathophysiology and treatment of heparin induced thrombocytopenia. *Thromb J* 2013;11.
- [252] Rauova L, Hirsch JD, Greene TK, Zhai L, Hayes VM, Kowalska MA et al. Monocyte-bound PF4 in the pathogenesis of heparin-induced thrombocytopenia. *Blood* 2010;116:5021-31.
- [253] Cornelis JJ, Lang SI, Stroh-Dege AY, Balboni G, Dinsart C, Rommelaere J. Cancer gene therapy through autonomous parvovirus-mediated gene transfer. *Curr Gene Ther* 2004;4:249-61.
- [254] Hagedorn M, Zilberberg L, Lozano RM, Cuevas P, Canron X, Redondo-Horcajo M et al. A short peptide domain of platelet factor 4 blocks angiogenic key events induced by FGF-2. *FASEB J* 2001;15:550-2.

- [255] Vandercappellen J, Liekens S, Bronckaers A, Noppen S, Ronsse I, Dillen C et al. The COOH-terminal peptide of platelet factor-4 variant (CXCL4L1/PF-4var47-70) strongly inhibits angiogenesis and suppresses B16 melanoma growth in vivo. *Mol Cancer Res* 2010;8:322-34.
- [256] Wang P, Yang X, Xu W, Li K, Chu Y, Xiong S. Integrating individual functional moieties of CXCL10 and CXCL11 into a novel chimeric chemokine leads to synergistic antitumor effects: a strategy for chemokine-based multi-target-directed cancer therapy. *Cancer Immunol Immunother* 2010;59:1715-26.
- [257] van der Schaft DW, Dings RP, de Lussanet QG, van Eijk LI, Nap AW, Beets-Tan RG et al. The designer anti-angiogenic peptide anginex targets tumor endothelial cells and inhibits tumor growth in animal models. *FASEB J* 2002;16:1991-3.
- [258] Mayer L, Sandborn WJ, Stepanov Y, Geboes K, Hardi R, Yellin M et al. Anti-IP-10 antibody (BMS-936557) for ulcerative colitis: a phase II randomised study. *Gut* 2014;63:442-50.
- [259] Yellin M, Paliienko I, Balanescu A, Ter-Vartanian S, Tseluyko V, Xu LA et al. A phase II, randomized, double-blind, placebo-controlled study evaluating the efficacy and safety of MDX-1100, a fully human anti-CXCL10 monoclonal antibody, in combination with methotrexate in patients with rheumatoid arthritis. *Arthritis Rheum* 2012;64:1730-9.
- [260] Nagy B, Jr., Miszti-Blasius K, Kerenyi A, Clemetson KJ, Kappelmayer J. Potential therapeutic targeting of platelet-mediated cellular interactions in atherosclerosis and inflammation. *Curr Med Chem* 2012;19:518-31.
- [261] Lippi G, Favaloro EJ. Recombinant platelet factor 4: a therapeutic, anti-neoplastic chimera? *Semin Thromb Hemost* 2010;36:558-69.
- [262] Lazzeri E, Romagnani P. CXCR3-binding chemokines: novel multifunctional therapeutic targets. *Curr Drug Targets Immune Endocr Metabol Disord* 2005;5:109-18.
- [263] Christen U. Chemokines as drug targets in type 1 diabetes. *Endocr Metab Immune Disord Drug Targets* 2007;7:7-12.
- [264] De Sutter J, van de Veire NR, Struyf S, Philippe J, De Buyzere M, Van Damme J. PF-4var/CXCL4L1 predicts outcome in stable coronary artery disease patients with preserved left ventricular function. *Plos One* 2012;7:e31343.
- [265] Meuwis MA, Fillet M, Lutteri L, Maree R, Geurts P, de Seny D et al. Proteomics for prediction and characterization of response to infliximab in Crohn's disease: A pilot study. *Clin Biochem* 2008;41:960-7.
- [266] Trocme C, Marotte H, Baillet A, Pallot-Prades B, Garin J, Grange L et al. Apolipoprotein A-I and platelet factor 4 are biomarkers for infliximab response in rheumatoid arthritis. *Ann Rheum Dis* 2009;68:1328-33.
- [267] Romero AI, Lagging M, Westin J, Dhillon AP, Dustin LB, Pawlotsky JM et al. Interferon (IFN)-gamma-inducible protein-10: association with histological results, viral kinetics, and outcome during treatment with pegylated IFN-alpha 2a and ribavirin for chronic hepatitis C virus infection. *J Infect Dis* 2006;194:895-903.

- [268] Falconer K, Askarieh G, Weis N, Hellstrand K, Alaeus A, Lagging M. IP-10 predicts the first phase decline of HCV RNA and overall viral response to therapy in patients co-infected with chronic hepatitis C virus infection and HIV. *Scand J Infect Dis* 2010;42:896-901.
- [269] Casrouge A, Decalf J, Ahloulay M, Lababidi C, Mansour H, Vallet-Pichard A et al. Evidence for an antagonist form of the chemokine CXCL10 in patients chronically infected with HCV. *J Clin Invest* 2011;121:308-17.
- [270] Ragab D, Laird M, Duffy D, Casrouge A, Mamdouh R, Abass A et al. CXCL10 antagonism and plasma sDPPiV correlate with increasing liver disease in chronic HCV genotype 4 infected patients. *Cytokine* 2013;63:105-12.
- [271] Proost P, Schutyser E, Menten P, Struyf S, Wuyts A, Opdenakker G et al. Amino-terminal truncation of CXCR3 agonists impairs receptor signaling and lymphocyte chemotaxis, while preserving antiangiogenic properties. *Blood* 2001;98:3554-61.
- [272] Burns JM, Summers BC, Wang Y, Melikian A, Berahovich R, Miao Z et al. A novel chemokine receptor for SDF-1 and I-TAC involved in cell survival, cell adhesion, and tumor development. *J Exp Med* 2006;203:2201-13.
- [273] Chaudhuri A, Zbrzezna V, Polyakova J, Pogo AO, Hesselgesser J, Horuk R. Expression of the Duffy antigen in K562 cells. Evidence that it is the human erythrocyte chemokine receptor. *J Biol Chem* 1994;269:7835-8.
- [274] Gardner L, Patterson AM, Ashton BA, Stone MA, Middleton J. The human Duffy antigen binds selected inflammatory but not homeostatic chemokines. *Biochem Biophys Res Commun* 2004;321:306-12.
- [275] Wu Z, Han X, Yan J, Pan Y, Gong J, Di J et al. The prognostic significance of chemokine receptor CXCR3 expression in colorectal carcinoma. *Biomed Pharmacother* 2012;66:373-7.
- [276] Gupta SK, Hassel T, Singh JP. A potent inhibitor of endothelial cell proliferation is generated by proteolytic cleavage of the chemokine platelet factor 4. *Proc Natl Acad Sci U S A* 1995;92:7799-803.
- [277] Zipfel PF, Bialonski A, Skerka C. Induction of members of the IL-8/NAP-1 gene family in human T lymphocytes is suppressed by cyclosporin A. *Biochem Biophys Res Commun* 1991;181:179-83.
- [278] Struyf S, Burdick MD, Peeters E, Van den Broeck K, Dillen C, Proost P et al. Platelet factor-4 variant chemokine CXCL4L1 inhibits melanoma and lung carcinoma growth and metastasis by preventing angiogenesis. *Cancer Res* 2007;67:5940-8.
- [279] Lu XL, Jiang XB, Liu RE, Zhang SM. The enhanced anti-angiogenic and antitumor effects of combining flk1-based DNA vaccine and IP-10. *Vaccine* 2008;26:5352-7.
- [280] Keyser J, Schultz J, Ladell K, Elzaouk L, Heinzerling L, Pavlovic J et al. IP-10-encoding plasmid DNA therapy exhibits anti-tumor and anti-metastatic efficiency. *Exp Dermatol* 2004;13:380-90.
- [281] Sharpe RJ, Byers HR, Scott CF, Bauer SI, Maione TE. Growth inhibition of murine melanoma and human colon carcinoma by recombinant human platelet factor 4. *J Natl Cancer Inst* 1990;82:848-53.

- [282] Kolber DL, Knisely TL, Maione TE. Inhibition of development of murine melanoma lung metastases by systemic administration of recombinant platelet factor 4. *J Natl Cancer Inst* 1995;87:304-9.
- [283] Arenberg DA, White ES, Burdick MD, Strom SR, Strieter RM. Improved survival in tumor-bearing SCID mice treated with interferon-gamma-inducible protein 10 (IP-10/CXCL10). *Cancer Immunol Immunother* 2001;50:533-8.
- [284] Li G, Tian L, Hou JM, Ding ZY, He QM, Feng P et al. Improved therapeutic effectiveness by combining recombinant CXC chemokine ligand 10 with Cisplatin in solid tumors. *Clin Cancer Res* 2005;11:4217-24.
- [285] Wang YQ, Wada A, Ugai S, Tagawa M. Expression of the Mig (CXCL9) gene in murine lung carcinoma cells generated angiogenesis-independent antitumor effects. *Oncol Rep* 2003;10:909-13.
- [286] Zhang R, Tian L, Chen LJ, Xiao F, Hou JM, Zhao X et al. Combination of MIG (CXCL9) chemokine gene therapy with low-dose cisplatin improves therapeutic efficacy against murine carcinoma. *Gene Ther* 2006;13:1263-71.
- [287] Ruehlmann JM, Xiang R, Niethammer AG, Ba Y, Pertl U, Dolman CS et al. MIG (CXCL9) chemokine gene therapy combines with antibody-cytokine fusion protein to suppress growth and dissemination of murine colon carcinoma. *Cancer Res* 2001;61:8498-503.
- [288] Pan J, Burdick MD, Belperio JA, Xue YY, Gerard C, Sharma S et al. CXCR3/CXCR3 ligand biological axis impairs RENCA tumor growth by a mechanism of immunoangiostasis. *J Immunol* 2006;176:1456-64.
- [289] Walser TC, Ma X, Kundu N, Dorsey R, Goloubeva O, Fulton AM. Immune-mediated modulation of breast cancer growth and metastasis by the chemokine Mig (CXCL9) in a murine model. *J Immunother* 2007;30:490-8.
- [290] Hampl M, Tanaka T, Albert PS, Lee J, Ferrari N, Fine HA. Therapeutic effects of viral vector-mediated antiangiogenic gene transfer in malignant ascites. *Hum Gene Ther* 2001;12:1713-29.
- [291] Yang F, Gou M, Deng H, Yi T, Zhong Q, Wei Y et al. Efficient inhibition of ovarian cancer by recombinant CXC chemokine ligand 10 delivered by novel biodegradable cationic heparin-polyethyleneimine nanogels. *Oncol Rep* 2012;28:668-76.
- [292] Jiang XB, Lu XL, Hu P, Liu RE. Improved therapeutic efficacy using vaccination with glioma lysate-pulsed dendritic cells combined with IP-10 in murine glioma. *Vaccine* 2009;27:6210-6.
- [293] Liu TC, Zhang T, Fukuhara H, Kuroda T, Todo T, Martuza RL et al. Oncolytic HSV armed with platelet factor 4, an antiangiogenic agent, shows enhanced efficacy. *Mol Ther* 2006;14:789-97.
- [294] Enderlin M, Kleinmann EV, Struyf S, Buracchi C, Vecchi A, Kinscherf R et al. TNF-alpha and the IFN-gamma-inducible protein 10 (IP-10/CXCL-10) delivered by parvoviral vectors act in synergy to induce antitumor effects in mouse glioblastoma. *Cancer Gene Ther* 2009;16:149-60.

- [295] Antoszczyk S, Spyra M, Mautner VF, Kurtz A, Stemmer-Rachamimov AO, Martuza RL et al. Treatment of orthotopic malignant peripheral nerve sheath tumors with oncolytic herpes simplex virus. *Neuro Oncol* 2014;16:1066.
- [296] Chen P, Yang LL, Yang HS, Wang YS, Li G, Wu Y et al. Synergistic antitumor effect of CXCL10 with hyperthermia. *J Cancer Res Clin Oncol* 2008;134:679-87.
- [297] Giese NA, Raykov Z, DeMartino L, Vecchi A, Sozzani S, Dinsart C et al. Suppression of metastatic hemangiosarcoma by a parvovirus MVMp vector transducing the IP-10 chemokine into immunocompetent mice. *Cancer Gene Ther* 2002;9:432-42.
- [298] Sgadari C, Farber JM, Angiolillo AL, Liao F, Teruya-Feldstein J, Burd PR et al. Mig, the monokine induced by interferon-gamma, promotes tumor necrosis in vivo. *Blood* 1997;89:2635-43.
- [299] Hensbergen PJ, Wijnands PG, Schreurs MW, Scheper RJ, Willemze R, Tensen CP. The CXCR3 targeting chemokine CXCL11 has potent antitumor activity in vivo involving attraction of CD8+ T lymphocytes but not inhibition of angiogenesis. *J Immunother* 2005;28:343-51.