ALPERT MULTIWAVELETS AND LEGENDRE-ANGELESCO MULTIPLE ORTHOGONAL POLYNOMIALS*

JEFFREY S. GERONIMO[†], PLAMEN ILIEV[†], AND WALTER VAN ASSCHE[‡]

Abstract. We show that the multiwavelets, introduced by Alpert in 1993, are related to type I Legendre-Angelesco multiple orthogonal polynomials. We give explicit formulas for these Legendre-Angelesco polynomials and for the Alpert multiwavelets. The multiresolution analysis can be done entirely using Legendre polynomials, and we give an algorithm, using Cholesky factorization, to compute the multiwavelets and a method, using the Jacobi matrix for Legendre polynomials, to compute the matrices in the scaling relation for any size of the multiplicity of the multiwavelets.

Key words. Alpert multiwavelets, Legendre-Angelesco polynomials, Legendre polynomials, hypergeometric functions

AMS subject classifications. 42C40, 65T60, 33C20, 33C45

1. Introduction. Wavelets are a powerful way for approximating functions and have been very successful in harmonic analysis, Fourier analysis, signal and image processing and approximation theory [4], [17]. Multiresolution analysis, as introduced by Mallat and Meyer [19, 20], is a useful way to describe functions in $L^2(\mathbb{R})$ in a nested sequence of subspaces, and wavelets are very well suited to move from one resolution to higher resolutions. Multiwavelets are an extension of wavelets where instead of using a single scaling function one uses a vector scaling function, which allows to describe the spaces in the multiresolution analysis in terms of translates of linear combinations of the functions in the scaling vector. Multiwavelets came up in a natural way in fractal interpolation and iterated functions theory [5, 6, 7, 12] and were also put forward by Goodman [10, 11] and Hervé [13]. A good description of multiwavelets can be found in [17, part II] and [18]. In this paper we will investigate the multiwavelets described by Alpert [1]. Previously Geronimo, Marcellán and Iliev investigated a modification of Alpert's multiwavelets in [9] and [8], but they used an orthogonal basis for the wavelets with fewer zero moments than proposed by Alpert in [1]. In this paper we will investigate the wavelets with the maximal number of zero moments, as originally proposed by Alpert.

The paper is organized as follows. We give some information on Alpert's multiwavelets in section 2, we make the connection with type I multiple orthogonal polynomials in section 3 and give explicit formulas for the Legendre-Angelesco multiple orthogonal polynomials in section 4. A formula for Alpert's multiwavelets in terms of the Legendre-Angelesco polynomials is given in section 5 including explicit expressions for their Fourier transforms. An algorithm to expand the multiwavelets in terms of Legendre polynomials is given in section 6.

2. Alpert multiwavelets. Let $n \in \mathbb{N} = \{1, 2, 3, \dots\}$ be fixed and consider the functions

$$\phi_j(x) = \sqrt{2j-1} P_{j-1}(2x-1) \chi_{[0,1)}(x), \quad 1 \leq j \leq n,$$

*Version of October 17, 2016

[†]School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332-0160 (geronimo@math.gatech.edu, iliev@math.gatech.edu).

[‡]Department of Mathematics, KU Leuven, Celestijnenlaan 200B box 2400, BE-3001 Leuven, Belgium (walter@wis.kuleuven.be).

where P_j ($0 \leq j \leq n-1$) are the Legendre polynomials on $[-1, 1]$ and $\chi_{[0,1]}$ is the characteristic function of the interval $[0, 1)$. If we set

$$\Phi_n(x) = \begin{pmatrix} \phi_1(x) \\ \vdots \\ \phi_n(x) \end{pmatrix},$$

then Φ_n is a vector of compactly supported L^2 -functions with

$$\int_0^1 \Phi_n(x) \Phi_n^T(x) dx = \mathbb{I}_n,$$

where $\mathbb{I}_n$ is the identity matrix of order n . Introduce the linear space

$$V_0 = \text{cl}_{L^2(\mathbb{R})} \text{span}\{\phi_i(\cdot - j), 1 \leq i \leq n, j \in \mathbb{Z}\},$$

then V_0 is a space of piecewise polynomials of degree $\leq n-1$ and it is a finitely generated shift invariant space, for which $\{\phi_i(x-j), 1 \leq i \leq n, j \in \mathbb{Z}\}$ is an orthonormal basis. For every $k \in \mathbb{Z}$ we define $V_k = \{\phi(2^k \cdot), \phi \in V_0\}$, so that V_k contains functions in $L^2(\mathbb{R})$ at different resolutions. Clearly the spaces $(V_k)_{k \in \mathbb{Z}}$ are nested

$$\cdots \subset V_{-2} \subset V_{-1} \subset V_0 \subset V_1 \subset V_2 \subset \cdots \subset L^2(\mathbb{R}),$$

and they form a multiresolution analysis (MRA) of $L^2(\mathbb{R})$. We can write

$$V_{k+1} = V_k \oplus W_k, \quad k \in \mathbb{Z},$$

and the spaces $(W_k)_{k \in \mathbb{Z}}$ are called the wavelet spaces. If $\psi_1, \dots, \psi_n$ generate a shift-invariant basis for W_0 , then these functions are known as multiwavelets when $n > 1$. For $n = 1$ one retrieves the MRA associated with the Haar wavelet. Multiwavelets for $n > 1$ were investigated by Alpert [1], Geronimo and Marcellán [9], Geronimo and Iliev [8].

In [1] Alpert constructs multiwavelet functions $f_1, \dots, f_n$ supported on $[-1, 1]$ with the following properties:

1. The restriction of f_i to $(0, 1)$ is a polynomial of degree $n-1$;
2. $f_k(-t) = (-1)^{k+n-1} f_k(t)$ for $t \in (0, 1)$;
3. $f_1, \dots, f_n$ have the orthogonality property

$$(1) \quad \int_{-1}^1 f_i(t) f_j(t) dt = \delta_{i,j}, \quad 1 \leq i, j \leq n;$$

4. The function f_k has vanishing moments

$$(2) \quad \int_{-1}^1 f_k(t) t^i dt = 0, \quad i = 0, 1, \dots, k+n-2.$$

Our observation is the following

PROPOSITION 1. *The last function f_n is, up to a normalization factor, the type I Legendre-Angelesco multiple orthogonal polynomial*

$$f_n(x) = A_{n,n}(x) \chi_{[-1,0)} + B_{n,n}(x) \chi_{[0,1]}.$$

Of course we need to explain what the type I Legendre-Angelesco polynomial is. In the next section we explain what multiple orthogonal polynomials are and in particular we define the type I and type II Legendre-Angelesco polynomials. **Proposition 1** then follows immediately from the definition of type I Legendre Angelesco polynomials. We will then also explain how the other multiwavelets $f_1, \dots, f_{n-1}$ can be expressed in terms of the Legendre-Angelesco polynomials.

3. Multiple orthogonal polynomials. Let $r \in \mathbb{N}$ and $\vec{n} = (n_1, \dots, n_r) \in \mathbb{N}^r$ with size $|\vec{n}| = n_1 + n_2 + \dots + n_r$. Multiple orthogonal polynomials are polynomials in one variable that satisfy orthogonality conditions with respect to r measures $\mu_1, \dots, \mu_r$ on the real line. There are two types: the type I multiple orthogonal polynomials for the multi-index $\vec{n}$ are $(A_{\vec{n},1}, \dots, A_{\vec{n},r})$, where the degree of $A_{\vec{n},j}$ is $\leq n_j - 1$ and

$$\sum_{j=1}^r \int x^k A_{\vec{n},j}(x) d\mu_j(x) = 0, \quad 0 \leq k \leq |\vec{n}| - 2,$$

with the normalization

$$\sum_{j=1}^r \int x^{|\vec{n}|-1} A_{\vec{n},j}(x) d\mu_j(x) = 1,$$

and the type II multiple orthogonal $P_{\vec{n}}$ is the monic polynomial of degree $|\vec{n}|$ for which

$$\int x^k P_{\vec{n}}(x) d\mu_j(x) = 0, \quad 0 \leq k \leq n_j - 1,$$

for $1 \leq j \leq r$ (see, e.g., [14, Chapter 26]). If the measures are supported on disjoint intervals, then the system $\mu_1, \dots, \mu_r$ is called an Angelesco system. In this paper we will deal with two measures ($r = 2$) with μ_1 the uniform measure on $[-1, 0]$ and μ_2 the uniform measure on $[0, 1]$, and the corresponding multiple orthogonal polynomials are called Legendre-Angelesco polynomials. These polynomials were introduced by Angelesco in 1918–1919 and investigated in detail by Kalyagin and Ronveaux [16, 15].

The type I Legendre-Angelesco polynomials $(A_{n,m}, B_{n,m})$ are thus defined by $\deg A_{n,m} = n - 1$, $\deg B_{n,m} = m - 1$ and the orthogonality conditions

$$(3) \quad \int_{-1}^1 \left(A_{n,m}(x) \chi_{[-1,0]} + B_{n,m}(x) \chi_{[0,1]} \right) x^k dx = 0, \quad 0 \leq k \leq n + m - 2,$$

and the normalization

$$(4) \quad \int_{-1}^1 \left(A_{n,m}(x) \chi_{[-1,0]} + B_{n,m}(x) \chi_{[0,1]} \right) x^{n+m-1} dx = 1.$$

These polynomials are uniquely determined by these conditions. Note that the orthogonality (3) corresponds to the condition for the vanishing moments (condition 4) for $k = n$, hence proving Proposition 1. The normalization, however, is different since one uses condition 3 instead of (4). Note that the type I Legendre-Angelesco polynomials are piecewise polynomials on $[-1, 1]$ but their restrictions to $[-1, 0]$ and $[0, 1]$ are respectively the polynomials $A_{n,m}$ (of degree $n - 1$) and $B_{n,m}$ (of degree $m - 1$).

The remaining multiwavelets $f_1, \dots, f_{n-1}$ can also be expressed in terms of Legendre-Angelesco orthogonal polynomials. Denote the type I multiple orthogonal polynomials by

$$Q_{n,m}(x) = A_{n,m}(x) \chi_{[-1,0]} + B_{n,m}(x) \chi_{[0,1]}.$$

As mentioned before, these are piecewise polynomials on $[-1, 1]$ as are the multiwavelets.

PROPOSITION 2. *Let $(f_1, \dots, f_n)$ be the Alpert multiwavelets of multiplicity n . Then*

$$f_k(x) = \begin{cases} \sum_{j=\frac{k+n}{2}}^n c_j Q_{j,j}(x), & \text{if } k+n \text{ is even,} \\ \frac{1}{2} \sum_{j=\frac{k+n+1}{2}}^n a_j d_{j-1} Q_{j,j}(x) + \sum_{j=\frac{k+n-1}{2}}^{n-1} d_j Q_{j+1,j}(x), & \text{if } k+n \text{ is odd,} \end{cases}$$

where the coefficients a_j, c_j, d_j are real numbers. The a_j are independent of k and n , but the c_j, d_j depend on k and n .

Proof. The type I multiple orthogonal polynomials $\{Q_{j,j}, 1 \leq j \leq n\}$ on the diagonal, together with $\{Q_{j+1,j}, 0 \leq j \leq n-1\}$ are a basis for the piecewise polynomials $p_{n-1}\chi_{[-1,0]} + q_{n-1}\chi_{[0,1]}$, where $p_{n-1}, q_{n-1} \in \mathbb{P}_{n-1}$ are polynomials of degree at most $n-1$. There is a biorthogonality relation for the multiple orthogonal polynomials [14, §23.1.3]

$$\int_{-1}^1 P_{n,m}(x) Q_{k,\ell}(x) dx = \begin{cases} 0 & \text{if } k \leq n \text{ and } \ell \leq m, \\ 0 & \text{if } n+m \leq k+\ell-2, \\ 1 & \text{if } n+m = k+\ell-1, \end{cases}$$

where $P_{n,m}$ are the type II Legendre-Angelesco multiple orthogonal polynomials, for which

$$\begin{aligned} \int_{-1}^0 P_{n,m}(x) x^k dx &= 0, & 0 \leq k \leq n-1, \\ \int_0^1 P_{n,m}(x) x^k dx &= 0, & 0 \leq k \leq m-1. \end{aligned}$$

The biorthogonality gives for the multi-indices near the diagonal

$$\int_{-1}^1 P_{j,j}(x) Q_{k,k}(x) dx = 0 = \int_{-1}^1 P_{j+1,j}(x) Q_{k+1,k}(x) dx, \quad j, k \in \mathbb{N},$$

$$\int_{-1}^1 P_{j,j}(x) Q_{k+1,k}(x) dx = \delta_{k,j} = \int_{-1}^1 P_{j,j-1}(x) Q_{k,k}(x) dx.$$

So if we expand f_k in the basis of type I Legendre-Angelesco polynomials, then

$$(5) \quad f_k(x) = \sum_{j=1}^n c_j Q_{j,j}(x) + \sum_{j=0}^{n-1} d_j Q_{j+1,j}(x),$$

and the biorthogonality gives

$$(6) \quad c_j = \int_{-1}^1 f_k(x) P_{j,j-1}(x) dx, \quad d_j = \int_{-1}^1 f_k(x) P_{j,j}(x) dx.$$

The conditions imposed by Alpert give some properties for these coefficients. The symmetry property 2 gives (change variables $x \rightarrow -x$)

$$d_j = \int_{-1}^1 f_k(-x) P_{j,j}(-x) dx = (-1)^{k+n+1} \int_{-1}^1 f_k(x) P_{j,j}(x) dx = (-1)^{k+n+1} d_j,$$

so that $d_j = 0$ whenever $k + n$ is even. We have used that the type II multiple orthogonal polynomial $P_{j,j}$ is an even function. For c_j we have (with the same change of variables)

$$c_j = \int_{-1}^1 f_k(-x) P_{j,j-1}(-x) dx = (-1)^{k+n} \int_{-1}^1 f_k(x) P_{j-1,j}(x) dx,$$

where we used that $P_{j,j-1}(-x) = -P_{j-1,j}(x)$, which can easily be seen from the orthogonality conditions. The type II multiple orthogonal polynomials satisfy the nearest neighbor recurrence relations

$$\begin{aligned} xP_{n,m}(x) &= P_{n+1,m}(x) + c_{n,m}P_{n,m}(x) + a_{n,m}P_{n-1,m}(x) + b_{n,m}P_{n,m-1}(x), \\ xP_{n,m}(x) &= P_{n,m+1}(x) + d_{n,m}P_{n,m}(x) + a_{n,m}P_{n-1,m}(x) + b_{n,m}P_{n,m-1}(x). \end{aligned}$$

Subtracting both relations gives

$$P_{n+1,m}(x) - P_{n,m+1}(x) = (d_{n,m} - c_{n,m})P_{n,m}(x),$$

so that $P_{j,j-1}(x) - P_{j-1,j}(x) = a_j P_{j-1,j-1}(x)$, where $a_j = d_{j-1,j-1} - c_{j-1,j-1}$. Using this we find

$$c_j = (-1)^{k+n} \int_{-1}^1 f_k(x) (P_{j,j-1}(x) - a_j P_{j-1,j-1}(x)) dx = (-1)^{k+n} (c_j - a_j d_{j-1}),$$

and hence we have that $2c_j = a_j d_{j-1}$ whenever $k + n$ is odd. So the expansion (5) reduces to

$$(7) \quad f_k(x) = \begin{cases} \sum_{j=1}^n c_j Q_{j,j}(x), & \text{if } k + n \text{ is even,} \\ \frac{1}{2} \sum_{j=1}^n a_j d_{j-1} Q_{j,j}(x) + \sum_{j=0}^{n-1} d_j Q_{j+1,j}(x), & \text{if } k + n \text{ is odd.} \end{cases}$$

Alpert's moment condition 4 also shows that

$$\begin{cases} c_j = 0, & \text{if } 2j - 1 \leq k + n - 2, \\ d_j = 0, & \text{if } 2j \leq k + n - 2, \end{cases}$$

so that the expansion (7) reduces to the one given in the proposition. $\square$

4. Legendre-Angelesco polynomials. It is now important to find the Legendre-Angelesco polynomials. We introduce two families of polynomials

$$(8) \quad p_n(x) = \sum_{k=0}^n \binom{n}{k} \binom{n + \frac{k}{2}}{n} (-1)^{n-k} x^k,$$

$$(9) \quad q_n(x) = \sum_{k=0}^n \binom{n}{k} \binom{n + \frac{k-1}{2}}{n} (-1)^{n-k} x^k.$$

Observe that p_n and q_n are polynomials of degree n with positive leading coefficient. These two families are immediately related to the last two multiwavelets, and this holds for every multiplicity n . We will need the Mellin transform of these polynomials to make the connection.

PROPOSITION 3. *The Mellin transforms of the polynomials p_n and q_n on $[0, 1]$ are given by*

$$(10) \quad \int_0^1 p_n(x) x^s dx = (-1)^n \frac{(\frac{1}{2} - \frac{s}{2})_n}{(s+1)_{n+1}},$$

and

$$(11) \quad \int_0^1 q_n(x) x^s dx = (-1)^n \frac{(-\frac{s}{2})_n}{(s+1)_{n+1}}.$$

Proof. The proof is along the lines of [8, Lemma 4] and [23, Thm. 2.2]. We will compute the Mellin transform of p_n , which by integrating (8) is

$$\int_0^1 p_n(x) x^s dx = \sum_{k=0}^n \binom{n}{k} \binom{n + \frac{k}{2}}{n} \frac{(-1)^{n-k}}{k+s+1}.$$

This is the partial fraction decomposition of a rational function $T(s)/R(s)$ with poles at the integers $\{-k-1, 0 \leq k \leq n\}$, so $R(s) = (s+1)_{n+1}$ and T is a polynomial of degree $\leq n$ with residue $(-1)^{n-k} \binom{n}{k} \binom{n+\frac{k}{2}}{n}$ for the pole at $-k-1$, hence

$$\frac{T(-k-1)}{R'(-k-1)} = \binom{n}{k} \binom{n + \frac{k}{2}}{n} (-1)^{n-k}.$$

Observe that $R'(-k-1) = (-1)^k k!(n-k)!$ so that

$$T(-k-1) = (-1)^n \binom{n + \frac{k}{2}}{n} n! = (-1)^n \left(\frac{k}{2} + 1\right)_n, \quad 0 \leq k \leq n.$$

The polynomial $T(s) = (-1)^n (\frac{1}{2} - \frac{s}{2})_n$ of degree n has precisely these values at $-k-1$, and hence (10) follows. Observe that the Mellin transform of p_n is a rational function of s and that the numerator has zeros when $s = 2k-1$ ($1 \leq k \leq n$), so that all the odd moments $\leq 2n-1$ of p_n on $[0, 1]$ vanish. The Mellin transform of q_n can be obtained easily in a similar way. From (11) one sees that q_n has vanishing even moments $\leq 2n-2$. $\square$

The last two multiwavelets f_n and f_{n-1} are now easily given in terms of these polynomials.

PROPOSITION 4. *Let $f_1, \dots, f_n$ be Alpert's multiwavelets of multiplicity n . Then for $x \in [0, 1]$*

$$f_n(x) = c_{n,0} p_{n-1}(x), \quad f_{n-1}(x) = d_{n,0} q_{n-1}(x),$$

where $c_{n,0}$ and $d_{n,0}$ are normalizing constants given by

$$\frac{1}{c_{n,0}^2} = 2 \int_0^1 p_{n-1}^2(x) dx, \quad \frac{1}{d_{n,0}^2} = 2 \int_0^1 q_{n-1}^2(x) dx.$$

Proof. We need to verify the moment condition (2) for f_n . Observe that

$$\begin{aligned} \int_{-1}^1 f_n(x) x^k dx &= \int_0^1 f_n(x) x^k dx + \int_{-1}^0 f_n(x) x^k dx \\ &= (1 + (-1)^{k+1}) \int_0^1 f_n(x) x^k dx, \end{aligned}$$

so that all the even moments are already zero. So we only need to concentrate on the odd moments of p_{n-1} . The Mellin transform (10) for p_{n-1} shows that the odd moments $\leq 2n-3$ vanish, and since all the even moments for f_n are already zero, it must follow from (2) that f_n is equal to p_{n-1} up to a normalizing factor.

The proof for f_{n-1} is similar with a few changes. The symmetry $f_{n-1}(-x) = f_{n-1}(x)$ now gives

$$\int_{-1}^1 f_{n-1}(x)x^k dx = (1 + (-1)^k) \int_0^1 f_{n-1}(x)x^k dx,$$

so that all the odd moments vanish. We only need to check that the even moments vanish. The orthogonality of f_n and f_{n-1} will automatically be true because of the symmetry:

$$\int_{-1}^1 f_n(x)f_{n-1}(x) dx = \int_0^1 f_n(x)f_{n-1}(x) dx - \int_0^1 f_n(x)f_{n-1}(x) dx = 0.$$

For q_{n-1} we have from (11) that all the even moments $\leq 2n-4$ vanish, so that together with the vanishing odd moments, the moment condition (2) holds and f_{n-1} must be proportional to q_{n-1} . $\square$

An explicit formula for the normalizing constants $c_{n,0}$ and $d_{n,0}$ will be given in Proposition 8.

4.1. Type I Legendre-Angelesco polynomials. The type I Legendre-Angelesco polynomials near the diagonal can be expressed explicitly in terms of the functions p_n and q_n in (8)–(9):

PROPOSITION 5. *Let $(A_{n,m}, B_{n,m})$ be the type I Legendre-Angelesco polynomial for the multi-index (n, m) . Then on the diagonal*

$$(12) \quad B_{n+1,n+1}(x) = \frac{1}{2} \frac{(3n+2)!}{n!(2n+1)!} p_n(x), \quad A_{n+1,n+1}(x) = -B_{n+1,n+1}(-x).$$

and for $|n-m|=1$

$$(13) \quad b_n B_{n+1,n}(x) = \binom{n+\frac{n}{2}}{n} q_n(x) - \binom{n+\frac{n-1}{2}}{n} p_n(x),$$

$$(14) \quad b_n B_{n,n+1}(x) = \binom{n+\frac{n}{2}}{n} q_n(x) + \binom{n+\frac{n-1}{2}}{n} p_n(x),$$

and

$$(15) \quad A_{n+1,n}(x) = B_{n,n+1}(-x), \quad A_{n,n+1}(x) = B_{n+1,n}(-x),$$

where the normalizing constant is given by $b_n = 2(\frac{n}{2} + 1)_n \frac{(2n)!}{(3n+1)!}$.

Proof. The result in (12) follows from Proposition 1 and Proposition 4. The constant can be found by putting $s = 2n+1$ in the Mellin transform, since that moment has to be 1. For (13)–(15) one can check that the vanishing moment conditions (3) hold and that the polynomials are of the correct degree. The normalizing constant b_n can be obtained by computing the $2n$ th moment of q_n , i.e., putting $s = 2n$ in (11), and the using the normalization in (4). $\square$

A consequence of the Mellin transform (10) is the Rodrigues type formula

$$p_n(x) = \frac{(-1)^n}{2^n n!} (D^*)^n x^{2n} (1-x)^n, \quad D^* = \frac{1}{x} \frac{d}{dx},$$

which follows by using

$$(s - 2k - 1)\hat{f}(s) = -\widehat{D_k f}(s), \quad D_k f(x) = x^{-2k-1} \frac{d}{dx} x^{2k+2} f(x),$$

where $\hat{f}$ is the Mellin transform of f

$$\hat{f}(s) = \int_0^\infty f(x) x^s dx,$$

(see, e.g., [23, §2.2]). In a similar way the Mellin transform (11) implies the Rodrigues type formula

$$q_n(x) = \frac{(-1)^n}{2^n n!} x (D^*)^n x^{2n-1} (1-x)^n.$$

The p_n and q_n can be written as linear combinations of two hypergeometric functions

$$p_n(x) = (-1)^n {}_3F_2 \left(\begin{matrix} n+1, -\frac{n}{2}, -\frac{n}{2} + \frac{1}{2} \\ \frac{1}{2}, 1 \end{matrix} ; x^2 \right) - (-1)^n \frac{(\frac{3}{2})_n}{(n-1)!} x {}_3F_2 \left(\begin{matrix} n + \frac{3}{2}, -\frac{n}{2} + \frac{1}{2}, -\frac{n}{2} + 1 \\ \frac{3}{2}, \frac{3}{2} \end{matrix} ; x^2 \right),$$

and

$$q_n(x) = (-1)^n \frac{(\frac{1}{2})_n}{n!} {}_3F_2 \left(\begin{matrix} n + \frac{1}{2}, -\frac{n}{2}, -\frac{n}{2} + \frac{1}{2} \\ \frac{1}{2}, \frac{1}{2} \end{matrix} ; x^2 \right) - (-1)^n n x {}_3F_2 \left(\begin{matrix} n+1, -\frac{n}{2} + \frac{1}{2}, -\frac{n}{2} + 1 \\ 1, \frac{3}{2} \end{matrix} ; x^2 \right).$$

There is also a system of recurrence relations.

PROPOSITION 6. *One has*

$$(16) \quad (3n-1)x p_{n-1}(x) = 2n q_n(x) + (2n-1)q_{n-1}(x),$$

$$(17) \quad (3n-2)x q_{n-1}(x) = \frac{2}{3} \left(n p_n(x) + (2n-1)p_{n-1}(x) + (n-1)p_{n-2}(x) \right).$$

Proof. We will prove this by comparing coefficients. For (16) the coefficient of $(-x)^k$ on the right hand side is

$$(-1)^n \left(2n \binom{n}{k} \binom{n + \frac{k-1}{2}}{n} - (2n-1) \binom{n-1}{k} \binom{n-1 + \frac{k-1}{2}}{n-1} \right).$$

This can easily be seen to be equal to

$$(-1)^n \frac{(1 + \frac{k-1}{2})_{n-1}}{k!(n-k)!} \left(2n(n + \frac{k-1}{2}) - (2n-1)(n-k) \right).$$

The factor between brackets is $k(3n-1)$, hence together this gives

$$(-1)^n(3n-1)\frac{(1+\frac{k-1}{2})_{n-1}}{(k-1)!(n-k)!} = (-1)^n(3n-1)\binom{n-1}{k-1}\binom{n-1+\frac{k-1}{2}}{n-1},$$

which is the coefficient of $(-x)^k$ of the left hand side of (16). The proof is similar for (17), except that one now has to combine three terms on the right hand side. $\square$

These recurrence relations can be simplified in the following way. Introduce one more sequence of polynomials

$$(18) \quad r_n(x) = \sum_{k=0}^n \binom{n}{k} \binom{n+\frac{k+1}{2}}{n} (-1)^{n-k} x^k,$$

then

$$\begin{aligned} x r_{n-1}(x) &= \frac{2}{3} (p_{n-1}(x) + p_n(x)), \\ (3n-1)x p_{n-1}(x) &= 2n q_n(x) + (2n-1) q_{n-1}(x), \\ (3n+1) q_n(x) &= (n+1) r_n(x) + n r_{n-1}(x), \end{aligned}$$

or in matrix form

$$\begin{pmatrix} p_n(x) \\ q_n(x) \\ r_n(x) \end{pmatrix} = \begin{pmatrix} -1 & 0 & \frac{3}{2}x \\ \frac{3n-1}{2n}x & -\frac{2n-1}{2n} & 0 \\ \frac{(3n+1)(3n-1)}{2n(n+1)}x & -\frac{(3n+1)(2n-1)}{2n(n+1)} & -\frac{n}{n+1} \end{pmatrix} \begin{pmatrix} p_{n-1}(x) \\ q_{n-1}(x) \\ r_{n-1}(x) \end{pmatrix}.$$

4.2. Type II Legendre-Angelesco polynomials. The type II Legendre-Angelesco polynomials $P_{n,n}(x)$ and $P_{n+1,n}(x)$ also have nice Mellin transforms.

PROPOSITION 7. *One has*

$$(19) \quad \int_0^1 P_{n,n}(x) x^s dx = C_n \frac{(-s)_n}{(\frac{1}{2} + \frac{s}{2})_{n+1}}$$

and

$$(20) \quad \int_0^1 [P_{n+1,n}(x) + P_{n,n+1}(x)] x^s dx = D_n \frac{(-s)_n}{(1 + \frac{s}{2})_{n+1}},$$

where C_n and D_n are constants.

Proof. The orthogonality conditions for $P_{n,n}$ are

$$\int_{-1}^0 P_{n,n}(x) x^k dx = 0 = \int_0^1 P_{n,n}(x) x^k dx, \quad 0 \leq k \leq n-1.$$

The symmetry $P_{n,n}(-x) = P_{n,n}(x)$ implies that $P_{n,n}(x)$ is an even polynomial and we only need to consider the orthogonality conditions on $[0, 1]$. Let

$$P_{n,n}(x) = \sum_{j=0}^n a_j x^{2j},$$

then the Mellin transform is

$$\int_0^1 P_{n,n}(x) x^s dx = \sum_{j=0}^n \frac{a_j}{2j + s + 1}.$$

This is a rational function of s with poles at the odd negative integers $-1, -3, -5, \dots, -2n - 1$, hence the denominator is $(\frac{1}{2} + \frac{s}{2})_{n+1}$. The orthogonality conditions on $[0, 1]$ imply that the numerator has zeros at $0, 1, \dots, n - 1$, hence the numerator is proportional to $(-s)_n$. This gives (19), where the constant C_n has to be determined so that $P_{n,n}$ is a monic polynomial.

The orthogonality conditions for $P_{n+1,n}$ and $P_{n,n+1}$ are

$$\int_{-1}^0 P_{n+1,n}(x) x^k dx = 0 = \int_0^1 P_{n,n+1}(x) x^k dx, \quad 0 \leq k \leq n,$$

and

$$\int_0^1 P_{n+1,n}(x) x^k dx = 0 = \int_{-1}^0 P_{n,n+1}(x) x^k dx, \quad 0 \leq k \leq n - 1.$$

The difference $P_{n+1,n}(x) - P_{n,n+1}(x)$ is a polynomial of degree $2n$ which is proportional to $P_{n,n}$, and $P_{n+1,n}(-x) = -P_{n,n+1}(x)$. The sum $P_{n+1,n} + P_{n,n+1}$ is therefore an odd polynomial and we can write

$$P_{n+1,n}(x) + P_{n,n+1}(x) = x \sum_{j=0}^n b_j x^{2j}.$$

The Mellin transform of this sum is of the form

$$\int_0^1 [P_{n+1,n}(x) + P_{n,n+1}(x)] x^s dx = \sum_{j=0}^n \frac{b_j}{2j + s + 2},$$

and this is a rational function with poles at the even negative integers $-2, -4, -6, \dots, -2n - 2$. Hence the denominator is $(1 + \frac{s}{2})_{n+1}$. The orthogonality conditions on $[0, 1]$ imply that the numerator vanishes at $0, 1, \dots, n - 1$, hence it is proportional to $(-s)_n$. This gives (20), where the constant D_n has to be determined so that the leading coefficient is 2. $\square$

The Mellin transform (19) implies the Rodrigues formula

$$P_{n,n}(x) = (-1)^n \frac{(2n)!}{(3n)!} \frac{d^n}{dx^n} x^n (1 - x^2)^n,$$

if we take into account that

$$\int_0^1 (1 - x^2)^n x^s dx = \frac{1}{2} \frac{n!}{(\frac{1}{2} + \frac{s}{2})_{n+1}}.$$

This Rodrigues formula was already known, e.g., [14, §23.3] [16], and it gives the explicit expression

$$P_{n,n}(x) = \frac{n!(2n)!}{(3n)!} \sum_{k=0}^n \binom{n}{k} \binom{n+2k}{2k} (-1)^{n-k} x^{2k}.$$

This Rodrigues was already given by Appell in 1901 without any mentioning of the property of multiple orthogonality, see [2] [3, p. 433].

5. Alpert multiwavelets and Legendre-Angelesco polynomials. **Proposition 4** already gives the last two multiwavelets f_n and f_{n-1} in terms of the polynomials p_{n-1} and q_{n-1} . In order to be able to work with multiwavelets of various multiplicity, we will change the notation somewhat. If $n \in \mathbb{N}$ we will denote the multiwavelets of multiplicity n by $(f_1^n, f_2^n, \dots, f_n^n)$, and these correspond to what we so far denoted as $(f_1, f_2, \dots, f_n)$. **Proposition 4** then says that for $x \in [0, 1]$

$$f_{n+1}^{n+1}(x) = c_{n+1,0}p_n(x), \quad f_n^{n+1}(x) = d_{n+1,0}q_n(x).$$

To find the other multiwavelets, we proceed as follows. Every f_{n+1-2k}^{n+1} is a linear combination of $p_n, p_{n-1}, \dots, p_{n-k}$ in such a way that the orthogonality (1) holds. This means that we start from the sequence $(p_n, p_{n-1}, \dots, p_{n-k})$ and we use the Gram-Schmidt process to obtain $(f_{n+1}^{n+1}, f_{n-1}^{n+1}, \dots, f_{n+1-2k}^{n+1})$. In a similar way every f_{n-2k}^{n+1} is a linear combination of q_n, q_{n-1}, q_{n-k} in such a way that the orthogonality (1) holds. Hence we use Gram-Schmidt on $(q_n, q_{n-1}, \dots, q_{n-k})$ to obtain $(f_n^{n+1}, f_{n-2}^{n+1}, \dots, f_{n-2k}^{n+1})$. Observe that the Gram-Schmidt process is applied to the polynomials $(p_n)_n$ and $(q_n)_n$ starting from degree n and going down in the degree, which is different from the usual procedure where one starts from the lowest degree, going up to the highest degree. If we denote by $\langle f, g \rangle$ the integral of fg over the interval $[0, 1]$, then an explicit formula for $x \in (0, 1]$ is

$$(21) \quad f_{n+1-2k}^{n+1}(x) = c_{n+1,k} \begin{vmatrix} \langle p_n, p_n \rangle & \langle p_n, p_{n-1} \rangle & \cdots & \langle p_n, p_{n-k} \rangle \\ \langle p_{n-1}, p_n \rangle & \langle p_{n-1}, p_{n-1} \rangle & \cdots & \langle p_{n-1}, p_{n-k} \rangle \\ \vdots & \vdots & \cdots & \vdots \\ \langle p_{n-k+1}, p_n \rangle & \langle p_{n-k+1}, p_{n-1} \rangle & \cdots & \langle p_{n-k+1}, p_{n-k} \rangle \\ p_n(x) & p_{n-1}(x) & \cdots & p_{n-k}(x) \end{vmatrix},$$

where $c_{n+1,k}$ is such that the norm of f_{n+1-2k}^{n+1} is one. Similarly

$$(22) \quad f_{n-2k}^{n+1}(x) = d_{n+1,k} \begin{vmatrix} \langle q_n, q_n \rangle & \langle q_n, q_{n-1} \rangle & \cdots & \langle q_n, q_{n-k} \rangle \\ \langle q_{n-1}, q_n \rangle & \langle q_{n-1}, q_{n-1} \rangle & \cdots & \langle q_{n-1}, q_{n-k} \rangle \\ \vdots & \vdots & \cdots & \vdots \\ \langle q_{n-k+1}, q_n \rangle & \langle q_{n-k+1}, q_{n-1} \rangle & \cdots & \langle q_{n-k+1}, q_{n-k} \rangle \\ q_n(x) & q_{n-1}(x) & \cdots & q_{n-k}(x) \end{vmatrix},$$

where the constant $d_{n+1,k}$ normalizes the function to have norm one. For the normalizing constant $c_{n+1,k}$ we define

$$(23) \quad \Delta_{n,k}^p = \begin{vmatrix} \langle p_n, p_n \rangle & \langle p_n, p_{n-1} \rangle & \cdots & \langle p_n, p_{n-k} \rangle \\ \langle p_{n-1}, p_n \rangle & \langle p_{n-1}, p_{n-1} \rangle & \cdots & \langle p_{n-1}, p_{n-k} \rangle \\ \vdots & \vdots & \cdots & \vdots \\ \langle p_{n-k}, p_n \rangle & \langle p_{n-k}, p_{n-1} \rangle & \cdots & \langle p_{n-k}, p_{n-k} \rangle \end{vmatrix},$$

then it follows from (21) that $\langle f_{n+1-2k}^{n+1}, p_{n-j} \rangle = 0$ for $0 \leq j < k$, since this is equal to a determinant with two equal rows, and $\langle f_{n+1-2k}^{n+1}, p_{n-k} \rangle = c_{n+1,k} \Delta_{n,k}^p$. Now on $[0, 1]$ we have $f_{n+1-2k}^{n+1}(x) = c_{n+1,k} \Delta_{n,k}^p p_{n-k}(x) + \cdots$, and by the symmetry properties of the functions f_k^{n+1}

$$\int_{-1}^1 [f_{n+1-2k}^{n+1}(x)]^2 dx = 2 \int_0^1 [f_{n+1-2k}^{n+1}(x)]^2 dx,$$

hence

$$\frac{1}{2} = \langle f_{n+1-2k}^{n+1}, f_{n+1-2k}^{n+1} \rangle = c_{n+1,k}^2 \Delta_{n,k}^p \Delta_{n,k-1}^p,$$

so that

$$(24) \quad c_{n+1,k} = \frac{1}{\sqrt{2\Delta_{n,k}^p \Delta_{n,k-1}^p}}.$$

In a similar way we find

$$(25) \quad d_{n+1,k} = \frac{1}{\sqrt{2\Delta_{n,k}^q \Delta_{n,k-1}^q}},$$

where

$$(26) \quad \Delta_{n,k}^q = \det \begin{pmatrix} \langle q_n, q_n \rangle & \langle q_n, q_{n-1} \rangle & \cdots & \langle q_n, q_{n-k} \rangle \\ \langle q_{n-1}, q_n \rangle & \langle q_{n-1}, q_{n-1} \rangle & \cdots & \langle q_{n-1}, q_{n-k} \rangle \\ \vdots & \vdots & \cdots & \vdots \\ \langle q_{n-k}, q_n \rangle & \langle q_{n-k}, q_{n-1} \rangle & \cdots & \langle q_{n-k}, q_{n-k} \rangle \end{pmatrix}.$$

5.1. Hypergeometric functions. We will express the inner products $\langle p_n, p_k \rangle$ in terms of hypergeometric functions.

PROPOSITION 8. *The entries in the Gram-matrix for $k \leq n$ are given by*

$$(27) \quad \int_0^1 p_n(x) p_k(x) dx = (-1)^{n+k} \frac{\Gamma(n + \frac{1}{2})}{(n+1)! \sqrt{\pi}} {}_4F_3 \left(\begin{matrix} k+1, \frac{1}{2}, -\frac{k-1}{2}, -\frac{k}{2} \\ -n + \frac{1}{2}, \frac{n+2}{2}, \frac{n+3}{2} \end{matrix} ; 1 \right),$$

and

$$(28) \quad \int_0^1 q_n(x) q_k(x) dx = (-1)^{n+k} \frac{\Gamma(n - \frac{1}{2})k}{2(n+2)! \sqrt{\pi}} {}_4F_3 \left(\begin{matrix} k+1, \frac{3}{2}, -\frac{k-2}{2}, -\frac{k-1}{2} \\ -n + \frac{3}{2}, \frac{n+3}{2}, \frac{n+4}{2} \end{matrix} ; 1 \right).$$

Proof. For (27) we expand p_k using (8) to find

$$\langle p_n, p_k \rangle = (-1)^k \sum_{j=0}^k \binom{k}{j} \binom{k + \frac{j}{2}}{k} (-1)^j \int_0^1 x^j p_n(x) dx.$$

The integral can be evaluated by using the Mellin transform (10), which gives

$$\int_0^1 x^j p_n(x) dx = (-1)^n \frac{(\frac{1}{2} - \frac{j}{2})_n}{(1+j)_{n+1}},$$

and this is zero whenever j is odd and $1 \leq j \leq 2n-1$. So we only need to consider even terms in the sum. This leaves

$$\begin{aligned} \langle p_n, p_k \rangle &= (-1)^{n+k} \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} \binom{k}{2j} \binom{k+j}{k} \frac{(\frac{1}{2} - \frac{j}{2})_n}{(1+2j)_{n+1}} \\ &= (-1)^{n+k} \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} \frac{(k+j)! (\frac{1}{2} - j)_n}{(k-2j)! j! (2j+n+1)!}. \end{aligned}$$

Now we can write

$$\left(\frac{1}{2} - j\right)_n = \left(\frac{1}{2}\right)_j \frac{\left(\frac{1}{2}\right)_n}{\left(-n + \frac{1}{2}\right)_j}$$

and $(k + j)! = (k + 1)_j k!$, so that

$$\langle p_n, p_k \rangle = (-1)^{n+k} \left(\frac{1}{2}\right)_n k! \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} \frac{(k + 1)_j \left(\frac{1}{2}\right)_j}{(k - 2j)! (2j + n + 1)! \left(-n + \frac{1}{2}\right)_j j!}.$$

Next, we have

$$(k - 2j)! = \frac{k!}{(-k)_{2j}} = \frac{k!}{2^{2j} \left(-\frac{k}{2}\right)_j \left(-\frac{k-1}{2}\right)_j},$$

and $(2j + n + 1)! = 2^{2j} \left(\frac{n+2}{2}\right)_j \left(\frac{n+3}{2}\right)_j (n + 1)!$, so that

$$\langle p_n, p_k \rangle = (-1)^{n+k} \frac{\left(\frac{1}{2}\right)_n}{(n + 1)!} \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} \frac{(k + 1)_j \left(-\frac{k}{2}\right)_j \left(-\frac{k-1}{2}\right)_j \left(\frac{1}{2}\right)_j}{\left(-n + \frac{1}{2}\right)_j \left(\frac{n+2}{2}\right)_j \left(\frac{n+3}{2}\right)_j j!},$$

and this coincides with the hypergeometric expression in (27). The computations for $\langle q_n, q_k \rangle$ are similar, starting from the expansion (9) and using the Mellin transform (11). $\square$

Observe that both (27) and (28) are terminating hypergeometric series.

5.2. Fourier transform. We will now compute the Fourier transform of the multiwavelets f_k^{n+1} for $k = n + 1$ and $k = n$.

PROPOSITION 9. *One has*

$$\hat{f}_{n+1}^{n+1}(t) = \int_{-1}^1 f_{n+1}^{n+1}(x) e^{ixt} dx = 2ic_{n+1,0} t^{2n+1} \frac{(-1)^n n!}{(3n + 2)!} {}_1F_2 \left(\begin{matrix} n + 1 \\ \frac{3n+3}{2}, \frac{3n+4}{2} \end{matrix} ; -\frac{t^2}{4} \right),$$

and

$$\hat{f}_n^{n+1}(t) = \int_{-1}^1 f_n^{n+1}(x) e^{ixt} dx = 2d_{n+1,0} t^{2n} \frac{(-1)^n n!}{(3n + 1)!} {}_1F_2 \left(\begin{matrix} n + 1 \\ \frac{3n+2}{2}, \frac{3n+3}{2} \end{matrix} ; -\frac{t^2}{4} \right).$$

Proof. We have

$$\begin{aligned} \int_{-1}^1 f_{n+1}^{n+1}(x) e^{ixt} dx &= \int_0^1 f_{n+1}^{n+1}(x) e^{ixt} dx - \int_0^1 f_{n+1}^{n+1}(x) e^{-ixt} dx \\ &= 2i \int_0^1 f_{n+1}^{n+1}(x) \sin(xt) dx. \end{aligned}$$

If we use the Taylor series expansion of the sin function, then the Fourier transform is

$$\hat{f}_{n+1}^{n+1}(t) = 2i \sum_{k=0}^{\infty} (-1)^k \frac{t^{2k+1}}{(2k + 1)!} \int_0^1 x^{2k+1} f_{n+1}^{n+1}(x) dx.$$

On $(0, 1]$ we have $f_{n+1}^{n+1}(x) = c_{n+1,0} p_n(x)$, so we can use the Mellin transform in (10) to evaluate the integral:

$$\hat{f}_{n+1}^{n+1}(t) = (-1)^n 2ic_{n+1,0} \sum_{k=0}^{\infty} (-1)^k \frac{t^{2k+1}}{(2k + 1)!} \frac{(-k)_n}{(2k + 2)_{n+1}}.$$

The first n terms in the series are zero, so if we set $k = j + n$, then

$$\hat{f}_{n+1}^{n+1}(t) = 2ic_{n+1,0}t^{2n+1} \sum_{j=0}^{\infty} (-1)^j \frac{t^{2j}}{(2j+2n+1)!} \frac{(-j-n)_n}{(2j+2n+2)_{n+1}}.$$

Now use

$$(2j+2n+1)!(2j+2n+2)_{n+1} = (2j+3n+2)! = (3n+2)! 2^{2j} \left(\frac{3n+3}{2}\right)_j \left(\frac{3n+4}{2}\right)_j,$$

and $(-j-n)_n = (-1)^n n!(n+1)_j/j!$ to find

$$\hat{f}_{n+1}^{n+1}(t) = 2ic_{n+1,0}t^{2n+1} \frac{(-1)^n n!}{(3n+2)!} \sum_{j=0}^{\infty} (-1)^j \frac{t^{2j}}{2^{2j}} \frac{(n+1)_j}{j! \left(\frac{3n+3}{2}\right)_j \left(\frac{3n+4}{2}\right)_j},$$

which can be identified in terms of the hypergeometric series.

The proof for the second Fourier transform is similar. We now have

$$\hat{f}_n^{n+1}(t) = 2 \int_0^1 f_n^{n+1}(x) \cos(xt) dx = 2 \sum_{k=0}^{\infty} (-1)^k \frac{t^{2k}}{(2k)!} \int_0^1 x^{2k} f_n^{n+1}(x) dx.$$

Now use $f_n^{n+1}(x) = d_{n+1,0}q_n(x)$ and the Mellin transform (11) for q_n to find

$$\hat{f}_n^{n+1}(t) = (-1)^n 2d_{n+1,0} \sum_{k=0}^{\infty} (-1)^k \frac{t^{2k}}{(2k)!} \frac{(-k)_n}{(2k+1)_{n+1}}.$$

Again the first n terms in the series are zero, so the change $k = j + n$ gives

$$\hat{f}_n^{n+1}(t) = 2d_{n+1,0}t^{2n} \sum_{j=0}^{\infty} (-1)^j \frac{t^{2j}}{(2j+2n)!} \frac{(-j-n)_n}{(2j+2n+1)_{n+1}}.$$

In this case we need to use

$$(2j+2n)!(2j+2n+1)_{n+1} = (2j+3n+1)! = (3n+1)! 2^{2j} \left(\frac{3n+2}{2}\right)_j \left(\frac{3n+3}{2}\right)_j,$$

and some elementary work gives the required hypergeometric function. $\square$

The Fourier transform of the other multiwavelets f_k^{n+1} can be obtained by taking the sin-transform in (21) or the cos-transform in (22). Both will give a determinant containing hypergeometric functions.

6. Alpert multiwavelets and Legendre polynomials. In order to compute the multiwavelets, we introduce the notation

$$\Psi_n(x) = \begin{pmatrix} f_1^n(2x-1) \\ f_2^n(2x-1) \\ \vdots \\ f_n^n(2x-1) \end{pmatrix},$$

and the idea is to write

$$\Psi_n\left(\frac{x+1}{2}\right) = D_{-1}^n \Phi_n(x+1) + D_1^n \Phi_n(x)$$

where

$$\Phi_n(x) = \begin{pmatrix} \ell_0(x) \\ \ell_1(x) \\ \vdots \\ \ell_{n-1}(x) \end{pmatrix} \chi_{[0,1]},$$

and $\ell_n(x)$ are the orthonormal Legendre polynomials on $[0, 1]$ with χ_A the indicator function for the set A . The scaling relation is

$$\Phi_n\left(\frac{t+1}{2}\right) = C_{-1}^n \Phi_n(t+1) + C_1^n \Phi_n(t),$$

where C_{-1}^n and C_1^n are $n \times n$ matrices (which are lower triangular). The orthogonality of the Legendre polynomials gives

$$\int_{-1}^1 \Phi_n\left(\frac{x+1}{2}\right) \Phi_n^T\left(\frac{x+1}{2}\right) dx = 2\mathbb{I}_n,$$

where $\mathbb{I}_n$ is the identity matrix of order n , so that

$$C_{-1}^n (C_{-1}^n)^T + C_1^n (C_1^n)^T = 2\mathbb{I}_n.$$

The orthogonality of Alpert's multiwavelets gives

$$\int_{-1}^1 \Psi_n\left(\frac{x+1}{2}\right) \Psi_n^T\left(\frac{x+1}{2}\right) dx = \mathbb{I}_n,$$

so that

$$D_{-1}^n (D_{-1}^n)^T + D_1^n (D_1^n)^T = \mathbb{I}_n.$$

Furthermore, the multiwavelets are orthogonal to polynomials of degree $\leq n$, hence

$$\int_{-1}^1 \Psi_n\left(\frac{x+1}{2}\right) \Phi_n^T\left(\frac{x+1}{2}\right) dx = \mathbb{O}_n,$$

so that

$$D_{-1}^n (C_{-1}^n)^T + D_1^n (C_1^n)^T = \mathbb{O}_n.$$

Let n be even, then we write

$$\Psi_n^e(x) = \begin{pmatrix} f_2(x) \\ f_4(x) \\ \vdots \\ f_n(x) \end{pmatrix}, \quad \Psi_n^o(x) = \begin{pmatrix} f_1(x) \\ f_3(x) \\ \vdots \\ f_{n-1}(x) \end{pmatrix}.$$

From (21) and (22) we find that

$$\Psi_n^e(x) = G_n^p \begin{pmatrix} p_{n/2}(x) \\ \vdots \\ p_{n-1}(x) \end{pmatrix}, \quad \Psi_n^o(x) = G_n^q \begin{pmatrix} q_{n/2}(x) \\ \vdots \\ q_{n-1}(x) \end{pmatrix},$$

where G_n^p and G_n^q are $\frac{n}{2} \times \frac{n}{2}$ upper triangular matrices. The orthogonality of the multiwavelets and the symmetry gives

$$\mathbb{I}_{n/2} = 2 \int_0^1 \Psi_n^e(x) \Psi_n^e(x)^T dx = 2G_n^p \int_0^1 \begin{pmatrix} p_{n/2}(x) \\ \vdots \\ p_{n-1}(x) \end{pmatrix} (p_{n/2}(x) \cdots p_{n-1}(x)) (G_n^p)^T,$$

so that

$$(G_n^p)^{-1}(G_n^{pT})^{-1} = 2 \left(\langle p_i, p_j \rangle \right)_{i,j=n/2}^{n-1},$$

and hence $(G_n^p)^{-1}(G_n^{pT})^{-1}$ is the Cholesky factorization of the matrix $2 \left(\langle p_i, p_j \rangle \right)_{i,j=n/2}^{n-1}$. Note however that this is an UL-factorization rather than an LU-factorization. In a similar way, $(G_n^q)^{-1}(G_n^{qT})^{-1}$ is the UL-Cholesky factorization of the positive definite matrix $2 \left(\langle q_i, q_j \rangle \right)_{i,j=n/2}^{n-1}$. Next, we expand the p_k and q_k functions in terms of Legendre polynomials as

$$\begin{pmatrix} p_0(x) \\ p_1(x) \\ \vdots \\ p_{n-1}(x) \end{pmatrix} = A_n \Phi_n(x), \quad \begin{pmatrix} q_0(x) \\ q_1(x) \\ \vdots \\ q_{n-1}(x) \end{pmatrix} = B_n \Phi_n(x),$$

where A_n and B_n are lower triangular $n \times n$ matrices. The orthonormality of the Legendre polynomials gives

$$\int_0^1 \begin{pmatrix} p_0(x) \\ p_1(x) \\ \vdots \\ p_{n-1}(x) \end{pmatrix} \begin{pmatrix} p_0(x) & p_1(x) & \cdots & p_{n-1}(x) \end{pmatrix} dx = A_n A_n^T,$$

so that $A_n A_n^T$ is the Cholesky factorization of the matrix $\left(\langle p_i, p_j \rangle \right)_{i,j=0}^{n-1}$ and similarly $B_n B_n^T$ is the Cholesky factorization of $\left(\langle q_i, q_j \rangle \right)_{i,j=0}^{n-1}$. Note that the positivity of the diagonal elements of the Cholesky factorization reflects that the leading coefficients of p_k and q_k are positive. Combining this gives

$$\Psi_n^e(x) = \left(\mathbb{O}_{\frac{n}{2}, \frac{n}{2}} \ G_n^p \right) A_n \Phi_n(x), \quad \Psi_n^o(x) = \left(\mathbb{O}_{\frac{n}{2}, \frac{n}{2}} \ G_n^q \right) B_n \Phi_n(x),$$

where $\mathbb{O}_{n,m}$ is a $n \times m$ matrix containing only zeros, and D_1^n is obtained by intertwining Ψ_n^e and Ψ_n^o , i.e.,

$$D_1^n = \left(\mathbb{O}_{n, \frac{n}{2}} \ \hat{G}_n^p \right) A_n + \left(\mathbb{O}_{n, \frac{n}{2}} \ \hat{G}_n^q \right) B_n,$$

where $\hat{G}_n^p$ is a stretched version of G_n^p where zero rows have been inserted between every row, starting from the first row, and $\hat{G}_n^q$ is obtained from G_n^q by inserting zero rows starting from the second row:

$$\hat{G}_n^p = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 \\ 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix} G_n^p, \quad \hat{G}_n^q = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ \vdots & & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix} G_n^q.$$

For n odd we have to modify the method only slightly. We now use

$$\Psi_n^e(x) = \begin{pmatrix} f_2(x) \\ f_4(x) \\ \vdots \\ f_{n-1}(x) \end{pmatrix}, \quad \Psi_n^o(x) = \begin{pmatrix} f_1(x) \\ f_3(x) \\ \vdots \\ f_n(x) \end{pmatrix},$$

so that Ψ_n^e has $(n-1)/2$ elements and Ψ_n^o has $(n+1)/2$ elements. From (21) and (22) we now find

$$\Psi_n^e(x) = G_n^q \begin{pmatrix} q_{(n+1)/2}(x) \\ \vdots \\ q_{n-1}(x) \end{pmatrix}, \quad \Psi_n^o(x) = G_n^p \begin{pmatrix} p_{(n-1)/2}(x) \\ \vdots \\ p_{n-1}(x) \end{pmatrix},$$

where G_n^p is a $\frac{n+1}{2} \times \frac{n+1}{2}$ and G_n^q a $\frac{n-1}{2} \times \frac{n-1}{2}$ upper triangular matrix. As before, $(G_n^p)^{-1}(G_n^{pT})^{-1}$ is the UL-Cholesky factorization of the matrix $2\left(\langle p_i, p_j \rangle\right)_{i,j=(n-1)/2}^{n-1}$ and $(G_n^q)^{-1}(G_n^{qT})^{-1}$ is the UL-Cholesky factorization of the positive definite matrix $2\left(\langle q_i, q_j \rangle\right)_{i,j=(n+1)/2}^{n-1}$. The lower triangular matrices A_n and B_n are the same as before. The final result is that

$$D_1^n = \left(\mathbb{O}_{n, \frac{n-1}{2}} \hat{G}_n^p\right) A_n + \left(\mathbb{O}_{n, \frac{n+1}{2}} \hat{G}_n^q\right) B_n,$$

where the stretched matrices are now given by

$$\hat{G}_n^p = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & \\ \vdots & & & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix} G_n^p, \quad \hat{G}_n^q = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 \\ 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & & & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix} G_n^q.$$

Appendix A. An alternative proof of Proposition 8. The Mellin transforms (10)–(11) will be useful, if we combine this with Parseval's formula for the Mellin transform

$$(29) \quad \int_0^\infty f(x)g(x) dx = \frac{1}{2\pi} \int_{-\infty}^\infty \hat{f}\left(-\frac{1}{2} + it\right) \overline{\hat{g}\left(-\frac{1}{2} + it\right)} dt,$$

where

$$\hat{f}(s) = \int_0^\infty f(x)x^s dx, \quad \hat{g}(s) = \int_0^\infty g(x)x^s dx.$$

By using Parseval's formula and (10) we have

$$\int_0^1 p_n(x)p_k(x) dx = (-1)^{n+k} \frac{1}{2\pi} \int_{-\infty}^\infty \frac{\Gamma(\frac{3}{4} + n - \frac{it}{2})\Gamma(\frac{3}{4} + k + \frac{it}{2})|\Gamma(\frac{1}{2} + it)|^2}{\Gamma(\frac{3}{2} + n + it)\Gamma(\frac{3}{2} + k - it)|\Gamma(\frac{3}{4} + \frac{it}{2})|^2} dt.$$

The arguments $it/2$ are unusual, so to get rid of them we change the variable $t = 2s$ to find

$$\langle p_n, p_k \rangle = (-1)^{n+k} \frac{1}{\pi} \int_{-\infty}^\infty \frac{\Gamma(\frac{3}{4} + n - is)\Gamma(\frac{3}{4} + k + is)|\Gamma(\frac{1}{2} + 2is)|^2}{\Gamma(\frac{3}{2} + n + 2is)\Gamma(\frac{3}{2} + k - 2is)|\Gamma(\frac{3}{4} + is)|^2} ds.$$

Now we use Legendre's duplication formula for the Gamma function

$$\Gamma(2z) = \frac{2^{2z-1}}{\sqrt{\pi}} \Gamma(z) \Gamma(z + \frac{1}{2})$$

to find

$$\begin{aligned} \langle p_n, p_k \rangle &= (-1)^{n+k} 2^{-(n+k+1)} \\ &\times \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\Gamma(\frac{3}{4} + n - is) \Gamma(\frac{3}{4} + k + is) |\Gamma(\frac{1}{4} + is)|^2}{\Gamma(\frac{n}{2} + \frac{3}{4} + is) \Gamma(\frac{n}{2} + \frac{5}{4} + is) \Gamma(\frac{k}{2} + \frac{3}{4} - is) \Gamma(\frac{k}{2} + \frac{5}{4} - is)} ds. \end{aligned}$$

If we now change the variable is to y , then we get a Mellin-Barnes integral defining a Meijer G-function

$$(30) \quad \int_0^1 p_n(x) p_k(x) dx = (-1)^{n+k} 2^{-(n+k+1)} G_{4,4}^{2,2} \left(1; \begin{matrix} -k + \frac{1}{4}, \frac{3}{4}, \frac{k}{2} + \frac{3}{4}, \frac{k}{2} + \frac{5}{4} \\ n + \frac{3}{4}, \frac{1}{4}, -\frac{n}{2} + \frac{1}{4}, -\frac{n}{2} - \frac{1}{4} \end{matrix} \right),$$

where the Meijer G-function is defined as

$$G_{p,q}^{m,n} \left(z; \begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix} \right) = \frac{1}{2\pi i} \int_{\Gamma} \frac{\prod_{j=1}^m \Gamma(b_j - s) \prod_{j=1}^n \Gamma(1 - a_j + s)}{\prod_{j=m+1}^q \Gamma(1 - b_j + s) \prod_{j=n+1}^p \Gamma(a_j - s)} z^s ds,$$

where the path Γ separates the poles of $\Gamma(b_j - s)$ from the poles of $\Gamma(1 - a_j + s)$ [21, 22, §16.17]. In our case we have $m = n = 2$, $p = q = 4$ and

$$\begin{aligned} b_1 &= n + \frac{3}{4}, \quad b_2 = \frac{1}{4}, \quad b_3 = -\frac{n}{2} + \frac{1}{4}, \quad b_4 = -\frac{n}{2} - \frac{1}{4}, \\ a_1 &= -k + \frac{1}{4}, \quad a_2 = \frac{3}{4}, \quad a_3 = \frac{k}{2} + \frac{3}{4}, \quad a_4 = \frac{k}{2} + \frac{5}{4}. \end{aligned}$$

This Meijer G-function can be expressed as a linear combination of two hypergeometric functions

$$\begin{aligned} G_{4,4}^{2,2} \left(z; \begin{matrix} a_1, a_2, a_3, a_4 \\ b_1, b_2, b_3, b_4 \end{matrix} \right) &= \frac{z^{b_1} \Gamma(b_2 - b_1) \Gamma(1 + b_1 - a_1) \Gamma(1 + b_1 - a_2)}{\Gamma(1 + b_1 - b_3) \Gamma(1 + b_1 - b_4) \Gamma(a_3 - b_1) \Gamma(a_4 - b_1)} \\ &\times {}_4F_3 \left(\begin{matrix} 1 + b_1 - a_1, 1 + b_1 - a_2, 1 + b_1 - a_3, 1 + b_1 - a_4 \\ 1 + b_1 - b_2, 1 + b_1 - b_3, 1 + b_1 - b_4 \end{matrix} ; z \right) \\ &+ \frac{z^{b_2} \Gamma(b_1 - b_2) \Gamma(1 + b_2 - a_1) \Gamma(1 + b_2 - a_2)}{\Gamma(1 + b_2 - b_3) \Gamma(1 + b_2 - b_4) \Gamma(a_3 - b_2) \Gamma(a_4 - b_2)} \\ &\times {}_4F_3 \left(\begin{matrix} 1 + b_2 - a_1, 1 + b_2 - a_2, 1 + b_2 - a_3, 1 + b_2 - a_4 \\ 1 + b_2 - b_1, 1 + b_2 - b_3, 1 + b_2 - b_4 \end{matrix} ; z \right). \end{aligned}$$

We need the case $z = 1$. Note that the ${}_4F_3$ hypergeometric functions are balanced [21, 22, §16.4 (i)] : for the first ${}_4F_3$ the sum of the parameters in the numerator is $4n + 4$ and the sum of the denominator parameters is $4n + 5$, for the second ${}_4F_3$ the numerator parameters add up to 2 and the denominator parameters to 3. Observe that $a_3 - b_1 = \frac{k}{2} - n$ and $a_4 - b_1 = \frac{k+1}{2} - n$, hence one of these two is a negative integer, and the Gamma function $\Gamma(a_3 - b_1)$ or $\Gamma(a_4 - b_1)$ has a pole, so that the first term vanishes. This means that (27) follows.

In a similar way one can also obtain the following expression for the integrals involving the q_n polynomials

$$(31) \quad \int_0^1 q_n(x) q_k(x) dx = (-1)^{n+k} 2^{-(n+k+1)} G_{4,4}^{2,2} \left(1; \begin{matrix} -k + \frac{3}{4}, \frac{1}{4}, \frac{k}{2} + \frac{3}{4}, \frac{k}{2} + \frac{5}{4} \\ n + \frac{1}{4}, \frac{3}{4}, -\frac{n}{2} + \frac{1}{4}, -\frac{n}{2} - \frac{1}{4} \end{matrix} \right).$$

which simplifies to (28).

Acknowledgments. The research of the first author was partially supported by Simons Foundation Grant 210169. The research of the second author was partially supported by Simons Foundation Grant 280940. The third author was supported by FWO research projects G.0934.13 and G.0864.16 and KU Leuven research grant OT/12/073. This work was done while the third author was visiting Georgia Institute of Technology and he would like to thank FWO-Flanders for the financial support of his sabbatical and the School of Mathematics at GaTech for their hospitality.

REFERENCES

- [1] B. K. ALPERT, *A class of bases in L^2 for the sparse representation of integral operators*, SIAM J. Math. Anal., 24 (1993), pp. 246–262, doi:10.1137/0524016.
- [2] P. APPELL, *Sur une suite de polynômes ayant toutes leurs racines réelles*, Arch. Math. Phys. (3), 1 (1901), pp. 69–71.
- [3] A. APTEKAREV, *Multiple orthogonal polynomials*, J. Comput. Appl. Math., 99 (1998), pp. 423–447, doi:10.1016/S0377-0427(98)00175-7.
- [4] I. DAUBECHIES, *Ten Lectures on Wavelets*, vol. 61 of CBMS-NSF Regional Conference Series in Applied Mathematics, SIAM, Philadelphia, PA, 1992, doi:10.1137/1.9781611970104.
- [5] G. C. DONOVAN, J. S. GERONIMO, AND D. P. HARDIN, *Intertwining multiresolution analyses and the construction of piecewise-polynomial wavelets*, SIAM J. Math. Anal., 27 (1996), pp. 1791–1815, doi:10.1137/S0036141094276160.
- [6] G. C. DONOVAN, J. S. GERONIMO, D. P. HARDIN, AND P. R. MASSOPUST, *Construction of orthogonal wavelets using fractal interpolation functions*, SIAM J. Math. Anal., 27 (1996), pp. 1158–1192, doi:10.1137/S0036141093256526.
- [7] J. S. GERONIMO, D. P. HARDIN, AND P. R. MASSOPUST, *Fractal functions and wavelet expansions based on several scaling functions*, J. Approx. Theory, 78 (1994), pp. 373–401, doi:10.1006/jath.1994.1085.
- [8] J. S. GERONIMO AND P. ILIEV, *A hypergeometric basis for the Alpert multiresolution analysis*, SIAM J. Math. Anal., 47 (2015), pp. 654–668, doi:10.1137/140963923.
- [9] J. S. GERONIMO AND F. MARCELLÁN, *On Alpert multiwavelets*, Proc. Amer. Math. Soc., 143 (2015), pp. 2479–2494, doi:10.1090/S0002-9939-2015-12493-8.
- [10] T. GOODMAN AND S. LEE, *Wavelets of multiplicity r* , Trans. Amer. Math. Soc., 342 (1994), pp. 307–324, doi:10.2307/2154695.
- [11] T. GOODMAN, S. LEE, AND W. TANG, *Wavelets in wandering subspaces*, Trans. Amer. Math. Soc., 338 (1993), pp. 639–654, doi:10.1090/S0002-9947-1993-1117215-0.
- [12] D. P. HARDIN, B. KESSLER, AND P. R. MASSOPUST, *Multiresolution analysis and fractal functions*, J. Approx. Theory, 71 (1992), pp. 104–120, doi:10.1016/0021-9045(92)90134-A.
- [13] L. HERVÉ, *Multi-resolution analysis of multiplicity d : applications to dyadic interpolation*, Appl. Comput. Harm. Anal., 1 (1994), pp. 299–315, doi:10.1006/acha.1994.1017.
- [14] M. E. ISMAIL, *Classical and Quantum Orthogonal Polynomials in One Variable*, vol. 98 of Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2009 (paperback).
- [15] V. KALIAGUINE AND A. RONVEAUX, *On a system on “classical” polynomials of simultaneous orthogonality*, J. Comput. Appl. Math., 67 (1996), pp. 207–217, doi:10.1016/0377-0427(94)00129-4.
- [16] V. KALYAGIN, *On a class of polynomials defined by two orthogonality relations*, Mat. Sbornik, 110(152) (1979), pp. 609–627, doi:10.1070/SM1981v038n04ABEH001465. Translated in Math. USSR Sbornik 38 (1981), nr. 4, 563–580.
- [17] F. KEINERT, *Wavelets and Multiwavelets*, Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [18] F. KEINERT, *Multiwavelets*, in Encyclopedia of Complexity and Applied Systems Science, R. A. Meyers, ed., Springer-Verlag, New York, 2009, pp. 5841–5858.
- [19] S. G. MALLAT, *Multiresolution approximations and wavelet orthonormal bases of $L^2(\mathbb{R})$* , Trans. Amer. Math. Soc., 315 (1989), pp. 69–87, doi:10.2307/2001373.
- [20] Y. MEYER, *Ondelettes et Opérateurs, I*, Actualités Mathématiques, Hermann, Paris, 1990.
- [21] *NIST Digital Library of Mathematical Functions*, <http://dlmf.nist.gov>. Release 1.0.10 of 2015-08-07, online companion to [22].
- [22] F. W. OLVER, D. W. LOZIER, R. F. BOISVERT, AND C. W. CLARK, *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, NY, 2010.

- [23] C. SMET AND W. VAN ASSCHE, *Mellin transforms for multiple Jacobi-Piñeiro polynomials and a q -analogue*, J. Approx. Theory, 162 (2010), pp. 782–806, doi:[10.1016/j.jat.2009.09.004](https://doi.org/10.1016/j.jat.2009.09.004).