Citation	Jenniskens M, Langouche L, (2017), Cholestatic alterations in the critically ill: some new light on an old problem Chest. 2017 Aug 25 (epub ahead of print)
Archived version	Author manuscript: the content is identical to the content of the published paper, but without the final typesetting by the publisher
Published version	http://dx.doi.org/ 10.1016/j.chest.2017.08.018
Journal homepage	http://www.chestnet.org/Publications
Author contact	greet.vandenbergh@med0kuleuven.be + 32 (0)16 344021
IR	https://lirias.kuleuven.be/handle/123456789/590821

(article begins on next page)

CHOLESTATIC ALTERATIONS IN THE CRITICALLY ILL: SOME NEW LIGHT ON AN OLD PROBLEM

Running title

CHOLESTATIC ALTERATIONS IN THE CRITICALLY ILL

Jenniskens M¹, MD; Langouche L¹, MSc, PhD; Van den Berghe G¹, MD, PhD

¹ Clinical Division and Laboratory of Intensive Care Medicine, Department of Cellular and Molecular Medicine, KU Leuven, Belgium.

Corresponding author: Greet Van den Berghe, Clinical Division and Laboratory of Intensive Care Medicine, KU Leuven, Herestraat 49, B-3000 Leuven, Belgium. E-mail: greet.vandenberghe@med.kuleuven.be.

Source funding: Methusalem Program of the Flemish Government to GVdB and LL via the KU Leuven University (METH/08/07); ERC Advanced grant (AdvG-2012-321670) to GVdB from the Ideas Program of the European Union 7th framework program; Research Foundation Flanders Grant to GVdB and LL (G.0C78.17N.)

Disclosure: The authors have no conflicts of interest to disclose.

Key words: critical illness, sepsis, cholestasis, BA, bilirubin, liver

Abbreviations

ALP	Alkaline phosphatase
ALT	Alanine aminotransferase
AST	Aspartate aminotransferase
BA	Bile acids
FXR	Farnesoid-X-Receptor
γ GT	Gamma-glutamyltranspeptidase
GLDH	Glutamate dehydrogenase
ICU	Intensive Care Unit
RXR	Retinoid-X-Receptor

Abstract

Hepatic dysfunction and jaundice are traditionally viewed as late features of sepsis and other critical illnesses, and are associated with a complicated ICU stay. However, recent studies suggest that cholestatic alterations already occur early in the course of critical illnesses, perceived only as minor abnormalities in routinely used biochemical liver tests. Inflammation-induced alterations in the transport of bile acids appears to drive bile acids and bilirubin towards the systemic circulation. Ongoing bile acid synthesis with an, at least partial, loss of feedback inhibition further contributes to elevated circulating bile acids and bilirubin. To what extent these changes reflect a biochemical epiphenomenon, true illness-induced hepatic dysfunction or a beneficial and adaptive response to illness should be further investigated.

Because of the lack of specificity of standard laboratory tests, especially in the context of a complex systemic condition as critical illness, identifying true cholestatic liver dysfunction remains a great challenge. However, very high levels of cholestatic markers that are sustained in prolonged critically ill patients almost always indicate a complicated illness course and should be monitored closely. Prevention of cholestatic liver dysfunction comprises minimizing inflammation and hypoxia in the liver and preventing hyperglycemia, avoiding early use of parenteral nutrition and reducing the administration of avoidable drugs.

Future research on the effects of bile acids and on modulating underlying drivers of critical illness-induced cholestasis is warranted as this could open perspectives for a targeted diagnostic approach and ultimately for novel therapies to improve outcome.

Introduction

During sepsis and other critical illnesses, the liver is a fundamental player in the host defense mechanisms as it regulates a wide range of metabolic, endocrine and immunological processes^{1,2}. However, inflammation and hypoxemia severely challenge the hepatic microcirculation and function. Abnormal liver tests, such as hyperbilirubinemia, elevated gamma glutamyl transpeptidase (γ -GT), alkaline phosphatase (ALP), alanine transaminase (ALT) and aspartate transaminase (AST) are reported in up to 61% of patients admitted to the ICU³⁻⁵. In fact, sepsis is reported as the second leading cause of jaundice in patients presenting at the emergency department⁶. Strongly elevated values, several fold the upper limit of normality, are associated with distinct hepatic pathology such as hepatic ischemia or (iatrogenic) pharmacological effects and indicative of poor outcome⁷. However, the large majority of deranged liver tests are only slightly outside the normal range. Nonetheless, also low-grade hyperbilirubinemia and mild rises in plasma concentrations of liver enzymes are associated with a more complicated ICU course, a longer duration of ICU stay and a higher risk of death^{3,4,8-10}. While obstructive cholestasis due to gallstones or by inflammation-driven narrowing of the bile ducts could induce these biochemical liver test abnormalities, it is rarely the cause of hyperbilirubinemia in critical illness¹¹⁻¹⁴. Indeed, the precise etiology of mild alterations in liver tests is not completely understood and it remains unclarified whether they clinically reflect true hepatic dysfunction or are part of an adaptive stress response and reflect severity of illness.

In this review, we summarize the state of the art knowledge, integrating some novel insights, about mild and severe cholestatic alterations in intensive care patients, discuss the epidemiology along with the relationship with outcome, the clinical diagnostic approach, the underlying (patho)physiological pathways and possible therapeutic implications.

Definitions, incidence and prognostic value of liver test alterations in critical illness

Liver test abnormalities are often found in the critically ill patient, the majority of which can be classified in two major pathological entities: “hypoxic hepatitis” and “critical illness-induced cholestasis” (Figure 1)¹⁵. Hypoxic hepatitis, or shock liver, presents as an early and severe rise in hepatic lysis enzymes ALT and AST, which reflects diffuse hepatic injury as a consequence of failed oxygen delivery and ischemia of the liver, due to any cause of severe shock. Mortality of patients diagnosed with hypoxic hepatitis is

directly dependent to the severity of the underlying condition, and overall high mortality rates are documented^{16,17}. The typical time course shows a rapid, steep increase in plasma ALT or AST concentrations reaching at least 20x the upper limit of normality^{15,17}. Hypoxic hepatitis presents in up to 10% of ICU patients, of which around at least 75% fulfill these diagnostic criteria already upon admission^{16,17}. Importantly, one-third of patients suffering from hypoxic hepatitis will also develop jaundice¹⁸.

More frequently observed in the ICU department are mild liver test abnormalities with predominantly cholestatic features, which is currently labeled “critical illness-induced cholestasis” (Figure 1). Cholestasis is clinically defined as decreased or absent bile-flow towards the duodenum, caused by either impaired bile formation or an inability to excrete bile through the biliary system. This manifestation is traditionally considered a late event in sepsis and multi-organ failure, although recent findings indicate that such signs can also be observed in the more early stages of severe illness^{3,8,19}. Whereas hypoxic hepatitis evidently relates to patients with severe shock and liver hypoxia, critical illness-induced cholestasis appears to be hallmarked by inflammation-driven cellular alterations resulting in a transient accumulation of bile acids (BA) and bilirubin within the liver and in the systemic circulation. In daily clinical practice, the most often used diagnostic criterion for critical illness-induced cholestasis is a total plasma bilirubin concentration higher than 2 or 3 mg/dl (34-41mmol/L) often combined with a maximally two or threefold increase in ALP and γ -GT levels. Accordingly, up to 20% of ICU patients fulfill these diagnostic criteria of critical illness-induced cholestasis at admission^{3,8,19}. However, cholestatic alterations may be more prevalent, considering that most patients display milder derangements of cholestatic markers that do not exceed these cutoff levels²⁰. The occurrence of critical illness-induced cholestasis has been related to the use of parenteral nutrition, antibiotics or other medications and to mechanical ventilation with high levels of PEEP^{8,21,22}. Other typical risk factors for onset of critical illness-induced cholestasis are sepsis with gram-negative bacteria, major surgery, severe trauma or other severe shock states^{14,15}. In addition, pre-existing (subclinical) presence of alcoholic, viral or toxin-induced hepatitis might further accentuate liver test abnormalities during critical illness²³.

Diagnostic tools for critical illness-induced liver dysfunction and cholestasis

The liver plays a pivotal role in key processes of metabolism, synthesis of acute phase proteins and coagulation components and excretion of toxic compounds. However, no single biomarker currently reflects all these functional activities (Table 1). In current daily practice, bilirubin is the most widely used biomarker in the assessment of liver function and has been incorporated in several organ dysfunction scores^{24,25}. In addition, elevated circulating levels of ALT, AST, ALP and γ -GT can further indicate liver injury and cholestasis^{5,8}. However, because of the lack of specificity of these standard laboratory tests, identifying true (cholestatic) liver dysfunction remains a great challenge⁸.

Bilirubin is the routinely used marker in the assessment of excretory dysfunction. However, the reliability of hyperbilirubinemia as a marker of cholestasis in critically ill patients may be questioned, since multiple ICU-associated factors can influence levels of circulating bilirubin, such as changes in red blood cell turnover and blood transfusions²⁶. Furthermore, bile excretion is a secretory process that depends largely on the osmotic drive induced by BA^{27,28}. If bile-flow is hampered as a consequence of retained BA, bilirubin will also accumulate, but essentially reflects a biochemical epiphenomenon. Emerging evidence indicates that elevated BA levels reflect critical illness-induced cholestasis with a higher sensitivity and specificity, independently from bilirubin²⁹. Furthermore, BA could potentially be useful as disease severity marker. Indeed, in a cohort of critically ill patients, total bile acids predicted 28-day mortality independently of severity of illness, gender, age and serum bilirubin concentration³⁰. Moreover, molecular alterations in intrahepatic BA regulation appear to precede biochemical laboratory abnormalities³¹. However, BA isoforms including the conjugated fractions are difficult to measure and not readily available in clinical practice.

Increased γ -GT and ALP are very sensitive indicators for the diagnosis of liver and bile duct diseases³². Their synthesis and systemic release are observed during cholestasis with increased intra-biliary BA³³. Increased levels of γ -GT and ALP are observed in 20-30% of ICU patients at admission and decrease thereafter^{8,34}. Surprisingly, elevated levels of γ -GT and ALP are not associated with complicated clinical events, such as mechanical ventilation, hypoxia, higher APACHE II scores, or antibiotic use⁸. Furthermore, both markers have a high sensitivity but low specificity, making them weak indicators when used for the individual assessment of hepatic dysfunction in a single patient³².

The release of ALT and AST in the circulation is largely dependent on cellular necrosis and evidently related to severe hypoxia (Figure 1). However, levels may be mildly elevated during critical illness, as

inflammatory signals increase cellular permeability and could promote enzyme leakage. The enzyme glutamate dehydrogenase (GLDH) is predominantly localized in the centrolobular region, which is more susceptible to hypoxia, and may potentially point to acute liver injury with more accuracy³⁵. However, the use of GLDH in the context of critical illness has not been systematically investigated.

Next to static laboratory measurements, dynamic tests such as indocyanine green can be used to assess excretory function of the liver. Indocyanine green is an organic anion that is exclusively eliminated by the liver. In a recent study, dynamic measurement of hepatic biotransformation and excretion was investigated in critically ill animals using the clearance rate of indocyanine green²⁹. In these critically ill mice, dye was rapidly cleared from the circulation and taken up by liver cells, but the appearance in bile was delayed, indicating that defective cellular function primarily involves excretion of transformed molecules from the hepatocyte into the biliary canaliculi rather than the uptake in liver cells²⁹. Furthermore, in a retrospective analysis from 336 critically ill patients, plasma disappearance rate of indocyanine green correlated with mortality with comparable sensitivity and specificity as the SAPS and APACHE II scores³⁶. Moreover, in patients with hypoxic hepatitis, dynamics of plasma disappearance rate of indocyanine green over time demonstrated that plasma clearance did not increase over time in patients who died within 28-days on the ICU, whereas a clear improvement was observed in patients who survived³⁷. Noteworthy, plasma disappearance rate of indocyanine green is greatly depending on liver perfusion and impaired clearance rates may reflect both liver hypoperfusion and reduced excretory function³⁸. Furthermore, redistribution in extravascular tissues, protein binding and complex pharmacokinetics further complicate the interpretation of the plasma disappearance rate of indocyanine green³⁹. Hence, indocyanine green should be used with caution in the complex clinical context of critical illness³⁹.

Outside the context of critical illness, hepatic synthesis capacity can be estimated by quantifying serum concentrations of proteins that are exclusively produced by the liver, such as albumin. Albumin is the most abundant circulating protein and accounts for up to 50% of hepatic protein synthesis. However, serum protein concentrations do not directly reflect protein synthesis, as breakdown and distribution to the extravascular compartment also play a role. During critical illness, low albumin can reflect blood losses, hemodilution, and extravascular albumin leakage, rather than hepatic synthesis dysfunction^{40,41}. Also, availability of coagulation factors, indirectly estimated via measuring standardized prothrombin time (INR),

is used to assess the synthetic capacity of the liver. The INR is used to calculate the Model for End stage Liver Disease-score (MELD), in turn used to prioritize end stage liver disease patients for liver transplant²⁵. However, in the context of critical illness, other factors, such as bleeding, hemodilution, disseminated intravascular coagulation and the administration of coagulation factors or fresh frozen plasma, can alter the INR⁴¹. It is clear that neither albumin nor INR are preferred markers to test synthetic function of the liver in critically ill patients. However, observational data suggest that the hepatocellular synthesis performance could be estimated with use of serum cholinesterase^{42,43}. Indeed, serum cholinesterase activity is lower than normal in patients with burns and septic shock⁴⁴ and correlates with the severity of the injury⁴². However, because normal values span over a wide range, clear diagnostic cutoffs for cholinesterase activity to diagnose liver dysfunction in critical illness are lacking. Furthermore, healthy volunteers showed both increased and decreased cholinesterase activity after an endotoxin challenge⁴⁵. Possibly, in the context of critical illness, cholinesterase activity reflects the systemic cholinergic status rather than the synthetic capacity of the liver^{42,45}.

Critically ill patients also often develop post-hepatic biliary abnormalities. Most frequently, biliary sludge is observed during critical illness, with a prevalence of approximately 50-60% in patients requiring more than 5 days of intensive care¹⁹. True mechanical biliary obstruction can be easily and reliably assessed with use of abdominal sonography. However, mechanical biliary obstruction is rarely the cause of deranged cholestatic liver parameters in the context of critical illness, which argues against a routine sonographic screening for calculous cholecystitis^{11,13}. However, decreased blood supply via the hepatic artery can impair the microcirculation of the gallbladder wall and lead to acalculous cholecystitis, inflammation and necrosis. This rare but severe condition is associated with a complicated ICU stay and can be lethal¹². The use of ultrasound to detect such acalculous cholecystitis is only of value in combination with a high level of clinical suspicion⁴⁶. Also the use of CT imaging for acalculous cholecystitis has previously shown to be too aspecific, as up to 96% of patients displayed gallbladder abnormalities⁴⁷. Importantly, persisting cholestatic serum abnormalities may indicate the development of secondary cholestatic pathologies, such as secondary sclerosing cholangitis, in turn associated with poor outcome^{48,49}. For patients who continue to reveal cholestatic serum abnormalities, also after clinical improvement, further clinical follow up with biliary imaging is advisable.

As to date, no specific liver test for critical illness-induced cholestasis is available, it remains a 'diagnosis of exclusion'. In absence of a recognized hypoxic or pharmacotoxic insult, obstruction or pre-existing liver disease, abnormal liver tests more likely indicate critical illness-induced cholestasis.

Underlying pathophysiological pathways of critical illness-induced cholestasis

Critical illness-induced cholestasis appears to be the result of functional intracellular alterations in BA homeostasis, with changes observed in intrahepatic BA synthesis, feedback-regulation, conjugation and BA transportation^{31,50}. BA are actively produced in the liver from its precursor cholesterol through an extensive enzymatic cascade⁵¹. Key proteins in BA production are CYP7A1 and CYP27A1, the rate limiting enzymes of the normal and alternative synthesis pathway respectively^{51,52}. BA can suppress their own synthesis through binding with its regulating nuclear receptors Farnesoid-X-Receptor (FXR) and the Retinoid-X-Receptor (RXR). Upon binding with BA, FXR-RXR dimers translocate to the hepatocyte nucleus and regulate DNA transcription of Small Heterodimer Partner (SHP), which inhibits CYP7A1 and CYP27A1⁵². In an observational study of non-surviving ICU patients, circulating BA levels were 11-fold increased as compared with elective surgery patients, but in the liver biopsies, protein expression of CYP7A1 was not suppressed. Furthermore, hepatocytic nuclear content of the key sensors and transcriptional regulators of BA synthesis and intrahepatic transport, FXR and RXR, was remarkably low, suggesting an at least partial loss of feedback-inhibition in high risk critically ill patients⁴.

Bilirubin and BA are normally conjugated within the hepatocyte to detoxify and convert these molecules to less toxic forms^{51,52}. While in a lethal animal model of sepsis, a BA pattern indicative of a conjugation defect was documented²⁹, data from non-surviving ICU patients suggested that conjugation was unaffected by critical illness⁴. In an observational study of 280 critically ill patients, conjugation of primary BA even increased over time in ICU²⁰

Hepato-biliary BA transporters are the final critical determinant of BA homeostasis and control intrahepatic and circulating levels of BA^{27,28}. In normal hepato-biliary transport, BA enter the hepatocyte from the portal circulation via Na⁺-Taurocholate Cotransporting Polypeptide (NCTP) at the basolateral membrane. In addition, several multispecific anion transporters (OATPs) facilitate transport of BA, bilirubin and other hormones and compounds. The canalicular membrane transporter bile salt export pump (BSEP)

regulates the excretion of BA into the bile duct lumen. The multispecific export transporters multidrug resistance-associated protein 2 (MRP2) and multidrug resistance proteins (MDR1, 2, 3) mediate excretion of BA, bilirubin and various xenobiotic compounds. Alternatively, BA can be excreted into the systemic circulation via multidrug resistance-associated protein 3 and 4 (MRP3, MRP4). In liver biopsies harvested from critically ill patients who died in the ICU, expression of hepatic BA transporters was altered in comparison with liver biopsies from patients with elective non liver related surgery, suggesting a diverted export of bilirubin and BA to the systemic circulation⁵³. Gene expression of hepatic basolateral uptake transporters NTCP and OATP's was suppressed, as well as apical exporter BSEP gene expression. Interestingly, export pumps MRP3 and MRP4 were strongly upregulated, which may serve as alternative escape routes from hepatocytes into the sinusoidal blood⁴. Inevitably, this postmortem study was restricted to the most severe phenotype of ICU patients. It is not clear whether such reversal in BA transport is a general phenomenon occurring in all types of critical illnesses. However, also acute animal models of sepsis have shown changes in the maximal transport velocity of BA and organic anions along with downregulation in gene and protein levels of BA transporters^{29,54}.

In vitro and acute models of sepsis suggest cytokines as possible drivers of cholestatic alterations in critical illness. Indeed, cell culture models showed similar nuclear export of RXR after administration of IL-1 β and TNF α ⁵⁵. In addition, in acute models of cytokine-induced sepsis, signs of diverted transport were observed^{56,57}. However, there was no association between markers of inflammation and markers of cholestasis in liver biopsies from ICU-nonsurvivors, suggesting that inflammation may not be the main contributor to cholestatic phenotype in prolonged critical illness⁴. In addition, disturbances in the microcirculation of the biliary system could play a part in the development of hyperbilirubinemia in critical illness. Ischemic cholangiopathy, secondary to severely impaired biliary blood supply, can contribute to bile stasis and the formation of biliary casts, and can promote the development of clinical jaundice^{58,59}.

Clinical implications of critical illness-induced cholestasis

The current novel insights from studies on bilirubin and BA synthesis and transport in the critically ill, and their regulation by nuclear receptors do not suffice to conclude whether they represent a "failure" of liver functions or instead reflect a beneficial compensatory response within the hepatocyte in response to

critical illness. Thus, whether the typical mild cholestatic alterations observed in most if not all ICU patients reflect an adaptive response or indicate impaired organ function is still unclear. The diverted hepato-biliary transport could indeed reflect an adaptive escape mechanism for the hepatocyte to reduce the high energy demanding transport of bile components against the concentration gradient into the biliary system. Alternatively, in order to increase favorable effects of systemic bilirubin and BA signaling, ongoing BA production and reversal of intrahepatic transport towards the circulation might also reflect an adaptive and beneficial process. Indeed, via a positive feedforward mechanism, even small amounts of bilirubin may exert known antioxidative and cytoprotective roles⁶⁰. Bilirubin has shown to be protective in various animal models, including sepsis⁶⁰⁻⁶³. In an animal model of LPS-induced sepsis, bilirubin treated rats showed attenuated hepatic organ damage and had improved survival, at least partially mediated by inhibited hepatic iNOS expression⁶³. In addition to bilirubin, metabolic and endocrine effects of BA could play a protective role in the hypermetabolic and inflammatory response of acute critical illness. BA are currently being investigated as promising new drug targets for metabolic diseases as they play a role in glucose, cholesterol, triglyceride and energy homeostasis⁶⁴⁻⁶⁷. Furthermore, animals with reduced BA signaling are more susceptible to liver injury in LPS-induced sepsis, and show higher cytokine mRNA levels, more pronounced inflammatory infiltrates and an increased rate of apoptotic cells⁶⁸. In addition, injection of (conjugated or unconjugated) BA have shown to increase circulating cortisol levels in rats by reducing the expression and activity of cortisol metabolizing enzymes⁶⁹. A positive correlation between the rise in BA and plasma cortisol has also been described in critically ill patients, suggesting that the reduced breakdown of cortisol, recently documented as a universal phenomenon in the critically ill patient, may be at least partially explained by increased circulating levels of BA in response to critical illness. BA could thereby contribute to elevated plasma cortisol concentrations and higher cortisol availability in those vital organs and tissues that express the cortisol-metabolizing enzymes, thus playing a key role in the inflammatory and metabolic stress response⁷⁰.

In contrast, since BA play a vital role in intestinal integrity and reduce bile-flow in the intestine, absence of flow may promote translocation of the gut microbiome and elevate levels of LPS in the portal circulation⁷¹⁻⁷³. This might result in sustained inflammatory signals as is often observed in prolonged critical illness^{71,72}. In addition, prolonged lack of BA in the intestine can cause malabsorption of lipids and vitamins.

In addition, beneficial effects of other cholestatic markers, such as ALP, have been suggested. ALP has shown to elicit favorable effects in animal experiments and in two small phase II clinical trials of critically ill patients suffering from acute kidney injury⁷⁴⁻⁷⁶. Administration of recombinant ALP improved markers of kidney function and inhibited the upregulation of renal inducible NO synthase, leading to subsequently reduced NO metabolite production⁷⁵. Furthermore, experimental data from animal studies suggest that ALP is able to detoxify LPS and may provide a new strategy to attenuate LPS-induced diseases^{77,78}. In addition, next to the detoxification of LPS, presence of ALP prevented bacterial invasion across the intestinal mucosal barrier in an ischemia/reperfusion mouse model⁷⁹.

Importantly, the clinical impact of an altered hepatic transport on compounds normally excreted by the hepatocyte, such as toxins, drugs and other metabolic substances remains uncertain. The exogenous administration of antibiotics and other drugs can accumulate due to decreased bile excretion. Furthermore, certain drugs could amplify the underlying cholestatic alterations as several drugs routinely used in the ICU are themselves associated with the occurrence of cholestasis^{29,80}.

Therapeutic implications

Because critical illness-induced cholestasis is primarily a result of functional alterations at the hepatocyte and bile duct level, it is usually reversible and there is no clear argument for attempting to treat this condition. Up until today, there is in fact no treatment available for critical illness-induced cholestasis, but the general management of the ICU patient focusses on avoiding additional inflammatory stimuli and hypoxia in the liver microenvironment. Prevention however appears possible to a certain extent. Indeed, optimizing supportive care and reducing risk factors that could provoke severe cholestasis, such as the excessive use of antibiotics and other drugs, preventing hyperglycemia and not using parenteral nutrition during the first week of critical illness have shown to prevent cholestatic features observed during critical illness^{19,81,82}.

Adequately optimizing hemodynamic perfusion is an essential step in preventing liver dysfunction^{83,84}. Importantly, experimental data suggest that some catecholamines can also exacerbate hepatic dysfunction observed during sepsis by altering the biotransformation of drugs⁸⁵. As hepatic biotransformation and biliary export are the primary mechanisms for clearance of most drugs and toxins,

routinely used drugs in intensive care medicine can accumulate in the critically ill patient^{80,86}. Compounds that need to be metabolized in the liver and those that need to be excreted via the bile are known to have hepatic adverse effects⁸⁷. Commonly used drugs used in daily critical care practice, such as the majority of antibiotics, anesthetics and sedatives therefore form an additional threat for the hepatocyte. Although insights in individualized therapy are increasing rapidly, there are currently no commonly used strategies to adjust doses for decreased biliary excretion in the critically ill.

Lowering blood glucose to healthy fasting ranges, thereby avoiding hyperglycemia, in adult critically ill patients also reduced the occurrence of cholestasis and biliary sludge, as revealed by 2 randomized controlled trials^{19,81,82}. Also the impact of early initiation of parenteral nutrition, as compared with accepting a large macronutrient deficit for one week in the ICU when only using enteral feeds, on the occurrence of cholestasis has been investigated. For hemodynamically stable patients with a functioning gastrointestinal tract, early enteral feeding is currently the recommended standard of care. Enteral nutrition promotes the intestinal reuptake of BA and recovery of the enterohepatic cycle⁸⁸. Furthermore, restoration of the enterohepatic cycle by enteral nutrition improves gut oxygenation and barrier function, and may reduce microbial translocation and inflammatory signaling in the liver^{89,90}. The use of enteral nutrition compared to parenteral nutrition has been associated with fewer infections and reduced occurrence of metabolic complications in patients with liver disease and after liver transplantation⁹¹. However, enteral feeding is often not well tolerated by ICU patients. A large randomized controlled trial assessed the impact of early versus late initiation of parenteral nutrition in adults and found an important effect on liver function markers⁹². Indeed, this study showed that not using early parenteral nutrition not only shortened ICU and hospital length of stay and reduced the number of new infections, it also improved laboratory markers of hepatobiliary damage (maximum levels of ALT, γ -GT and ALP)^{92,93} and lowered the incidence of biliary sludge²⁰. Surprisingly, levels of bilirubin were slightly but significantly higher in the group not receiving early parenteral nutrition²⁰. Similar results were observed in the pediatric equivalent of this study⁹³. Underlying mechanisms of the beneficial effects of delaying parenteral nutrition have been investigated in a prolonged critical illness rabbit model⁵³. In this study, withholding parenteral nutrition reduced markers of hepatocyte injury and further accentuated the reversed BA transport towards the systemic circulation, suggesting that fasting by itself could partially initiate some “cholestatic” alterations during critical illness⁵³.

Conclusions

Hepatic dysfunction and jaundice are traditionally viewed as late features of sepsis and as part of multi-organ failure. However, recent studies suggest that minor “cholestatic” alterations occur already early in the course of critical illness, but these would not be considered as a clinically relevant problem according to advised thresholds for normality of standard laboratory tests. Inflammation- and hypoxia-driven alterations in the synthesis and transport of BA and bilirubin explain these early and often mild alterations. Elevated values of routinely used laboratory tests appear to be brought about by altered BA and organic anion transport machineries at the basolateral and biliary pole that appear to drive bile components towards the systemic circulation. To what extent these changes reflect a biochemical epiphenomenon, true illness-induced hepatic dysfunction or a beneficial and adaptive response to illness should be further investigated. It is known that BA and bilirubin play important roles in various metabolic and inflammatory pathways and mild elevation of these compounds could attenuate the severe burden of critical illness. Possibly, in the context of a complex systemic condition as critical illness, routinely used liver tests may not necessarily represent similar processes than in patients with primary liver disease. However, very high levels of cholestatic markers that are sustained in prolonged critically ill patients almost always indicate a complicated illness course and should be monitored more closely. The fascinating possibility of beneficial effects of increased BA availability during critical illness opens perspectives for future research in a targeted diagnostic approach and, ultimately, in the development of novel therapies aimed to improve outcome.

Acknowledgements

Source funding: Methusalem Program of the Flemish Government to GVdB and LL via the KU Leuven University (METH/08/07); ERC Advanced grant (AdvG-2012-321670) to GVdB from the Ideas Program of the European Union 7th framework program; Research Foundation Flanders Grant to GVdB and LL (G.0C78.17N.)

Disclosure: The authors have no conflicts of interest to disclose.

References

1. Szabo G, Romics L, Frendl G. Liver in sepsis and systemic inflammatory response syndrome. *Clin Liver Dis.* 2002;6(4):1045-1066, x.
2. Dhainaut JF, Marin N, Mignon A, Vinsonneau C. Hepatic response to sepsis: interaction between coagulation and inflammatory processes. *Crit Care Med.* 2001;29(7 Suppl):S42-47.
3. Kramer L, Jordan B, Druml W, Bauer P, Metnitz PGH, Austrian Epidemiologic Study on Intensive Care ASG. Incidence and prognosis of early hepatic dysfunction in critically ill patients--a prospective multicenter study. *Crit Care Med.* 2007;35(4):1099-1104.
4. Vanwijngaerden YM, Wauters J, Langouche L, et al. Critical illness evokes elevated circulating bile acids related to altered hepatic transporter and nuclear receptor expression. *Hepatology.* 2011;54(5):1741-1752.
5. Kortgen A, Paxian M, Werth M, et al. Prospective assessment of hepatic function and mechanisms of dysfunction in the critically ill. *Shock.* 2009;32(4):358-365.
6. Whitehead MW, Hainsworth I, Kingham JG. The causes of obvious jaundice in South West Wales: perceptions versus reality. *Gut.* 2001;48(3):409-413.
7. Harbrecht BG, Zenati MS, Doyle HR, et al. Hepatic dysfunction increases length of stay and risk of death after injury. *J Trauma.* 2002;53(3):517-523.
8. Thomson SJ, Cowan ML, Johnston I, Musa S, Grounds M, Rahman TM. 'Liver function tests' on the intensive care unit: a prospective, observational study. *Intensive Care Med.* 2009;35(8):1406-1411.
9. Nessler N, Launey Y, Aninat C, et al. Liver Dysfunction Is Associated with Long-Term Mortality in Septic Shock. *Am J Respir Crit Care Med.* 2016;193(3):335-337.
10. Zhai R, Sheu CC, Su L, et al. Serum bilirubin levels on ICU admission are associated with ARDS development and mortality in sepsis. *Thorax.* 2009;64(9):784-790.
11. Murray FE, Stinchcombe SJ, Hawkey CJ. Development of biliary sludge in patients on intensive care unit: results of a prospective ultrasonographic study. *Gut.* 1992;33(8):1123-1125.
12. Kalliafas S, Ziegler DW, Flancbaum L, Choban PS. Acute acalculous cholecystitis: incidence, risk factors, diagnosis, and outcome. *Am Surg.* 1998;64(5):471-475.
13. Boland GW, Slater G, Lu DS, Eisenberg P, Lee MJ, Mueller PR. Prevalence and significance of gallbladder abnormalities seen on sonography in intensive care unit patients. *AJR Am J Roentgenol.* 2000;174(4):973-977.
14. Molenat F, Boussuges A, Valantin V, Sainty JM. Gallbladder abnormalities in medical ICU patients: an ultrasonographic study. *Intensive Care Med.* 1996;22(4):356-358.
15. Horvatits T, Trauner M, Fuhrmann V. Hypoxic liver injury and cholestasis in critically ill patients. *Curr Opin Crit Care.* 2013;19(2):128-132.
16. Fuhrmann V, Kneidinger N, Herkner H, et al. Hypoxic hepatitis: underlying conditions and risk factors for mortality in critically ill patients. *Intensive Care Med.* 2009;35(8):1397-1405.
17. Fuhrmann V, Kneidinger N, Herkner H, et al. Impact of hypoxic hepatitis on mortality in the intensive care unit. *Intensive Care Med.* 2011;37(8):1302-1310.
18. Jager B, Drolz A, Michl B, et al. Jaundice increases the rate of complications and one-year mortality in patients with hypoxic hepatitis. *Hepatology.* 2012;56(6):2297-2304.
19. Mesotten D, Wauters J, Van den Berghe G, Wouters PJ, Milants I, Wilmer A. The effect of strict blood glucose control on biliary sludge and cholestasis in critically ill patients. *J Clin Endocrinol Metab.* 2009;94(7):2345-2352.
20. Vanwijngaerden Y-M, Langouche L, Brunner R, et al. Withholding parenteral nutrition during critical illness increases plasma bilirubin but lowers the incidence of biliary sludge. *Hepatology.* 2013.

21. Grau T, Bonet A, Rubio M, et al. Liver dysfunction associated with artificial nutrition in critically ill patients. *Crit Care*. 2007;11(1):R10.
22. Brienza N, Dalfino L, Cinnella G, Diele C, Bruno F, Fiore T. Jaundice in critical illness: promoting factors of a concealed reality. *Intensive Care Med*. 2006;32(2):267-274.
23. Moreau R, Jalan R, Gines P, et al. Acute-on-chronic liver failure is a distinct syndrome that develops in patients with acute decompensation of cirrhosis. *Gastroenterology*. 2013;144(7):1426-1437, 1437 e1421-1429.
24. Vincent JL, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med*. 1996;22(7):707-710.
25. Kamath PS, Wiesner RH, Malinchoc M, et al. A model to predict survival in patients with end-stage liver disease. *Hepatology*. 2001;33(2):464-470.
26. Bauer M, Huse K, Settmacher U, Claus RA. The heme oxygenase – carbon monoxide system: regulation and role in stress response and organ failure. *Intensive care medicine*. 2008;34(4):640-648.
27. Trauner M, Boyer JL. Bile salt transporters: molecular characterization, function, and regulation. *Physiol Rev*. 2003;83(2):633-671.
28. Trauner M, Meier PJ, Boyer JL. Molecular pathogenesis of cholestasis. *N Engl J Med*. 1998;339(17):1217-1227.
29. Recknagel P, Gonnert FA, Westermann M, et al. Liver dysfunction and phosphatidylinositol-3-kinase signalling in early sepsis: experimental studies in rodent models of peritonitis. *PLoS Med*. 2012;9(11):e1001338.
30. Horvatits T, Drolz A, Rutter K, et al. Circulating bile acids predict outcome in critically ill patients. *Annals of intensive care*. 2017;7(1):48.
31. Jenniskens M, Langouche L, Vanwijngaerden Y-M, Mesotten D, Van den Berghe G. Cholestatic liver (dys)function during sepsis and other critical illnesses. *Intensive Care Med*. 2016;42(1):16-27.
32. Giannini EG, Testa R, Savarino V. Liver enzyme alteration: a guide for clinicians. *Cmaj*. 2005;172(3):367-379.
33. Accatino L, Figueroa C, Pizarro M, Solis N. Enhanced biliary excretion of canalicular membrane enzymes in estrogen-induced and obstructive cholestasis, and effects of different bile acids in the isolated perfused rat liver. *J Hepatol*. 1995;22(6):658-670.
34. Sanfilippo F, Veenith T, Santonocito C, Vrettou CS, Matta BF. Liver function test abnormalities after traumatic brain injury: is hepato-biliary ultrasound a sensitive diagnostic tool? *Br J Anaesth*. 2014;112(2):298-303.
35. Schomaker S, Warner R, Bock J, et al. Assessment of emerging biomarkers of liver injury in human subjects. *Toxicol Sci*. 2013;132(2):276-283.
36. Sakka SG, Reinhart K, Meier-Hellmann A. Prognostic value of the indocyanine green plasma disappearance rate in critically ill patients. *Chest*. 2002;122(5):1715-1720.
37. Horvatits T, Kneidinger N, Drolz A, et al. Prognostic impact of ICG-PDR in patients with hypoxic hepatitis. *Annals of intensive care*. 2015;5.
38. Uusaro A, Ruokonen E, Takala J. Estimation of splanchnic blood flow by the Fick principle in man and problems in the use of indocyanine green. *Cardiovascular research*. 1995;30(1):106-112.
39. Stehr A, Ploner F, Traeger K, et al. Plasma disappearance of indocyanine green: a marker for excretory liver function? *Intensive Care Med*. 2005;31(12):1719-1722.
40. Nicholson JP, Wolmarans MR, Park GR. The role of albumin in critical illness. *Br J Anaesth*. 2000;85(4):599-610.

41. Lescot T, Karvellas C, Beaussier M, Magder S. Acquired liver injury in the intensive care unit. *Anesthesiology*. 2012;117(4):898-904.
42. Ba L, Wu DQ, Qian AY, Zhang M, Xiong B. Dynamic changes of serum cholinesterase activity after severe trauma. *Journal of Zhejiang University. Science. B*. 2014;15(12):1023-1031.
43. Kamolz LP, Andel H, Greher M, Ploner M, Meissl G, Frey M. Serum cholinesterase activity reflects morbidity in burned patients. *Burns : journal of the International Society for Burn Injuries*. 2002;28(2):147-150.
44. al-Kassab AS, Vijayakumar E. Profile of serum cholinesterase in systemic sepsis syndrome (septic shock) in intensive care unit patients. *European journal of clinical chemistry and clinical biochemistry : journal of the Forum of European Clinical Chemistry Societies*. 1995;33(1):11-14.
45. Ofek K, Krabbe KS, Evron T, et al. Cholinergic status modulations in human volunteers under acute inflammation. *Journal of molecular medicine (Berlin, Germany)*. 2007;85(11):1239-1251.
46. Myrianthefs P, Evodia E, Vlachou I, et al. Is routine ultrasound examination of the gallbladder justified in critical care patients? *Critical care research and practice*. 2012;2012:565617.
47. Ahvenjarvi L, Koivukangas V, Jartti A, et al. Diagnostic accuracy of computed tomography imaging of surgically treated acute acalculous cholecystitis in critically ill patients. *J Trauma*. 2011;70(1):183-188.
48. Jaeger C, Mayer G, Henrich R, et al. Secondary sclerosing cholangitis after long-term treatment in an intensive care unit: clinical presentation, endoscopic findings, treatment, and follow-up. *Endoscopy*. 2006;38(7):730-734.
49. Kulaksiz H, Heuberger D, Engler S, Stiehl A. Poor outcome in progressive sclerosing cholangitis after septic shock. *Endoscopy*. 2008;40(3):214-218.
50. Trauner M, Fickert P, Stauber RE. Inflammation-induced cholestasis. *J Gastroenterol Hepatol*. 1999;14(10):946-959.
51. Russell DW. The enzymes, regulation, and genetics of bile acid synthesis. *Annu Rev Biochem*. 2003;72:137-174.
52. Chiang JYL. Bile acids: regulation of synthesis. *J Lipid Res*. 2009;50(10):1955-1966.
53. Vanwijngaerden Y-M, Langouche L, Derde S, et al. Impact of parenteral nutrition versus fasting on hepatic bile acid production and transport in a rabbit model of prolonged critical illness. *Shock*. 2014;41(1):48-54.
54. Cherrington NJ, Slitt AL, Li N, Klaassen CD. Lipopolysaccharide-mediated regulation of hepatic transporter mRNA levels in rats. *Drug Metab Dispos*. 2004;32(7):734-741.
55. Zimmerman TL, Thevananther S, Ghose R, Burns AR, Karpen SJ. Nuclear export of retinoid X receptor alpha in response to interleukin-1beta-mediated cell signaling: roles for JNK and SER260. *J Biol Chem*. 2006;281(22):15434-15440.
56. Andrejko KM, Raj NR, Kim PK, Cereda M, Deutschman CS. IL-6 modulates sepsis-induced decreases in transcription of hepatic organic anion and bile acid transporters. *Shock*. 2008;29(4):490-496.
57. Kim PK, Chen J, Andrejko KM, Deutschman CS. Intraabdominal sepsis down-regulates transcription of sodium taurocholate cotransporter and multidrug resistance-associated protein in rats. *Shock*. 2000;14(2):176-181.
58. Ruemmele P, Hofstaedter F, Gelbmann CM. Secondary sclerosing cholangitis. *Nat Rev Gastroenterol Hepatol*. 2009;6(5):287-295.
59. Rady MY, Kodavatiganti R, Ryan T. Perioperative predictors of acute cholecystitis after cardiovascular surgery. *Chest*. 1998;114(1):76-84.
60. Baranano DE, Rao M, Ferris CD, Snyder SH. Biliverdin reductase: a major physiologic cytoprotectant. *Proc Natl Acad Sci U S A*. 2002;99(25):16093-16098.

61. Maruhashi T, Soga J, Fujimura N, et al. Hyperbilirubinemia, augmentation of endothelial function, and decrease in oxidative stress in Gilbert syndrome. *Circulation*. 2012;126(5):598-603.
62. Castilho ÁF, Aveleira CA, Leal EC, et al. Heme oxygenase-1 protects retinal endothelial cells against high glucose- and oxidative/nitrosative stress-induced toxicity. *PLoS One*. 2012;7(8):e42428.
63. Wang WW, Smith DLH, Zucker SD. Bilirubin inhibits iNOS expression and NO production in response to endotoxin in rats. *Hepatology*. 2004;40(2):424-433.
64. Houten SM, Watanabe M, Auwerx J. Endocrine functions of bile acids. *The EMBO journal*. 2006;25(7):1419-1425.
65. Keitel V, Kubitz R, Haussinger D. Endocrine and paracrine role of bile acids. *World J Gastroenterol*. 2008;14(37):5620-5629.
66. Watanabe M, Houten SM, Matak C, et al. Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation. *Nature*. 2006;439(7075):484-489.
67. Porez G, Prawitt J, Gross B, Staels B. Bile acid receptors as targets for the treatment of dyslipidemia and cardiovascular disease. *J Lipid Res*. 2012;53(9):1723-1737.
68. Wang YD, Chen WD, Yu D, Forman BM, Huang W. The G-protein-coupled bile acid receptor, Gpbar1 (TGR5), negatively regulates hepatic inflammatory response through antagonizing nuclear factor kappa light-chain enhancer of activated B cells (NF-kappaB) in mice. *Hepatology*. 2011;54(4):1421-1432.
69. McNeilly AD, Macfarlane DP, O'Flaherty E, et al. Bile acids modulate glucocorticoid metabolism and the hypothalamic-pituitary-adrenal axis in obstructive jaundice. *J Hepatol*. 2010;52(5):705-711.
70. Boonen E, Vervenne H, Meersseman P, et al. Reduced cortisol metabolism during critical illness. *N Engl J Med*. 2013;368(16):1477-1488.
71. Hofmann AF, Eckmann L. How bile acids confer gut mucosal protection against bacteria. *Proc Natl Acad Sci U S A*. 2006;103(12):4333-4334.
72. Pastores SM, Katz DP, Kvetan V. Splanchnic ischemia and gut mucosal injury in sepsis and the multiple organ dysfunction syndrome. *Am J Gastroenterol*. 1996;91(9):1697-1710.
73. Zahiri HR, Perrone EE, Strauch ED. Bile salt supplementation acts via the farnesoid X receptor to alleviate lipopolysaccharide-induced intestinal injury. *Surgery*. 2011;150(3):480-489.
74. Su F, Brands R, Wang Z, et al. Beneficial effects of alkaline phosphatase in septic shock. *Crit Care Med*. 2006;34(8):2182-2187.
75. Heemskerk S, Masereeuw R, Moesker O, et al. Alkaline phosphatase treatment improves renal function in severe sepsis or septic shock patients. *Crit Care Med*. 2009;37(2):417-423, e411.
76. Pickkers P, Heemskerk S, Schouten J, et al. Alkaline phosphatase for treatment of sepsis-induced acute kidney injury: a prospective randomized double-blind placebo-controlled trial. *Crit Care*. 2012;16(1):R14.
77. Bentala H, Verweij WR, Huizinga-Van der Vlag A, van Loenen-Weemaes AM, Meijer DK, Poelstra K. Removal of phosphate from lipid A as a strategy to detoxify lipopolysaccharide. *Shock*. 2002;18(6):561-566.
78. van Veen SQ, van Vliet AK, Wulferink M, Brands R, Boermeester MA, van Gulik TM. Bovine intestinal alkaline phosphatase attenuates the inflammatory response in secondary peritonitis in mice. *Infection and immunity*. 2005;73(7):4309-4314.
79. Goldberg RF, Austen WG, Jr., Zhang X, et al. Intestinal alkaline phosphatase is a gut mucosal defense factor maintained by enteral nutrition. *Proc Natl Acad Sci U S A*. 2008;105(9):3551-3556.
80. Gonnert FA, Recknagel P, Hilger I, Claus RA, Bauer M, Kortgen A. Hepatic excretory function in sepsis: implications from biophotonic analysis of transcellular xenobiotic transport in a rodent model. *Crit Care*. 2013;17(2):R67.

81. Van den Berghe G, Wilmer A, Hermans G, et al. Intensive insulin therapy in the medical ICU. *N Engl J Med*. 2006;354(5):449-461.
82. Van den Berghe G, Wouters P, Weekers F, et al. Intensive insulin therapy in critically ill patients. *N Engl J Med*. 2001;345(19):1359-1367.
83. Dellinger RP, Levy MM, Rhodes A, et al. Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock, 2012. *Intensive Care Med*. 2013;39(2):165-228.
84. Guo K, Ren J, Wang G, et al. Early Liver Dysfunction in Patients With Intra-Abdominal Infections. *Medicine (Baltimore)*. 2015;94(42):e1782.
85. Aninat C, Seguin P, Descheemaeker PN, Morel F, Malledant Y, Guillouzo A. Catecholamines induce an inflammatory response in human hepatocytes. *Crit Care Med*. 2008;36(3):848-854.
86. Oude Elferink RP, Jansen PL. The role of the canalicular multispecific organic anion transporter in the disposal of endo- and xenobiotics. *Pharmacol Ther*. 1994;64(1):77-97.
87. Lammert C, Bjornsson E, Niklasson A, Chalasani N. Oral medications with significant hepatic metabolism at higher risk for hepatic adverse events. *Hepatology*. 2010;51(2):615-620.
88. de Vree JM, Romijn JA, Mok KS, et al. Lack of enteral nutrition during critical illness is associated with profound decrements in biliary lipid concentrations. *The American journal of clinical nutrition*. 1999;70(1):70-77.
89. Braga M, Gianotti L, Gentilini O, Parisi V, Salis C, Di Carlo V. Early postoperative enteral nutrition improves gut oxygenation and reduces costs compared with total parenteral nutrition. *Crit Care Med*. 2001;29(2):242-248.
90. Hu QG, Zheng QC. The influence of Enteral Nutrition in postoperative patients with poor liver function. *World J Gastroenterol*. 2003;9(4):843-846.
91. Hegazi RA, Wischmeyer PE. Clinical review: Optimizing enteral nutrition for critically ill patients--a simple data-driven formula. *Crit Care*. 2011;15(6):234.
92. Casaer MP, Mesotten D, Hermans G, et al. Early versus late parenteral nutrition in critically ill adults. *N Engl J Med*. 2011;365(6):506-517.
93. Favez T, Kerklaan D, Mesotten D, et al. Early versus Late Parenteral Nutrition in Critically Ill Children. *N Engl J Med*. 2016;374(12):1111-1122.

Table 1: Diagnostic tools for critical illness induced cholestasis

Parameter	Description and rationale	Advantage/disadvantage as diagnostic marker for ICU cholestasis
Total Bilirubin	Catabolic end-product of senescent red blood cells. Elevated levels indicate high secretion, intrahepatic dysfunction or post-hepatic obstruction	<ul style="list-style-type: none"> + Partially indicates hepatic excretory function + Correlates with ICU outcome - Biliary excretion dependent on BA-flow - Differentiation in pre/intra/post hepatic origin difficult
Direct Bilirubin	Conjugated isoform of bilirubin. In addition to total bilirubin, elevated levels indicate a maintained conjugation function	<ul style="list-style-type: none"> + Indicates detoxification/conjugation function + Partially indicates hepatic excretory function - Biliary excretion dependent on BA flow - Differentiation in pre/intra/post hepatic origin difficult
Bile acids (BA)	Intrahepatic synthesized drivers of bile-flow. Elevated levels indicate high secretion, intrahepatic dysfunction or post-hepatic obstruction	<ul style="list-style-type: none"> + Indicates ongoing synthesis and impaired feedback + Indicates excretory function - Differentiation in pre/intra/post hepatic origin difficult - Conjugated/unconjugated fraction difficult to measure
Aspartate Transaminase (AST)	Intracellular enzyme mainly located in liver, heart, skeletal muscle, kidney, brain. Serum activity of AST increases after hypoxia with liver parenchymal cell loss	<ul style="list-style-type: none"> + Indicates parenchymal cell loss - Parenchymal cell loss not primary present - Does not indicate liver (dys)function
Alanine Transaminase (ALT)	Intracellular enzyme mainly located in liver. Serum activity of ALT increases after hypoxia with liver parenchymal cell loss	<ul style="list-style-type: none"> + Indicates parenchymal cell loss - Parenchymal cell loss not primary present - Does not indicate liver (dys)function
Gamma-glutamyl transpeptidase (γ -GT)	Intracellular enzyme mainly located in liver, bile ducts, kidney and bone. Cholestasis enhances ALP synthesis and release to the circulation	<ul style="list-style-type: none"> + High sensitivity - Low specificity
Alkaline phosphatase (ALP)	Intracellular enzyme mainly located in liver, bile ducts, kidney and bone. Biliary 'stasis' enhances ALP synthesis and release to the circulation	<ul style="list-style-type: none"> + High sensitivity - Low specificity - Also increased in bone/skeletal disease
glutamate dehydrogenase (GLDH)	Intracellular enzyme mainly located in the (oxygen poor) region of the liver. Serum activity of GLDH increases earlier after hypoxia with liver parenchymal cell loss	<ul style="list-style-type: none"> + May detect early hypoxic injury with higher accuracy - Does not indicate (dys)liver function - Use in ICU patients not well investigated
Indocyanine green plasma disappearance rate	Cyanine dye that is removed from the circulation exclusively by the liver. Decreased levels indicate impaired splanchnic blood flow or excretory function	<ul style="list-style-type: none"> + Partially indicates excretory function + Correlates with mortality and severity of illness - Confounded by hepatic blood flow
Cholinesterase activity	Intrahepatic synthesized esterase that lyses choline-based esters. Decreased levels indicate impaired hepatic synthetic capacity	<ul style="list-style-type: none"> + Might indicate lower synthetic function of the liver - Use in ICU patients not well investigated - No clear diagnostic criteria - Confounded by drugs

Figure 1: Schematic overview of hepatic alterations and serum markers in hypoxic hepatitis and critical illness induced cholestasis. A: In hypoxic hepatitis, oxygen supply to the liver is impaired, resulting in cellular necrosis of hepatocytes. Accordingly, ALT and AST, and to a lesser degree ALP and γ -GT are released into the circulation. Excretory function measured by bilirubin and BA may be mildly impaired. In the recovery phase, oxygen supply to the liver is restored and hepatocytes regenerate. In 30% of patients, clinical jaundice develops after hypoxic hepatitis. B: During ICU cholestasis, intrahepatic alterations in the liver transport machinery results in higher circulating bilirubin and BA. Biliary stasis promotes release of ALP and γ -GT from cholangiocytes. Mild elevation of ALT and AST may also be present. Abbreviations: Bili: Bilirubin, BA: bile acids, ALT: alanine transaminase, AST: aspartate transaminase, ALP: alkaline phosphatase, γ -GT: gamma glutamyl transpeptidase, ULN: Upper limit of normality.

A. Hypoxic Hepatitis

B. Critical illness induced cholestasis