WILEY

Online Proofing System Instructions

The Wiley Online Proofing System allows authors and proof reviewers to review PDF proofs, mark corrections, respond to queries, upload replacement figures, and submit these changes directly from the PDF proof from the locally saved file or while viewing it in your web browser.

1. For the best experience reviewing your proof in the Wiley Online Proofing System please ensure you are connected to the internet. This will allow the PDF proof to connect to the central Wiley Online Proofing System server. If you are connected to the Wiley Online Proofing System server you should see the icon with a green check mark above in the yellow banner.

Connected

Disconnected

2. Please review the article proof on the following pages and mark any corrections, changes, and query responses using the Annotation Tools outlined on the next 2 pages.

3. To save your proof corrections, click the “Publish Comments” button appearing above in the yellow banner. Publishing your comments saves your corrections to the Wiley Online Proofing System server. Corrections don’t have to be marked in one sitting, you can publish corrections and log back in at a later time to add more before you click the “Complete Proof Review” button below.

4. If you need to supply additional or replacement files bigger than 5 Megabytes (MB) do not attach them directly to the PDF Proof, please click the “Upload Files” button to upload files:
5. When your proof review is complete and you are ready to submit corrections to the publisher, please click the “Complete Proof Review” button below:

IMPORTANT: Do not click the “Complete Proof Review” button without replying to all author queries found on the last page of your proof. Incomplete proof reviews will cause a delay in publication.

IMPORTANT: Once you click “Complete Proof Review” you will not be able to publish further corrections.

Once you have Acrobat Reader open on your computer, click on the [Comment](#) tab at the right of the toolbar:

This will open up a panel down the right side of the document. The majority of tools you will use for annotating your proof will be in the [Annotations](#) section, pictured opposite. We've picked out some of these tools below:

1. Replace (Ins) Tool – for replacing text.

Strikes a line through text and opens up a text box where replacement text can be entered.

How to use it

- Highlight a word or sentence.
- Click on the [Replace \(Ins\)](#) icon in the Annotations section.
- Type the replacement text into the blue box that appears.

standard framework for the analysis of microeconomic behavior. Nevertheless, it also led to the development of a new class of strategic form games. The number of competitors in the industry is that the structure of the game is a main component. At the micro level, are exogenous variables and important works on entry by firms (M. Henceforth) we open the 'black b

2. Strikethrough (Del) Tool – for deleting text.

Strikes a red line through text that is to be deleted.

How to use it

- Highlight a word or sentence.
- Click on the [Strikethrough \(Del\)](#) icon in the Annotations section.

there is no room for extra profits as mark-ups are zero and the number of firms (set) values are not determined by Blanchard and Kiyotaki (1987), perfect competition in general equilibrium of aggregate demand and supply in the classical framework assuming monopoly between an exogenous number of firms

3. Add note to text Tool – for highlighting a section to be changed to bold or italic.

Highlights text in yellow and opens up a text box where comments can be entered.

How to use it

- Highlight the relevant section of text.
- Click on the [Add note to text](#) icon in the Annotations section.
- Type instruction on what should be changed regarding the text into the yellow box that appears.

dynamic responses of mark-ups consistent with the VAR evidence

sation by Markov processes. The number of competitors and the impact of demand shocks on the demand-

4. Add sticky note Tool – for making notes at specific points in the text.

Marks a point in the proof where a comment needs to be highlighted.

How to use it

- Click on the [Add sticky note](#) icon in the Annotations section.
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the yellow box that appears.

and supply shocks. Most of the standard framework for the analysis of microeconomic behavior. The number of competitors and the impact of demand shocks on the demand-

5. **Attach File** Tool – for inserting large amounts of text or replacement figures.

Inserts an icon linking to the attached file in the appropriate place in the text.

How to use it

- Click on the **Attach File** icon in the Annotations section.
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.

END

6. **Drawing Markups** Tools – for drawing shapes, lines and freeform annotations on proofs and commenting on these marks.

Allows shapes, lines and freeform annotations to be drawn on proofs and for comment to be made on these marks.

How to use it

- Click on one of the shapes in the Drawing Markups section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, move the cursor over the shape until an arrowhead appears.
- Double click on the shape and type any text in the red box that appears.

ORIGINAL ARTICLE

Economic evaluation of the randomized European Achalasia trial comparing pneumodilation with Laparoscopic Heller myotomy

A. Moonen¹ | O. Busch² | M. Costantini³ | E. Finotti⁴ | J. Tack¹ | R. Salvador³ |
 1 G. Boeckxstaens¹ | G. Zaninotto⁵

¹Department of Gastroenterology, Catholic University of Leuven, Leuven, Belgium

²Gastrointestinal Surgery, Academic Medical Center, Amsterdam, The Netherlands

³Department of Surgical Oncological and Gastroenterological Sciences, University of Padua, Padua, Italy

⁴Department of Surgery, Policlinico di Abano Terme, Abano, Italy

⁵Department of Academic Surgery, St Mary's Hospital, Imperial College, London, UK

Correspondence

G.E. Boeckxstaens, Department of Gastroenterology, Translational Research Center for Gastrointestinal Disorders (TARGID), University Hospital Leuven, Catholic University Leuven, Leuven, Belgium. Email: Guy.boeckxstaens@med.kuleuven.be and

G. Zaninotto, Department of Academic Surgery, St Mary's Hospital, Imperial College, London, UK. Email: g.zaninotto@imperial.ac.uk

Abstract

Background: A recent multicenter randomized trial in achalasia patients has shown that pneumatic dilation resulted in equivalent relief of symptoms compared to laparoscopic Heller myotomy. Additionally, the cost of each treatment should be also taken in consideration. Therefore, the aim of the present study was to perform an economic analysis of the European achalasia trial.

Methods: Patients with newly diagnosed achalasia were enrolled from 2003 to 2008 in 14 centers in five European countries and were randomly assigned to either pneumatic dilation (PD) or laparoscopic Heller (LHM). The economic analysis was performed in the three centers in three different countries where most patients were enrolled (Amsterdam [NL], Leuven, [B] and Padova [I]) and then applied to all patients included in the study. The total raw costs of the two treatments per patient include the initial costs, the costs of complications, and the costs of retreatments.

Results: Two hundred and one patients, 107 (57 males and 50 females, mean age 46 CI: 43-49 years) were randomized to LHM and 94 (59 males and 34 females, mean age 46 CI 43-50 years) to PD. The total cost of PD per patient was quite comparable in the three different centers; €3397 in Padova, €3259 in Amsterdam and €3792 in Leuven. For LHM, the total costs per patient were highest in Amsterdam: €4488 in Padova, €6720 in Amsterdam, and €5856 in Leuven.

Conclusion: In conclusion, the strategy of treating achalasia starting with PD appears the most economic approach, independent of the health system.

KEYWORDS

Achalasia, costs, economic, laparoscopic Heller myotomy, pneumatic dilation

1 | INTRODUCTION

Esophageal achalasia is a relatively uncommon disease, characterized by an impairment of smooth muscle activity and unrelaxing Lower Esophageal Sphincter.¹ These abnormalities lead to impaction of the

bolus at the level of the cardia inducing the sensation of dysphagia. Other typical symptoms include regurgitation, cough especially at night and chest pain.² The etiology of achalasia remains elusive, but the medical community learnt how to manage the disease, or better, how to effectively palliate its symptoms since many decades, by decreasing the obstruction outflow at the cardia either by forceful disruption with a dilating balloon or by dividing surgically the LES muscle fibers.³ The

G. Boeckxstaens and G. Zaninotto shared senior authorship

Journal Name		NMO	
Manuscript No.		13115	
WILEY		No. of pages: 7	
Dispatch: 10-5-2017		CE: Viswanathan	
PE: Deivapriyadharsini D.			

latter has been performed in the last twenty years through a minimal invasive approach and has gained a vast popularity.⁴⁻⁶ Nevertheless, over the years, firm opinions have been expressed in favor or against each treatment and the competition between the two therapies is not yet settled.^{7,8} The choice of the treatment for achalasia is still left to the patient or to his/her physician and it is based on patient's preference, patient's surgical risk, age, or availability of local expertise in one of the techniques. A recent multicenter randomized trial has shown that a series of pneumatic dilation (PD) resulted in equivalent relief of achalasia symptoms compared to laparoscopic Heller (LHM) myotomy, both at mid-term (2 years) and long-term (5 years) follow-up.^{9,10} In this scenario and given the increasing attention to the economic health aspects, the cost of each treatment should also be taken into consideration.

The few economic analyses since now performed on achalasia treatments were mainly based on models derived from data obtained by the medical literature or by panels of experts or even by assumptions in the absence of sound data.¹¹⁻¹³ Moreover, only two of these studies were settled within the "European health-care system" with a mix of social and private medicine, while other studies refer to the United States mainly private health-care system.^{11,14}

The aim of the present study was to perform an economic analysis of the most recent randomized trial on achalasia treatment available, considering the initial cost of each therapy, the cost of the complications and that of retreatments performed in patients in whom the primary therapy failed. These costs were calculated in three of the Hospitals in whom most patients were enrolled: the University Hospital Leuven, Belgium, the Academic Medical Center, Amsterdam, the Netherlands and the Padova University Hospital, Italy and then applied to the whole patients group.

2 | MATERIAL AND METHODS

2.1 | Patients

Symptomatic patients with a newly diagnosed achalasia based on clinical, radiological, and manometric criteria were enrolled from 2003 to 2008 in 14 Centers in five European countries and were randomly assigned to either PD or LHM.⁹

Patients' symptoms (dysphagia, regurgitation and chest pain) were scored according to Eckardt (a score of 0 indicates the absence of symptoms, 1 indicates occasional symptoms, 2 daily symptoms and 3 indicates symptoms at each meal) and weight loss (0 no weight loss, 1 indicating a loss <5 Kg, 2 indicating a loss of 5-10 kg, and 3 indicating a loss >10 Kg). The maximum Eckardt score is 12 and an Eckardt score greater than 3 was assumed as the threshold to define the failure of treatment.

The data for this study ie, the duration of hospitalization of each patient, the number and type of complication, the treatment received, the number of failures and the type of retreatment performed in patients who failed, were obtained by reviewing the case report files stored in the coordination center and, when necessary, by interviewing directly the data manager of each center involved. The data were

Keypoints

- Pneumatic dilation and laparoscopic Heller myotomy have similar success rates after 5 years of follow-up, but economic aspects of the treatment are not taken into account in the treatment strategy of achalasia
- Treating achalasia starting with PD appears the most economic, independent of the health system, even when costs of complications and retreatment of failures are taken into account.
- Economic aspects should be added to the decision tree when new guidelines for the treatment of achalasia will be developed.

updated until December 31, 2013. The institutional review board at each hospital approved the study protocol and written informed consent was obtained from all patients. The study was registered in the "Netherlands trial register" (NTR37) and in the Current Controlled Trials registry (ISRCTN56304564).

2.2 | Interventions

2.2.1 | Pneumatic dilation

Pneumatic dilation was performed using a Rigidflex Balloon dilating the sphincter at a pressure of 5 and 8 psi per one minute each, starting with a 30 mm diameter balloon. The first dilation was followed within 3 weeks by a second dilation using a larger balloon (35 mm). All patients therefore received at least two dilations. If the Eckardt score was greater than 3, a third dilation with a 40 mm Balloon was performed. Patients with a recurrence of symptoms within the follow-up period were allowed to receive a second series of dilation using a 35 mm and—if necessary—40 mm balloon. A third and final series of dilation was allowed only if symptoms recurred more than 2 years after the latter series of dilations. If symptoms recurred within 2 years from the second series, they were considered failures.²

In Leuven, the two dilations were performed on two consecutive days with an overnight stay.

In the Padova University Hospital, patients after dilation were admitted as day-hospital patients. If overnight stay was necessary, the patients were admitted to the surgical ward. Amsterdam patients were treated as "day-hospital" patients, but if in case they required an overnight stay they were kept in the same low-intensity ward.

2.2.2 | Laparoscopic Heller myotomy with Dor antireflux procedure

The distal esophagus was mobilized on the lateral and anterior side and a myotomy was performed extending at least 6 cm above the gastroesophageal junction and at least 1 to 1.5 cm over the stomach. Thereafter, an anterior 180-degree fundoplication according to the

method of Dor was performed. If symptoms recurred after surgery, ie, an Eckardt score > 3, the patient was considered a treatment failure. In all three centers, a naso-gastric tube was left in place for the first 24 hours. In the first p.o. day, a gastrograffin swallow was given and—if no leakage from the myotomy was documented—the patient was allowed to drink. Soft food was given in 2nd p.o. day. Timing of patient discharge was determined by the treating physician and based on the patients' medical condition and social situation.

2.3 | Study outcomes

The primary aim of the study was to establish the cost to treat a patient following one or the other of the two therapies including the cost of complications and retreatment.

2.4 | Retreatment

Patients who failed were advised to have pneumatic dilation if they were in the surgical arm or LM if they were in the endoscopic dilation arm. As failures were—by definition—excluded from the study, these were only recommendations and the final decision on the type of retreatment was taken by the patient himself and/or by his/her general practitioner.

2.5 | Economic analysis

In this study, the cost analysis was performed in the three centers where most patients were enrolled (Amsterdam, Leuven and Padova) and then applied to all patients included in the study for each center separately, based on the total number of successful treatments, failures, retreatment of failures and complications. Indirect costs, such as non-medical services relating to the treatment (eg, transportation and family care, and indirect morbidity costs such as absence from work for hospitalization and convalescence) were not assessed. Costs of treatments were calculated using the Activity-Based Costing (ABC) method, which is a full costing system that considers treatment as a production process as a whole (eg, the costs of a single blood test, disposable instruments, operating room time, costs of medical and non-medical personnel, etc.). The cost of personnel (medical doctors and nurses) was based on the standard cost indicated in the hospital budget per person per year and divided by the time required for the treatments. The cost of hospital stay (both for a full admission or "day-hospital" admission) was provided by the Hospital Administrations. The cost for treating a complication was calculated by adding the cost of any re-operation, and the cost of the prolonged hospital stay (the cost of the prolonged hospital stay in all three hospitals included also the cost of extra medication and diagnostic.). The cost of any retreatments for failure of the primary therapy was calculated assuming that was the same as that of the primary treatment (PD or LHM) with the same duration of hospital stay. The final cost of each treatment was calculated adding the cost of the primary procedure (multiple PDs or LM), the cost of the complications and the cost of the retreatments, according to the following formula:

Costs of a treatment in center A = (Cost of primary treatments in center A × total No. of patients in the study) + (Cost of treatments for complications in center A × total No. of complications in the study) + (Cost of retreatments of failures in center A × total No. of failures in the study)/total No. of patients in the study.

All costs were rounded to unit (<50 cents = 0; >50 cents = 1 €).

2.6 | Statistical analysis

Continuous variables are presented as means with 95% confidence interval and were compared using the Student's *t* test. A *P* < .05 was considered statistically significant.

3 | RESULTS

The first thirteen patients of the 214 randomized patients had pneumatic dilation with a 35 mm Rigidflex balloon (as per initial protocol). Perforations occurred in four of them, representing a much higher percentage than reported in the medical literature. This led to a modification of the protocol and the first dilation was performed with a smaller diameter balloon (30 mm). These 13 patients were excluded from the original study, as well as from the economic audit. In addition, two patients were randomized to PD but erroneously treated by LHM. These patients were analyzed according the treatment received. Thus, the final study group comprised 201 patients, 107 (57 males and 50 females, mean age 46 CI: 43-49 years) were randomized to LHM and 94 (59 males and 34 females, mean age 46 CI: 43-50 years) to PD.

3.1 | Uncomplicated patients

3.1.1 | Pneumatic dilation

In eighty-nine patients PD was uneventful. Sixty-three uncomplicated patients had their PD in a day-hospital setting, and 26 uncomplicated patients stayed in the hospital more than 1 day. The mean hospital stay of uncomplicated patients after PD was 2.8, (CI: 2.4-3.1). Twenty-four patients needed retreatment after the initial PD, 21 patients needed only one series of redilation. Five patients were retreated twice (Figure 1: Panel A).

The costs of a single dilation (inclusive of the access in the day-hospital suite) were €1261 in Padova, €734 in Amsterdam, and €981 in Leuven. The costs in Padova included the presence of an extra nurse and an anesthesiologist (required by internal regulations for administration of Propofol) and the regular use of a gastrograffin swallow study after each dilation. In Leuven, dilations were mostly performed with the patient hospitalized for 2 days. The cost of a 2-day hospitalization including dilation was €1550. The cost per each extra day of hospital admission after a dilation was €271 in Padova, €571 in Amsterdam, and €437 in Leuven. The total cost of a successful pneumodilation, including redilation, was therefore, €3060 in Padova, €2049 in Amsterdam, and €2569 in Leuven (Table 1).

FIGURE 1 A, Flow chart of pneumodilation treatment. B, Flow chart of laparoscopic Heller myotomy treatment

TABLE 1 Cost of successful PD (including redilations) in the three different centers. Data are expressed in euro per patient

	Costs of first series of PD	Costs of retreatment	Total costs
Padova	2435	2320	3060
Amsterdam	1930	1790	2412
Leuven	2064	1873	2569

3.1.2 | Laparoscopic Heller myotomy

Ninety-two patients had an uneventful recovery after LHM; their mean hospital stay was 4.8 days (CI: 4.0-5.3). The cost of the surgical procedure was €2811, €3525, and €3392 for Padova, Amsterdam, and Leuven respectively. Also, the cost of hospital admission was different in the different centers: €270/d in Padova; €539/d in Amsterdam, and €401/d in Leuven. Given a mean hospital stay after LHM of 4.8 days, the costs for hospital stay were €1296 in Padova, €2587 in Amsterdam, and €1923 in Leuven. Therefore, the total cost

of LHM including surgical procedure and hospital stay were €4107 in Padova, €6112 in Amsterdam, and €5315 in Leuven.

3.2 | Complications

3.2.1 | Pneumatic dilation

After adaptation of the dilation protocol (starting with 30 mm balloon instead of 35 mm balloon), five perforations occurred (5.3%) during the entire study. Three were treated conservatively, ie, patients were fastened (nothing by mouth) and were treated with antibiotics, and they received nasogastric suction and total parenteral nutrition. Two patients underwent surgery during which the perforation was sutured and a myotomy was added. The median hospital stay after a perforation was 13.4 (CI: 3.8-23) days. Patients who were treated conservatively stayed on average 16±6 days in hospital, compared to 9.5±7.5 days for patients who were operated ($P = .4$). The costs of a complicated patient were €11008 in Padova, €9224 in Amsterdam, and €11627 in Leuven.

3.2.2 | Laparoscopic Heller myotomy

Thirteen of the 107 patients (12%) had a mucosal tear during the myotomy. In twelve patients, the tear was sutured during laparoscopy; in one patient required a conversion to open approach. Patients who had a mucosal tear had a prolonged hospital stay (mean: 8.9 days [CI: 5.7-12.1] compared to 4.7 [CI 4.0-5.3], $P < .0001$; Student's *t* test). One additional patient had a myocardial infarction and was treated with a coronary stent. He remained hospitalized during 31 days. One patient remained hospitalized during 5 days because of a postoperative bleeding that was treated conservatively with a drain. Taken all complications into account, the average costs of complications per patient was €5542 in Padova, €8976 in Amsterdam, and €7444 in Leuven.

3.3 | Retreatment of failures

3.3.1 | Pneumatic dilation

Until December 2013, 15 patients were considered as treatment failures. Eight patients because of recurrent symptoms, while seven patients refused further dilation(s). PD treatment failures were most often treated with LHM ($n=12$), additional pneumodilations occurred in two patients and botox injection was given in one patient. No complications were observed in this group. The costs of retreatment per patient were €3723 in Padova, €5192 in Amsterdam, and €4578 in Leuven.

3.3.2 | Laparoscopic Heller myotomy

At December 2013, 21 patients experienced recurrence of symptoms after LHM. Ten patients were treated with PD. Six patients required only one session of PD, two patients had two sessions and the other two patients had three and five dilations respectively. (Figure 1: Panel B). One patient had a pneumatic dilation and a redo myotomy (through a laparoscopic approach), six patients refused further treatment, and four patients were lost at follow-up. No complications were reported in this group. The costs of retreatment per patient were €1428 in Padova, €1050 in Amsterdam, and €1236 in Leuven.

3.4 | Total costs

The total raw costs of the two treatments per patient include the initial costs, the costs of complications, and the costs of retreatments and is calculated based on the flow chart shown in Figure 1. The total cost

of PD per patient was quite comparable in the three different centers; €3997 in Padova, €3258 in Amsterdam, and €3792 in Leuven. For LHM, the total costs per patient were highest in Amsterdam: €4488 in Padova, €6720 in Amsterdam, and €5856 in Leuven (Table 2).

4 | DISCUSSION

Until a few years ago, the rarity of achalasia and the lack of randomized clinical trials confronting different therapeutic options, led to a customer based (patient or physician) choice of treatment. The latter was largely determined by factors such as the availability of skilled surgeons or endoscopists performing the dilation procedure, the subjective preference of the treating physician, or the patient's propensity toward a more or less invasive procedure. Treatment costs, although a relevant issue, probably had only a marginal influence on this choice, given the low number of patients treated and the few studies reporting on this topic. In the current study, we present the economic analysis of the European Achalasia trial, a large multicenter randomized trial comparing LHM (with Dor's fundoplication) with pneumatic dilation as first treatment for achalasia. Our present economic analysis indicates that initial treatment with PD is cheaper than initial treatment with LHM, even when costs of complications and retreatment of failure are included.

In the present study, the costs of LHM and PD were compared based on data collected during the European Achalasia Trial. The latter is a large multicenter randomized trial comparing LHM with pneumatic dilation as a treatment for achalasia with long-term follow-up (at least 5 years). The European achalasia trial was started in 2003 and showed that both treatments are equally effective.^{9,10} The trial was performed in 14 centers scattered around Europe in countries with different health system, significantly complicating the economic analysis. Hence, data analysis was limited to the three centers that randomized most of the patients (47%), ie, Padova, (Italy), Amsterdam (the Netherlands), and Leuven (Belgium). The strength of our study however is that, to date, it is the only comparative health economic analysis with respect to the management of newly diagnosed esophageal achalasia carried out within the framework of an adequately sized RCT, and most importantly, which is based on actual costs. The data² used in our study were indeed extracted from the RCT files of the European Achalasia Trial, and thus reflect better "the real world" of treating achalasia patients in specialized units. Our study contrasts with previous economic studies that used "ideal data" to build the models mainly conducted on data subtracted from the literature. In

TABLE 2 Overview of costs of LHM compared to PD Cost of successful initial uncomplicated treatment, complications, retreatment after failure and total costs with PD and LHM are given for the three centers

	Success		Complications		Retreatment after failure		Total cost	
	PD	LHM	PD	LHM	PD	LHM	PD	LHM
Padova	3060	4107	11008	5542	3956	1428	4072	4488
Amsterdam	2412	9159	11360	12134	8195	1437	4110	9858
Leuven	2569	5315	11627	7444	4578	126	3792	5856

addition, a mixture of surgical procedures was included in these studies, ie, using both laparoscopic and open surgery with and without antireflux procedure.¹²⁻¹⁵ In the most recent analysis, the probability of each event and its plausible range was even estimated by a panel of five experts, rather than based on real data.¹¹ Moreover graded distention for PD was not included, which led clearly to an overestimation of the failure rate of PD. Only one previous study has been based on results of a real life scenario within the framework of a randomized controlled trial.¹⁴ In this study, 51 patients were randomized to LHM or PD with a graded distention protocol. In males, dilation was started at 35 mm, consequently an unacceptable high perforation rate was noted (8%). Moreover, all perforations were treated surgically, which is not the usual policy, and follow-up lasted only 12 months. Although the data reported so far are of great interest and importance, they are not really based on daily clinical practice and comprise only a short period of follow-up. A limitation of our study however is that the cost of complications was mainly based on the duration of hospital stay and not on the cost of each single patient's treatment.

The main finding of our study is that a treatment strategy of idiopathic achalasia starting with PD is less costly than a strategy starting with LHM, even when cost of complications, redilation, and retreatment of failures are included. LHM was more costly in all three centers. In Padova, the costs of LHM were only 9% (€4488) higher than PD, in Leuven and Amsterdam the difference was 35% (€5856) and 58% (€9858) respectively. This result is in line with the model-based study of Karanicolas et al.¹¹ This study, using a model in which recurrence of symptoms after PD was managed with LHM, showed that the strategy starting with LHM was more costly than the strategy with PD (€7300 vs €4500 after 5 years). Also, the study based on a smaller RCT of Kostic et al. came to the same conclusion, although follow-up was very short.¹⁴ The total median costs, including follow-up costs were €1969 for PD and €7545 for myotomy after a follow-up period of 12 months. A major difference though between our study and previous model-based reports is that the length of hospital stay in our study is longer for both pneumatic dilation and surgery. This is explained by the fact that especially in Italy, patients live at a relevant distance from the often high-volume hospital where treatment is provided. As a consequence, patients need to be admitted 1 day before treatment, and likewise, they can only be discharged safely the day after the procedure. Moreover, treating physicians were likely more conservative in discharging patients as treatment was performed within the framework of a clinical trial, which is reflected in the fact that 20% of uncomplicated PD patients were admitted to the hospital. Other possible explanations for the increased hospital stay might be a reimbursement model linked to the duration of hospitalization in some countries. Be as it may, it is evident that the cost can be reduced by shortening the hospital stay, but this applies for both procedures implying that the difference in costs will remain largely in favor of PD.

Indirect costs were not taken into account in our study. However, as the hospitalization and recovery period is longer after LHM than after PD, including the indirect morbidity costs due to work absence, would only increase the difference in costs between LHM and PD.

We found relevant differences in the costs of the procedures for the raw hospital costs for both LHM and PD in the different countries evaluated. Raw costs of LHM in Italy and Belgium were one-third less than those in the Netherlands. In Padova, the costs of LHM were only €4488 compared to €5856 in Leuven and €6720 in Amsterdam. The costs of PD, including costs of redilation, were comparable in all three centers: €3997 in Padova, €3259 in Amsterdam, and €3792 in Leuven. PD was relatively more expensive in Italy due to differences in health-care regulations (the need of an anesthesiologist and extra nurse in Italy). The reason for these variations between centers is mainly that these three countries have a different health system based largely on private insurances in the Netherlands, completely subsidized by the state in Italy and with a mixed component of insurances and state reimbursement in Belgium which makes comparison between countries very difficult. For the same reason, it is difficult to generalize the results to other continents. In years in which the free circulation of patients is easier, this phenomenon of different costs for the same procedures should be studied and monitored with attention.

In conclusion, the raw hospital costs of a strategy starting from pneumatic dilation are less expensive in all three countries, although the differences in Italy were less pronounced. Therefore, at least in the framework of the European achalasia trial, the strategy of treating achalasia starting with PD appears the most economically convenient, independently of the health-care system. This new information should be added to the decision tree when new guidelines for the treatment of achalasia will be developed.

DISCLOSURE

The authors have nothing to disclose.

AUTHORS CONTRIBUTION

This investigator initiated trial was conceived by the last authors (GB and GZ). All the authors had substantial roles in the design of the study, the interpretation of the data, and the writing of the manuscript. The first author and last authors wrote the first and final versions of the manuscript and, in consultation with the other authors, made the decision to submit the manuscript for publication. No commercial entity had any role in the study. All the authors vouch for the completeness and accuracy of the data.

REFERENCES

1. Moonen AJ, Boeckxstaens GE. Management of achalasia. *Gastroenterol Clin North Am*. 2013;42:45-55.
2. Boeckxstaens GE, Zaninotto G, Richter JE. Achalasia. *Lancet*. 2014;383:83-93.
3. Moonen A, Boeckxstaens G. Current diagnosis and management of achalasia. *J Clin Gastroenterol*. 2014;48:484-490.
4. Oelschlager BK, Chang L, Pellegrini CA. Improved outcome after extended gastric myotomy for achalasia. *Arch Surg*. 2003;138:490-495.
5. Perrone JM, Frisella MM, Desai KM, Soper NJ. Results of laparoscopic Heller-Toupet operation for achalasia. *Surg Endosc*. 2004;18:1565-1571.

6. Schuchert MJ, Luketich JD, Landreneau RJ, et al. Minimally-invasive esophagomyotomy in 200 consecutive patients: factors influencing postoperative outcomes. *Ann Thorac Surg*. 2008;85:1729-1734.
7. Rohof WO, Moonen A, Boeckxstaens GE. Reply to Massey. *Gastroenterology*. 2013;144:e18.
8. Richter JE, Boeckxstaens GE. Management of achalasia: surgery or pneumatic dilation. *Gut*. 2011;60:869-876.
9. Boeckxstaens GE, Annese V, des Varannes SB, et al. Pneumatic dilation versus laparoscopic Heller's myotomy for idiopathic achalasia. *N Engl J Med*. 2011;364:1807-1816.
10. Moonen A, Annese V, Belmans A, et al. Long-term results of the European achalasia trial: a multicenter randomised controlled trial comparing pneumatic dilation versus laparoscopic Heller myotomy. *Gut*. 2015;???-???-???
11. Karanicolas PJ, Smith SE, Inculet RI, et al. The cost of laparoscopic myotomy versus pneumatic dilatation for esophageal achalasia. *Surg Endosc*. 2007;21:1198-1206.
12. O'Connor JB, Singer ME, Imperiale TF, Vaezi MF, Richter JE. The cost-effectiveness of treatment strategies for achalasia. *Dig Dis Sci*. 2002;47:1516-1525.
13. Imperiale TF, O'Connor JB, Vaezi MF, Richter JE. A cost-minimization analysis of alternative treatment strategies for achalasia. *Am J Gastroenterol*. 2000;95:2737-2745.
14. Kostic S, Johnsson E, Kjellin A, et al. Health economic evaluation of therapeutic strategies in patients with idiopathic achalasia: results of a randomized trial comparing pneumatic dilatation with laparoscopic cardiomyotomy. *Surg Endosc*. 2007;21:1184-1189.
15. Parkman HP, Reynolds JC, Ouyang A, Rosato EF, Eisenberg JM, Cohen S. Pneumatic dilation or esophagomyotomy treatment for idiopathic achalasia: clinical outcomes and cost analysis. *Dig Dis Sci*. 1993;38:75-85.

How to cite this article: Moonen A, Busch O, Costantini M, et al. Economic evaluation of the randomized European Achalasia trial comparing pneumodilation with Laparoscopic Heller myotomy. *Neurogastroenterol Motil*. 2017;00:e13115. <https://doi.org/10.1111/nmo.13115>

Graphical Abstract

The contents of this page will be used as part of the graphical abstract of html only. It will not be published as part of main.

(A)

(B)

Treating achalasia starting with PD appears the most economic, independent of the health system, even when costs of complications and retreatment of failures are taken into account. Economic aspects should be added to the decision tree when new guidelines for the treatment of achalasia⁵ will be developed.

Author Query Form

Journal: NMO

Article: 13115

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.

Many thanks for your assistance.

Query reference	Query	Remarks
1	AUTHOR: Please confirm that given names (red) and surnames/family names (green) have been identified correctly.	
2	AUTHOR: Define RCT	
3	AUTHOR: Please provide the volume number, page range for reference [10].	
4	AUTHOR: Please approve or update the conflicts of interest statement.	
5	AUTHOR: Please check and confirm if the copy-edited Graphical Abstract caption is fine.	