To cite this article:

Cabrieto, J., Adolf, J., Tuerlinckx, F., Kuppens, P., & Ceulemans, E. (in press). An objective, comprehensive and flexible statistical framework for detecting early warning signs of mental health problems. *Psychotherapy and Psychosomatics*.

An objective, comprehensive and flexible statistical framework for detecting early warning signs of mental health problems

Jedelyn Cabrieto^{1*†}, Janne Adolf^{1†}, Francis Tuerlinckx¹, Peter Kuppens¹, Eva Ceulemans¹

¹ Research Group of Quantitative Psychology and Individual Differences, KU Leuven - University of Leuven, Leuven, Belgium

* Corresponding author

Jedelyn Cabrieto

Research Group of Quantitative Psychology and Individual Differences

KU Leuven - University of Leuven

Tiensestraat 102

3000 Leuven, Belgium

Tel: +32 16 32 03 74

E-mail: jed.cabrieto@kuleuven.be

† These authors contributed equally to this work

1. Text

The early detection of mental health problems, such as depression or psychosis, is of great scientific and societal interest as it allows for prevention or early intervention. Fueled by dynamical systems theory and grounded in developmental theories [1], research is increasingly focusing on the identification of early warning signs (EWS [2]) preceding the onset of psychopathology (also referred to as prodromal symptoms [3]). The key idea is that transitions between global states, e.g. from not-being to being depressed, are foreshadowed by subtle changes in the daily fluctuations and dynamics of individual symptoms [4]. Initial evidence for this EWS hypothesis has been found for depression [5]. Detecting EWS is however not straightforward, and most statistical approaches so far have been mostly univariate, descriptive and rather ad hoc. However, given what is at stake, for the at-risk individual and society, comprehensive, objective and yet flexible, statistical tools for accurately detecting EWS are urgently needed.

In this paper, we propose a statistical framework, Kernel change point detection (KCP; [6-7]), that detects EWS by testing whether summary statistics of repeatedly measured symptoms (e.g., means, variances, correlations, auto-correlations) change across time. The method, a detailed description of which can be found in the Supplementary Material,

- is flexible, in that it can be easily adapted to any summary statistic that may potentially yield EWS
- is comprehensive, in that multiple variables can be monitored simultaneously
- is objective in that it implements formal decision procedures, such as a kernel-based algorithm to search for optimal change point locations, a permutation-based significance-test to decide whether at least one of the located change points is statistically significant, and a penalty approach to decide on the total number of change points

We demonstrate the large potential of KCP for studying EWS by re-analyzing experience sampling data that stem from an anti-depressant dose reduction experiment tracking a single participant diagnosed with major depression for 239 days over five experimental phases [5]. Momentary mood ratings were collected via a smartphone that beeped the participant 10 times a day at random occasions. Importantly, the patient relapsed into depression in the course of the experiment.

Given that Wichers et al. [5] describe changes in the variances, auto-correlations and correlations of these variables before relapse, we monitor the corresponding running statistics (computed in a moving window of size 25; see Supplementary Material) with KCP to test for significant change points before relapse. As these statistics quantify affective dynamics linked to psychopathology [8-9], it makes indeed sense to expect that change points in them can serve as EWS of an upcoming depressive episode.

To enhance specificity of results (i.e., the probability with which changes are attributable to the running statistic monitored) we usually recommend to first implement KCP on the running means and filter the data for significant mean changes, before other changes are searched for. Since the present data are already detrended (see Wichers et al. [5]), we skip this step here and run three separate KCP analyses, on the running variances, auto-correlations and correlations (yielding KCP-var, KCP-ac, and KCP-corr). Note that we replaced values deviating from the mean by more than 3 SDs with maximum values within 3 SDs. Moreover, we removed observations more than one lag apart and overnight lags to keep time intervals invariant for the accurate computation of auto-correlations.

Our results show that each monitored running statistic contains at least one significant change point (for the exact p-values and multiple testing corrections used, see Supplementary Material). The penalty approach further reveals that, for the running auto-correlation, there is one change point observed at Day 86 (41 days before relapse), with all variables exhibiting an increase in auto-

correlation (max $\Delta_AC = .20$ for worry). For the running variance, a major change point occurs at Day 130, which is already 3 days after relapse. However, if a second change point is searched for, we detect it 53 days before relapse, at Day 74. This second change point is mainly characterized by an elevation in the variance of mental unrest ($\Delta_var = .58$), while the change point at Day 130 is marked by drastic variance jumps for worry ($\Delta_var = 1.01$) and suspicion ($\Delta_var = .97$). Lastly, for the running correlation, we find a change point at Day 116, characterized by all pairwise correlations being strengthened except for the one between PA and mental unrest. The largest correlation shifts occur for PA and Worry ($\Delta_corr = .39$), PA and suspicion ($\Delta_corr = .39$), NA and worry ($\Delta_corr = .38$), and NA and suspicion ($\Delta_corr = .38$).

As we are interested in EWS, we repeated the analyses excluding all time points after relapse. This yields very similar evidence. That is, although no significant change point was detected for the running auto-correlation, significant change points were flagged for the running variance at Day 88 and the running correlation at Day 116. These change points are consistently characterized by upward-shifts in the statistics' values.

Summarizing, the proposed KCP approach allows to monitor *different* summary statistics of *multiple* variables simultaneously for change points. The method is statistically sound, well-suited for explorative usage and has been shown to identify valid change points in a large variety of applications [6-7]. Here, we were able to reveal change points in a number of running statistics of affective symptoms, detecting EWS before relapse into depression. Specifically, we see that correlation changes, occur relatively late, whereas variance and auto-correlation changes occur early (11 days vs. more than a month before relapse). Thus, monitoring these statistics might be more informative and can benefit timely intervention and enable prevention.

2. Statements

2.1. Acknowledgement

The authors would like to thank Marieke Wichers and Peter Groot for making the data of the anti-depressant dose reduction experiment publicly available.

2.2. Disclosure Statement

The authors have no conflicts of interest to declare.

2.3. Funding Sources

The research leading to the results reported in this paper was sponsored by a research grant from the Fund for Scientific Research-Flanders (FWO, Project No. G.0582.14 awarded to Eva Ceulemans, Peter Kuppens and Francis Tuerlinckx) and by the Research Council of KU Leuven (GOA/15/003).

3. References (Numerical)

1. Nelson B, McGorry PD, Wichers M, Wigman JT, Hartmann J A. Moving from static to dynamic models of the onset of mental disorder: A review. *JAMA Psychiatry* 2017 May;74(5):528.
2. Scheffer M, Bascompte J, Brock W, Brovkin V, Carpenter S, Dakos, V. Early-warning signals for critical transitions. *Nature* 2009 Sept;461:53-59.
3. Fava G. Subclinical symptoms in mood disorders: pathophysiological and therapeutic implications. *Psychol Med* 1999 Jan;29:47-61.
4. Nesselroade JR. The warp and the woof of the developmental fabric. In Downs RM, Liben LS, Palermo DS, editors. *Visions of aesthetics, the environment, and development: The legacy of Joachim F. Wohlwill*. Hillsdale, NJ: Erlbaum; 1991; pp. 213-240.
5. Wichers M, Groot PC, Psychosystems, ESM Group, EWS Group. Critical slowing down as a personalized early warning signal for depression. *Psychother Psychosom* 2016 Feb;85(2):114-116.
6. Cabrieto J, Tuerlinckx F, Kuppens P, Borbála H, Ceulemans E. Testing for the presence of correlation changes in a multivariate time series: A permutation based approach. *Sci Rep* 2018 Jan;8(1):769.
7. Cabrieto J, Tuerlinckx F, Kuppens P, Wilhelm FH, Liedlgruber M, Ceulemans E. Capturing correlation changes by applying kernel change point detection on the running correlations. *Information Sciences* 2018 Jun;447:117-139.
8. Kuppens P, Verduyn P. Emotion dynamics. *Curr Opin Psychol* 2017 Oct;17:22-26.
9. Trull TJ, Lane SP, Koval P, Ebner-Priemer UW. Affective dynamics in psychopathology. *EMR* 2015 Jul;7(4):355-361.

4. Figure Legends

Fig. 1. Running statistics and the corresponding change points. The first and second panel exhibit the running variances and auto-correlations for the 5 symptoms, while the last panel displays the running correlations for the 10 pairwise correlations. All running statistics are computed using a moving window of 25 beeps. The experimental phases are labeled in the first panel and are indicated by background shading across all panels. The relapse to depression at Day 127 is marked by black vertical lines, while the change points in the running statistics are indicated by red vertical lines.

Running Var

Running AR

Running Corr

Supplementary Material for

An objective, comprehensive and flexible statistical framework for detecting early warning signs of mental health problems

Jedelyn Cabrieto^{1*†}, Janne Adolf^{1†}, Francis Tuerlinckx¹, Peter Kuppens¹, Eva Ceulemans¹

¹ Research Group of Quantitative Psychology and Individual Differences, KU Leuven - University of Leuven, Leuven, Belgium

* Corresponding author

† These authors contributed equally to this work

I. KCP on the running statistics

Step 1. Searching for change points.

The similarity measure

After obtaining the running statistic of interest (e.g., running mean, running variance, running autocorrelation or running correlation) by sliding a time window across the time series and computing the statistic in each window, KCP looks for change points in this running statistic, by examining the similarities between windows. The key idea is straightforward: When there is no change point, there will only be small fluctuations in the similarities. However, when statistics in subsequent windows are very similar, while those that are distant in time are dissimilar, there is more evidence for (a) possible change point(s). KCP employs a kernel function to quantify these similarities. Specifically, we use the Gaussian kernel, such that if RS_i and RS_j denote the running statistic computed in windows i and j , respectively, we compute the following similarity,

$$Gk(RS_i, RS_j) = \exp\left(\frac{-\|RS_i - RS_j\|^2}{2h_{RS}^2}\right), \quad (1)$$

where h_{RS} is the median Euclidean distance between all the RS_i values. The similarity value ranges from 0 to 1, with 0 implying extreme dissimilarity and 1 indicating complete similarity.

Since the goal is to analyze all variables in the time series simultaneously (cfr. comprehensive analysis), our approach is multivariate such that RS_i is a vector of the running statistic for all the V variables in the system. Specifically, for univariate statistics such as the mean, variance and

autocorrelation, the corresponding $\mathbf{RS}_i$ will consist of V elements as each variable yields one running statistic. For correlations however, the corresponding $\mathbf{RS}_i$ will have more elements since there will be $\frac{V(V-1)}{2}$ pairwise correlations to monitor. Regarding the window size in extracting the running statistic, $\mathbf{RS}_i$, we recommend to use $w=25$ as our previous studies [1-2] showed that in comparison to larger window sizes, this leads to more power in detecting a change, as well as less biased change point estimates. We emphasize, though, that prior information can help in making this choice. For instance, one can set the window size equal to the length of the shortest expected event. If no prior information is available as is often the case, employing different window sizes to check the stability of the obtained change points can be helpful.

The variance criterion to optimize change point locations

To locate the change points, KCP optimizes a variance criterion based on the intra-phase scatter,

$$\hat{V}_{\tau_1, \tau_2, \dots, \tau_K, m} = (\tau_m - \tau_{m-1}) - \frac{1}{\tau_m - \tau_{m-1}} \sum_{i=\tau_{m-1}+1}^{\tau_m} \sum_{j=\tau_{m-1}+1}^{\tau_m} Gk(\mathbf{RS}_i, \mathbf{RS}_j), \quad (2)$$

where the indices, $\tau_1, \tau_2, \dots, \tau_K$, are the phase boundaries, m is the current phase number, and τ_{m-1} and τ_m are the first and last time points of this phase. The more similar the values of the running statistic are inside a phase, the lower is the intra-phase scatter, as $\hat{V}_m$'s rightmost term will become more negative due to the high similarities. If the user requests to locate K change points, the variance criterion,

$$\hat{R}(\tau_1, \tau_2, \dots, \tau_K) = \frac{1}{W} \sum_{m=1}^{K+1} \hat{V}_{\tau_1, \tau_2, \dots, \tau_K, m}. \quad (3)$$

which is simply the sum of all $K+1$ intra-phase scatters divided by the total number of windows, is minimized to ensure optimal homogeneity (i.e., high similarity) of the running statistic within a phase.

Let's consider the simplest case where only one change point ($K=1$) has to be estimated. Two intra-phase scatters will be optimized: $\hat{V}_{\tau_1, 1}$ and $\hat{V}_{\tau_1, 2}$. The phase containing the first observations until τ_1 will determine $\hat{V}_{\tau_1, 1}$, and the phase comprising all remaining observations will determine $\hat{V}_{\tau_1, 2}$. The goal therefore is to search among all $\tau_1 \in \{2, 3, \dots, w-1\}$ the location $\hat{\tau}_1$ that yields the lowest value of the variance criterion, $\hat{R}(\tau_1) = \frac{1}{w} (\hat{V}_{\tau_1, 1} + \hat{V}_{\tau_1, 2})$, and to optimally segment the time series into phases within which the values of the running statistic are very similar. The change point is then

set at $\hat{\tau}_1 + 1$. Generalizing the procedure, the optimal K change points, $\hat{\tau}_1 + 1, \hat{\tau}_2 + 1, \dots, \hat{\tau}_K + 1$, are obtained by finding

$$\hat{\tau}_1, \hat{\tau}_2, \dots, \hat{\tau}_K = \arg \min \hat{R}(\tau_1, \tau_2, \dots, \tau_K). \quad (4)$$

For ease of notation, in the following steps, we will denote the optimized variance criterion for a given K as $\hat{R}_{min,K}$.

Step 2. Selecting the number of change points.

In practice, users often do not know the expected number of change points and should infer this from the data itself. We therefore employ a two-step procedure to choose the optimal number of change points.

Step 2.1 Testing if there is at least one significant change point.

First, a significance test is performed to decide on the presence of at least one change point in the running statistic. The test compares the variance criterion $\hat{R}_{min,K}$ of the running statistic computed from the original data to that of the running statistic from the permuted data (in which time points are reshuffled). Two subtests are implemented: the variance and the variance drop test.

Variance test. The idea of the variance test is that when the running statistic contains at least one change point, their overall variance will be large. This is not expected if the running statistic is obtained from permuted data, as possible changes across time are effectively removed via reshuffling. The variance test therefore compares the overall variance (i.e., $\hat{R}_{min,K=0}$, the variance criterion value when no change point is extracted from the running statistic) to the distribution of $\hat{R}_{min,K=0,perm}$, which is obtained by computing the same overall variance from a large number of permuted versions of the data. The p-value of the variance test is therefore given by,

$$p_{variance\ test} = \frac{\#(\hat{R}_{min,K=0,perm} > \hat{R}_{min,K=0})}{B} \quad (5)$$

where B denotes the number of permutations carried out. In this paper, B was always set to 1000.

We note that this test is not applied to the running autocorrelation as it leads to an inflated Type 1 error rate.

Variance drop test. The variance drop test looks at the extent to which the variance criterion improves due to extracting change points. Recall that the variance criterion, $\hat{R}_{min,K}$, improves (i.e.,

decreases) when more change points are allowed for. However, if there are no true change points present in the data, this improvement will not be substantial and will be comparable to that of permuted data. Thus, in the variance drop test, we compare the drop in the variance criterion, $\hat{R}_{min,K} - \hat{R}_{min,K-1}$ to a distribution of variance drops obtained after reshuffling the data. We only use the strongest evidence for change, by retaining the maximum $\hat{R}_{min,K} - \hat{R}_{min,K-1}$ among all considered K 's. The variance drop test therefore yields a p-value equal to

$$p_{variance\ drop\ test} = \frac{\#(\max variance\ drop_{perm} > \max variance\ drop)}{B} \quad (6)$$

where, B , again indicates the number of reshuffled data sets on which the permuted distribution is based.

Combining the sub-tests. Since two sub-tests are carried out, we correct for multiple testing by adopting a corrected significance level of $\frac{\alpha}{2}$ for each test. If at least one of the sub-tests yield a significant result, we declare that there is at least one change point in the running statistics. For the running autocorrelation, we only employ the variance drop test, and thus, in this case, its significance level is set to α .

Step 2.2 If there is at least one change point ($K > 0$), choose the optimal number of K via penalization.

If the permutation test above yields a significant result, we employ the penalty term $pen_K = C \frac{V_{max}(K+1)}{w} \left[1 + \log \left(\frac{w}{K+1} \right) \right]$ proposed by Arlot et al. [3], to aid in choosing a K that balances fit, as evidenced by a reduction in the variance criterion, $\hat{R}_{min,K}$, and complexity, by penalizing too large K -values that would lead to unnecessary change points. The idea is to pick the number of change points that minimizes the sum of the variance criterion and the penalty term:

$$\hat{K} = \arg \min \hat{R}_{min,K} + pen_K. \quad (7)$$

The penalty coefficient C can be tuned by the user to adjust the impact of the penalty term. Small C -values imply a weak penalty for extra change points in the solution and will therefore tend to favor higher K -values (many change points). Large C -values, on the other hand, more strongly penalize adding more change points and therefore would favor small K -values (less change points). The remaining term, V_{max} , is obtained by computing the trace of the empirical covariance matrix of the first and last 5% elements of the running statistic, retaining whichever is larger among the two.

Since the choice of the penalty coefficient, C , highly influences the result of the penalization scheme, Cabrieto et al.[4] proposed to tune this coefficient through a grid search so that the choice will not solely depend on a single choice for C . The procedure starts by first setting C equal to 1, which will always yield K_{max} (the largest K) as a solution because the penalty is not strong. Moving on, we increase C by incrementing it linearly. This step increases the influence of the penalty term, and thus lower K -values are eventually favored. The grid search terminates once C becomes too large such that the K -value returned is zero. The whole procedure, at the end, outputs the most stable K , that is, the most frequently returned K -value across all considered C -values.

The final KCP change point solution then consists of the change point locations that correspond to the most stable K .

II. Multiple testing correction for monitoring several statistics

When multiple statistics are tracked, as is done in this paper, a multiple testing correction can be used in Step 2.1. Since we monitored three running statistics (i.e., running variance, running correlation, and running autocorrelation) in three separate KCP-analyses, we maintained the overall type 1 error rate of .05 by allotting a corrected α -level of .017 (i.e., .05 divided by 3) to each analysis and thus each running statistic. This implies that for the running autocorrelation, where only the variance drop test is employed, the corrected α -level is .017. For the running variance and running correlation where two sub-tests are employed, the α -level for each sub-test is corrected such that the per statistic α -level (.017) is maintained. This means that each subtest will have an alpha level of .008 (i.e., per statistic α -level divided by 2). In Table 1, we present this multiple testing plan and the corresponding KCP permutation test results for the depression data.

We remark that using such a Bonferroni correction is a conservative way of controlling for multiple testing. Less conservative methods have been proposed, for instance a non-parametric combination of dependent permutation tests [5]. However, to the best of our knowledge, this combined test is an Omnibus test, postulating the global null hypothesis that none of the running statistics contains a change point and the corresponding alternative hypothesis that at least one running statistic shows at least one change point. In contrast, we are interested in local effects, in that we aim to test for each running statistic separately whether it contains at least one change point. Nevertheless, exploring the potential of a non-parametric combined test for increasing the power of KCP-analyses, might be a useful direction for future research.

Table 1. Corrected α -levels and observed p-values for the KCP permutation tests employed to determine whether the 3 running statistics (i.e., running autocorrelation, running variance and running correlation) contain at least one change point.

Data	Running Statistic	Tests	Corrected α -level	p-value
Full	Running autocorrelation*	Variance	.017	.01
		Variance drop	.008	0
	Running variance*	Variance	.008	.016
		Variance drop	.008	0
		Variance drop	.008	0
Before Relapse	Running autocorrelation	Variance	.017	.221
		Variance drop	.008	0
	Running variance*	Variance	.008	0
		Variance drop	.008	0
		Variance drop	.008	.148

* Contains at least one significant change point

References

1. Cabrieto J, Tuerlinckx F, Kuppens P, Borbála H, Ceulemans E. Testing for the presence of correlation changes in a multivariate time series: A permutation based approach. *Sci Rep* 2018 Jan;8(1):769.
2. Cabrieto J, Tuerlinckx F, Kuppens P, Wilhelm FH, Liedlgruber M, Ceulemans E. Capturing correlation changes by applying kernel change point detection on the running correlations. *Information Sciences* 2018 Jun;447:117-139.
3. Arlot S, Celisse A, Harchaoui Z. Kernel change-point detection. <https://arxiv.org/pdf/1202.3878.pdf>, 2012.
4. Cabrieto J, Tuerlinckx F, Kuppens P, Grassmann M, Ceulemans E. Detecting correlation changes in multivariate time series: A comparison of four non-parametric change point detection methods. *Behav Res Methods* 2017 Jun;49(2):988-1005.
5. Pesarin F, Salmaso S. *Permutation Tests for Complex Data: Theory, Applications and Software*. Chichester, UK: Wiley & Sons; 2010. 450 p.