Abstract

Objective: Media stories on how to cope with suicidal crises have been shown to reduce suicidal ideation, but studies investigating if effects differ depending on delivery by individuals with or without personal experience of suicidality are lacking. The present study aimed to examine effects of newspaper articles featuring interviews with experts with vs. without personal experience of suicidal ideation.

Method: A total of 527 adults were randomized to read a newspaper article featuring an interview with a suicide expert disclosing personal experience of suicidal ideation, the same article without disclosure of personal experience, or an article unrelated to suicide in a web-based randomized controlled trial. The primary outcome was suicidal ideation; the secondary outcome was suicide-prevention-related knowledge. Data were collected using online questionnaires before and after exposure.

Results: Participants in the two intervention groups reported a decrease in suicidal ideation (Group #1: $P < .001$, $d = -0.16$, 95% CI -0.25 to -0.07; Group #2: $P < .001$, $d = -0.25$, 95% CI -0.33 to -0.16) and an increase in suicide-prevention-related knowledge (Group #1: $P < .001$, $d = 0.72$, 95% CI 0.64 to 0.81; Group #2: $P < .001$, $d = 0.70$, 95% CI 0.62 to 0.79) after article exposure. There were no differences between the two intervention groups.

Conclusions: Newspaper articles educating about how to cope with suicidality which are delivered by experts with or without personal experience of suicidality appear to have similar suicide-protective effects.

Clinical Trial Registration: American Economic Association's RCT registry, www.socialscienceregistry.org, identifier: RCT-ID: 0002100.

Keywords: Suicide, media, Papageno effect, personal experience of suicidal ideation, randomized controlled trial

Suicide Education Delivered by Individuals with vs. without Personal Experience of Suicidal Ideation: Randomized Controlled Trial

The way how mass media portray suicide influences suicidal behaviour in the population.^{1,2} According to the Werther effect,³ sensationalist suicide reports in newspapers can trigger so-called imitational suicides.^{2,4} In contrast, the potential of some media portrayals highlighting how to cope with suicidal ideation to reduce suicidal behaviour has been labelled Papageno effect.^{5,6} In the first study on this phenomenon, Niederkrotenthaler and colleagues⁵ found a decrease of suicides in Austria following newspaper reports focusing on constructive ways of coping with adverse circumstances. Recent randomized controlled trials found further support for this effect, demonstrating a reduction of suicidal ideation or an increase in protective factors after exposure to newspaper articles,⁷ fictional films,⁸ or educative websites⁹ featuring individual accounts of how to cope with suicidality⁷⁻⁹ and general educative information on suicide prevention.⁹ These effects appeared most pronounced among individuals with some degree of vulnerability to suicide, as indicated by their baseline suicidal ideation scores.^{8,9} So far, it remains unknown if the beneficial effect of such stories depend upon the status of the featured character as having personal experience with coping with suicidal ideation versus not having or disclosing such experience. In recent years, individuals with personal experience of suicidal ideation have become an increasingly acknowledged and organized proponents in suicide prevention, and involvement of suicide attempt survivors and other individuals with personal experience of suicidal ideation or loss to suicide has been described as “the way forward” for suicide prevention.¹⁰ Consistent with this development, the media increasingly cover individuals with personal experience of suicidal ideation who share their stories in order to help prevent suicide.^{11,12} So far, however, it remains unknown whether stories featuring individuals with and without personal experience of suicidal ideation differ with regard to their impact on the audience.

To bridge this research gap, we conducted a web-based randomized controlled trial to examine the effects of suicide-prevention-related newspaper reports compared to a control group. The reports featured interviews with suicide experts who differed with regard to their disclosure of personal experience with suicidal ideation. We evaluated the articles' impact on suicidal ideation (the primary outcome) and suicide-prevention-related knowledge (secondary outcome).

Hypotheses

We hypothesized that both suicide prevention articles will reduce suicidal ideation and increase suicide-related knowledge among audiences compared to the control group (hypotheses 1 and 2). Based on previous findings,^{8,9} we also predicted that the reduction in suicidal ideation will be most pronounced among individuals with higher baseline suicidal ideation (hypothesis 3). Because no previous study has explored the differential influence of messages by an individual with or without personal experience of suicidal ideation, we investigated whether there are differences between the two intervention groups (research question 1).

Method

Participants

A web-based double-blinded randomized controlled trial was conducted in April and May 2017. We recruited 656 German-speaking individuals aged 18 or above who had registered on SoSci Survey—a non-commercial online panel¹³—via e-mail. In total, 527 individuals (80.3%) were included in the final analysis. Of the included 479 individuals who completed measures on socio-demographics, 272 were women (56.8%) and 207 men (43.2%), and mean age was 41.3 ($SD = 14.9$). Among them, 82 participants (17.15%) indicated that they had no high school diploma, 137 (28.6%) reported high school as highest completed level of education, and 260 (54.3%) indicated having a college degree.

Power analysis

Based on the results of a power analysis with G*Power 3.1.9.2,¹⁴ an ANOVA model with two repeated measures, six groups (three study groups, low vs. high baseline suicidal ideation), and an assumed correlation of .79 among repeated measures^{8,9} would require a total sample size of 270 participants to detect an effect of .21 found in related previous studies.^{8,9} Based on a less conservative sample size calculation with GLIMMPSE 2.2.4¹⁵ it is estimated that an equivalent linear mixed model would require a minimum sample size of 168 participants.

Materials and Procedure

We conducted a web-based randomized controlled trial with two intervention groups and one control group. Participants were informed that they would participate in a study on the impact of health-related awareness material without further specification. At the beginning of the trial, we measured participants' suicidal ideation and suicide-prevention-related knowledge. After completing the two measures, the participants were automatically randomly allocated to one of three groups based on an allocation ratio of 1:1:1 to achieve equal group sizes.

Participants of Group #1 were exposed to a newspaper article featuring an interview with a suicide expert on suicide prevention. In this interview, the expert states that suicidal ideation affects many people in society, offers advice on how to cope with suicidal thoughts, explains how families and friends can support individuals in crises, and debunks the most common misconceptions about suicide. The article consisted of two pages, was headlined *Coping with suicidal crises*, and included two pull quotes ("Suicide announcements should always be taken seriously", "If you suspect someone may be thinking of suicide, it is important to ask them directly about it") that highlighted the interview's key messages. Participants of Group #2 were exposed to the same newspaper article, however in this version the expert reported personal experience when asked about ever having a suicidal crises herself. She tells a story about coping with a suicidal crisis as a young adult by contacting a

crisis-intervention centre. This version of the article also had one additional pull quote saying “One evening I wanted to take my own life. I can still hear the Samaritan’s ‘Don’t do it’ in my ear”. Participants of the control group read a newspaper article of similar layout and style featuring an expert interview unrelated to suicide or mental health. In this article, the expert talked about prevention strategies for seasonal influenza.

After reading the articles, we measured suicidal ideation and suicide-prevention-related knowledge again along socio-demographics. We also administered one item to assess whether or not participants were currently having suicidal thoughts, two items for manipulation check, and one item to assess blinding success. On the last page of the online survey, we provided contact information of organizations providing help for individuals in crisis. Participants who discontinued their participation before randomization was completed ($n = 111$) or took less than 30 seconds to read their respective article ($n = 18$) were excluded from the study. Thus, a total of 527 participants were included in the final statistical analysis (Group #1: $n = 173$; Group #2: $n = 174$; Control group: $n = 180$). Figure 1 shows the study flowchart. There were no differences in suicidal ideation ($t(18.54) = -0.09, P = .93$) or suicide-prevention-related knowledge ($t(604) = -1.91, P = .06$) before reading the articles between those excluded and included in the final statistical analyses.

The study was approved by the Research Ethics Board at the Medical University of Vienna (study protocol 1285/2017, date: 2017-04-25). The trial was registered with the American Economic Association’s registry for randomized controlled trials (www.socialscienceregistry.org) as RCT-ID: 0002100.

Primary outcome measure

RFL Survival Coping Beliefs subscale. Suicidal ideation was assessed with the 23-item Survival Coping Beliefs subscale of the Reasons for Living Inventory¹⁶ (e.g., “I am afraid of the actual ‘act’ of killing myself”). This subscale has been shown to be particularly sensitive to media-induced changes of suicidal ideation.⁹ Respondents rated their beliefs and

expectations for *not* dying by suicide on a scale ranging from 1 (*not at all important*) to 6 (*extremely important*). Scale scores were reverse-coded, resulting in higher scores indicating higher suicidal ideation.

Secondary outcome measure

Questionnaire on Suicide-Prevention-Related Knowledge. To determine participants' suicide-prevention-related knowledge, we compiled a collection of 10 statements related to suicide prevention adopted from previous questionnaires.¹⁷⁻²⁰ All items were answerable based on the information provided in the articles of the intervention groups. We administered the statements (e.g., "Most suicidal individuals do not want any help") to the participants and asked them to rate the accuracy of the respective statement (*true*, *false*, or *don't know*). We coded whether participants correctly (coded as 1) or incorrectly (coded as 0) rated each of the statements as true or false. Respondents received one point for each correct answer. Higher scores indicate more suicide-prevention-related-knowledge.

Additional measures

Current suicidal thoughts. Respondents were asked to indicate whether or not they currently have suicidal thoughts with one single self-report item (*yes* = 1, *no* = 0).

Manipulation Check. In order to assess if the experimental manipulation was successful, we asked the respondents to indicate whether or not the expert in the respective article (1) was an expert on suicide prevention (*true*, *false*, or *don't know*) and (2) has ever experienced a suicidal crisis in her life (*true*, *false*, or *don't know*). The manipulation check was performed after the measurement of the repeated measurement of the main outcomes of interest to the study.

Blinding success. In order to assess blinding success, we asked respondents to indicate what group they think they have been allocated to (*intervention group*, *control group*, or *don't know*) at the very end of the study, before the debriefing.^{21,22}

Data analysis

For each measure, negative items were reversed and mean scores across all items of the respective questionnaire were calculated for each participant. In order to test hypothesis 3, it was necessary to compare the impact of article exposure on participants with lower vs. higher baseline suicidal ideation. Since our measure of suicidal ideation does not have a validated cut-off score,¹⁶ we applied a median split of the sample, using the score for suicidal ideation ($Md = 2.30$) observed prior to article exposure, and stratified the sample into two groups: one group including participants with lower baseline suicidal ideation (i.e., suicidal ideation scores < 2.30) ($n = 263$, $Md = 1.70$, $IQR = 0.61$, $Min = 1.00$, $Max = 2.26$) and the other group including those with higher baseline suicidal ideation (i.e., suicidal ideation scores ≥ 2.30) ($n = 264$, $Md = 2.89$, $IQR = 0.68$, $Min = 2.30$, $Max = 5.78$). The same approach was also used in previous research.^{8,9}

The scores for the primary outcome variable (i.e., suicidal ideation) were subjected to a study group (Group #1, Group #2, Control group) $\times$ test condition (pre-exposure, post-exposure) $\times$ baseline suicidal ideation (baseline suicidal ideation below vs. above the median) analysis using linear mixed models with individual group differences tested with Bonferroni-corrected contrast tests. Due to their ability to account for correlations between repeated measures of the same subject and missing data, linear mixed models have been used increasingly in recent years replacing traditional repeated measures ANOVA models in psychiatric research.^{9,23} The scores for the secondary outcome variable, suicide-prevention-related knowledge, were subjected to a study group (Group #1, Group #2, Control group) $\times$ test condition (pre-exposure, post-exposure) analysis using linear mixed models along with Bonferroni-corrected contrast tests.

Results

Of the 480 participants who completed the items to assess blinding success and for manipulation check, 133 participants (27.2%) correctly guessed their group allocation, whereas 127 participants (26.5%) were incorrect and 220 (45.8%) were unsure. We can

assume that blinding was successful because the number of unsure participants was relatively high,²¹ correct and incorrect guesses were balanced,²¹ and the distribution was similar to other published randomized controlled trials.²² $N = 404$ participants (76.7%) answered the two manipulation check items correctly, and there were no differences between the three study groups in terms of successful manipulation check ($\chi^2 = 4.23$, $df = 2$, $P = .12$). See Table 1 for an overview of the demographic characteristics of participants in each group. There were no significant differences in terms of the participants' gender, age, highest completed education, or baseline suicidal ideation across the three groups, as indicated by χ^2 and ANOVA tests. Table 2 displays the means and corresponding 95% confidence intervals for the two outcome variables before and after article exposure. The mean values and their corresponding 95% confidence intervals for suicidal ideation before and after article exposure stratified by baseline suicidal ideation below and above the sample median are provided in Table 3.

Suicidal ideation

The analysis of the impact of the newspaper articles revealed a significant study group $\times$ test condition interaction for the primary outcome, suicidal ideation. The contrast tests indicated that participants exposed to one of the two newspaper articles on suicide prevention experienced a small reduction of suicidal ideation after the exposure (Group #1: contrast test after vs. before exposure: Bonferroni-corrected $P < .001$, $d = -0.16$, 95% CI -0.25 to -0.07; Group #2: contrast test after vs. before exposure: Bonferroni-corrected $P < .001$, $d = -0.25$, 95% CI -0.33 to -0.16), whereas suicidal ideation did not change in the control group (contrast test after vs. before exposure: Bonferroni-corrected $P = .62$, $d = -0.02$, 95% CI -0.11 to 0.07). A difference between the two intervention groups was not observed (contrast test Group #2 vs. Group #1: Bonferroni-corrected $P = 1.00$, $d = 0.03$, 95% CI -0.07 to 0.14). There was no significant interaction with baseline suicidal ideation, and the same effects emerged when baseline suicidal ideation was not entered into the statistical model. Table 4 gives an overview

of the results of the linear mixed models. Based on these results, hypothesis 1 was supported, whereas hypothesis 3 was rejected.

Suicide-prevention-related knowledge

There was a significant study group $\times$ test condition interaction in terms of the secondary outcome of suicide-prevention-related knowledge. Participants in the two intervention groups reported a large increase of suicide-prevention-related knowledge after reading the article (Group #1: contrast test after vs. before exposure: Bonferroni-corrected $P < .001$, $d = 0.72$, 95% CI 0.64 to 0.81; Group #2: contrast test after vs. before exposure: Bonferroni-corrected $P < .001$, $d = 0.70$, 95% CI 0.62 to 0.79), whereas a small increase of suicide-prevention-related knowledge was found in the control group (contrast test after vs. before exposure: Bonferroni-corrected $P < .05$, $d = 0.09$, 95% CI 0.01 to 0.18). This increase was significantly stronger in the two intervention groups than in the control group (contrast test control group vs. Group #1: Bonferroni-corrected $P < .001$, $d = -0.23$, 95% CI -0.34 to -0.12; contrast test control group vs. Group #2: Bonferroni-corrected $P < .001$, $d = -0.30$, 95% CI -0.40 to -0.19), and there was no difference observed between the two intervention groups (contrast test Group #2 vs. Group #1: Bonferroni-corrected $P = .41$, $d = 0.07$, 95% CI -0.04 to 0.17). Based on these results, hypothesis 2 was supported.

Impact on audience members with current suicidal thoughts

Of the 503 participants who completed the single item assessing current suicidal thoughts, only 11 (2.2%) indicated that they were currently experiencing suicidal thoughts (Group #1: $n = 3$, Group #2: $n = 5$, control group: $n = 3$). Due to the low frequencies, we were unable to include this variable into inferential statistical analyses. The patterns were generally consistent with the findings of the total sample (not shown).

Discussion

This study indicates that educative newspaper articles featuring suicide experts had a beneficial impact on readers in terms of a reduction in suicidal ideation and increase in

suicide-prevention-related knowledge. The effect was present independent of the personal overtly-expressed experience status of the featured suicide expert. Our findings are consistent with previous research on the Papageno effect, which describes a suicide-protective effect of media stories of how to cope with adverse circumstances.⁵⁻⁹ Consistent with this effect, previous studies have shown that media stories focusing on how to cope with suicidal ideation have the potential to reduce suicidal ideation and increase suicide-prevention-related knowledge.⁷⁻⁹

In the current study, the expert interviews featured in the newspaper articles may have empowered participants to modify their perceptions regarding their ability to cope constructively with suicidal ideation, which has also been found previously with educative websites for suicide prevention.⁹ The present findings also extend previous research on the Papageno effect. Evidence to date suggests that the suicide-protective effect is present for individual narrations on coping with adverse circumstances,⁵⁻⁸ as well as for educative narratives including both perspectives of personal experience and other expert insight.⁹ The results of the present study suggest that the effect is present on a general level for media content on how to cope with suicidal crises, independent of the personal experience status of the featured character. The general focus of such stories on what can be done to cope with suicidal ideation may be the key prerequisite of a suicide-protective effect of these newspaper stories. Further studies are warranted to investigate if particular story elements have different effects in different subpopulations.

While previous studies have found a media-induced reduction in suicide risk predominantly among vulnerable individuals,^{8,9} baseline suicidal ideation did not moderate the article effects in the current study. This may be due to differences in sample characteristics (the current study sample was older, more educated, and slightly less suicidal than in previous studies^{8,9}). The present study also had a considerably larger sample size as compared to previous works in the area.^{8,9}

Limitations

With the study sample being recruited from the general population and not from a clinical sample, it remains unknown whether and how our findings generalize to individuals with suicide risk. The current sample, included only 11 participants reporting current suicidal thoughts, and the impact of the articles in this particular subsample appeared to be consistent with the findings in the overall sample. However, more studies that investigate specifically the effects on individuals at risk of suicide as a function of the personal experience of a featured character are warranted. A further limitation of the study was the use of a convenience sample not representative of the population in terms of age and education. Finally, the Cronbach α estimating the reliability of the items measuring suicide-prevention related knowledge was relatively low (see Table 1), which is a known psychometric limitation of comparable scales in this field of research.^{9,20} With $\alpha = .62$, however, the reliability of the current measure was considerably higher than comparable scales in previous studies.^{9,20}

Implications

Interviews with suicide experts debunking suicide myths and highlighting that suicidal ideation affects many people in society and does not necessarily mean that the person would eventually die from suicide and emphasizing what can be done specifically to cope if experiencing suicidal ideation appear to be effective in educating the public and reducing suicidal ideation. Health professionals with and without personal experience of suicidal ideation appear to be similarly effective when targeting primarily non-suicidal individuals such as those included in the present study. The present study highlights the relevance of national and international recommendations to include expert interviews in suicide-related reporting.^{2,24,25} Previous studies have shown, however, that expert interviews are sometimes used in sensationalist contexts that are not consistent with media recommendations,⁵ we recommend that experts should always ask journalists and editors to consider current media

recommendations for suicide reporting,² explain their practical importance, and also offer to read a draft of the article before publication.

References

1. Niederkrotenthaler T, Reidenberg D, Till B, et al. Increasing help-seeking and referrals for individuals at risk for suicide by decreasing stigma: The role of mass media. *Am J Prev Med.* 2014;47:235-243.
2. World Health Organization. *Preventing suicide. A resource for media professionals: Update 2017.* Geneva, Switzerland: WHO; 2017.
3. Phillips DP. The influence of suggestion on suicide: substantive and theoretical implications of the Werther effect. *Am Sociol Rev.* 1974;39:340–354.
4. Niederkrotenthaler T, Till B, Kapusta ND, et al. Copycat effects after media reports on suicide: A population-based ecologic study. *Soc Sci Med.* 2009;69:1085-1090.
5. Niederkrotenthaler T, Voracek M, Herberth A, et al. The role of media reports in completed and prevented suicide – Werther versus Papageno effects. *Br J Psychiatry.* 2010;197:234-243.
6. Niederkrotenthaler T, Voracek M, Herberth A, et al. Papageno v Werther effect. *BMJ.* 2010;341:5841.
7. Arendt F, Till B, Niederkrotenthaler T. Effects of suicide awareness material on implicit suicide cognition: A laboratory experiment. *Health Commun.* 2016;31:18-26.
8. Till B, Strauss M, Sonneck G, et al. Determining the effects of films with suicidal content: a laboratory experiment. *Br J Psychiatry.* 2015;207:72-78.
9. Till B, Tran US, Voracek M, et al. Papageno vs. Werther Effect online: Randomized controlled trial of beneficial and harmful impacts of educative suicide prevention websites. *Br J Psychiatry.* 2017;211:109-115.
10. National Action Alliance for Suicide Prevention: Suicide Attempt Survivors Task Force. *The way forward: Pathways to hope, recovery, and wellness with insights from lived experiences.* Washington, DC: Author; 2014.

11. Hines K. After my suicide attempt, I made this plan to stay alive and well. Huffington Post. www.huffingtonpost.com/kevin-hines/after-my-suicide-attempt-i-made-this-plan-to-stay-aliveand-well_b_9058776.html. Accessed September 29, 2017.
12. Substance and Mental Health Service Administration (SAMHSA). Stories of hope and recovery: A video guide for suicide attempt survivors. https://www.youtube.com/playlist?list=PLBXgZMI_zqfR4dvBdX7XHD-fjgoehFM_9&feature=plcp. Accessed June 24, 2014.
13. Leiner D. *SoSci Panel: The noncommercial Online Access Panel*. Poster presented at the GOR 2012, 6th March, Mannheim, Germany. <https://www.soscisurvey.de/panel/download/SoSciPanel.GOR2012.pdf>. Accessed September 29, 2017.
14. Faul F, Erdfelder E, Lang AG, et al. G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. 2007;39:175-191.
15. Kreidler SM, Barón AE, Glueck DH. *GLIMMPSE Tutorial: Selecting a test*. Denver, CO: University of Colorado; 2012.
16. Linehan MM, Goodstein JL, Nielsen SL, et al. Reasons for staying alive when you are thinking of killing yourself: The Reasons for Living Inventory. *J Consult Clin Psychol*. 1983;51:276-286.
17. Domino G. Popular misconceptions about suicide: How popular are they? *Omega-J Death Dying*. 1990;21:167-175.
18. Schurtz DR, Cerel J, Rodgers P. Myths and facts about suicide from individuals involved in suicide prevention. *Suicide Life Threat Behav*. 2010;40:346-352.
19. Sonneck G, Kapusta N, Tomandl G, et al. (Eds.) *Krisenintervention und Suizidverhütung*. [Crisis intervention and suicide prevention.] Vienna, Austria: Facultas; 2016.

20. Voracek M, Tran US, Sonneck G. Psychometric properties of the Revised Facts On Suicide Quiz in Austrian medical and psychology undergraduates. *Death Stud.* 2008;32:937-950.
21. Kolahi J, Bang H, Park J. Towards a proposal for assessment of blinding success in clinical trials: up-to-date review. *Community Dent Oral Epidemiol.* 2009;37:477-484.
22. Lao L, Bergman S, Hamilton GR, et al. Evaluation of acupuncture for pain control after oral surgery: A placebo-controlled trial. *Arch Otolaryngol Head Neck Surg.* 1999;125:567-572.
23. Gueorguieva R, Krystal JH. Move over ANOVA. Progress in analyzing repeated-measures data and its reflection in papers published in the Archives of General Psychiatry. *Arch Gen Psychiatry.* 2004;61:310-317.
24. U.S. Department of Health and Human Services. *National strategy for suicide prevention: Goals and objectives for action.* Rockville, MD: Author; 2001.
25. World Health Organization. *The World health report 2001. Mental Health: New understanding, new hope.* Geneva, Switzerland: WHO; 2001.

Figure 1

Study flowchart *Figure 1. Study flowchart.*

Table 1

Descriptive Demographic Statistics (Sex, Age, and Highest Completed Education) and Baseline Suicidal Ideation across All Three Groups

Variable	Group #1: No personal experience	Group #2: Per- sonal experience	Control group	χ^2/F
Females n (%)	91 (58.3%)	84 (54.2%)	97 (57.7%)	0.64 ¹
Age M (SD)	41.4 (15.3)	41.2 (14.3)	41.3 (15.1)	0.00 ²
College n (%)	85 (54.5%)	89 (57.4%)	86 (50.9%)	1.40 ¹
High school n (%)	43 (27.6%)	41 (26.5%)	54 (32.0%)	1.35 ¹
Below high school n (%)	28 (17.9%)	25 (16.1%)	29 (17.2%)	0.18 ¹
Baseline suicidal ideation M (SD)	2.36 (0.9)	2.32 (0.8)	2.39 (0.8)	0.35 ³

Note. Values are frequencies (n), percentages (%), means (M), and standard deviations (SD) provided for each group, as well as χ^2 values from χ^2 tests and F values from analyses of variance (ANOVAs) testing group differences.

¹ χ^2 test result, $df = 2$

²ANOVA result, $df_1 = 2$, $df_2 = 477$

³ANOVA result, $df_1 = 2$, $df_2 = 524$

Table 2

Study Variables in Audience Members Before and After Article Exposure

Study variable	Group	Before	After
Primary outcome: Suicidal ideation			
Suicidal ideation ($\alpha = .93$)	#1: No personal	2.33	2.24
	experience	(2.19, 2.48)	(2.09, 2.40)
	#2: Personal experience	2.36	2.22
		(2.23, 2.48)	(2.09, 2.34)
	Control	2.39	2.38
		(2.26, 2.52)	(2.23, 2.52)
Secondary outcome: Suicide-Prevention-Related Knowledge			
Suicide-prevention-related knowledge ($\alpha = .62$)	#1: No personal	0.59	0.84
	experience	(0.55, 0.63)	(0.81, 0.87)
	#2: Personal experience	0.62	0.86
		(0.58, 0.65)	(0.84, 0.88)
	Control	0.59	0.62
		(0.55, 0.62)	(0.58, 0.65)

Note. Values are means with 95% confidence intervals given in parentheses for both outcome variables before and after article exposure as well as lower-bound (Cronbach α) sample reliabilities.

Table 3

Suicidal Ideation in Audience Members Before (T1) and After (T2) Article Exposure Stratified for Baseline Suicidal Ideation Below and Above Sample Median

Study variable	Group	Baseline Suicidal Ideation Below the Median		Baseline Suicidal Ideation Above the Median	
		T1	T2	T1	T2
Suicidal ideation	G1	1.64	1.50	3.09	3.04
		(1.55, 1.72)	(1.41, 1.58)	(2.94, 3.24)	(2.87, 3.22)
	G2	1.74	1.63	2.98	2.81
		(1.66, 1.81)	(1.54, 1.71)	(2.86, 3.11)	(2.66, 2.96)
	CG	1.69	1.68	3.04	3.02
		(1.60, 1.77)	(1.57, 1.79)	(2.90, 3.17)	(2.85, 3.18)

Note. Values are means with 95% confidence intervals given in parentheses for suicidal ideation before (T1) and after article exposure (T2).

G1 = Group #1: No personal experience, G2 = Group #2: Personal experience, CG = Control group

Table 4

Findings from Linear Mixed Models for Suicidal Ideation and Suicide-Prevention-Related Knowledge

Study Variable		Group	Time	Baseline Suicidal Ideation	Group × Time	Group × Baseline Suicidal Ideation	Baseline Suicidal Ideation × Time	Group × Baseline Suicidal Ideation × Time
Primary outcome								
Suicidal ideation	<i>F</i> (<i>df</i> ₁ , <i>df</i> ₂)	0.81 (2, 520.87)	31.27 (1, 474.03)	849.99 (1, 520.89)	6.71 (2, 474.03)	2.59 (2, 520.87)	0.04 (1, 474.03)	2.35 (2, 474.03)
	<i>P</i>	.446	<.001	<.001	.001	.076	.836	.096
Secondary outcome								
Suicide-prevention- related knowledge	<i>F</i> (<i>df</i> ₁ , <i>df</i> ₂)	25.09 (2, 524.43)	418.97 (1, 515.96)	---	72.67 (2, 515.91)	---	---	---
	<i>P</i>	<.001	<.001	---	<.001	---	---	---

Note. Values are *F* and *P* values with degrees of freedom (*df*₁, *df*₂) given in parentheses from linear mixed models representing the change of the respective outcome variable with regard to group (intervention vs. control group), time (T1, T2), baseline suicidal ideation (low vs. high suicidal ideation/vulnerability), and interactions between these factors. Significant *p* values (< 0.05) are in bold.