Cross-country validation of the association between oral health and general health in community-dwelling older adults

Keywords: oral health; general health, bivariate autoregressive models, Bayesian models, cross-country validation

de Almeida Mello Johanna, Ms Econ, PhD¹

Trung Dung Tran, MSc²

Stefanie Krausch-Hofmann, MSc³

Brigette Meehan, PhD⁵

Hein van Hout, PhD⁶

Luke Turcotte, MSc⁷

Henriette G. van der Roest, PhD⁶

Vjenka Garms-Homolová, PhD⁸

Pálmi Jónsson, MD, PhD⁹

Graziano Onder, MD, PhD¹⁰

Harriet Finne-Soveri, MD, PhD¹¹

Jan De Lepeleire, MD, PhD⁴

Dominique Declerck, PhD³

Emmanuel Lesaffre, Dr. Sc.²

Joke Duyck, PhD³

Anja Declercq, Ms Econ, Ms Soc, PhD¹,

1 KU Leuven LUCAS - Centre for Care Research and Consultancy, Leuven, Belgium

2 KU Leuven Biostatistics and Statistical Bioinformatics Centre (L-BioStat) - Department of Public Health and Primary Care, Leuven, Belgium

- 3 KU Leuven Population Studies in Oral Health - Department of Oral Health Sciences, Leuven, Belgium
- 4 KU Leuven Academic Centre for General Practice - Department of Public Health and Primary Care, Leuven, Belgium
- 5 Technical Advisory Services (TAS), interRAI New Zealand
- 6 Department of General Practice and Elderly Care Medicine, Amsterdam Public Health research institute, Amsterdam UMC, VU University medical center, Amsterdam, The Netherlands.
- 7 School of Public Health and Health Systems, University of Waterloo, Waterloo, Canada
- 8 Department of Economics and Law, HTW, Berlin University of Applied Sciences, Berlin, Germany
- 9 Faculty of Medicine, University of Iceland and Department of Geriatrics, Landspítali National University Hospital, Reykjavik, Iceland.
- 10 Fondazione Universitaria Policlinico A. Gemelli, IRCCS, Università Cattolica del Sacro Cuore, Italy
- 11 Helsinki City – Department for Social Services and Health Care, Finland

Abstract

Objective: Oral health is known to be associated with general health, but longitudinal relationships between oral health and general health indicators have not yet been fully explored in international research.

Setting and Participants: The sample consisted of three longitudinal databases: a sample from Belgium from the Protocol 3 project (n=8359), a combined sample from six European countries (n=2501) from the iBenC study (Belgium, Finland, Iceland, Germany, Italy and the Netherlands), and a sample from New Zealand (n=15012). All clients were 65 years or older and received long-term home care services.

Methods: Bayesian models were used to analyze the associations between three oral health indicators (chewing difficulty, non-intact teeth and dry mouth) and four aspects of general health (activities of daily living functioning, cognition, depression and health instability). In addition, the models explored the associations between current oral health and general health status and future oral health and general health status.

Results Clients who had poorer oral health had a higher risk of suffering from poor general health. Especially chewing difficulty was associated with all general health indicators in all datasets (odds ratios > 1). Dry mouth and non-intact teeth showed significant associations with almost all general health indicators. Additionally, having poor oral health (respectively general health) was predictive of poor general health (respectively oral health) at future assessments (significant cross-lagged parameters).

Conclusions and Implications The results point out the need of the inclusion of oral health assessment and advice from dentists or oral health practitioners into the multidisciplinary conversation. In addition, identifying older people with oral health problems is essential in order to provide treatment and monitoring. Raising awareness for oral health is important and policy makers should foster oral health promotion and care for older adults in order to keep them in good health.

Cross-country validation of the association between oral health and general health in community-dwelling older adults

Introduction

International scientific literature shows that oral health is associated with general health outcomes, functional performance and well-being¹⁻⁶. Several studies identify poor oral health as a major risk factor of aspiration pneumonia⁷⁻⁹. Reduction in saliva flow, which is often a side effect from medication use, leads to a higher risk of oral infections^{10,11} such as caries or periodontitis, which in turn is associated with increased risk for diabetes and cardiovascular disease^{12,13}. Also impaired chewing function can affect a person's general health and well-being^{14,15}.

A recently published Belgian longitudinal study showed higher risk of poor general health status for older people having poor oral health¹⁶. This study showed evidence of the predictive association of oral health for general health (and vice versa). This second study aims at examining and comparing the association between oral health and general health in three cross-country databases of older people receiving long-term home care services. Another objective is to examine the predictive associations between the current oral health and general health status and their future status.

Methods

Sample selection

Three longitudinal databases were included in the study: 1) a sample from Belgium consisting of participants in the Protocol 3 project with data collection from 2010 until 2016; 2) a combined sample from six European countries participating in the iBenC study (Finland, Iceland, Germany, Italy, the Netherlands and an additional sample from Belgium), with data collected between 2014 and 2016; and 3) a sample from New Zealand, where the interRAI Home Care (HC) instrument was implemented

nationally. Details about the study design and data collection of the three samples can be found in de Almeida Mello et al. (Protocol 3 project in Belgium)¹⁷, Garms-Homolová et al. (IBenC project Europe)¹⁸ and interRAI New Zealand (<http://www.interrai.co.nz>)¹⁹.

All clients were 65 years or older and received long-term home care services. Assessments were performed every six months by registered health or care professionals delivering home care. They filled out the interRAI-HC instrument, an internationally validated comprehensive assessment compiling information on several domains (functional, clinical, psychosocial, environmental and social)^{20,21}. Only complete assessments for all oral health items were selected from the three databases and these assessments were performed at baseline and every six months thereafter (longitudinal evaluation). Missing assessments for oral health were not included, but have been studied previously²². If a client died, or home care services stopped before the six month period, the client's assessment was not included in the longitudinal analysis. On average, clients kept receiving care for 14 months in the Belgian sample, for 12 months in the European sample and for 18 months in the sample from New Zealand.

Ethics approval

The data collection from Belgium was approved by the Belgian Privacy Commission and by the Ethics committees of the KULeuven and Université Catholique de Louvain. The data from iBenC was approved from authorized medical ethical committees according to local country regulations. In New Zealand, national data access protocols ensured appropriate ethical clearance. Written consent was sought from all participants from all countries.

Outcome measures

Three items from the interRAI-HC indicating oral health status were used: non-intact teeth (NT), chewing difficulty (CD), and dry mouth (DM). 'Non-intact teeth' means that the client has broken, fragmented or loose natural teeth. The item about 'dry mouth' indicates that the client does not make enough saliva to make keep the mouth wet or that there is difficulty in moving a food bolus in the mouth due to dryness. The item about chewing difficulty indicates that the client is unable to chew food easily and without pain or difficulties, regardless of cause (e.g., the person uses ill-fitting dentures, or has a neurologically impaired chewing mechanism, temporomandibular joint pain, or a painful tooth). These three items are scored by asking the clients whether these conditions are present, by observations during meals or clients also spontaneously report the problems.

In order to measure general health, four interRAI validated scales were calculated: the Activities of Daily Living Hierarchy scale (ADL scale), the Cognitive Performance Scale (CPS scale with five items), the Depression Rating Scale (DRS), and the Changes in Health, End-Stage Disease, Signs, and Symptoms Scale (CHESS)²³⁻²⁶. The interRAI ADL and CPS scales measure ADL and cognitive function and have a 0-6 range. The interRAI DRS (0-14 range) measures the presence of depressive symptoms. The CHESS (0-5 range) identifies clients at risk of adverse outcomes and with health instability. For all these scales, higher values indicate greater severity. In addition, the following characteristics were included in the analysis as covariates: age, gender, 'Living situation' (alone or not), and 'Having at least one informal caregiver' (yes or no). In the sample from Belgium a covariate was added to account for the type of innovative home care project older people were enrolled in (case management, occupational therapy at home, night care, etc.). In Belgium, older people were receiving home care interventions by nurses, occupational therapists, physiotherapists as well as other professional caregivers. In the European sample and in the sample from New Zealand, clients were receiving nursing care as usual

Analyses

All analyses were performed in the Bayesian framework²⁷. The models were implemented in R (version 3.2.5) using `runjags`²⁸ and `rstan` packages²⁹. Four chains were run and convergence was checked by examining the trace-plots. The proportional odds model was applied to analyze the associations between general health indicators and oral health indicators. In addition, 12 modified bivariate autoregressive models were fit for the pairs between the oral health indicators (3 items) and the general health indicators (4 items).³⁰

The first analysis explored the association between the items from the oral health section of the interRAI-HC and interRAI scales. Associations were expressed as odds ratios. The second analysis examined the predictive association of a given oral health (or general health) status combined with a general health (or oral health) status to predict general health (or oral health) at a time-point in the future.

Results

Table 1 presents descriptive characteristics of the individuals in the samples. The mean age of clients was similar; however, female clients were less frequent in the New Zealand sample. The median scores on the interRAI scales did not differ for ADL. For the other scales, the differences were of one point. Almost half of the clients in the Belgian and in the iBenC project samples showed moderate to severe ADL-impairment while 26% in the New Zealand sample. Cognitive impairment was the highest in the iBenC sample (34.6%) and very similar between the samples from Belgium and New Zealand (about 30%). In the Belgian sample about 28% of the clients showed depressive symptoms, followed by the clients in the iBenC sample and New Zealand. The largest difference emerged from the CHESS. About 28% of the clients in the New Zealand sample showed health instability, while only about 10% in the

samples from Belgium and iBenC project. People receiving full interRAI-HC assessments in New Zealand seem to have more clinical issues than in the European countries. This can be explained by the fact that formal home care provided by nurses is more available in the European countries than in New Zealand. In the European countries, nurses also give all types of ADL help, while in New Zealand this task is mostly provided by home support workers (health aides). Nurses focus on more clinical tasks and on incontinence care. In addition, clients in the New Zealand sample were the least likely to live alone (31%). Clients in the Belgian sample were the least likely to have an informal caregiver (79%). Between 10% and 15% of the clients presented oral health problems and the most common problem was dry mouth.

Table 1 - Descriptive characteristics of the three samples

Median, mean or percentage	Belgium N= 8359	IBenC N=2501	New Zealand N=15012
Age (mean (SD))	81.2 (7.0)	82.7 (7.3)	83.0 (7.4)
Gender (% female)	68.3%	67.9 %	59.4 %
ADLH (0-6) (median) ADLH 3-6 (%)	2 45.70 %	2 44.66 %	2 25.98 %
CPS (0-6) (median) CPS 3-6 (%)	1 29.56 %	1 34.56 %	2 29.98 %
DRS (0-14) (median) DRS 3-14 (%)	1 28.16 %	0 23.50 %	0 16.96 %
CHESS (0-5) (median) CHESS 3-5 (%)	1 9.69 %	1 9.80 %	2 27.96 %
Non-intact teeth (%)	10.9 %	12.1 %	10.3 %
Chewing difficulty (%)	9.3 %	10.3 %	10.0 %
Dry mouth (%)	11.4 %	14.1 %	10.7 %
Living alone (%)	55.1 %	59.1 %	30.8 %
Has at least one informal caregiver (%)	78.6 %	93.4 %	97.0 %

Association between oral health and general health

Table 2 shows the associations (odds ratios - O.R.) from the proportional odds model between oral health status and general health indicators. Most of the significant odds ratios were higher than 1. This means that clients with oral health problems were more likely to suffer from poor general health, compared with older persons without these problems. Among the three oral health items, chewing difficulty was highly associated with all general health items in all three samples. In the sample from Belgium, the O.R. for chewing difficulty and CPS was five times higher than in the other samples. In addition, non-intact teeth was associated with CPS and DRS in the three samples. In the samples from iBenC project and New Zealand, non-intact teeth was also associated with ADL functioning and in the iBenC sample, the O.R. was almost 4 times higher than in the sample from New Zealand. Moreover, non-intact teeth was associated with CHES for Belgium and for the iBenC sample, but not for New Zealand.

Dry mouth was significantly associated with DRS and CHES in all three samples, but this indicator was only associated with ADL impairment in the Belgian dataset. In the other samples, there was no association between dry mouth and ADL function. A significant inverted association between dry mouth and cognitive function (CPS) was found in the samples from iBenC project and New Zealand. An invert association means that people with dry mouth showed less cognitive problems. For Belgium the results pointed in the same direction although the association was not significant.

Table 2 - Odds ratios of ADL, CPS, DRS, and CHES for clients with oral health problems compared to individuals without.

		Odds Ratios (O.R.)			Credible Interval 95% (C.I.)					
		Belgium (N=8359)			IBenC (N=2501)			New Zealand (N=15012)		
NT (yes=1)	ADLH	1.19	0.92	1.56	4.32	2.49	7.45	1.14	1.03	1.26
CD (yes=1)		3.45	2.63	4.55	2.12	1.18	3.76	1.90	1.72	2.09
DM (yes=1)		1.39	1.09	1.80	0.76	0.45	1.29	1.03	0.93	1.14
NT (yes=1)	CPS	2.38	1.63	3.46	2.84	1.67	4.87	1.17	1.02	1.35
CD (yes=1)		10.88	7.35	16.29	2.06	1.17	3.61	1.75	1.53	2.00
DM (yes=1)		0.97	0.68	1.37	0.60	0.37	0.99	0.58	0.51	0.66
NT (yes=1)	DRS	1.71	1.29	2.26	1.62	1.04	2.56	1.15	1.01	1.29
CD (yes=1)		3.70	2.77	5.04	3.60	2.24	5.84	1.69	1.49	1.92
DM (yes=1)		3.71	2.85	4.86	2.54	1.72	3.75	2.46	2.18	2.76
NT (yes=1)	CHES	1.29	1.07	1.55	1.56	1.09	2.24	1.01	0.92	1.11
CD (yes=1)		3.10	2.52	3.83	2.54	1.68	3.84	1.72	1.57	1.89
DM (yes=1)		2.69	2.23	3.23	3.65	2.55	5.19	2.27	2.08	2.48

NT: Non-intact teeth CD: Chewing difficulty DM: Dry mouth

Controlled for age, gender, living situation and presence of at least one informal caregiver

Predictive association of oral health and general health items

Figures 1, 2 and 3 show the significant pairs of associations between oral health and general health indicators. The arrows were drawn from the estimates for the cross-lagged parameters (Supplementary material Tables A1, A2 and A3 - gamma estimates: γ_{12} and γ_{21}). A full arrow means a prediction with a positive association, showing that the presence of a current health problem is associated with an increased likelihood of the presence, or worsening of another health problem at a future assessment. Therefore, a full arrow starting at an oral health item and pointing to a general health item indicates a positive association between these indicators, meaning that given a certain general health status, the presence of an oral health problem is associated with greater severity or impairment on the general health status at a future assessment. A bi-directional arrow means that the association is reciprocal, so it is significant in both directions, from an oral health item to a general health item and vice versa.

A dotted arrow means a prediction with an inverse association, showing that the presence of a current problem is associated with higher odds of an improvement or absence of another problem in the future.

The numbers on the arrows indicate the estimates for the cross-lagged parameters (gamma estimates: γ_{12} and γ_{21}). As the highest part of the prediction is made by the parameters γ_{11} and γ_{22} , which are the parameters related to the situation of the same indicator in the previous time period, these values are often close to 1. The values of the cross-lagged parameters are normally much lower than 1 as most of the prediction is made by the parameters γ_{11} and γ_{22} . The cross-lagged parameter γ_{12} (respectively γ_{21}) is an indication of the amount of extra information that the current oral health (respectively general health) indicator provides to the future general health (respectively oral health) indicator. In case the cross-lagged parameters are significant, like all the estimates shown in the figure, this means that they are able to add to the prediction in the model.

Figure 1 shows the significant associations between oral health and general health indicators in the Belgian sample. The results show that, given current ADL, current chewing status is predictive of ADL status at the next time-point. The bi-directional arrow shows that current ADL status is also predictive of chewing difficulty at the next time-point. In summary, the estimate for the cross-lagged parameters indicates that given the current status of ADL (respectively chewing difficulty), the current chewing difficulty (respectively ADL) provides a significant amount of information in predicting the future value of ADL (respectively chewing difficulty). Chewing difficulty was also predictive of health instability and cognitive impairment. The bi-directional arrow also shows that CPS, unlike CHES, was predictive of chewing difficulty at the next time-point. Non-intact teeth were predictive of CHES and ADL at the next time-point, but only CPS was predictive of non-intact teeth. Dry mouth was predictive of DRS at the next time-point and vice-versa, showing that current dry mouth status was associated with more depressive symptoms at follow-up and vice versa. Dry mouth was also predictive of ADL at the next time-point. In addition, given the current health instability status, dry mouth was predictive of health instability at the next time-point.

Figure 1. Significant associations of the current oral health (respectively general health) status to the future general health (respectively oral health) in the Belgian dataset.

NT: Non-intact teeth CD: Chewing difficulty DM: Dry mouth

The results for the pairs of associations between oral health and general health indicators from the iBenC dataset can be found in Figure 2. In this model, given current DRS and CHESS, current dry mouth predicts both DRS and CHESS at a time-point in the future. DRS and CHESS can also predict dry mouth at the next time-point. In addition, DRS is predictive of chewing difficulty. The results also show that CPS and ADL are both predictive of non-intact teeth.

The dotted arrow from non-intact teeth to DRS shows a significant inverse association between these indicators, as well as the dotted arrow from CPS to dry mouth. This means that a decrease in cognitive problems is associated with increased odds for dry mouth in the future and that presence of non-intact teeth is associated with less depressive symptoms in the future.

Figure 2. Significant associations of the current oral health (respectively general health) status to the future general health (respectively oral health) in the iBenC dataset.

NT: Non-intact teeth CD: Chewing difficulty DM: Dry mouth

Figure 3 shows the associations between oral health and general health indicators from the New Zealand sample. The results point out that given a current ADL status, chewing difficulty is predictive of ADL. The dotted line in the opposite direction shows a significant inverse association between ADL and chewing difficulty. In addition, given current CHES, chewing difficulty is predictive of CHES in the future. Dry mouth is predictive of DRS and CHES at a time-point in the future, but no associations were found between non-intact teeth and the health indicators.

The dotted lines show a significant inverse association between ADL and dry mouth, as well as between CPS and dry mouth and vice versa.

Figure 3. Significant associations of the current oral health (respectively general health) status to the future general health (respectively oral health) in dataset from New Zealand

NT: Non-intact teeth CD: Chewing difficulty DM: Dry mouth

In summary, the predictive association between dry mouth and health instability emerged in all three samples, as well as between dry mouth and depressive symptoms. An inverse predictive association was found between dry mouth and CPS in the samples from iBenC and New Zealand but not in the Belgian sample. For CPS and the prediction of non-intact teeth, the association emerged for the samples from Belgium and iBenC, but not for New Zealand, as well as the prediction between non-intact teeth and ADL. Chewing difficulty showed predictive association for ADL and CHESS for Belgium and New Zealand.

Discussion

This research paper shows evidence of the association between oral health and general health using large cross-country databases: Belgium, iBenC project and New Zealand. When comparing the odds ratios for oral health and general health for the three study populations, most cross-sectional associations were similar. The associations between chewing difficulty and ADL, cognition, depressive symptoms and health instability were confirmed. Moreover, the associations between non-intact teeth and cognition, as well as non-intact teeth and depressive symptoms were verified, as well as the associations between dry mouth and depressive symptoms and dry mouth and health instability. The results show the relevance of oral health in relation to general health, and vice versa. Strong significant associations between oral health and general health were confirmed in the three datasets, providing evidence of robust interrelation between oral health and general health.

In practice, the results indicate that an oral health assessment and the treatment of oral health problems should be fostered, and that this can help older people to remain in good health, especially when complemented with tailored nutritional advice. By solving oral health problems, nutritional intake deficiencies may be avoided, which in turn can prevent general health problems³¹⁻³³. In addition, associations show that if older people have good general health, they seem to have better oral health.

Especially chewing difficulty was associated with each of the general health indicators. As chewing difficulty indicates difficulty to masticate regardless of cause (e.g., ill-fitting dentures, neurological impairment, temporomandibular joint pain, painful tooth, etc.), it is not possible to explore the direct cause of the problem within the interRAI assessment. Further examination is thus needed in order to be able to advise adequate treatment^{34,35}. Dry mouth and non-intact teeth showed significant associations with almost all general health indicators in the three countries. The association between non-intact teeth and general health is not well-documented in the scientific literature and this paper can trigger further research. Non-intact teeth, however, is a rather vague description of a dental condition and can imply both infectious (caries) as well as non-infectious (e.g. attrition, erosion) dental conditions. The mechanism underlying the association between non-intact teeth and general health conditions depends on the specific etiology of the dental disease.

The results also suggest that dry mouth is associated with better cognitive function. The latter finding may be explained by the fact that drugs for dementia often induce dry mouth and that the drug-induced cognitive improvement is thereby associated with dry mouth^{36,37}.

In the population of our study, we suspect that the highest contributor to dry mouth is medication use, seen by the prevalence of polypharmacy in older adults³⁸. On the other hand, functional loss or cognitive problems are most likely the contributors to non-intact teeth and chewing difficulty as older people lose their capacity to perform self-care sufficiently. Addressing dry mouth, as well as non-intact teeth and chewing difficulty will require different approaches. A review of the medication could be a possible strategy to treat dry mouth, as well as non-drug treatments for depression and anxiety^{36,39-41}. For non-intact teeth and chewing difficulty, dental check-ups and care, as well as support with daily oral self-care could be effective^{42,43}.

The predictions from the second analysis differed in some cases but the model provided validation for the ability to predict outcomes and the results were significant. This finding is important in order to work on prevention of oral health and general health problems. The inclusion of oral health indicators

in longitudinal research of health outcomes, although important, is rarely done. Raising awareness for oral health assessment and treatment is advisable. Policy makers should foster oral health promotion and care in order to keep older adults in good health.

Although the interRAI assessments are comprehensive and have proven validity for many aspects of general health (cognition, functional performance, depressive symptoms, health instability, etc.), some authors point out that the items about oral health from the previous version of the interRAI instruments did not detect all oral health problems present, contesting their validity⁴⁴⁻⁴⁶. This is due to lack of completeness and unclear wording of the items, or due to lack of training. In our study, professional caregivers received training to fill out the items in general, and the assessment was not only based on self-report. This may have mitigated the problem, but further research is needed to evaluate whether these items should be adapted or replaced by more valid oral health items.

Strengths and limitations

A strength of the present study was the availability of large longitudinal databases. The tested Bayesian models can be considered more appropriate than regression analysis due to the complexity of the data. Further, this approach was chosen for the more flexible modelling tools in Bayesian software such as Stan.

A limitation of the study was the selection bias, as only clients receiving home care were included. The population in the study is thus not representative of a general population of older people in the community. Due to the General Data Protection Regulation⁴⁷, the three datasets could not be merged and therefore, parameter differences between the samples and the models could not be tested. In addition, the datasets could not be combined to construct one large model. In addition, no data on medication use or diagnoses could be included in the analysis.

Conclusions and Implications

The results point out the need of the inclusion of oral health assessment and advice from dentists or oral health practitioners into the multidisciplinary conversation. This would allow for the development of care plans focusing also on preventing oral health problems, with the broader goal of improving general health. The interRAI assessments can be used to trigger decision-support algorithms called Client Assessment Protocols (CAPs)²¹. An oral health CAP has not yet been developed but it would be an asset in order to identify clients who have current oral health problems and clients who are at risk of developing them.

Conflict of interests

The authors declare no conflict of interests.

References

1. Tsakos G, Quiñonez C, Sabbah W, et al. A sober look at the links between oral and general health. *J Epidemiol Community Health*. 2013; 67(5):381-382. doi:10.1136/jech-2013-202481.
2. Kimura Y, Ogawa H, Yoshihara A, et al. Evaluation of chewing ability and its relationship with activities of daily living, depression, cognitive status and food intake in the community-dwelling elderly. *Geriatr Gerontol Int*. 2013;13(3):718-725.
3. Locker D. Concepts of oral health, disease and the quality of life. In: Slade GD, editor. *Measuring oral health and quality of life*. Chapel Hill: University of North Carolina, Dental Ecology. 1997; 11-23.
4. Welmer AK, Rizzuto D, Parker M G, Xu, W. Impact of tooth loss on walking speed decline over time in older adults: a population-based cohort study. *Aging Clin Exp Res*. 2017; 29, 793–800.
5. AlJameel AH, Watt RG, Brunner EJ, Tsakos G. Earlier depression and later-life self-reported chewing difficulties: Results from the Whitehall II study. *J Oral Rehabil*. 2015;42, 98–104.
6. Onder G, Liperoti R, Soldato M et al. Chewing problems and mortality in older adults in home care: results from the aged in home care study. *J Am Geriatr Soc*, 2007; 55, 1961-1966.
7. Oh E, Weintraub N, Dhanani S. Can we prevent aspiration pneumonia in the nursing home? *J Am Med Dir Assoc* 2005;6(3 Suppl):S76-80.
8. James KH Luk, FHKCP, FHKAM, Daniel KY Chan, MD, FACP (2014). Preventing aspiration pneumonia in older people: do we have the 'know-how'? *Hong Kong Med J* 2014; Oct;20(5):421–7.
9. Müller, F. Oral Hygiene Reduces the Mortality from Aspiration Pneumonia in Frail Elders. *Journal of Dental Research*, 2015;94(3 Suppl), 14S–16S.
10. Pedersen AML, Sørensen CE, Proctor GB, Carpenter GH, Ekström J. Salivary secretion in health and disease. *J Oral Rehabil*. Sep 2018;45(9):730-746.
11. Napeñas JJ, Rouleau TS. Oral complications of Sjögren's syndrome. *Oral Maxillofac Surg Clin North Am*. 2014; 26(1):55-62.

12. Graziani F, Gennai S, Solini A, Petrini M. A systematic review and meta-analysis of epidemiologic observational evidence on the effect of periodontitis on diabetes: An update of the EFP-AAP review. *J Clin Periodontol*. 2018;45(2):167-187.
13. Lamster IB, Pagan M. Periodontal disease and the metabolic syndrome. *Int Dent J*. 2017 Apr;67(2):67-77.
14. Teixeira FB, Pereira Fernandes L de M, Tavares Noronha PA, et al. Masticatory deficiency as a risk factor for cognitive dysfunction. *Int J Med Sci*. 2014;11(2):209-214.
15. Tada A, Miura H. Association between mastication and cognitive status: A systematic review. *Arch Gerontol Geriatr*. 2017; 70:44-53.
16. Trung DT, Krausch-Hofmann S, Duyck J, et al. Association between oral health and general health indicators in older adults. *Scientific Reports*; 2018; 8; iss. 1.
17. de Almeida Mello J, Van Durme T, Macq J, Declercq A. Interventions to delay institutionalization of frail older persons: design of a longitudinal study in the home care setting. *BMC Public Health*. 2012;12:615.
18. Garms-Homolová V, Notthoff N, Declercq A et al. Social and functional health of home care clients with different levels of cognitive impairments. *Aging Ment Health* 2017;21:18–23
19. New Zealand Technical Advisory Services (TAS), New Zealand. Ad-hoc dataset provided from interRAI New Zealand data warehouse, Request number 2017-328, April 2018. www.interrai.co.nz
20. Landi F, Tua E, Onder G et al. The Minimum Data Set for Home Care: a valid instrument to assess frail older people living in community. *Med Care*. 2000;30: 1184-1190.
21. Carpenter GI, Hirdes JP Using interRAI assessment systems to measure and maintain quality of long-term care in OECD/European Commission: A Good Life in Old Age? Monitoring and Improving Quality in Long-term Care, OECD Health Policy Studies, OECD Publishing 2013;93-139.
22. Krausch-Hofmann, S., Bogaerts, K., Hofmann, M., de Almeida Mello, J., Fávaro Moreira, N. C., Lesaffre, E., ... Duyck, J. (2015). Missing Oral Health-Related Data in the interRAI-HC - Associations with Selected Variables of General Health and the Effect of Multiple Imputation on the Relationship between Oral and General Health. *Plos One*, 10(12), e0146065.
23. Morris JN, Fries BE, Morris S. Scaling ADLs within the MDS. *J Gerontol A Biol Sci Med Sci*.

- 1999;54(11):M546-M553.
24. Morris JN, Fries BE, Mehr DR, et al. MDS Cognitive Performance Scale. *J Gerontol.* 1994;49(4):M174-M182.
 25. Burrows AB, Morris JN, Simon SE et al. Development of a minimum data set-based depression rating scale for use in nursing homes. *Age Ageing.* 2000;29(2):165-172.
 26. Hirdes JP, Frijters DH, Teare GF. The MDS-CHESS Scale: A new measure to predict mortality in institutionalized older people. *J Am Geriatr Soc.* 2003;51(1):96-100.
 27. Lesaffre E, Lawson AB. *Bayesian Biostatistics.* Wiley. <https://books.google.be/books?id=WV7KVjEQnJMC>. Series - Statistics in Practice 2012. ISBN: 978-0-470-01823-1
 28. Denwood MJ. Runjags: An R Package Providing Interface Utilities, Model Templates, Parallel Computing Methods and Additional Distributions for MCMC Models in JAGS. *J Stat Softw.* 2016;71(9):1-25.
 29. Stan Development Team. RStan: the R interface to Stan. Version 2.16, <http://mc-stan.org/>
 30. Johnson VE, Albert JH. *Ordinal Data Modeling.* Springer New York; 2000. <https://books.google.be/books?id=6-Y8OL3sW14C>.
 31. Sheiham A, Steele J. Does the condition of the mouth and teeth affect the ability to eat certain foods, nutrient and dietary intake and nutritional status amongst older people? *Public Health Nutrition* 2001;4:797-803.
 32. Ono S, Ishimaru M, Yamana H, Morita K, Ono Y, Matsui H, Yasunaga H. Enhanced Oral Care and Health Outcomes Among Nursing Facility Residents: Analysis Using the National Long-Term Care Database in Japan. *J Am Med Dir Assoc.* 2017 Mar 1;18 (3):277.e1-277.e5.
 33. Kossioni A. The Association of Poor Oral Health Parameters with Malnutrition in Older Adults: A Review Considering the Potential Implications for Cognitive Impairment *Nutrients* 2018, 10(11), 1709; <https://doi.org/10.3390/nu10111709>.

34. Woo J, Tong C, Yu R. Chewing Difficulty Should be Included as a Geriatric Syndrome. *Nutrients*. 2018;10(12):1997. Published 2018 Dec 17. doi:10.3390/nu10121997
35. Takagi D, Watanabe Y, Eda Hiro A, et al. Factors affecting masticatory function of community-dwelling older people: Investigation of the differences in the relevant factors for subjective and objective assessment. *Gerodontology*. 2017;34:357–364.
36. Tan CKE, Lexomboon D, Sandborgh-Englund G et al. Medications that cause dry mouth as an adverse effect in older people: A Systematic Review and Metaanalysis. *J Am Geriatr Soc*. 2018;66:76–84.
37. Gil-Montoya JA, Barrios R, Sánchez-Lara I, et al. Prevalence of drug-induced xerostomia in older adults with cognitive impairment or dementia: an observational study. *Drugs Aging*. 2016;33(8):611.
38. Stewart D, Mair A, Wilson M et al. SIMPATHY consortium Guidance to manage inappropriate polypharmacy in older people: systematic review and future developments. *Expert Opin Drug Saf*. 2017;16(2):203-213.
39. Altamimi AM. Update knowledge of dry mouth- a guideline for dentists. *African Health Sciences*. 2014; 14(3):736–742.
40. Takiguchi T, Yoshihara A, Takano N , Miyazaki H. Oral health and depression in older Japanese people. *Gerodontology*. 2016;33,439–446.
41. Dudas R, Malouf R, McCleery J, and Denning T. Antidepressants for treating depression in dementia. *Cochrane Database of Systematic Reviews*. John Wiley & Sons, Ltd. 2018.
42. Razak PA, Richard KMJ, Thankachan RP, et al. Geriatric Oral Health: A Review Article. *Journal of International Oral Health JIOH*, 2014;6(6), 110–116.
43. Tomata Y, Watanabe T, Sugiyama K et al. Effects of a Community-Based Program for Oral Health and Nutrition on Cost-Effectiveness by Preventing Disability in Japanese Frail Elderly: A Quasi-Experimental Study Using Propensity Score Matching. *J Am Med Dir Assoc*. 2017 Aug 1;18(8):678-685.

44. Hoben M, Poss JW, Norton PG, Estabrooks CA. Oral/dental items in the resident assessment instrument - minimum Data Set 2.0 lack validity: results of a retrospective, longitudinal validation study. *Popul Health Metr.* 2016;14:36. Published 2016 Oct 21.
45. Cohen-Mansfield J, Lipson S. The underdetection of pain of dental etiology in persons with dementia. *Am J Alzheimers Dis Other Demen.* 2002;17:249–53.
46. Folse GJ. National MDS and dental deficiency data reported by the US Health Care Financing Administration (HCFA) Spec Care Dentist. 2001;21:37–8.
47. European Commission: 2018 reform of EU data protection rules:
https://ec.europa.eu/commission/priorities/justice-and-fundamental-rights/data-protection/2018-reform-eu-data-protection-rules_en Accessed on November 26 2018.

Supplementary material:

Table S1- Parameter estimates and 95% credible intervals (CI) for all γ 's for all pairs of one oral health and one general health indicator
Belgium

Pair	γ_{11}			γ_{12}			γ_{21}			γ_{22}		
	Est.	95% CI		Est.	95% CI		Est.	95% CI		Est.	95% CI	
ADL and NT	0.83	0.82	0.84	0.03	0.001	0.06	0.00	-0.02	0.03	0.98	0.98	0.99
ADL and CD	0.83	0.82	0.84	0.08	0.04	0.11	0.03	0.002	0.06	0.97	0.95	0.98
ADL and DM	0.83	0.82	0.84	0.07	0.04	0.11	0.00	-0.03	0.03	0.96	0.95	0.97
CPS and NT	0.93	0.93	0.94	0.02	0.00	0.04	0.03	0.01	0.06	0.98	0.98	0.99
CPS and CD	0.93	0.93	0.94	0.03	0.01	0.05	0.05	0.02	0.08	0.96	0.95	0.97
CPS and DM	0.93	0.93	0.94	0.02	-0.01	0.04	0.02	-0.01	0.06	0.96	0.95	0.97
DRS and NT	0.88	0.86	0.89	0.01	-0.02	0.03	0.01	-0.02	0.03	0.98	0.98	0.99
DRS and CD	0.87	0.86	0.88	0.02	-0.01	0.05	0.01	-0.02	0.04	0.97	0.95	0.98
DRS and DM	0.87	0.86	0.88	0.04	0.01	0.07	0.05	0.01	0.08	0.96	0.94	0.97
CHESS and NT	0.57	0.54	0.60	0.09	0.05	0.14	0.02	-0.01	0.04	0.98	0.98	0.99
CHESS and CD	0.56	0.53	0.59	0.18	0.13	0.23	0.01	-0.02	0.05	0.97	0.96	0.98
CHESS and DM	0.55	0.52	0.59	0.23	0.18	0.28	0.00	-0.03	0.03	0.96	0.95	0.97

NT: Non-intact teeth CD: Chewing difficulty DM: Dry mouth

Table S2- Parameter estimates and 95% credible intervals (CI) for all γ 's for all pairs of one oral health and one general health indicator

IBenC sample

Pair	γ 11			γ 12			γ 21			γ 22		
	Est.	95% CI		Est.	95% CI		Est.	95% CI		Est.	95% CI	
ADL and NT	0.93	0.93	0.94	0.02	0.00	0.04	0.05	0.02	0.08	0.98	0.98	0.99
ADL and CD	0.93	0.93	0.94	0.00	-0.02	0.02	0.01	-0.01	0.04	0.98	0.98	0.99
ADL and DM	0.93	0.93	0.94	-0.01	-0.03	0.01	-0.02	-0.05	0.00	0.98	0.97	0.98
CPS and NT	0.93	0.93	0.94	0.01	-0.01	0.03	0.02	0.002	0.05	0.98	0.98	0.99
CPS and CD	0.93	0.93	0.93	0.03	0.003	0.05	0.00	-0.03	0.03	0.98	0.98	0.99
CPS and DM	0.93	0.93	0.94	0.01	-0.01	0.03	-0.05	-0.08	-0.03	0.97	0.96	0.98
DRS and NT	0.89	0.89	0.91	-0.03	-0.06	0.001	0.02	0.00	0.04	0.98	0.98	0.99
DRS and CD	0.89	0.89	0.91	0.01	-0.02	0.04	0.05	0.02	0.08	0.98	0.97	0.99
DRS and DM	0.89	0.89	0.90	0.04	0.01	0.06	0.07	0.04	0.10	0.97	0.96	0.98
CHESS and NT	0.73	0.71	0.75	0.02	-0.02	0.07	0.03	0.00	0.05	0.98	0.98	0.99
CHESS and CD	0.73	0.70	0.75	0.05	0.00	0.09	0.02	-0.01	0.05	0.98	0.98	0.99
CHESS and DM	0.73	0.70	0.75	0.07	0.03	0.12	0.05	0.02	0.08	0.97	0.96	0.98

NT: Non-intact teeth CD: Chewing difficulty DM: Dry mouth

Table S3- Parameter estimates and 95% credible intervals (CI) for all γ 's for all pairs of one oral health and one general health indicator

New Zealand

Pair	γ 11			γ 12			γ 21			γ 22		
	Est.	95% CI		Est.	95% CI		Est.	95% CI		Est.	95% CI	
ADL and NT	0.24	0.20	0.28	0.01	-0.05	0.07	0.02	-0.02	0.06	0.96	0.95	0.97
ADL and CD	0.23	0.20	0.27	0.13	0.07	0.19	-0.05	-0.09	-0.01	0.93	0.91	0.95
ADL and DM	0.24	0.20	0.28	0.05	-0.01	0.10	-0.05	-0.10	-0.01	0.92	0.89	0.93
CPS and NT	0.66	0.63	0.68	-0.04	-0.09	0.01	0.00	-0.04	0.03	0.96	0.95	0.97
CPS and CD	0.66	0.63	0.68	0.01	-0.04	0.06	-0.01	-0.05	0.03	0.92	0.90	0.94
CPS and DM	0.66	0.63	0.68	-0.06	-0.10	-0.01	-0.11	-0.15	-0.06	0.91	0.88	0.93
DRS and NT	0.62	0.59	0.64	0.00	-0.05	0.05	-0.03	-0.07	0.02	0.96	0.95	0.97
DRS and CD	0.62	0.59	0.64	0.03	-0.03	0.08	0.00	-0.05	0.04	0.93	0.90	0.94
DRS and DM	0.61	0.59	0.64	0.09	0.05	0.14	0.03	-0.02	0.07	0.92	0.89	0.93
CHESS and NT	0.30	0.26	0.33	-0.05	-0.11	0.01	0.00	-0.03	0.04	0.96	0.95	0.97
CHESS and CD	0.29	0.26	0.33	0.10	0.05	0.16	-0.02	-0.06	0.02	0.93	0.90	0.94
CHESS and DM	0.29	0.26	0.33	0.12	0.06	0.17	-8E-04	-0.04	0.04	0.92	0.90	0.94

NT: Non-intact teeth CD: Chewing difficulty DM: Dry mouth