Associative fear learning to interoceptive stimuli features in theoretical accounts of panic disorder. In a similar vein, interoceptive exposure therapy is based on the assumption that interoceptive fear is learned and can be extinguished again. Nonetheless, fear conditioning research is dominated by exteroceptive paradigms that score low on their ecological validity for panic disorder. From a functional perspective, panic-relevant conditioning paradigms ought to use panicogenic unconditional stimuli (US), conditioned stimuli with a natural relationship to such US, and study panic-relevant response systems. We review the empirical work with respiratory USs, discuss it from such functional perspective, and outline some potential avenues for future research.

INTEROCEPTIVE FEAR CONDITIONING

EARLY CAREER AWARD, 2013

Interoception, Conditioning, and Fear: a potent Threesome with special Relevance to Panic Disorder

Ilse Van Diest

Health, Behavior & Psychopathology

Faculty of Psychology & Educational Sciences

University of Leuven, Belgium

Running title: Interoceptive fear conditioning**Corresponding author:**

Ilse Van Diest, Ph.D.

Health Psychology, University of Leuven, Tiensestraat 102, 3000 Leuven, Belgium. Telephone:

0032 16 32 60 29. Email: ilse.vandiest@kuleuven.be

INTEROCEPTIVE FEAR CONDITIONING

1. Interoception: its definition, its functions, and two of its troubles

The concept of interoception has not been consistently defined in the scientific literature, leading to a lack of clarity about what can and cannot be considered 'interoception' or 'interoceptive'. Various different views and definitions have been proposed (Khalsa & Lapidus, 2016), ranging from the very restrictive (e.g., only visceral) to the very inclusive (incorporating touch and gustation; for more details, see Cameron, 2001; Ceunen, Vlaeyen, & Van Diest, 2016) – but these are based on disparate principles, and many people seem to default to a more intuitive sense of what constitutes 'interoception'. Despite the fluctuating boundaries of the word's meaning, most authors in the field would agree with Bud Craig's tenet that interoception informs the brain about the physiological condition of the body and serves important regulatory functions (Craig, 2002). Consistent with this, a consensus was recently reached by a group of interoception researchers (Khalsa et al., 2017, p 501), who defined interoception as "*a process by which the nervous system senses, interprets, and integrates signals originating from within the body, providing a moment-by-moment mapping of the body's internal landscape across conscious and unconscious levels.*" This definition emphasizes that very different levels and layers of the central nervous system can be involved in interoception, ranging from local regulatory circuits to central, higher-order circuits involved in the awareness and perception of bodily functions and states. As an example, respiration can be sensed with a variety of specialized interoceptors (e.g., pulmonary stretch receptors, chemoreceptors), and the vital function of breathing is usually regulated unconsciously as respiratory sensory input, though accessible to awareness, does not capture attention or lead to a conscious percept. When respiratory functions are threatened and automatic regulatory systems within the body fail, respiratory perception and discomfort typically arise, which engages the behavioral system. The perception of breathlessness (or dyspnea), in particular, motivates one to reduce the aversive state of breathlessness, whether it is by reducing ongoing or planned physical activity, gasping, opening the window and get some fresh air, seeking distraction

INTEROCEPTIVE FEAR CONDITIONING

1 from the aversive experience by starting in an engaging activity, using one's inhaler, breathing
2 more deeply or rapidly, or seeking medical help. This behavioral layer in the homeostatic circuit
3 allows for flexible selection of the most adaptive and appropriate response for a certain situation at
4 a certain time (Schroijen et al., in press). The degrees of freedom in behavioral responding may
5 come with a cost, however, as less adaptive behaviors that may maintain or even worsen the
6 situation can also be selected. To complicate things further, behavior that may be adaptive at one
7 level (e.g., temporary relief of dyspneic feelings by sighing) is not necessarily beneficial at another
8 level (e.g., excessive sighing inducing a decrease in the arterial pressure of CO₂).

9 The above example from the respiratory domain illustrates that interoception is indeed
10 multifaceted and relies on diverse functions and structures of the nervous system that emerged and
11 were preserved throughout evolution. As such, interoception seems to contribute to a wide range of
12 behavioral and psychological aspects of human life. Apart from helping to maintain the body's
13 homeostasis, interoception is involved in emotions, emotional regulation, eating behavior, sexual
14 functioning, craving and addiction, symptom perception (e.g., pain, dyspnea and fatigue),
15 perception and control of bodily urges and functions (e.g., micturition and defecation, hunger and
16 thirst, perception of cardiorespiratory and gastro-intestinal activity), self-awareness and
17 consciousness, time perception, decision making and empathy (Ceunen et al., 2016; Khalsa et al.,
18 2017). Given the role of interoception in so many, apparently diverse functions, it is not surprising
19 to see that interoceptive malfunctions seem to play a role in many major psychopathologies
20 (Khalsa et al., 2017; Petzschner et al., 2017). Perhaps the psychopathologies most studied in this
21 respect are panic disorder and psychosomatic disorders (Cameron, 2001). Two interoceptive
22 hallmarks of panic disorder are the sudden and unexpected occurrence of panic attacks, and
23 excessive and disabling fear of bodily sensations of arousal, sometimes referred to as interoceptive
24 fear or 'fear of fear'. Psychosomatic disorders, on the other hand, are characterized by the
25 disabling experience of bodily symptoms without any identifiable medical cause. If chronic, such

INTEROCEPTIVE FEAR CONDITIONING

symptoms are often referred to as psychosomatic, functional or (medically) unexplained symptoms, because central processes - rather than peripheral afferent sensory input - are assumed to generate them (Fava & Sonino, 2010).

The present paper focuses on one such process. Specifically, our goal is to illustrate and review some of the evidence for how associative learning involving interoceptive stimuli, or ‘interoceptive conditioning’, can elicit changes in physiological, emotional and potentially also perceptual outcomes. We first outline a functional perspective on Pavlovian conditioning and provide the reader with a general introduction and historical background on interoceptive conditioning as a special instance of Pavlovian conditioning. Next, we discuss biological and learning accounts of panic disorder and analyze which stimuli and responses may be most useful for modelling learning that is related to panic disorder. Finally, we review and discuss studies on interoceptive fear conditioning with respiratory stimuli, and outline future directions.

2. Pavlovian conditioning

2.1 A functional perspective

Pavlovian (or classical) conditioning refers to associative learning between stimuli or events, meaning that one stimulus/event evokes the mental representation or expectation of the other stimulus/event. In the prototypical laboratory paradigm, only one of the two stimuli is significant (US, or unconditional stimulus), meaning that it unconditionally (and prior to any learning experience) evokes an observable response (unconditioned response, UR). Typically, the other stimulus (conditioned stimulus, CS) is initially ineffective in triggering a response relevant to the US and is therefore labeled ‘neutral’. That associative learning between CS and US has occurred can be deduced from the observation that the initially ineffective stimulus (CS) starts to elicit a response (conditioned response, CR) after its contingent pairing with the US. Importantly, the prior ‘neutrality’ of the CS is not a strict necessary condition for the occurrence of associative learning, it

INTEROCEPTIVE FEAR CONDITIONING

only facilitates an unconfounded observation of a developing CR in the laboratory and makes it easier to conclude that learning has taken place.

Functionally, the conditioned response (CR) prepares the organism to deal with the upcoming US and typically engages the same response system as the US, though the CR is not necessarily identical to and can sometimes even oppose the UR (Krause & Domjan, 2017). Drug tolerance, for example, is often explained by the formation of a CR (to cues associated with drug administration) that counteracts the drug effect (UR) (Ramsay & Woods, 1997; Siegel, 1989).

Pavlov and colleagues were likely the first to realize and provide empirical evidence that associative learning between stimuli is a potentially powerful mechanism to support physiological and behavioral regulation by learning to respond (CR) to an anticipated, biologically relevant stimulus (US). Indeed, already starting to salivate (CR) in response to a bell (CS) that announces upcoming food (US) will ultimately facilitate the dog's digestion of that food. Likewise, male Japanese quails ejaculated more sperm and mounted the female quail more quickly after a conditioned visual stimulus announced the presence of a receptive female quail, compared to males who copulated with an 'unannounced' female (Krause & Domjan, 2017; Domjan, Blesbois, & Williams, 1998). Numerous other examples can be found in the literature - from different species (even those with very simple nervous systems) and involving diverse physiological or behavioral regulatory systems, each with their specific underlying neural circuit. All this suggests strongly that Pavlovian conditioning is a fundamental principle that is preserved across species, which can occur at many levels of the nervous system and imply diverse response and memory systems (Fanselow & Poulos, 2005). In humans, for example, Pavlovian conditioning has been demonstrated to affect physiological regulation (e.g., Dworkin, 1993; Stockhorst et al., 2000), sensory processing and perception (e.g., Miskovic & Keil, 2012; Rossi et al., 2017; Zaman et al., 2016), emotions and their associated motivated behaviors (e.g., Fannes et al., 2008), as well as declarative knowledge and contingency learning (e.g., De Houwer & Beckers, 2002).

INTEROCEPTIVE FEAR CONDITIONING

This ubiquity of Pavlovian conditioning likely explains why the dominant focus in the research on associative learning in Western psychology is on identifying general principles of associative learning. Typically, the response system under study is relatively arbitrary in this line of research. As its primary interest is on the formation of associations and changes in associative strength (between CS and US), it tends to focus on the occurrence and intensity, rather than on the nature or direction, of the CR. Most work from this tradition uses CSs and a US that allow maximal experimental control, and focuses on responses that are convenient to observe and quantify. This type of work is often characterized by excellent internal validity and has yielded important insights in the principles of associative learning that otherwise would have remained undiscovered or poorly understood. A potential downside of the approach, however, may be that the external validity is less obvious, thereby carrying the risk that the translation of the research to increase the understanding and malleability of responses and behaviors that occur in natural settings - and that matter in real life - is poor. As an example, prototypical human fear conditioning paradigms with visual/auditory CSs and an electrical shock or a loud noise as the US have offered only limited insights into whether and how associative learning can establish the interoceptive fear that is so characteristic of panic disorder, or how it can modulate the nature and occurrence of panic attacks. Paradigms with biologically significant USs and response systems, and the selection of CSs that show a natural relationship with the US, may offer an interesting solution to this problem. In this respect, an inspirational area of research is that on interoceptive conditioning.

2.2 Interoceptive conditioning

As many interoceptive events occur sequentially and take a predictable course, associative learning between them can arise easily. For example, whenever skeletal muscles start working harder, the body will increase its production of CO₂, which reflexively augments ventilation through a chemoreceptor feedback loop supporting homeostatic mechanisms that strive for a

INTEROCEPTIVE FEAR CONDITIONING

constant and stable pressure of CO₂ in the arterial blood. Paradoxically, however, ventilation already increases at the onset of exercise, *prior* to any rise in arterial CO₂-pressure, a phenomenon called 'exercise hyperpnea'. A prevailing hypothetical explanation is that Pavlovian conditioning may have installed a feedforward loop for the regulation of respiratory activity. Specifically, cues associated with the onset of exercise (CSs) are repeatedly followed by an increase in CO₂ production (US), which may lead to associative learning between them. As a result of such learning, a conditioned anticipatory response to the cues (increase in ventilation, the CR) is established, which helps to deal with the upcoming homeostatic disturbance caused by the US (Poon, Tin, & Yu, 2007). Another respiratory example, but now at the behavioral level, is an asthma patient who learns to recognize early respiratory sensations of upcoming asthma attacks and develops anticipatory anxiety. Such learning may actually also affect respiratory sensation and perception, as will be discussed later in this paper.

Associative learning involving interoceptive stimuli, such as in the above examples, is referred to as 'interoceptive conditioning'. Although interoceptive conditioning is far less studied than exteroceptive conditioning, it seems to contribute substantially to physiological regulation (Dworkin, 1993; Somjen, 1992) and to emerge ontogenetically even prior to exteroceptive conditioning (Haroutounian & Campbell, 1979). The literature often distinguishes between three subtypes of interoceptive conditioning, dependent on whether the CS, the US or both are of an interoceptive nature - these are referred to as extero-interoceptive, intero-exteroceptive and intero-interoceptive types of conditioning, respectively (Razran, 1961). Interoceptive conditioning has a history that dates back to the old Russian physiological school. Although in the past decades Pavlovian conditioning on human animals has been heavily skewed towards extero-exteroceptive pairings (visual or auditory CSs paired with electrocutaneous shocks in particular), Pavlov himself was primarily interested in visceral learning and regulation and used biologically significant USs in many of his studies (Tansey, 2006). It was also Pavlov who emphasized that interoceptive events

INTEROCEPTIVE FEAR CONDITIONING

1 can also serve as a CS. Therefore, in its infancy, the study of classical conditioning was closely
2 connected to the topic of interoception and adopted a functional perspective that nowadays often
3 seems lost.

4 A Russian monograph by Bykov, describing work with interoceptive CSs, was translated
5 into English by H.W. Gantt (Bykov, 1957). Razran (1961) brought interoceptive conditioning to
6 the attention of Western Psychology by reviewing and discussing 14 of the Russian experiments on
7 interoceptive conditioning, as well as a few clinical observations. In animals, visceral CSs used in
8 the Russian experiments described by Razran (1961) included distentions using water/air inflatable
9 balloons inserted in the animals' uterus, bladder, or at distinct places along the gastro-intestinal
10 tract; chemical or electrical stimulation of viscera; rhythmic 'scratching' of the visceral mucosa.
11 These CSs were paired with one or more of the following USs: inhalation of CO₂-enriched air;
12 electrical shocks in the rectum or on the dog's leg; balloon distentions in the rectum; infusion of a
13 salty solution in the stomach. Distentions and thermal stimulation of the viscera were also applied
14 in human subjects, either by having the participant swallowing a balloon to which electrodes could
15 be attached to stimulate the stomach, or by studying patients with pre-existing urinary bladder
16 fistulae. Using these stimuli, conditioning effects were reported not only for visceromotor,
17 skeletal, vasomotor and respiratory reactions, but also for the urges to urinate or defecate, and for
18 visceral perception and the discrimination of visceral perceptions from one another.

19 Based on his reading of the Russian experiments, Razran (1961) formulated six
20 propositions on interoceptive condition (see box 1) that have since prevailed in the literature (e.g.,
21 Bouton et al., 2001; Ceunen, 2016; Cameron, 2001; Pappens et al., 2013). The propositions that
22 interoceptive conditioning (1) occurs largely unconsciously, and (2) is more resistant to extinction
23 than extero-exteroceptive conditioning, are especially well known and have regularly been
24 mentioned in more recent papers.

25

INTEROCEPTIVE FEAR CONDITIONING

----box 1 with 6 proposition (p97 in Razran, 1961)-----

- "1. Unlike the continuum of exteroceptive stimulation which is the body material of all our conscious experiences, the continuum of interoceptive stimulation leads largely to unconscious reactions.
2. Interoceptive conditioning, whether involving conditioned or unconditioned interoceptive stimuli, is readily obtainable and is by its very nature largely unconscious in character.
3. While interoceptive conditioning is by its nature more limited than is exteroceptive conditioning with respect to total kinds and variety of stimulations, the interoceptive kinds of stimulations are, on the other hand, by their very nature much more recurrent, periodic, and organism-bound, making interoceptive conditioning an almost built-in function that is constantly generated and regenerated in the very process of living and acting.
4. Interoceptive conditioning is somewhat slower in formation than is exteroceptive, but, once conditioned, it is more fixed and irreversible (less readily extinguished).
5. When equal but opposing interoceptively produced and exteroceptively produced reactions are juxtaposed, the interoceptive reactions dominate the exteroceptive ones, with the final result that preceding exteroceptive stimuli become conditioned stimuli for succeeding interoceptive reactions whereas preceding interoceptive reactions become strengthened by exteroceptive reactions succeeding them.
6. The juxtaposition of conditioned interoceptive and exteroceptive stimuli of the same conditioned reaction, unlike similar juxtapositions of exteroceptive stimuli of different modalities, produces a certain amount of conflict and decrementation of conditioning."

The Russian work on interoceptive conditioning clearly laid the foundation for later work that provided more convincing evidence for associative learning involving interoceptive stimuli. The research questions and responses studied in these early studies have been recognized by many as highly relevant to the understanding and malleability of responses and behaviors that occur in natural settings and that matter in real life. The next paragraphs will focus on interoceptive conditioning of fear in the context of panic disorder, although interoceptive conditioning has been studied in relation to many outcomes other than fear, some examples of which can be found in box 2.

----box 2: Examples of interoceptive conditioning in response systems other than fear-

Especially extero-interoceptive pairings with biologically significant USs have been commonly used, generating evidence that such learning can affect a diverse range of response systems. A few major examples are:

- (1) Baroreceptor stimulation as the US has been shown to generate conditioned cardiovascular responses and cortical inhibition (Dworkin & Dworkin, 1995; Rau, Elbert, Lutzenberger, & Eves, 1988).

INTEROCEPTIVE FEAR CONDITIONING

(2) A range of interoceptive USs causing sickness (e.g., liposacharide, lithiumchloride) install conditioned aversion, anticipatory nausea and hyperalgesia (Garcia, Hankins, & Rusiniak, 1974; Limebeer et al., 2008; Wirtelak et al., 1994). Immunosuppressive drugs (US) paired with a certain taste (CS) were found to generate conditioned immunosuppression (e.g., Wirth et al., 2011).

(3) Several other types of drugs as the US (e.g., opiates, alcohol, nicotine) have been applied in Pavlovian conditioning procedures to study anticipatory reactions to the drug in response to cues associated with drug intake (CSs); such learning contributes to drug tolerance, craving and addiction (e.g., Siegel, 1989; Skinner & Aubin, 2010; Ramsay & Woods, 1997).

(4) Water loads have served as a US in studies on conditioned diuresis in humans (Hofer & Hinkle, 1964).

(5) Appetite, flavor preference, food intake and blood glucose regulation can be modulated by Pavlovian conditioning procedures with nutrients or insuline as USs (e.g., Dworkin, 1993; Sclafanik & Ackroff, 2012; Uematsu et al., 2010; Woods & Ramsay, 2000).

(6) Conditioned respiration has been documented in studies using inhalation of CO₂-enriched, hypoxia or loaded breathing as USs (Fannes et al., 2008; Gallego & Perruchet, 1991; Zaman, Van den Bergh, Fannes, & Van Diest, 2014). CO₂-inhalation, hyperventilation and loaded breathing as a US can generate also conditioned stress-induced analgesia (Mongeluzi, Rosellini, Leu, Caldarone, & Stock, 2003) as well as placebo effects in self-reported respiratory symptoms (e.g., De Peuter, et al., 2005; Van Diest et al., 2006; Vlemincx, Sprenger & Büchel, 2017).

3. Integrating biological and learning accounts of panic disorder

Two key characteristics of panic disorder are (1) recurrent, unexpected bouts of extreme fear (i.e., panic attacks), and (2) substantial concerns and debilitating anxiety about such attacks. Whereas studies in the field of psychiatry have typically sought to identify biological causes of panic attacks, psychological research has been directed relatively more at the origins and consequences of, and treatment options for anticipatory anxiety in the context of panic attacks.

3.1 Panic attacks

A panic attack can be conceptualized as a hard-wired, defensive alarm reaction. In the case of spontaneous panic attacks, this alarm is apparently triggered without any obvious reason and is therefore sometimes referred to as a 'false alarm' (Barlow, 1988). Panic attacks are very common in the general population (Kessler et al., 2006) and are not considered pathological by definition. How panic attacks come about has puzzled many researchers over the past decades and is currently still a matter of debate. Perhaps because many symptoms of a panic attack (e.g., shortness of breath, dizziness, cardiorespiratory arousal, fear) can be reproduced by elevating levels of CO₂ in the

INTEROCEPTIVE FEAR CONDITIONING

1 blood (hypercapnia), researchers have hypothesized that an excessive sensitivity to CO₂ and
2 concomitant pH changes in the brain constitute the neurobiological basis of panic attacks (e.g.,
3 Woods et al., 1986; Klein, 1993). In support of this idea, persons suffering from panic disorder as
4 well as their healthy first-degree relatives do indeed seem very sensitive to laboratory challenges
5 that disrupt the acid-base balance (Perna, Gocchi, Bertani, Arancio, & Bellodi, 1995; van Beek, &
6 Griez, 2000), although the specificity and sensitivity of such challenges are a classical matter of
7 debate (e.g., Rappaport et al., 2017). In the past decade, work in animals has convincingly
8 demonstrated that changes in central CO₂ and pH can be sensed in several brain structures other
9 than those involved in brainstem ventilatory control. The baso-lateral amygdala of mice has
10 abundant acid-sensing ion channels that are crucially involved in both conditioned and innate fear
11 (Ziemann et al., 2009, Leibold et al. 2015), and the bed nucleus of the stria terminalis of mice is
12 also crucially involved in CO₂-evoked anxiety behaviors (Taughner et al., 2014). In humans, genetic
13 variations in the sensitivity of such ion channels have been found to relate to differential sensitivity
14 to CO₂ challenges (Leibold et al., 2017). Other evidence suggests that CO₂ hypersensitivity can be
15 distinguished from general trait anxiety and may result from gene-environment interactions
16 associated with early life stress (Battaglia, 2017; Giannese et al., 2018). These and other findings
17 have sparked a renewed interest in the hypothesis that acid-base dysregulations may in fact
18 underlie the occurrence of apparently spontaneous panic attacks and may even be key to panic
19 disorder (e.g., Esquivel, Schruers, Maddock, Colasanti, Griez, 2010; Vollmer, Strawn & Sah, 2015,
20 Leibold et al. 2015).

3.2 Learning about panic attacks

22 A change in pH that unconditionally activates the fear circuitry and can generate panic
23 qualifies as a biologically relevant event with excellent US-UR properties that may easily become
24 involved in associative learning processes. After two decades characterized by a predominantly
25 cognitive account of panic disorder,, a theoretical review paper by Bouton, Barlow and Mineka

INTEROCEPTIVE FEAR CONDITIONING

(2001) re-introduced and further elaborated on the hypothesis that conditioning plays a pivotal, etiological role in panic disorder (see also Barlow, 1988; Goldstein & Chambless, 1978; Wolpe & Rowan, 1988). In essence, panic attacks would cause the conditioning of panic or anxiety to cues in the internal or external environment associated with the panic attack. Following such learning, anxiety occurs in the presence of the conditioned cues, which can motivate avoidance behavior and precipitate another panic attack. Whereas learned anxiety to exteroceptive cues likely underlies agoraphobic fear and avoidance, panic attacks that apparently come out of the blue may result from fear learning to interoceptive cues. Fear of bodily sensations of anxiety/arousal is also sometimes referred to as 'fear of fear' (Goldstein & Chambless, 1978). The hypothesis is that sensations of arousal that precurse a panic attack can act as CSs that are associated with more intense arousal, fear and even panic (Bouton et al., 2001; Cameron, 2001). In line with this, Meuret and colleagues (2011) documented altered respiratory rhythms and changes in CO₂ pressure prior to the onset of 'spontaneous' panic attacks in patients with panic disorder, suggesting that attacks that seem to come 'out of the blue' on an experiential level are in fact preceded and may potentially be triggered by interoceptive, respiratory events.

It has been hypothesized that 'conditioned' panic attacks are an example of intra-event or homoreflexive learning, in which the early onset of an event becomes associated with later aspects of the same event (Bouton et al., 2001; Sinha et al., 2000). Homoreflexive learning has been demonstrated for outcomes other than fear and is generally thought to yield strong conditioning (Bouton et al., 2001), which is consistent with abundant evidence showing that associative learning occurs more rapidly to CSs that are natural precursors of the US than to CSs that lack an obvious connection to the US (Domjan, 2005). The hypothesis that interoceptive fear learning, as an instance of intra-event learning, may generate conditioned fear and panic has been featuring in learning accounts of panic disorder for quite some time, but attempts to empirically model such learning are relatively recent. In contrast, convenient, extero-exteroceptive paradigms have

INTEROCEPTIVE FEAR CONDITIONING

dominated the human fear conditioning literature for decades. In the next section, we discuss alternative responses and stimuli that could be used to study learning processes relevant to panic and bodily symptoms of arousal.

4. Towards panic-relevant conditioning paradigms

Some of the interesting alternative approaches to studying associative learning in the context of panic disorder include fear conditioning paradigms with biologically significant USs, the selection of CSs that show a natural relationship with such US, and a reasoned, a priori choice of the response system under study.

4.1 Conditioned Stimulus

As outlined above, interoceptive sensations originating from a cardiorespiratory activity have been hypothesized to be natural precursors of a panic attack (Bouton et al., 2001; Cameron, 2001). Evidence from ambulatory studies support this idea (Meuret et al., 2011). Furthermore, it is well described that respiratory distress is a prominent feature in (many) patients with panic disorder (e.g., Biber & Alkin, 1999; Telch, Jacquin, Smits, & Powers, 2003) and that this distress motivates them to avoid aerobic exercise (Broocks et al., 1997). Interoceptive exposure using respiratory challenges constitutes a major, effective component of cognitive-behavioral treatments for panic disorder (Brown & Craske, 2017; Deacon et al., 2013; Lee et al., 2006; Meuret et al., 2018).

Methods used to evoke respiratory sensations during interoceptive exposure can also be used to produce panic-relevant CSs in laboratory paradigms. Obviously, a certain level of experimental control over the induced respiratory sensation is required if one aims to implement it as an interoceptive CS, manipulate its pairing with a US, and measure psychophysiological responses to it. External respiratory resistances ('loads') are interesting in this respect. Adding external resistive loads to a breathing circuit introduces a calibrated level of resistance to breathing, meaning that more effort from the respiratory muscles is required to maintain flow rate and

INTEROCEPTIVE FEAR CONDITIONING

1 volume. Resistive loads can come in different intensities and can be applied for different durations.
2 Lower intensities (e.g., those just above perception threshold) applied for a short duration are
3 generally not experienced as unpleasant or arousing, whereas higher intensities for multiple breaths
4 are (Pappens et al., 2010; Alexander-Miller & Davenport, 2010). Another type of respiratory load
5 is the pressure threshold valve (Sapienza & Davenport, 2002), which has an adjustable spring that
6 remains open as long as sufficient pressure is produced to overcome the spring's strength. In other
7 words, the subject has to generate a minimum pressure throughout the entire breath to be able to
8 breathe. A potential and important caveat when applying respiratory loads to experimentally induce
9 respiratory sensations, is that the sensation inherently depends on the participant's ongoing
10 breathing behavior. As an example, when one breathes with great force through a minimal
11 pressure relief valve, the valve will open instantaneously and the resulting sensation will be
12 extremely weak or even absent. In the case of resistive loads, the respiratory sensation will become
13 more intense with an increased flow rate and will be absent during a pause in the respiratory cycle
14 (zero flow). Whenever it is important to standardize the intensity of the respiratory stimulus, it is
15 sensible to instruct participants to aim for a pre-defined target level of peak airflow during each
16 breath (e.g., Alexander-Miller & Davenport, 2010). As an alternative, spontaneous breathing
17 behavior can be monitored, measured and reported, which allows to document potential variations
18 in breathing patterns that could produce associated variations in the intensity of the respiratory
19 input stimulus.

4.2 Conditioned Response

21 Prototypical, extero-exteroceptive fear conditioning paradigms study a rather conventional
22 standard set of CRs, including the potentiated startle blink reflex, skin conductance responses, self-
23 reported fear, or an increased expectancy of the US occurrence (Lonsdorf et al., 2017). Although
24 this standard set of fear-related CRs certainly reflects important facets of (learned) fear, it is mainly
25 representative of the so-called 'post-encounter' phase on the continuum of defensive responding,

INTEROCEPTIVE FEAR CONDITIONING

during which an organism reduces physical activity and directs attention to an environmental stimulus (Lang & Bradley, 2013). Attentional responses are sometimes characterized by a decrease in cardiorespiratory activation (Denot-Ledunois, Vardon, Perruchet, & Gallego, 1998; Fokkema, 1999; Obrist, 1969; Stekelenburg & Van Boxtel, 2002; Van Diest et al., 2009). Panic attacks, in contrast, are thought to reflect acute, massive bursts cardiorespiratory arousal and energy mobilization to support overt, action-oriented defensive responses (fighting, fleeing) to imminent threat. To better understand how associative learning can modulate the occurrence of panic attacks, it is likely more fruitful to search for laboratory paradigms that generate and measure CRs that can be mapped onto the action-oriented, circa-strike mode of the defensive cascade. Interesting candidate CRs are an inhibition of the startle reflex, hyperventilation, and an increased cardiorespiratory activation (Krause et al., 2018; Löw, Weymar, & Hamm, 2015; Van Diest et al., 2009; Van Diest et al., 2001ab).

Relative to other CRs, fear-conditioned increases in respiration may be of particular relevance in the context of panic disorder. In the case of exteroceptive fears (e.g., spider phobia), the fear response (CR) will not facilitate the occurrence of the US (fear does not make the spider crawl towards and onto you). In contrast, fear-induced cardiorespiratory activation as a CR can elicit or aggravate the feared CSs (sensations of arousal) and make them culminate in a panic attack. Thus, in persons with interoceptive fear, a vicious circle of increasing fear and incremental cardiorespiratory sensations of arousal can easily occur. Consistent with the idea that respiratory activity plays a role in maintaining panic disorder, capnometry-assisted respiratory training is an effective treatment to reduce panic symptoms and restore respiratory dysregulation via respiratory-specific mechanisms (Han et al., 1996; Meuret, Rosenfield, Seidel, Bhaskara, & Hofmann, 2010).

4.3 Unconditional Stimulus -Unconditional Response

Relative to other psychiatric disorders, persons with panic disorder have a higher life-time and childhood prevalence of respiratory diseases and report more traumatic suffocation histories,

INTEROCEPTIVE FEAR CONDITIONING

such as near-drowning experiences or having had an object stuck in the throat causing fear of choking to death (Campbell, Sanderson, & Lavery, 1964; Zandbergen & Bright, 1991). This suggests that a suffocation experience may act as a US causing an immediate panic response (UR). Because adequate respiratory function is vital, multiple receptors and pathways can signal a threat to respiration and some of these receptors can be experimentally stimulated in the laboratory within the necessary ethical considerations and limitations. Several respiratory challenges with a panicogenic potential have been used in the laboratory, including hyperventilation, CO₂-inhalation (see box 3) and, more recently, mechanical interference with breathing. An element common to all these challenges is that they induce dyspnea (or breathlessness), which persons with panic disorder or increased anxiety sensitivity and/of fear of suffocation typically dislike and fear more than other persons do (Melzig, Holtz, Michalowski, & Hamm, 2011; Pappens, Smets, Van den Bergh, & Van Diest, 2012b; Van den Hout, van der Molen, Gries, & Lousberg, 1987; Sinha et al., 2012). Whether the fear reaction to this type of respiratory challenge is learned or whether it reflects an unconditional response (or a combination of both) is hard to disentangle but, at least for some of these challenges, neurobiological theories on panic disorder have claimed that the fear reaction is unconditional (e.g., Klein, 1993).

4.3.1 CO₂. Single or double inhalations of higher concentrations (e.g., 35%) of CO₂ are panicogenic in vulnerable persons (Schruiers et al., 2011), but require the active and voluntary compliance of the subject. Because high concentrations of CO₂ have an immediate and strong effect on the nervus trigeminus, causing a salient, aversive sensory experience (Gallego & Perruchet, 1991; Moessnang et al., 2013), a full inhalation can only be performed when participants actively overcome their reflexive tendency to withdraw from the mask or mouthpiece. This requires instructions to first exhale as deeply as possible, subsequently inhale deeply and hold their breath for a few seconds before exhaling again, before performing the full inhalation (e.g., De Cort et al., 2017). Lower concentrations of CO₂ do not generate panic acutely, but have the advantage

INTEROCEPTIVE FEAR CONDITIONING

that they can be applied for a longer time and more unobtrusively. The unconditional effects of CO₂-inhalation have been studied extensively and are described in more detail in box 3. In short, dose-response relationships have been documented for self-reported symptoms of general arousal, fear, panic, and for increases in ventilation and pCO₂, and for responses reflecting sympathetic activation including skin conductance levels and blood pressure (e.g., Leibold et al., 2013; Pappens, De Peuter, Vansteenwegen, Van den Bergh, & Van Diest, 2012a). Findings on startle blink EMG seem counterintuitive at first because, despite the presence of subjective fear, there is a lack of fear potentiated startle blink right after inhalation of CO₂, and a pronounced inhibition of startle during inhalation of CO₂ (Ceunen, Vlaeyen, & Van Diest, 2013; Pappens et al., 2012a; Pinkney, Wickens, Bamford, Baldwin, & Garner, 2014). Together, the described set of unconditional responses to CO₂-inhalation map well to the action-oriented, circa-strike side of the defensive continuum and support the idea that CO₂-induced changes in blood pH strongly and unconditionally activate the fear circuit in humans.

4.3.2 Obstructed breathing. Whereas CO₂-inhalation is a chemical stimulus acting upon chemoreceptors, respiration-related mechanoreceptors can also be experimentally stimulated to activate the fear circuit. Several studies have documented defensive activation (startle potentiation, cardiorespiratory activation, elevated skin conductance) *in anticipation of* obstructed breathing (e.g., Krause et al., 2017; Lang et al., 2011; Pappens et al., 2013; Schroyen et al., 2016). The actual application of resistive loads of a moderately high intensity or a complete breathing obstruction (infinite resistance) for a period of time induces subjective distress, fear and dyspnea and activates dyspnea-relevant sensorimotor and cortico-limbic brain regions (Alexander-Miller & Davenport, 2010; Benke, Hamm, Pané-Farré, 2017; Esser et al., 2017; Laviètes et al., 2000; Livermore et al., 2008; von Leupoldt et al., 2007). Pappens and colleagues (2010; 2011) were the first to investigate startle potentiation and SCR *during* loaded breathing. In two studies, they compared affective and psychophysiological responses during the viewing of aversive pictures with those evoked during

INTEROCEPTIVE FEAR CONDITIONING

breathing through two low-intensity inspiratory resistive loads applied for only one respiratory cycle. Findings of both studies indicated similar skin conductance responses (SCR) and low-moderate levels of subjective fear to both the loaded inspirations and the presentation of an unpleasant picture. Surprisingly, a paradoxical lack of startle potentiation during the most intense and unpleasant resistive load was observed in the first study, but not in the second study where participants concurrently engaged in an exteroceptive reaction time task. Therefore, fear-potentiated startle during loaded breathing seems to show up only when - despite the interoceptive experience -one keeps on directing attention to the external environment. This interpretation is supported by findings showing that the P300 to the acoustic startle probe is reduced during loaded compared to unloaded breathing (Alius et al., 2015). Other studies have replicated the lack of startle potentiation during loaded breathing and have documented similar findings with other types of aversive interoceptive stimulation (e.g., Alius et al., 2015; Ceunen, Vlaeyen, & Van Diest, 2013; Ceunen et al., 2015).

-----box 3: unconditional responses to CO₂ inhalation-----

Studies on acute fear and panic typically use higher concentrations of CO₂ (13% - 65%), whereas longer inhalations of lower concentrations (4-10%) have been applied in other studies, including on sustained arousal, stress and symptom perception (e.g., Rapee, Brown, Antony, & Barlow, 1992; Van den Bergh et al., 1995).

Experiential level. Depending on the concentration and the duration, CO₂ inhalation is known to escalate symptoms of arousal, partially mimicking the topography of human fear responses: deeper and/or faster breathing, dizziness, a sense of dyspnea (breathlessness), a faster and/or stronger heart beat, feeling of unreality, etc. (Pappens et al., 2012a). Responses to CO₂-inhalation show a large variability, not only between diagnostic groups, but also within groups and between similar groups in different studies. Baseline anxiety preceding the challenge is a relevant factor, as instructions can modulate reactivity to the challenge in non-clinical participants (Rapee, Mattick, Murell, 1986; Van der Molen & Van den Hout, 1988), and threatening instructions are sometimes even implemented to maximize the aversive value of the US (e.g., Forsyth, Eifert & Thompson, 1996; Forsyth & Eifert, 1998). Studies have demonstrated a relationship between physiological indices of baseline anxiety and subsequent subjective fear responses to biological challenges (Sinha et al., 2000). A range of trait-like fear-related variables have also been found to predict reactivity to CO₂-inhalation, including anxiety sensitivity, avoidant coping style, suffocation fear, and behavioral inhibition sensitivity (Blechert, Wilhelm, Meuret, Wilhelm, & Roth, 2013; Feldner, Zvolensky, Eifert, & Spira, 2003; McNally & Eke, 1996; Rassovsky, Kushner, Schwarze, & Wangenstein, 2000; Zinbarg, Brown, Barlow, & Rapee, 2001; Zvolensky, 2001). Another factor that probably contributes to

INTEROCEPTIVE FEAR CONDITIONING

ariability in subjective responses to CO₂-inhalation is the lack of a standardized set of stimuli (in terms of concentration, duration and method of application) or a standardized way to measure the responses, which often makes reports from different studies difficult to compare.

Respiration. Under normal circumstances, CO₂ is the main chemical drive to breathe, so ventilation increases reflexively during inhalation of CO₂. Initial increases in ventilation result mainly from an increase in tidal volume; an increase in breathing frequency typically occurs only later on, when a further increase in volume would become less cost-effective (Tipton et al., 2017). Inhalation of CO₂-enriched air may also affect physiological activity indirectly, via the fear circuit. Interestingly, evidence suggests that, during CO₂-inhalation or during exercise, the breathing frequency component makes a larger contribution to the total increase in ventilation in fearful subjects than in non-fearful subjects (Blechert et al., 2013; Masaoka & Homma, 1999; Pain, Biddle, & Tiller, 1988). This is consistent with findings that, compared to tidal volume, respiratory timing parameters are less tightly controlled by the chemical drive to breathe (Rafferty & Gardner, 1996) and states of fear and anxiety are known to be associated with more rapid breathing (Gardner, 1996; Masaoka & Homma, 1999; Tipton et al., 2017; Van Diest et al., 2001ab). It is as yet unclear whether the larger contribution of the breathing frequency component to the total ventilation of fearful subjects is an automatic response or an instance of fear-motivated operant behavior. Escape behavior occasionally occurs in the laboratory when participants stop the experiment and take off the mask. Attempts to avoid or minimize the inhalation have also been described. Specifically, it has been suggested that fearful subjects may try to limit the amount of inhaled CO₂ by deliberately breathing more shallowly and more rapidly (Sinha et al., 2000). Other findings suggest that some participants try to 'sabotage' the inhalation by deliberately creating leaks in the breathing circuit or by inhibiting their breathing when the CO₂ comes on (Fannes et al., 2008), which may lead to a 'paradoxical decrease in ventilation' (Rassovsky & Kushner, 2003).

Physiological responses other than respiration. Inhalation of CO₂-enriched air influences pH-regulated physiological processes in the body. For example, it is known to be a potent dilator of the cerebral arterioles and to cause a concomitant rise in cerebral blood flow. Relatively short inhalations (up to 2 min) at lower concentrations (7.5-20%) typically trigger increases in heart rate and skin conductance, but not necessarily in release of salivary cortisol (Coplan et al., 2002; Pappens et al., 2012a). Longer inhalations of 7.5% CO₂ (10-20 min) are associated with rises in systolic blood pressure, ventilation and heart rate, and with increased plasma levels of epinephrine, norepinephrine and 17-OH corticosteroids (Bailey, Argyopoulos, & Kendrick, 2005). A single vital capacity inhalation of 35%-CO₂-enriched air in healthy subjects induces significant bradycardia, sympathetic activation and, in some but not all studies, also increases in cortisol (e.g., Argyopoulos et al., 2002). With an increasing concentration of CO₂, double inhalations of 35% CO₂ are associated with a dose-dependent increase in diastolic and systolic blood pressure, but not in heart rate (Leibold et al., 2013). The eyeblink startle response is progressively inhibited during inhalation of 7.5% and 20% CO₂-enriched air, possibly indicating a sensory shut down at the action-oriented 'circa-strike' phase of the defensive continuum (Ceunen, et al., 2013; Krause et al., 2018; Pappens et al., 2012a).

5. Human interoceptive fear conditioning studies with panic-relevant stimuli: an overview

5.1 Studies with CO₂-inhalation as US

INTEROCEPTIVE FEAR CONDITIONING

Fear conditioning studies in humans with CO₂ as a US were identified based on a Pubmed search (April 2018) combining the following search terms in the title or abstract: (1) 'carbon dioxide' or 'CO₂' (2) 'conditioning', 'conditioned', 'learning' or 'learned', and (3) 'fear' or 'anxiety'. This search yielded 41 publications, from which 9 papers were judged to be relevant based on the following criteria: fear conditioning study on human subjects in which contingency between CS and US can derived only from experience (no instructed contingencies), including at least one physiological outcome as well as a self-reported fear-relevant outcome, CO₂-inhalation as the US. Three additional papers were identified by means of a snowball search strategy. From this list of 11 studies, 8 studies used exteroceptive CSs, whereas 3 studies interoceptive CSs.

5.1.1 Extero-interoceptive paradigms. Table 1 presents an overview of the studies that paired CO₂ as the US with exteroceptive CSs. All eight studies were performed on healthy volunteers and six studies used inhalations of 20% CO₂-enriched air as the US. In five of the latter studies, all by Forsyth and colleagues, video fragments, nonwords or geometric shapes served as CSs in various extero-interoceptive preparations; five or six seconds after onset of these CSs, the CO₂ was administered and both stimuli co-terminated 20 s later (Finlay & Forsyth, 2009; Forsyth & Eiffert, 1998; Forsyth et al., 1996; Valdivia-Salas, Forsyth, Berghoff, & Ritzert, 2014). Another study by Fannes et al. (2008) used odors as CSs in combination with 20% CO₂. The remaining two studies used mental imagery scripts as CSs, paired with either a concurrent inhalation of 5.5% CO₂ (Stegen, De Bruyne, Rasschaert, Van de Woestijne, & Van den Bergh, 1999) or a subsequent inhalation of 35% CO₂ (De Cort, Griez, Büchler, Schruers, 2012). Therefore, except for the study by De Cort et al. (2012), one could argue that the paradigms employed, though still using exteroceptive CSs, model some of the aspects of intra-event learning in which CSs and a panic-relevant US are close together in time and partially or completely overlap. Several insights emerged from this line of studies.

INTEROCEPTIVE FEAR CONDITIONING

----- Insert Table 1 around here-----

First, 20% CO₂ is a potent US that, when paired with exteroceptive CSs, reliably generates conditioned fear expressed in self-reported distress, panic symptoms, evaluative ratings, skin conductance and heart rate. The apparent potency of 20% CO₂ as a US to establish fear learning contrasts with the overall weak findings from the study with a single 35% CO₂ inhalation as the US (De Cort et al., 2012). In the latter study, differential conditioning could not be convincingly demonstrated, as only the between subject effects comparing responses to various CS+s provided some support for a selective conditioning effect to the fear-relevant interoceptive emotional imagery script. A potential explanation for the overall weak effects is that participants had to self-administer the 35% CO₂ US (by complying with an instructed inhalation maneuver, see box 4). Self-administration of the US may drastically reduce the surprise element (and prediction error) which, according to traditional associative theories (e.g., Rescorla & Wagner, 1972), is a major determinant of learning.

Second, although heart rate was no longer studied in the later studies of Forsyth and colleagues (Finlay & Forsyth, 2009; Valdivia-Salas et al., 2014), their initial two studies with 20% CO₂ as a US consistently observed a conditioned cardiac acceleration in response to the CS (Forsyth & Eiffert, 1998; Forsyth et al., 1996).

Third, all three studies that have looked at respiration as a CR observed significant conditioning effects in respiration (De Cort et al., 2012; Fannes et al., 2008; Stegen et al., 1999), thereby underscoring that conditioned respiration is a likely and 'natural' outcome in studies with CO₂ as a US. However, the direction and nature of the observed respiratory learning effects vary remarkably. Whereas Fannes et al. (2008) reported on a conditioned decrease in ventilation, Stegen et al. (1999) demonstrated fear conditioning and a conditioned increase in respiratory rate only to fear-relevant CSs. De Cort et al. (2012) reported a conditioned increase in tidal volume in their

INTEROCEPTIVE FEAR CONDITIONING

study. The inconsistent findings regarding changes in respiration as a conditioned response may result from the approach of averaging all respiration cycles across the entire CS window (typically 2 min), which may wipe out potentially opposing dynamic changes in breathing behavior over time. To explore this possibility, Fannes et al. (2008) studied changes in respiration over time in consecutive 10 s windows. Their results showed that a substantial number of participants learned to actively avoid the CO₂ by inhibiting their breathing behavior early on in the CS+ acquisition trials (Fannes et al., 2008). Although CO₂-enriched air is thought to be odor- and tasteless (especially at lower concentrations), the findings of this study strongly suggested that participants could taste/sense the CO₂ in the first few seconds of inhalation. Stegen et al. (1999), who observed not a conditioned decrease but a conditioned increase in ventilation, used a low concentration of CO₂ (5.5%), which is likely much harder to detect via taste. Therefore, whether and to which extent CO₂ can be tasted is an important methodological question that may shed light on the inconsistent pattern of findings in conditioned respiration.

Fourth, with CO₂ as a US, fear is more easily acquired and is more difficult to extinguish to fear-relevant than to fear-irrelevant CSs, and this is especially true for physiological CRs (Forsyth & Eifert, 1998; Forsyth et al., 1996). In a similar vein, selective conditioning effects have also been found in studies with different concentrations of CO₂ as US and mental imagery as the CSs (Stegen et al., 1999; De Cort et al., 2012 - Table 1). Fear conditioning in the latter studies could only be established to claustrophobic emotional scripts, but not to emotionally neutral or fear-irrelevant imagery scripts. All this is consistent with findings from extero-exteroceptive designs (e.g., Cook, Hodes, & Lang, 1986).

Fifth, Forsyth & Eifert (1998) found that conditioned heart rate acceleration and evaluative ratings -but not the other conditioned responses- showed more resistance to extinction when learned to an 'internal' CS depicting an arrhythmically beating human heart than when learned to an 'external' CS (video of a snake). Although speculative and hard to falsify, the latter effect can be

INTEROCEPTIVE FEAR CONDITIONING

understood by an enhanced 'belongingness' between the CS and the US-UR combination in case of the internal CS, which, as suggested by the authors, may even be more enhanced with 'true' interoceptive sensations as a CS.

5.1.2 Intero-interoceptive paradigms. Table 2 summarizes the three studies with CO₂ as a US that may qualify as intero-interoceptive studies. One study implemented a differential conditioning paradigm, in which the presence and absence of cardiac sensations of arousal caused by infusions of pentagastrine and saline, respectively, served as CSs (De Cort et al., 2017). These CSs were followed by a single inhalation of 35% CO₂ (US) or room air (no US). Conditioning effects were weak overall: they occurred for only one of the outcomes (expectancy ratings, not for self-reported anxiety and SCR), for only the pentagastrine CS+ (not for the saline CS+), and only in participants with lower levels of anxiety sensitivity. Thus, similar to De Cort et al. (2012), evidence for fear conditioning was also rather weak in this study, again perhaps because the US was a self-administered single inhalation of 35% CO₂. Another aspect that may have contributed to the weak findings is that the CSs used by De Cort et al. (2017) differed dramatically in salience: the absence of cardiac sensations of arousal is a less 'distinct' event than the sudden occurrence of such sensations (to 35% CO₂). Creating two comparable interoceptive stimuli as experimental CSs poses a methodological challenge, which may explain why most research on intero-interoceptive fear conditioning has turned to between subject designs, which are also more common in the animal literature.

----- Insert Table 2 around here-----

An interesting intero-interoceptive, homoreflexive, single-cue, between subject paradigm was first developed by Acheson and colleagues. Specifically, these authors used 20% CO₂ as both a CS (when applied for only 5 s) and a US (when applied for 15 s). In the first study (Acheson,

INTEROCEPTIVE FEAR CONDITIONING

Forsyth, Prenoveau & Bouton, 2007), three groups were run: one in which the CS was followed by the US (paired group), one in which the CS was never followed by the US (although the US was presented separately; explicitly unpaired group), and one group that received only CSs and no USs (CS-only group). Compared to the CS-only group, fear to the CS was greater in both the paired and unpaired groups in the later acquisition trials. The unpaired group displayed even greater fear to the CS than the paired group, and the enhanced fear responding to the CS was later extinguished in the paired, but not the unpaired group. To explain these unexpected findings, Acheson et al. (2007) argued that the unpaired procedure could actually be considered a partial reinforcement schedule in which CO₂ onset was 50% reinforced by a prolonged CO₂ inhalation. In other words, at CO₂ onset, participants were unable to discriminate whether the CO₂ would be brief and physiologically inert (inhalation of 5s; the CS) or sustained and physiologically prepotent (inhalation of 15s; US). In a follow-up study, Acheson, Forsyth and Moses (2012) replicated their 2007 study and extended it with a fourth, 'explicitly unpaired' group that received an additional exteroceptive cue, allowing participants to discriminate the CS from the US at CO₂ onset. The findings from the first study were replicated. Adding the discrimination cue in the explicitly unpaired group dramatically decreased fearful responding to the CS, thereby supporting the authors' hypothesis that fearful responding in the original unpaired group had been due to the participants' inability to discriminate between the CS and the US at CO₂ onset. An intriguing question, in this respect, is how participants could perceive the CS at all. As noted by Acheson and colleagues (2007), 5 s of 20% CO₂ is physiologically inert and therefore cannot produce perceptible interoceptive sensations. Possibly, participants could detect the CO₂ onset in their studies by its taste properties (see also Fannes et al., 2008), making the taste an *exteroceptive* cue that predicted the later interoceptive effects of CO₂-inhalation more reliably than any other CS that could be experimentally introduced.

5.1.3 Discussion. Consistent with the immediate threat to the body imposed by excessive levels of CO₂, the reviewed studies show that inhalation of CO₂ is a potent US to establish fear

INTEROCEPTIVE FEAR CONDITIONING

conditioning to a variety of CSs. Partial overlap between CS and US is a feature of intra-event learning that has been said to characterize learning processes in panic disorder (Bouton et al., 2001). Indeed, these studies suggest that fear learning seems to take place rather quickly (after only 3 acquisition trials in many of the studies, see Table 1), and in a diverse range of outcomes (including self-reports and physiological outcomes), to CSs that partially overlap with the CO₂ US. Notably, cardio-respiratory activation was also conditioned in several studies, which supports the idea that using 20% CO₂ as a US generates CRs that prepare for defensive action (heart rate acceleration), rather than a 'mere' attentional response to the CS (heart rate deceleration).

The consistent findings on conditioned heart rate acceleration with CO₂ as a US contrast with findings from more commonly used paradigms that use panic-irrelevant exteroceptive USs (e.g., shock, aversive noises). In the latter type of studies most participants show conditioned heart rate deceleration (e.g., Hodes, Cook, & Lang, 1985; Moratti, Keil, & Miller, 2006). Besides cardiorespiratory activation, an inhibition of the startle blink reflex is also thought to characterize defensive responding in the action-oriented phase of the defensive continuum. Interestingly, unconditioned responses to CO₂-inhalation involve an inhibition of the startle blink too (Pappens et al., 2012a). It is therefore surprising to see that none of the studies in Tables 1 and 2 has incorporated startle blink EMG responses as an outcome of fear learning. It remains thus to be seen whether the CO₂ paradigms are able to generate startle inhibition as a CR, which would lend further support to the idea that CO₂ as a US is likely to generate learned action-oriented defensive responses.

Several methodological considerations are relevant. A general challenge of implementing CO₂ inhalation in fear conditioning paradigms is the relative lack of experimental control. The time it takes for the CO₂ to reach the mouthpiece or face mask depends on the breathing equipment used (e.g., length and diameter of breathing tubes) and the actual breathing pattern of the participant. Even when the time when the CO₂ reaches the participant is known, the actual onset of CO₂-

INTEROCEPTIVE FEAR CONDITIONING

induced symptoms of arousal cannot be controlled and is highly variable, which makes it difficult to standardize the time between CS and US onset within and between participants.

Furthermore, fear learning can be expected to be maximal with a potent US and maximal prediction error, but there seems to be a trade-off between both aspects in the case of CO₂ as a US. Whereas the panicogenic properties of CO₂ rise with increasing concentrations of CO₂, successfully inducing panic-like symptoms 'out of the blue' (without taste cues preceding the symptoms, or without having to rely on self-administration and a voluntary inhalation maneuver) is very difficult with higher concentrations of CO₂. The problem of the taste properties of inhaling a prepared CO₂-enriched air mixture is also relevant if one aims to establish a 'truly' intero-interoceptive paradigm with CO₂ as the CS or US. Further work is needed to establish the concentrations of CO₂ for which people can detect the CO₂ via its taste. Until then, a promising alternative US may be re-breathing, in which one inhales and exhales the same air from a bag, leading to gradual build-up of CO₂ (Read, 1967) and a concomitant increase in breathlessness and fear (Wan et al., 2008). Another alternative US that circumvents the problematic taste properties of inhaled CO₂ may be a mechanical limitation to breathe (breathing obstruction).

5.2 Studies with obstructed breathing as US

Human fear conditioning studies that used mechanical limitation of breathing (obstructed breathing) as a US were identified based on a Pubmed search (November 2018) that combined the following search terms in the title or abstract: (1) 'respiration' or 'respiratory' (2) 'conditioning', 'conditioned', 'learning' or 'learned', (3) 'fear' or 'anxiety', (4) 'obstruction' or 'load'. This search yielded 9 publications, of which only 2 papers were judged to be relevant based on the following criteria: fear conditioning study on human subjects in which CS-US contingencies were not instructed, including at least one physiological outcome as well as a self-reported measure of fear; a limitation of breathing as the US. Seven additional reports were identified by means of a

INTEROCEPTIVE FEAR CONDITIONING

snowball search strategy, and one unpublished study was contributed directly by the author. From this list of 10 studies, 3 studies used exteroceptive, whereas 7 studies interoceptive CSs.

5.2.1 Extero-interoceptive paradigms. Table 3 presents an overview of the 3 studies that have paired auditory/visual CSs with a limitation/prevention of breathing as the US. The study of Campbell et al. (1964) applied transient respiratory muscle paralysis (inducing a total respiratory arrest) in alcohol-dependent patients. The procedure was introduced to the patients as a component of their therapy. Obviously, this paradigm and recruitment procedure would raise serious ethical concerns nowadays. The post-experimental verbal descriptions of how these patients experienced the US give a sense of the horrifying experience they must have gone through.

----- Insert Table 3 around here-----

A much weaker and acceptable US - breathing through a strong resistive load or a temporary total occlusion of the breathing circuit - was introduced more recently by Pappens and colleagues, who used it in healthy volunteers. The conditioned respiratory response in anticipation to these USs seemed to involve an increase of ventilation, established by increases in either respiratory rate (Pappens, Schroyen, Van den Bergh, & Van Diest, 2015a) or tidal volume (Pappens et al., 2013). The overall pattern of findings of the studies listed in Table 3 shows rapid fear learning in all studies for *all* of the studied outcomes, including self-reports, respiration, heart rate, startle blink EMG, and skin conductance. Mechanical limitation of breathing, even if only partial and shortlived, seems to be a potent US for establishing fear learning to exteroceptive cues. This is also reflected in the extinction findings. In the two studies with a between group design (Campbell et al., 1964; Pappens et al., 2013), there was no successful extinction for any of the physiological conditioned responses. In the study of Campbell et al. (1964) that used the traumatic and horrifying US, conditioned fear responses failed to extinguish, even after more than 100

INTEROCEPTIVE FEAR CONDITIONING

extinction trials. In the differential study of Pappens et al. (2015a), extinction occurred for all outcomes other than respiratory rate.

5.2.2 Intero-interoceptive paradigms. We found eight intero-interoceptive studies that had used mild respiratory loads as CSs, combined with partial or complete breathing obstruction as the US (see Table 4 for an overview). As both the CS and US are in the respiratory modality and differed only in terms of intensity and unpleasantness, these paradigms closely modeled interoceptive intra-event learning in the context of incremental sensations of breathlessness, as typically experienced during an asthma or panic attack.

----- Insert Table 4 around here-----

In the first type of intero-interoceptive studies, the respiratory load CS is paired with a more intense and aversive load of a relatively long duration as the US ('load-load' paradigm, see left pannel of Figure 1). In the first between subject study with this pairing, Pappens et al. (2013) observed a conditioned increase in startle blink EMG and respiratory rate, but not in self-reports or in SCR. These findings were successfully replicated by Pappens, Vandenbossche, Van den Bergh, & Van Diest (2015b). In the latter study, contingency awareness was also assessed and indicated that participants from the paired group were unaware of the contingencies between CS and US, despite having developed conditioned responses in respiration and startle blink EMG (Pappens et al., 2015b). Because the studies of Pappens et al. (2013; 2015b) demonstrate that interoceptive fear learning can take place without declarative knowledge of the CS-US contingencies, the load-load paradigm was put forward as a laboratory model to study "unexpected" panic attacks (Pappens et al., 2015b). Recently, Benke, Alius, Hamm & Pané-Farré (2018) introduced an interesting, within subject variation of the paradigm that allowed for the study of both cued and contextual conditioning. The relationship of the CS to the US-load was made dependent on variations in

INTEROCEPTIVE FEAR CONDITIONING

exteroceptive contexts. In the 'No threat' context, only the CS load was presented; in the 'Predictable' context, the US load followed the CS, and in the 'Unpredictable' context, the US was unpaired to the CS. These within subject conditions are commonly used in NPU (No/Predictable/Unpredictable threat) tasks, in which the predictable condition typically generates cued fear to the CS, whereas the unpredictable condition contextual fear to the context (e.g., Schmitz & Grillon, 2012; Schroyen et al., 2016). In contrast to the standard NPU task, however, participants were not instructed about the contingencies in the study of Benke et al. (2018). Fear learning to both the interoceptive cue and the exteroceptive context was established for startle blink EMG in the predictable and unpredictable contexts, respectively, but only in individuals scoring in the upper tertile on Fear of Suffocation of the Claustrophobia Questionnaire (Radomsky, Rachman, Thordarson, McIsaac, & Teachman, 2001). Extinction was not tested in this new type of within subject load-load paradigm.

-----insert Figure 1 around here-----

A second paradigm that also uses mild respiratory loads as CSs has used complete breathing obstructions as the US (load-occlusion paradigm, see lower panel of Figure 1). Typically, the duration of the obstruction is individually determined as a percentage of a person's maximal breath-holding time after expiration. Only one of four studies using this US implemented a differential paradigm in which two respiratory loads of a different intensity were used as CS+ and CS- (Schroyen et al., 2015). Findings from this differential study were complex and difficult to interpret because, similar to the study by De Cort et al. (2017), the two CSs were not comparable on

INTEROCEPTIVE FEAR CONDITIONING

several aspects (e.g., perceptual salience, similarity to the US) that may affect conditionability (see footnote 1). More convincing evidence for fear conditioning in the load-occlusion paradigm stems from the studies with a single-cue, between subject (paired/unpaired) design. In contrast to the load-load paradigm where fear learning occurred in only startle blink EMG and respiration (Pappens et al., 2013; 2015b; Benke et al., 2018), the load-occlusion paradigm also yielded conditioning effects in SCR and US-expectancy ratings (Pappens, Smets, Vansteenwegen, Van den Bergh, & Van Diest, 2012c; Pappens et al., 2014; Van Diest, 2017). In the load-occlusion paradigm, however, dissociation between CRs occurred during extinction (which was successful and complete for SCR and ratings), but not for startle blink EMG (Pappens et al., 2012c).

Interestingly, interindividual differences in short-term heart rate variability (HRV) during a resting baseline prior to conditioning were positively associated with extinction learning in startle blink EMG in the paired group (Pappens et al., 2014). Similar findings have been observed in a paradigm with exteroceptive stimuli (Wendt, Neubert, Koenig, Thayer, & Hamm, 2015) and suggest that HRV may index the strength or integrity of prefrontal mechanisms involved in inhibitory learning.

5.2.3 Discussion. Consistent with the direct and vital threat posed by mechanical limitation of breathing, findings from the reviewed studies demonstrate that obstructed breathing constitutes a potent US to establish fear learning to both exteroceptive (Table 3) and interoceptive (Table 4) CSs. Defensive CRs appear rapidly in both self-reports (US-expectancy, fear) and physiological outcomes (SCR, startle blink EMG, respiration). The variations in the paradigms used (see Tables 3-4) suggest that several factors co-determine the strength of conditioning.

First, a total breathing occlusion seems to be a more potent US than a partial breathing obstruction as imposed by a respiratory load. Indeed, in intero-interoceptive paradigms (Table 4), only total breathing occlusion established conditioning at all the levels of conditioned responding that were investigated. extinction is Also slower/less complete in the studies that use complete

INTEROCEPTIVE FEAR CONDITIONING

1 obstruction as the US. Given that a respiratory occlusion is an infinite resistance and, therefore, a
2 resistive load of the highest intensity, these findings are in line with the general principle that more
3 intense/aversive USs yield stronger conditioning.

4 Second, fear learning appears to be stronger and more resistant to extinction in single-cue,
5 between subject paradigms than in within subject / differential paradigms. A possible explanation
6 is that fear inhibition/inhibitory learning (to the CS-) is already active during acquisition in
7 differential paradigms, and facilitates subsequent extinction.

8 Third, conditioning seems to be more difficult to establish in paradigms where the CS is
9 perceptually less distinct from the US. In fact, although the load-load paradigm may model intra-
10 event learning most closely, it generally yields weaker fear learning than paradigms with a more
11 distinct CS, such as the load-occlusion paradigm (Table 4) or the paradigms that combine breathing
12 obstruction with a visual CS (Table 3). Using a CS that is perceptually less distinct from the US in
13 the load-load paradigm may downgrade the salience of the CS. Work in rodents has demonstrated
14 that CRs are less easily acquired with less salient CSs (Rescorla and Wagner, 1972), which may
15 explain why associative learning seems to be more difficult to establish in the load-load than in the
16 other paradigms.

17 Fourth, compared to the visual CSs (see Table 3), the respiratory load CS (see Table 4)
18 scores higher in fear-relevance/ 'preparedness' (Seligman, 1971), and seems to relate naturally to
19 the US (Garcia & Koelling, 1966). Such 'belongingness' between the respiratory load CS and the
20 occlusion US is present both at the sensory level (both are in the respiratory modality) and the
21 semantic level (the partial and benign breathing limitation imposed by the respiratory load
22 intrinsically refers to the inability to breathe). The relatively strong belongingness between the CS
23 and US may explain why, in the load-occlusion paradigm, the explicitly unpaired group fails to
24 learn that the respiratory load CS actually signals a relatively safe period. Indeed, US-expectancy
25 ratings of the unpaired group increased during the respiratory load CS, although to a lesser degree

INTEROCEPTIVE FEAR CONDITIONING

than for the paired group (Pappens et al., 2012c; 2014; see Table 4). Such 'failure' of the unpaired group to learn that the CS is actually safe was not observed if a visual CS was used in combination with an occlusion as the US. In the latter case, the unpaired group's US-expectancy ratings in response to CS decreased by the end of acquisition (Pappens et al., 2013; see Table 3).

Probably the most remarkable finding in the reviewed studies with limitation of breathing as a US is the reasonably consistent dissociation between CRs. In the load-load paradigm, acquisition can be established in startle blink EMG and in respiration, but not in self-reports and SCR (Table 4). In the load-occlusion paradigm (Table 4) as well as the paradigms with visual CSs (Table 3), extinction occurs more slowly (or not at all) in respiration and/or in startle blink EMG than in SCR and self-reports. That is, even though participants are unaware of the contingencies during acquisition (load-load paradigm), or, even though participants realize that the contingency no longer holds during extinction (load-occlusion paradigm), defensive responding to the respiratory CS (that was previously) paired with the US (breathing limitation) was present. Dissociations are not unique for interoceptive USs and have also been observed in more commonly used extero-exteroceptive fear conditioning paradigms. In the latter type of studies, dissociation between CRs seems to occur most typically when using prepared CSs (Öhman & Mineka, 2001), when using noradrenergic agents (e.g., Soeter & Kindt, 2010), or in persons with specific brain lesions (e.g., Bechara et al., 1995). Based on such findings, it has become progressively accepted that US-expectancy and SCR reflects declarative knowledge about the CS-US contingencies, whereas startle blink EMG reflects emotional learning that can occur independent of changes in declarative knowledge (e.g., Hamm & Vaitl, 1996; Sevenster, Beckers, & Kindt, 2014). Others have argued that dissociation between CRs reflects a differential responsiveness of CRs when expressing what is learned (Lovibond & Shanks, 2002). In either case, the load-load paradigm could form an experimental model of 'unexpected' fear/panic, as persons suffering from panic disorder also verbally report that some panic attacks are unpredictable although physiological

INTEROCEPTIVE FEAR CONDITIONING

recordings show that the attacks are actually preceded by changes in cardiorespiratory activation (Meuret et al., 2011).

In the case of a respiratory interoceptive CS and the demonstrated respiratory activation as a CR, such 'unconscious' defensive mobilization may lead to incremental arousal and facilitate a panic attack. Indeed, increased respiratory activation will intensify the experience of the respiratory CSs and make them increasingly more similar to the feared US. In this respect, it is relevant to note that a conditioned respiratory activation occurred in all 5 between subject studies that implemented respiratory measures, corroborating the idea that respiratory activation may indeed foster incremental arousal and even panic (Bouton et al., 2001). The findings on respiratory CRs also suggest that respiration may be the most natural response system to prepare for the aversive respiratory obstruction (US). Obviously, an increased respiration (CR) in anticipation of an upcoming breathing obstruction (US) is functionally adaptive. Cardiovascular CRs can be expected to covary with respiratory activation (Van Diest et al., 2009), but have not yet been investigated in paradigms that use obstructed breathing.

5.3 General conclusions

5.3.1 Respiratory USs are potent. The studies with panic-relevant, respiratory USs reviewed here generally confirm that both CO₂ inhalation and obstructed breathing are potent USs to rapidly establish fear learning in a range of different response systems. Indeed, whereas studies with electrical shocks as US typically implement 5-20 acquisition trials (Lonsdorf et al., 2017), most of the reviewed studies with respiratory USs use (and therefore seem to need) fewer CS-US pairings for fear learning to occur. Knowing that higher concentrations of CO₂-inhalation may not qualify as a purely 'interoceptive' stimulus, the apparent potency of reviewed respiratory USs is more likely to be explained by their biological significance and immediate vital threat than by their interoceptive nature.

INTEROCEPTIVE FEAR CONDITIONING

5.3.2 Differential interoceptive paradigms. Apart from a relatively quick fear learning, the *differential* paradigms using exteroceptive CSs in combination with a respiratory US (see Tables 1 and 3) generate a pattern of findings that is remarkably similar to the typical findings from more commonly used extero-exteroceptive fear conditioning paradigms in humans (e.g., tone - shock). Specifically, when respiratory USs are used in differential paradigms, both acquisition and extinction can be established to exteroceptive CSs for the majority of the outcomes (e.g., Fannes et al., 2008; Finlay & Forsyth, 2009; 1999; Pappens et al., 2015a; Valdivia-Salas et al., 2014), with superior conditioning and/or slower extinction for fear-relevant CSs (Forsyth & Eifert, 1996; 1998; Stegen et al., 1999). Cardiorespiratory activation as a CR may be more easily established with respiratory USs than with other USs, but this remains conjecture as no studies have directly compared cardiorespiratory CRs between different types of US (e.g., respiratory occlusion versus shock).

Differential, *intero*-interoceptive paradigms (see Tables 2 and 4) are rare and face the challenge of finding two interoceptive CSs that can be experimentally controlled and that are equivalent in a range of features that affect conditionability (De Cort et al., 2017; Schroyen et al., 2015). Most of the intero-interoceptive studies have therefore implemented single-cue, between subject paradigms. The following paragraphs will discuss some general findings apparent from these single-cue paradigms.

5.3.3 Functionally varying faces of fear. As outlined above, self-reports and SCR are consistently dissociated from startle blink EMG and respiration in single-cue intero-interoceptive paradigms (Table 4), during either acquisition (load-load paradigm) or extinction (load-occlusion paradigm). The consistency of this dissociation underscores that, as argued repeatedly in the literature (e.g., Hugdall, 1980; Lang, 1968; Ledoux, 2014; but see Fanselow & Pennington, 2018), fear is not a unitary concept. What is learned in a fear conditioning procedure can affect multiple, potentially independent but interacting memory and response systems. In line with a functional

INTEROCEPTIVE FEAR CONDITIONING

perspective on associative learning, the type of stimulus used may be essential to determine which memory and response systems are more or less engaged, and maybe even how they functionally connect to each other (Ferbinteanu, 2018). The reviewed findings strongly suggest that single-cue, intero-interoceptive paradigms favor the establishment of amygdala-based 'emotional' memories over declarative memories about the CS-US contingency, which may be related to the vital function of respiration and the likely existence of hard-wired defense circuits related to suffocation. Alternatively, the consistent dissociation between both CR types in the reviewed single-cue, intero-interoceptive paradigms may also be a typical feature of single-cue, between subject paradigms that model aspects of intra-event learning. A fruitful way to explore this question may be to study dissociation of CRs in between subject, intra-event paradigms with *exteroceptive* stimuli (e.g., neutral tone predicting an aversive tone; neutral tactile stimulus predicting a painful tactile stimulation). The use of single-cue, between subject paradigms in human fear conditioning would also facilitate closer comparisons with basic findings from animal studies (Lonsdorf et al., 2017).

5.3.4 Razran's propositions hold, but may be unfalsifiable. As outlined before, the propositions that interoceptive conditioning occurs largely unconsciously and that it is more resistant to extinction than exteroceptive conditioning were put forward many years ago (Razran, 1961). The findings discussed in the present review support these propositions a certain extent, as the load-load paradigm seems to generate 'unconscious' emotional learning, and the load-occlusion paradigm does not yield extinction in indices of emotional learning. However, even today, Razran's propositions seem extremely difficult to falsify. It is challenging to come up with operationalizations of 'consciousness' or 'awareness' that researchers would agree upon (see, for example, Lovibond & Shanks, 2002 for a related discussion on contingency awareness). Another difficulty is the fluctuating boundaries of how 'interoceptive' is defined, not to mention how to ensure that a CS is *truly* and *exclusively* 'interoceptive' and matched to an exclusively exteroceptive CS on other dimensions known to affect conditionability, such as perceptual salience, similarity, or

INTEROCEPTIVE FEAR CONDITIONING

belongingness to the US (Goddard, 1999). Rather than trying to infer differences between interoceptive and exteroceptive learning, it is likely more fruitful and parsimonious to assume that both interoceptive and exteroceptive learning obey the same basic learning principles, but are strongly modulated by 'pre-existing' relationships between the CS (whether interoceptive or exteroceptive) and the US, as well as the response system under study.

6. Future directions

Interoceptive conditioning has been hypothesized to play a role in panic and psychosomatic disorders. The present review indicates the availability of experimental evidence that associative learning involving interoceptive stimuli can indeed generate conditioned cardiorespiratory arousal and fear. This demonstration of interoceptive conditioning of fear and cardiorespiratory arousal supports its potential etiological role in both disorders. However, two aspects that may be more clinically and theoretically relevant to these disorders have not been extensively studied and warrant more attention in future research.

First, the reviewed work on interoceptive conditioning has focused mainly on the acquisition, rather than on the extinction of interoceptive fear. Although cognitive behavioral therapy including interoceptive exposure is an effective treatment strategy (Boettcher & Barlow, 2018; Craske et al., 2011; Flink, Nicholaoas, Boersma, & Linton, 2009; Pompoli et al., 2018; Zucker et al., 2018), there is considerable room for improvement (Freire, Zuglinai, Garcia, & Nardi, 2016; Mitte, 2005). We do not currently know how extinction of interoceptive fears may differ from extinction of exteroceptive fears, nor do we know how it could be facilitated. A promising tool for studying this may be transcutaneous vagal nerve stimulation as an add-on to interoceptive exposure / extinction learning (Burger et al., 2017).

Second, clinical studies have found that interoceptive exposure reduces not only fear, but also functional symptoms in persons suffering from irritable bowel syndrome (e.g.,

INTEROCEPTIVE FEAR CONDITIONING

1 Craske et al., 2011; Ljòtsson et al., 2013). In other words, extinction of interoceptive fear has
2 the potential to alter interoception itself. This is consistent with findings showing that fear
3 learning can influence a range of sensory and perceptual outcomes in exteroceptive
4 modalities (e.g., Miskovic & Keil, 2012; Resnik, Sobel & Paz, 2011; Rossi et al., 2017).
5 However, the relationship between conditioned interoceptive fear and interoception itself
6 remains understudied, despite its great potential to improve understanding and treatment of
7 the clinical problem of functional bodily symptoms (Zaman, Vlaeyen, Van Oudenhove,
8 Wiech, & Van Diest, 2015). In the visceral domain, we have found preliminary evidence
9 that fear learning to a visceral CS was associated with a subsequent lack of habituation of the
10 perceptual threshold of the visceral CS, as well as with a decreased performance in a visceral
11 discrimination task (Zaman et al., 2016). Whether such findings can be replicated and
12 extended to the cardiorespiratory domain, is currently unknown and subject to further
13 research.

14
15

Acknowledgements

Psychophysiology

INTEROCEPTIVE FEAR CONDITIONING

References

- Acheson, D. T., Forsyth, J. P., & Moses, E. (2012). Interoceptive fear conditioning and panic disorder: the role of conditioned stimulus–unconditioned stimulus predictability. *Behavior therapy*, 43(1), 174-189. doi:10.1016/j.beth.2011.06.001
- Acheson, D. T., Forsyth, J. P., Prenoveau, J. M., & Bouton, M. E. (2007). Interoceptive fear conditioning as a learning model of panic disorder: an experimental evaluation using 20% CO₂-enriched air in a non-clinical sample. *Behaviour research and therapy*, 45(10), 2280-2294. doi:10.1016/j.brat.2007.04.008
- Alexander-Miller, S., & Davenport, P. W. (2010). Perception of multiple-breath inspiratory resistive loads in males and females. *Biological psychology*, 84(1), 147-149. doi:10.1016/j.biopsycho.2010.03.004
- Alius, M. G., Pané-Farré, C. A., Löw, A., & Hamm, A. O. (2015). Modulation of the blink reflex and P3 component of the startle response during an interoceptive challenge. *Psychophysiology*, 52(1), 140-148. doi:10.1111/psyp.12295
- Argyropoulos, S. V., Bailey, J. E., Hood, S. D., Kendrick, A. H., Rich, A. S., Laszlo, G., Nash, J. R., Lightman, S., & Nutt, D. J. (2002). Inhalation of 35% CO₂ results in activation of the HPA axis in healthy volunteers. *Psychoneuroendocrinology*, 27, 715-729. doi:10.1016/s0306-4530(01)00075-0
- Bailey, J. E., Argyropoulos, S. V., Kendrick, A. H., & Nutt, D. J. (2005). Behavioral and cardiovascular effects of 7.5% CO₂ in human volunteers. *Depression and Anxiety*, 21, 18-25. doi:10.1002/da.20048
- Barlow, D. H. (1988). *Anxiety and its disorders: The nature and treatment of anxiety and panic*. New York: Guilford Press.

INTEROCEPTIVE FEAR CONDITIONING

- 1 Battaglia, M. (2017). Sensitivity to carbon dioxide and translational studies of anxiety
2 disorders. *Neuroscience*, 346, 434-436. doi:10.1016/j.neuroscience.2017.01.053
- 3 Bechara, A., Tranel, D., Damasio, H., Adolphs, R., Rockland, C., & Damasio, A. R. (1995).
4 Double dissociation of conditioning and declarative knowledge relative to the amygdala
5 and hippocampus in humans. *Science*, 269(5227), 1115-1118.
6 doi:10.1093/neucas/3.4.267-c
- 7 Benke, C., Alius, M. G., Hamm, A. O., & Pané-Farré, C. A. (2018). Cue and context
8 conditioning to respiratory threat: Effects of suffocation fear and implications for the
9 etiology of panic disorder. *International Journal of Psychophysiology*, 124, 33-42.
10 doi:10.1016/j.ijpsycho.2018.01.002
- 11 Benke, C., Hamm, A. O., & Pané-Farré, C. A. (2017). When dyspnea gets worse:
12 Suffocation fear and the dynamics of defensive respiratory responses to increasing
13 interoceptive threat. *Psychophysiology*, 54(9), 1266-1283. doi:10.1111/psyp.12881
- 14 Biber, B., & Alkln, T. (1999). Panic disorder subtypes: differential responses to CO2
15 challenge. *American Journal of Psychiatry*, 156(5), 739-744.
- 16 Blechert, J., Wilhelm, F. H., Meuret, A. E., Wilhelm, E. M., & Roth, W. T. (2013).
17 Experiential, autonomic, and respiratory correlates of CO2 reactivity in individuals with
18 high and low anxiety sensitivity. *Psychiatry research*, 209(3), 566-573.
19 doi:10.1016/j.psychres.2013.02.010
- 20 Boettcher, H., & Barlow, D. H. (2018). The unique and conditional effects of interoceptive
21 exposure in the treatment of anxiety: A functional analysis. *Behaviour research and*
22 *therapy*. doi:10.1016/j.brat.2018.12.002
- 23 Bouton, M. E., Mineka, S., & Barlow, D. H. (2001). A modern learning theory perspective
24 on the etiology of panic disorder. *Psychological Review*, 108(1), 4-32. doi:10.1037/0033-
25 295x.108.1.4

INTEROCEPTIVE FEAR CONDITIONING

- 1 Broocks, A., Meyer, T. F., Bandelow, B., George, A., Bartmann, U., Rther, E., & Hillmer-
2 Vogel, U. (1997). Exercise avoidance and impaired endurance capacity in patients with
3 panic disorder. *Neuropsychobiology*, 36(4), 182-187. doi:10.1159/000119381
- 4 Brown, L. A., & Craske, M. G. (2017). Panic Disorder and Agoraphobia. *Treatments for*
5 *Psychological Problems and Syndromes*, 34-55. doi:10.1002/9781118877142.ch4
- 6 Burger, A. M., Verkuil, B., Fenlon, H., Thijs, L., Cools, L., Miller, H. C., ... & Van Diest, I.
7 (2017). Mixed evidence for the potential of non-invasive transcutaneous vagal nerve
8 stimulation to improve the extinction and retention of fear. *Behaviour research and*
9 *therapy*, 97, 64-74. doi:10.1016/j.brat.2017.07.005
- 10 Bykov, K. M. (1957). The cerebral cortex and the inner organs. *Chemic. Publ., New York*.
- 11 Cameron, O. G. (2001). Interoception: the inside story—a model for psychosomatic
12 processes. *Psychosomatic medicine*, 63(5), 697-710. doi:10.1097/00006842-200109000-
13 00001
- 14 Campbell, D., Sanderson, R. E., & Laverty, S. G. (1964). Characteristics of a conditioned
15 response in human subjects during extinction trials following a single traumatic
16 conditioning trial. *The Journal of Abnormal and Social Psychology*, 68(6), 627–639.
17 doi:10.1037/h0042582
- 18 Ceunen, E., Vlaeyen, J. W., & Van Diest, I. (2013). Atypical modulation of startle in women
19 in face of aversive bodily sensations. *International journal of psychophysiology*, 88(2),
20 157-163. doi:10.1016/j.ijpsycho.2013.03.013
- 21 Ceunen, E., Vlaeyen, J. W., & Van Diest, I. (2016). On the origin of interoception. *Frontiers*
22 *in psychology*, 7, 743. doi:10.3389/fpsyg.2016.00743
- 23 Ceunen, E., Zaman, J., Herssens, N., Van Oudenhove, L., Bogaerts, K., Ly, H. G., ... & Van
24 Diest, I. (2015). Startle responding in the context of visceral pain. *International journal of*
25 *psychophysiology*, 98(1), 128-134. doi:10.1016/j.ijpsycho.2015.07.009

INTEROCEPTIVE FEAR CONDITIONING

- 1 Cook, E. W., Hodes, R. L., & Lang, P. J. (1986). Preparedness and phobia: effects of
2 stimulus content on human visceral conditioning. *Journal of abnormal psychology*, 95(3),
3 195. doi:10.1037/0021-843x.95.3.195
- 4 Coplan, J. D., Moreau, D., Chaput, F., Martinez, J. M., Hoven, C. W., Mandell, D. J.,
5 Gorman, J. M., & Pine, D. S. (2002). Salivary cortisol concentrations before and after
6 carbon-dioxide inhalations in children. *Biological Psychiatry*, 51(4), 326-333.
7 doi:10.1016/s0006-3223(01)01250-1
- 8 Craig, A. D. (2002). How do you feel? Interoception: the sense of the physiological
9 condition of the body. *Nature reviews neuroscience*, 3(8), 655. doi:10.1038/nrn894
- 10 Craske, M. G., Wolitzky-Taylor, K. B., Labus, J., Wu, S., Frese, M., Mayer, E. A., &
11 Naliboff, B. D. (2011). A cognitive-behavioral treatment for irritable bowel syndrome
12 using interoceptive exposure to visceral sensations. *Behaviour research and therapy*,
13 49(6-7), 413-21. doi:10.1016/j.brat.2011.04.001
- 14 Deacon, B., Kemp, J. J., Dixon, L. J., Sy, J. T., Farrell, N. R., & Zhang, A. R. (2013).
15 Maximizing the efficacy of interoceptive exposure by optimizing inhibitory learning: A
16 randomized controlled trial. *Behaviour Research and Therapy*, 51(9), 588-596.
17 doi:10.1016/j.brat.2013.06.006
- 18 De Cort, K., Griez, E., Büchler, M., & Schruers, K. (2012). The role of “interoceptive” fear
19 conditioning in the development of panic disorder. *Behavior therapy*, 43(1), 203-215.
20 doi:10.1016/j.beth.2011.06.005
- 21 De Cort, K., Schroijen, M., Hurlemann, R., Claassen, S., Hoogenhout, J., Van den Bergh, O.,
22 ... & Schruers, K. (2017). Modeling the development of panic disorder with interoceptive
23 conditioning. *European neuropsychopharmacology*, 27(1), 59-69.
24 doi:10.1016/j.euroneuro.2016.11.001

INTEROCEPTIVE FEAR CONDITIONING

- 1 Denot-Ledunois, S., Vardon, G., Perruchet, P., & Gallego, J. (1998). The effect of attentional
2 load on the breathing pattern in children. *International journal of psychophysiology*,
3 29(1), 13-21. doi:10.1016/s0167-8760(97)00086-x
- 4 De Houwer, J., & Beckers, T. (2002). A review of recent developments in research and
5 theories on human contingency learning. *The Quarterly Journal of Experimental*
6 *Psychology: Section B*, 55(4), 289-310. doi:10.1080/02724990244000034
- 7 De Peuter, S., Van Diest, I., Lemaigre, V., Li, W., Verleden, G., Demedts, M., & Van den
8 Bergh, O. (2005). Can subjective asthma symptoms be learned?. *Psychosomatic*
9 *Medicine*, 67(3), 454-461. doi:10.1097/01.psy.0000160470.43167.e2
- 10 Domjan, M. (2005). Pavlovian Conditioning: A Functional Perspective. *Annual Review of*
11 *Psychology*, 56(1), 179–206. doi:10.1146/annurev.psych.55.090902.141409
- 12 Domjan, M., Blesbois, E., & Williams, J. (1998). The adaptive significance of sexual
13 conditioning: Pavlovian control of sperm release. *Psychological Science*, 9(5), 411-415.
14 doi:10.1111/1467-9280.00077
- 15 Dworkin, B. R. (1993). *Learning and physiological regulation*. University of Chicago Press.
- 16 Dworkin, B. R., & Dworkin, S. (1995). Learning of physiological responses: II. Classical
17 conditioning of the baroreflex. *Behavioral Neuroscience*, 109(6), 1119–1136.
18 doi:10.1037/0735-7044.109.6.1119
- 19 Esquivel, G., Schruers, K., Maddock, R., Colasanti, A., & Griez, E. (2010). Review: Acids
20 in the brain: a factor in panic? *Journal of Psychopharmacology*, 24(5), 639–647.
21 doi:10.1177/0269881109104847
- 22 Esser, R. W., Stoeckel, M. C., Kirsten, A., Watz, H., Taube, K., Lehmann, K., ... & von
23 Leupoldt, A. (2017). Brain activation during perception and anticipation of dyspnea in
24 chronic obstructive pulmonary disease. *Frontiers in physiology*, 8, 617.
25 doi:10.3389/fphys.2017.00617

INTEROCEPTIVE FEAR CONDITIONING

- 1 Fanselow, M. S., & Pennington, Z. T. (2018). A return to the psychiatric dark ages with a
2 two-system framework for fear. *Behaviour Research and Therapy*, 100, 24-29.
3 doi:10.1016/j.brat.2017.10.012
- 4 Fanselow, M. S., & Poulos, A. M. (2005). The Neuroscience of Mammalian Associative
5 Learning. *Annual Review of Psychology*, 56(1), 207–234.
6 doi:10.1146/annurev.psych.56.091103.070213
- 7 Fannes, S., Van Diest, I., Meulders, A., De Peuter, S., Vansteenwegen, D., & Van den
8 Bergh, O. (2008). To inhale or not to inhale: Conditioned avoidance in breathing behavior
9 in an odor—20% CO₂ paradigm. *Biological Psychology*, 78(1), 87-92.
10 doi:10.1016/j.biopsycho.2008.01.003
- 11 Fava, G. A., & Sonino, N. (2010). Psychosomatic Medicine. *The International Journal of*
12 *Clinical Practice*, 64, 1155-1161. doi:10.1111/j.1742-1241.2009.02266.x
- 13 Feldner, M. T., Zvolensky, M. J., Eifert, G. H., & Spira, A. P. (2003). Emotional avoidance:
14 an experimental test of individual differences and response suppression using biological
15 challenge. *Behaviour Research and Therapy*, 41, 403-411. doi:10.1016/s0005-
16 7967(02)00020-7
- 17 Ferbinteanu, J. (2019). Memory systems 2018 – Towards a new paradigm. *Neurobiology of*
18 *Learning and Memory*, 157, 61–78. doi:10.1016/j.nlm.2018.11.005
- 19 Finlay, C. G., & Forsyth, J. P. (2009). Context and renewal of conditioned fear: An
20 experimental evaluation using 20% carbon dioxide-enriched air as an unconditioned
21 stimulus. *Journal of anxiety disorders*, 23(6), 737-745. doi:10.1016/j.janxdis.2009.02.014
- 22 Flink, I. K., Nicholas, M. K., Boersma, K., & Linton, S. J. (2009). Reducing the threat value
23 of chronic pain: A preliminary replicated single-case study of interoceptive exposure
24 versus distraction in six individuals with chronic back pain. *Behaviour research and*
25 *therapy*, 47(8), 721-728. doi:10.1016/j.brat.2009.05.003

INTEROCEPTIVE FEAR CONDITIONING

- 1 Fokkema, D. S. (1999). The psychobiology of strained breathing and its cardiovascular
2 implications: A functional system review. *Psychophysiology*, 36, 164-175.
3 doi:10.1111/1469-8986.3620164
- 4 Forsyth, J. P., & Eifert, G. H. (1998). Response intensity in content-specific fear
5 conditioning comparing 20% versus 13% CO₂-enriched air as unconditioned stimuli.
6 *Journal of Abnormal Psychology*, 107(2), 291-304. doi:10.1037/0021-843x.107.2.291
- 7 Forsyth, J. P., Eifert, G. H., & Thompson, R. N. (1996). Systemic alarms in fear
8 conditioning II: An experimental methodology using 20% carbon dioxide inhalation as an
9 unconditioned stimulus. *Behavior Therapy*, 27(3), 391-415. doi:10.1016/s0005-
10 7894(96)80024-4
- 11 Freire, R. C., Zugliani, M. M., Garcia, R. F., & Nardi, A. E. (2016). Treatment-resistant
12 panic disorder: a systematic review. *Expert opinion on pharmacotherapy*, 17(2), 159-168.
13 doi:10.1517/14656566.2016.1109628
- 14 Gallego, J., & Perruchet, P. (1991). Classical conditioning of ventilatory responses in
15 humans. *Journal of Applied Physiology*, 70(2), 676-682.
16 doi:10.1152/jappl.1991.70.2.676
- 17 Garcia, J., Hankins, W.G., & Rusiniak, K.W. (1974). Behavioral regulation of the milieu
18 interne in man and rat. *Science* 185, 824–831. doi:10.1126/science.185.4154.824
- 19 Garcia, J., & Koelling, R. A. (1966). Relation of cue to consequence in avoidance learning.
20 *Psychonomic science*, 4(1), 123-124. doi:10.3758/bf03342209
- 21 Gardner, W. N. (1996). The pathophysiology of hyperventilation disorders. *Chest*, 109(2),
22 516-534. doi:10.1378/chest.109.2.516
- 23 Giannese, F., Luchetti, A., Barbiera, G., Lampis, V., Zanettini, C., Knudsen, G. P., ... &
24 Battaglia, M. (2018). Conserved DNA Methylation Signatures in Early Maternal

INTEROCEPTIVE FEAR CONDITIONING

- 1 Separation and in Twins Discordant for CO₂ Sensitivity. *Scientific reports*, 8(1), 2258.
- 2 doi:10.1038/s41598-018-20457-3
- 3 Goddard, M. J. (1999). The role of US signal value in contingency, drug conditioning, and
- 4 learned helplessness. *Psychonomic bulletin & review*, 6(3), 412-423.
- 5 doi:10.3758/bf03210829
- 6 Goldstein, A. J., & Chambless, D. L. (1978). A reanalysis of agoraphobia. *Behavior*
- 7 *Therapy*, 9(1), 47-59. doi:10.1016/s0005-7894(78)80053-7
- 8 Hamm, A. O., & Vaitl, D. (1996). Affective learning: Awareness and aversion.
- 9 *Psychophysiology*, 33(6), 698-710. doi:10.1111/j.1469-8986.1996.tb02366.x
- 10 Han, J. N., Stegen, K., De Valck, C., Clement, J., & Van de Woestijne, K. P. (1996).
- 11 Influence of breathing therapy on complaints, anxiety and breathing pattern in patients
- 12 with hyperventilation syndrome and anxiety disorders. *Journal of psychosomatic*
- 13 *research*, 41(5), 481-493. doi:10.1016/s0022-3999(96)00220-6
- 14 Haroutunian, V., & Campbell, B. A. (1979). Emergence of interoceptive and exteroceptive
- 15 control of behavior in rats. *Science*, 205(4409), 927-929. doi:10.1126/science.472715
- 16 Hegel, M. T., & Ferguson, R. J. (1997). Psychophysiological assessment of respiratory
- 17 function in panic disorder: Evidence for a hyperventilation subtype. *Psychosomatic*
- 18 *Medicine*, 59, 224-230. doi:10.1097/00006842-199705000-00003
- 19 Hodes, R. L., Cook, E. W., & Lang, P. J. (1985). Individual differences in autonomic
- 20 response: conditioned association or conditioned fear? *Psychophysiology*, 22(5), 545-560.
- 21 doi:10.1111/j.1469-8986.1985.tb01649.x
- 22 Hofer, M. A., & Hinkle Jr., L. E. (1964). Conditioned diuresis in man: Effects of altered
- 23 environment, subjective state, and conditioning Experience. *Psychosomatic medicine*,
- 24 26(2), 108-124. doi:10.1097/00006842-196403000-0000

INTEROCEPTIVE FEAR CONDITIONING

- 1 Hugdahl, K. (1981). The three-systems-model of fear and emotion—a critical examination.
2 *Behaviour Research and Therapy*, 19(1), 75-83. doi:10.1016/0005-7967(81)90114-5
- 3 Kessler, R. C., Chiu, W. T., Jin, R., Ruscio, A. M., Shear, K., & Walters, E. E. (2006). The
4 epidemiology of panic attacks, panic disorder, and agoraphobia in the National
5 Comorbidity Survey Replication. *Archives of general psychiatry*, 63(4), 415-424.
6 doi:10.1001/archpsyc.63.4.415
- 7 Khalsa, S. S., Adolphs, R., Cameron, O. G., Critchley, H. D., Davenport, P. W., Feinstein, J.
8 S., ... & Meuret, A. E. (2017). Interoception and mental health: a roadmap. *Biological*
9 *Psychiatry: Cognitive Neuroscience and Neuroimaging*. Doi: 10.1016/j.bpsc.2017.12.004
- 10 Khalsa, S. S., & Lapidus, R. C. (2016). Can interoception improve the pragmatic search for
11 biomarkers in psychiatry?. *Frontiers in psychiatry*, 7, 121. doi:10.3389/fpsy.2016.00121
- 12 Klein, D. F. (1993). False suffocation alarms, spontaneous panics, and related conditions: an
13 integrative hypothesis. *Archives of general psychiatry*, 50(4), 306-317.
14 doi:10.1001/archpsyc.1993.01820160076009
- 15 Krause, E., Benke, C., Koenig, J., Thayer, J. F., Hamm, A. O., & Pané-Farré, C. A. (2018).
16 Dynamics of Defensive Response Mobilization to Approaching External Versus
17 Interoceptive Threat. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*,
18 3(6), 525–538. doi:10.1016/j.bpsc.2017.12.002
- 19 Krause, M. A., & Domjan, M. (2017). Ethological and evolutionary perspectives on
20 Pavlovian conditioning. *APA Handbook of Comparative Psychology. Vol 2. Perception,*
21 *Learning, and Cognition*, 247-266. doi:10.1037/0000012-012
- 22 Lang, P. J. (1968). Fear reduction and fear behavior: Problems in treating a construct. In
23 *Research in psychotherapy conference, 3rd, May-Jun, 1966, Chicago, IL, US*. American
24 Psychological Association. doi:10.1037/10546-004

INTEROCEPTIVE FEAR CONDITIONING

- 1 Lang, P. J., & Bradley, M. M. (2013). Appetitive and defensive motivation: Goal-directed or
2 goal-determined?. *Emotion Review*, 5(3), 230-234. doi:10.1177/1754073913477511
- 3 Lang, P. J., Wangelin, B. C., Bradley, M. M., Versace, F., Davenport, P. W., & Costa, V. D.
4 (2011). Threat of suffocation and defensive reflex activation. *Psychophysiology*, 48(3),
5 393-396. doi:10.1111/j.1469-8986.2010.01076.x
- 6 Lavietes, M. H., Sanchez, C. W., Tiersky, L. A., Cherniack, N. S., & Natelson, B. H. (2000).
7 Psychological profile and ventilatory response to inspiratory resistive loading. *American*
8 *journal of respiratory and critical care medicine*, 161(3), 737-744.
9 doi:10.1164/ajrccm.161.3.9810075
- 10 LeDoux, J. E. (2014). Coming to terms with fear. *Proceedings of the National Academy of*
11 *Sciences*, 111(8), 2871-2878. doi:10.1073/pnas.1400335111
- 12 Lee, K., Noda, Y., Nakano, Y., Ogawa, S., Kinoshita, Y., Funayama, T., & Furukawa, T. A.
13 (2006). Interoceptive hypersensitivity and interoceptive exposure in patients with panic
14 disorder: specificity and effectiveness. *BMC psychiatry*, 6(1), 32. doi:10.1186/1471-244x-
15 6-32
- 16 Leibold, N. K., van den Hove, D. L. A., Esquivel, G., De Cort, K., Goossens, L., Strackx, E.,
17 ... & Schruers, K. R. J. (2015). The brain acid–base homeostasis and serotonin: A
18 perspective on the use of carbon dioxide as human and rodent experimental model of
19 panic. *Progress in neurobiology*, 129, 58-78. doi:10.1016/j.pneurobio.2015.04.001
- 20 Leibold, N. K., van den Hove, D. L., Viechtbauer, W., Kenis, G., Goossens, L., Lange, I., ...
21 & Schruers, K. R. (2017). Amiloride-sensitive cation channel 2 genotype affects the
22 response to a carbon dioxide panic challenge. *Journal of Psychopharmacology*, 31(10),
23 1294-1301. doi:10.1177/0269881116686880
- 24 Leibold, N. K., Viechtbauer, W., Goossens, L., De Cort, K., Griez, E. J., Myin-Germeys, I.,
25 ... & Schruers, K. R. (2013). Carbon dioxide inhalation as a human experimental model of

INTEROCEPTIVE FEAR CONDITIONING

- 1 panic: the relationship between emotions and cardiovascular physiology. *Biological*
- 2 *psychology*, 94(2), 331-340. doi:10.1016/j.biopsycho.2013.06.004
- 3 Limebeer, C. L., Krohn, J. P., Cross-Mellor, S., Litt, D. E., Ossenkopp, K. P., & Parker, L.
- 4 A. (2008). Exposure to a context previously associated with nausea elicits conditioned
- 5 gaping in rats: a model of anticipatory nausea. *Behavioural brain research*, 187(1), 33-40.
- 6 doi:10.1016/j.bbr.2007.08.024
- 7 Livermore, N., Butler, J. E., Sharpe, L., McBain, R. A., Gandevia, S. C., & McKenzie, D. K.
- 8 (2008). Panic attacks and perception of inspiratory resistive loads in chronic obstructive
- 9 pulmonary disease. *American Journal of Respiratory and Critical Care Medicine*, 178(1),
- 10 7-12. doi:10.1164/rccm.200711-1700oc
- 11 Ljótsson, B., Hesser, H., Andersson, E., Lindfors, P., Hursti, T., Rück, C., ... & Hedman, E.
- 12 (2013). Mechanisms of change in an exposure-based treatment for irritable bowel
- 13 syndrome. *Journal of consulting and clinical psychology*, 81(6), 1113.
- 14 doi:10.1037/a0033439
- 15 Lonsdorf, T. B., Menz, M. M., Andreatta, M., Fullana, M. A., Golkar, A., Haaker, J., ... &
- 16 Drexler, S. M. (2017). Don't fear 'fear conditioning': Methodological considerations for
- 17 the design and analysis of studies on human fear acquisition, extinction, and return of
- 18 fear. *Neuroscience & Biobehavioral Reviews*, 77, 247-285.
- 19 doi:10.1016/j.neubiorev.2017.02.026
- 20 Lovibond, P. F., & Shanks, D. R. (2002). The role of awareness in Pavlovian conditioning:
- 21 empirical evidence and theoretical implications. *Journal of Experimental Psychology:*
- 22 *Animal Behavior Processes*, 28(1), 3. doi:10.1037/0097-7403.28.1.3
- 23 Löw, A., Weymar, M., & Hamm, A. O. (2015). When threat is near, get out of here:
- 24 Dynamics of defensive behavior during freezing and active avoidance. *Psychological*
- 25 *science*, 26(11), 1706-1716. doi:10.1177/0956797615597332

INTEROCEPTIVE FEAR CONDITIONING

- 1 Masaoka, Y., & Homma, I. (1999). Expiratory time determined by individual anxiety levels
2 in humans. *Journal of Applied Physiology*, 86(4), 1329-1336.
3 doi:10.1152/jappl.1999.86.4.1329
- 4 McNally, R. J., & Eke, M. (1996). Anxiety sensitivity, suffocation fear, and breath-holding
5 duration as predictors of response to carbon dioxide challenge. *Journal of Abnormal*
6 *Psychology*, 105(1), 146-149. doi:10.1037/0021-843x.105.1.146
- 7 Melzig, C. A., Holtz, K., Michalowski, J. M., & Hamm, A. O. (2011). Interoceptive threat
8 leads to defensive mobilization in highly anxiety sensitive persons. *Psychophysiology*,
9 48(6), 745-754. doi:10.1111/j.1469-8986.2010.01150.
- 10 Meuret, A. E., Ritz, T., Wilhelm, F. H., Roth, W. T., & Rosenfield, D. (2018).
11 Hypoventilation therapy alleviates panic by repeated induction of dyspnea. *Biological*
12 *Psychiatry: Cognitive Neuroscience and Neuroimaging* 3(6), 539–545.
13 doi:10.1016/j.bpsc.2018.01.010
- 14 Meuret, A. E., Rosenfield, D., Seidel, A., Bhaskara, L., & Hofmann, S. G. (2010).
15 Respiratory and cognitive mediators of treatment change in panic disorder: Evidence for
16 intervention specificity. *Journal of Consulting and Clinical Psychology*, 78(5), 691-704
17 doi:10.1037/a0019552
- 18 Meuret, A. E., Rosenfield, D., Wilhelm, F. H., Zhou, E., Conrad, A., Ritz, T., & Roth, W. T.
19 (2011). Do unexpected panic attacks occur spontaneously? *Biological psychiatry*, 70(10),
20 985-991. doi:10.1016/j.biopsych.2011.05.027
- 21 Miskovic, V., & Keil, A. (2012). Acquired fears reflected in cortical sensory processing: a
22 review of electrophysiological studies of human classical conditioning.
23 *Psychophysiology*, 49(9), 1230-1241. doi:10.1111/j.1469-8986.2012.01398.

INTEROCEPTIVE FEAR CONDITIONING

- 1 Mitte, K. (2005). A meta-analysis of the efficacy of psycho-and pharmacotherapy in panic
2 disorder with and without agoraphobia. *Journal of affective disorders*, 88(1), 27-45.
3 doi:10.1016/j.jad.2005.05.003
- 4 Moessnang, C., Pauly, K., Kellermann, T., Krämer, J., Finkelmeyer, A., Hummel, T., ... &
5 Habel, U. (2013). The scent of salience—Is there olfactory-trigeminal conditioning in
6 humans?. *Neuroimage*, 77, 93-104. doi:10.1016/j.neuroimage.2013.03.049
- 7 Mongeluzi, D. L., Rosellini, R. A., Ley, R., Caldarone, B. J., & Stock, H. S. (2003). The
8 conditioning of dyspneic suffocation fear: effects of carbon dioxide concentration on
9 behavioral freezing and analgesia. *Behavior modification*, 27(5), 620-636.
10 doi:10.1177/0145445503256316
- 11 Moratti, S., Keil, A., & Miller, G. A. (2006). Fear but not awareness predicts enhanced
12 sensory processing in fear conditioning. *Psychophysiology*, 43(2), 216-226.
13 doi:10.1111/j.1464-8986.2006.00386
- 14 Öhman, A., & Mineka, S. (2001). Fears, phobias, and preparedness: toward an evolved
15 module of fear and fear learning. *Psychological review*, 108(3), 483. doi:10.1037/0033-
16 295x.108.3.483
- 17 Pain, M. C. F., Biddle, N., & Tiller, J. W. G. (1988). Panic disorder, the ventilatory
18 response to carbon dioxide and respiratory variables. *Psychosomatic Medicine*, 50, 541-
19 548. doi:10.1097/00006842-198809000-00010
- 20 Pappens, M., De Peuter, S., Vansteenwegen, D., Van den Bergh, O., & Van Diest, I.
21 (2012a). Psychophysiological responses to CO2 inhalation. *International journal of*
22 *psychophysiology*, 84(1), 45-50. doi:10.1016/j.ijpsycho.2012.01.008
- 23 Pappens, M., Schroyen, M., Sütterlin, S., Smets, E., Van den Bergh, O., Thayer, J. F., & Van
24 Diest, I. (2014). Resting heart rate variability predicts safety learning and fear extinction

INTEROCEPTIVE FEAR CONDITIONING

- 1 in an interoceptive fear conditioning paradigm. *PloS one*, 9(9), e105054.
- 2 doi:10.1371/journal.pone.0105054
- 3 Pappens, M., Schroyen, M., Van den Bergh, O., & Van Diest, I. (2015a). Retention of
- 4 perceptual generalization of fear extinction. *International Journal of Psychophysiology*,
- 5 98(3), 520-528. doi:10.1016/j.ijpsycho.2015.01.007
- 6 Pappens, M., Smets, E., Bergh, O., & Van Diest, I. (2012b). Fear of suffocation alters
- 7 respiration during obstructed breathing. *Psychophysiology*, 49(6), 829-832.
- 8 doi:10.1111/j.1469-8986.2012.01355.
- 9 Pappens, M., Smets, E., Vansteenwegen, D., Van den Bergh, O., & Van Diest, I. (2012c).
- 10 Learning to fear suffocation: a new paradigm for interoceptive fear conditioning.
- 11 *Psychophysiology*, 49(6), 821-828. doi:10.1111/j.1469-8986.2012.01357
- 12 Pappens, M., Van den Bergh, O., De Peuter, S., Bresseleers, J., Vansteenwegen, D., & Van
- 13 Diest, I. (2010). Defense reactions to interoceptive threats: a comparison between loaded
- 14 breathing and aversive picture viewing. *Biological psychology*, 84(1), 98-103.
- 15 doi:10.1016/j.biopsycho.2010.02.006
- 16 Pappens, M., Van den Bergh, O., Vansteenwegen, D., Ceunen, E., De Peuter, S., & Van
- 17 Diest, I. (2013). Learning to fear obstructed breathing: comparing interoceptive and
- 18 exteroceptive cues. *Biological psychology*, 92(1), 36-42.
- 19 doi:10.1016/j.biopsycho.2011.05.004
- 20 Pappens, M., Van den Bergh, O., Vansteenwegen, D., & Van Diest, I. (2011).
- 21 Psychophysiological responses to inspiratory resistive loads. *International Journal of*
- 22 *Psychophysiology*, 80(2), 161-165. doi:10.1016/j.ijpsycho.2011.02.015
- 23 Pappens, M., Vandenbossche, E., Van den Bergh, O., & Van Diest, I. (2015b). Interoceptive
- 24 fear learning to mild breathlessness as a laboratory model for unexpected panic attacks.
- 25 *Frontiers in psychology*, 6, 1150. doi: 10.3389/fpsyg.2015.01150

INTEROCEPTIVE FEAR CONDITIONING

- 1 Perna G, Cocchi S, Bertani A, Arancio C, Bellodi L. Sensitivity to 35% CO₂ in healthy first-
2 degree relatives of patients with panic disorder (1995). *The American Journal of*
3 *Psychiatry*, 152, 623–625. doi:10.1176/ajp.152.4.623
- 4 Petzschner, F. H., Weber, L. A., Gard, T., & Stephan, K. E. (2017). Computational
5 psychosomatics and computational psychiatry: Toward a joint framework for differential
6 diagnosis. *Biological psychiatry*, 82(6), 421-430. doi:10.1016/j.biopsych.2017.05.012
- 7 Pinkney, V., Wickens, R., Bamford, S., Baldwin, D. S., & Garner, M. (2014). Defensive
8 eye-blink startle responses in a human experimental model of anxiety. *Journal of*
9 *Psychopharmacology*, 28(9), 874-880. doi:10.1177/0269881114532858
- 10 Pompili, A., Furukawa, T. A., Efthimiou, O., Imai, H., Tajika, A., & Salanti, G. (2018).
11 Dismantling cognitive-behaviour therapy for panic disorder: a systematic review and
12 component network meta-analysis. *Psychological medicine*, 48(12), 1945–1953.
13 doi:10.1017/s0033291717003919
- 14 Poon, C. S., Tin, C., & Yu, Y. (2007). Homeostasis of exercise hyperpnea and optimal
15 sensorimotor integration: the internal model paradigm. *Respiratory physiology &*
16 *neurobiology*, 159(1), 1-13. doi:10.1016/j.resp.2007.02.020
- 17 Radomsky, A. S., Rachman, S., Thordarson, D. S., McIsaac, H. K., & Teachman, B. A.
18 (2001). The claustrophobia questionnaire. *Journal of anxiety disorders*, 15(4), 287-297.
19 doi:10.1016/s0887-6185(01)00064-0
- 20 Rafferty, G. F., & Gardner, W. N. (1996). Control of the respiratory cycle in conscious
21 humans. *Journal of Applied Physiology*, 81(4), 1744-1753.
22 doi:10.1152/jappl.1996.81.4.1744
- 23 Ramsay, D. S., & Woods, S. C. (1997). Biological consequences of drug administration:
24 Implications for acute and chronic tolerance. *Psychological Review*, 104(1), 170-193.
25 doi:10.1037/0033-295x.104.1.170

INTEROCEPTIVE FEAR CONDITIONING

- 1 Rapee, R. M., Brown, T. A., Antony, M. M., & Barlow, D. H. (1992). Response to
2 hyperventilation and inhalation of 5.5% carbon dioxide-enriched air across the DSM-
3 III—R anxiety disorders. *Journal of Abnormal Psychology*, *101*(3), 538–552.
4 doi:10.1037/0021-843x.101.3.538.
- 5 Rapee, R., Mattick, R., Murrell, E. (1986). Cognitive mediation in the affective component
6 of spontaneous panic attacks. *Journal of Behavior Therapy and Experimental Psychiatry*,
7 *17*, 245–253. doi:10.1016/0005-7916(86)90059-5
- 8 Rappaport, L. M., Sheerin, C., Savage, J. E., Hettema, J. M., & Roberson-Nay, R. (2017).
9 Clinical characteristics of latent classes of CO₂ hypersensitivity in adolescents and young
10 adults. *Behaviour research and therapy*, *93*, 95-103. doi:10.1016/j.brat.2017.03.015
- 11 Rassovsky, Y., & Kushner, M. G. (2003). Carbon dioxide in the study of panic disorder:
12 issues of definition, methodology and outcome. *Anxiety Disorders*, *17*(1), 1-32.
13 doi:10.1016/s0887-6185(02)00181-0
- 14 Rassovsky, Y., Kushner, M. G., Schwarze, N. J., & Wangenstein, D. O. (2000).
15 Psychological and Physiological predictors of response to carbon dioxide challenge in
16 individuals with panic disorder. *Journal of Abnormal Psychology*, *109*(4), 616-623.
17 doi:10.1037/0021-843x.109.4.616
- 18 Rau, H., Elbert, T., Lutzenberger, W., Eves, F., et al. (1988). Pavlovian conditioning of
19 peripheral and central components of the baroreceptor reflex. *Journal of*
20 *Psychophysiology*, *2*(2), 119-127.
- 21 Razran, G. (1961). The observable and the inferable conscious in current Soviet
22 psychophysiology: Interoceptive conditioning, semantic conditioning, and the orienting
23 reflex. *Psychological review*, *68*(2), 81-147. doi:10.1037/h0039848
- 24 Read, D. J. C. (1967). A clinical method for assessing the ventilatory response to carbon
25 dioxide. *Australasian annals of medicine*, *16*(1), 20-32. doi:10.1111/imj.1967.16.1.20

INTEROCEPTIVE FEAR CONDITIONING

- 1 Rescorla, R. A., & Wagner, A. R. (1972). A theory of Pavlovian conditioning: Variations in
2 the effectiveness of reinforcement and nonreinforcement. *Classical conditioning II:
3 Current research and theory*, 2, 64-99.
- 4 Resnik, J., Sobel, N., & Paz, R. (2011). Auditory aversive learning increases discrimination
5 thresholds. *Nature neuroscience*, 14(6), 791. doi:10.1038/nn.2802
- 6 Rossi, V., Vanlessen, N., Bayer, M., Grass, A., Pourtois, G., & Schacht, A. (2017).
7 Motivational salience modulates early visual cortex responses across task sets. *Journal of
8 Cognitive Neuroscience*, 29(6), 968-979. doi:10.1162/jocn_a_01093
- 9 Sapienza, C. M., Davenport, P. W., & Martin, A. D. (2002). Expiratory muscle training
10 increases pressure support in high school band students. *Journal of Voice*, 16(4), 495-501.
11 doi:10.1016/s0892-1997(02)00125-x
- 12 Sclafani, A., & Ackroff, K. (2012). Role of gut nutrient sensing in stimulating appetite and
13 conditioning food preferences. *American Journal of Physiology-Regulatory, Integrative
14 and Comparative Physiology*, 302(10), R1119-R1133. doi:10.1152/ajpregu.00038.2012
- 15 Schmitz, A., & Grillon, C. (2012). Assessing fear and anxiety in humans using the threat of
16 predictable and unpredictable aversive events (the NPU-threat test). *Nature Protocols*,
17 7(3), 527–532. doi:10.1038/nprot.2012.001
- 18 Schroyen, M., Davenport, P., Van den Bergh, O., & Van Diest, I. (in press). Respiratory
19 perception: From sensation to symptom. In J. E. Cotes, J. E., R. Maynard, B. Nemery, P.
20 Wagner, B. Cooper (Eds.), *Lung function: physiology, measurement and application in
21 medicine, 7th edition*. Oxford: John Wiley & Sons.
- 22 Schroyen, M., Fantoni, S., Rivera, C., Vervliet, B., Schruers, K., Van den Bergh, O., & Van
23 Diest, I. (2016). Defensive activation to (un) predictable interoceptive threat: The NPU
24 respiratory threat test (NPUr). *Psychophysiology*, 53(6), 905-913.
25 doi:10.1111/psyp.12621

INTEROCEPTIVE FEAR CONDITIONING

- 1 Schroyen, M., Pappens, M., Schruers, K., Van den Bergh, O., Vervliet, B., & Van Diest, I.
2 (2015). Generalization of fear to respiratory sensations. *Behavior therapy*, 46(5), 611-
3 626. doi:10.1016/j.beth.2015.05.004
- 4 Schruers, K., Esquivel, G., van Duinen, M., Wichers, M., Kenis, G., Colasanti, A., ... &
5 Smeets, H. (2011). Genetic moderation of CO2-induced fear by 5-HTTLPR genotype.
6 *Journal of psychopharmacology*, 25(1), 37-42. doi:10.1177/0269881110372543
- 7 Seligman, M. E. (1971). Phobias and preparedness. *Behavior therapy*, 2(3), 307-320.
8 doi:10.1016/s0005-7894(71)80064-3
- 9 Sevenster, D., Beckers, T., & Kindt, M. (2014). Fear conditioning of SCR but not the startle
10 reflex requires conscious discrimination of threat and safety. *Frontiers in behavioral*
11 *neuroscience*, 8, 32. doi:10.3389/fnbeh.2014.00032
- 12 Siegel, S. (1989). Pharmacological conditioning and drug effects. In A. J. Goudie, & M. W.
13 Emmett-Oglesby, *Psychoactive drugs: Tolerance and sensitization* . Clifton, NJ: Humana
14 Press.
- 15 Skinner, M. D., & Aubin, H. J. (2010). Craving's place in addiction theory: contributions of
16 the major models. *Neuroscience & Biobehavioral Reviews*, 34(4), 606-623.
17 <https://doi.org/10.1016/j.neubiorev.2009.11.024>
- 18 Sinha, S., Papp, L. A., & Gorman, J. M. (2000). How study of respiratory physiology aided
19 our understanding of abnormal brain function in panic disorder. *Journal of Affective*
20 *Disorders*, 61, 191-200. doi:10.1016/s0165-0327(00)00337-2
- 21 Soeter, M., & Kindt, M. (2011). Noradrenergic enhancement of associative fear memory in
22 humans. *Neurobiology of learning and memory*, 96(2), 263-271.
23 doi:10.1016/j.nlm.2011.05.003
- 24 Somjen, G. G. (1992). The missing error signal—regulation beyond negative feedback.
25 *Physiology*, 7(4), 184-185. doi:10.1152/physiologyonline.1992.7.4.184

INTEROCEPTIVE FEAR CONDITIONING

- 1 Stegen, K., De Bruyne, K., Rasschaert, W., Van de Woestijne, K. P., & Van den Bergh, O.
2 (1999). Fear-relevant images as conditioned stimuli for somatic complaints, respiratory
3 behavior, and reduced end-tidal pCO₂. *Journal of Abnormal Psychology*, 108(1), 143-
4 152. doi:10.1037/0021-843x.108.1.143
- 5 Stekelenburg, J. J., & Van Boxtel, A. (2002). Pericranial muscular, respiratory, and heart
6 rate components of the orienting response. *Psychophysiology*, 39(6), 707-722.
7 doi:10.1111/1469-8986.3960707
- 8 Stockhorst, U., Steingrüber, H. J., & Scherbaum, W. A. (2000). Classically conditioned
9 responses following repeated insulin and glucose administration in humans. *Behavioural*
10 *brain research*, 110(1), 143-159. doi:10.1016/s0166-4328(99)00192-8
- 11 Tansey, E. M. (2006). Pavlov at home and abroad: His role in international physiology.
12 *Autonomic Neuroscience*, 125(1), 1-11. doi:10.1016/j.autneu.2006.01.013
- 13 Taugher, R. J., Lu, Y., Wang, Y., Kreple, C. J., Ghobbeh, A., Fan, R., ... & Wemmie, J. A.
14 (2014). The bed nucleus of the stria terminalis is critical for anxiety-related behavior
15 evoked by CO₂ and acidosis. *Journal of Neuroscience*, 34(31), 10247-10255.
16 doi:10.1523/jneurosci.1680-14.2014
- 17 Telch, M. J., Jacquin, K., Smits, J. A. J., & Powers, M. B. (2003). Emotional responding to
18 hyperventilation as a predictor of agoraphobia status among individuals suffering from
19 panic disorder. *Journal of Behavior Therapy*, 34, 161-170. doi:10.1016/s0005-
20 7916(03)00037-5
- 21 Tipton, M. J., Harper, A., Paton, J. F., & Costello, J. T. (2017). The human ventilatory
22 response to stress: rate or depth?. *The Journal of physiology*, 595(17), 5729-5752.
23 doi:10.1113/jp274596

INTEROCEPTIVE FEAR CONDITIONING

- 1 Uematsu, A., Tsurugizawa, T., Uneyama, H., & Torii, K. (2010). Brain–gut communication
2 via vagus nerve modulates conditioned flavor preference. *European Journal of*
3 *Neuroscience*, 31(6), 1136-1143. doi:10.1111/j.1460-9568.2010.07136.x
- 4 Valdivia-Salas, S., Forsyth, J. P., Berghoff, C. R., & Ritzert, T. R. (2014). Using
5 panicogenic inhalations of carbon dioxide enriched air to induce attentional bias for
6 threat: Implications for the development of anxiety disorders. *Cognition and Emotion*,
7 28(8), 1474-1482. doi:10.1080/02699931.2014.883964
- 8 van Beek, N., & Griez, E. (2000). Reactivity to a 35% CO₂ challenge in healthy first-degree
9 relatives of patients with panic disorder. *Biological Psychiatry*, 47(9), 830-835.
10 doi:10.1016/s0006-3223(99)00265-6
- 11 Van den Bergh, O., Kempynck, P. J., Van De Woestijne, K. P., Baeyens, F., & Eelen, P.
12 (1995). Respiratory learning and somatic complaints: a conditioning approach using
13 CO₂-enriched air inhalation. *Behaviour research and therapy*, 33(5), 517-527.
14 doi:10.1016/0005-7967(94)00080-4
- 15 Van den Hout, M. A., van der Molen, G. M., Griez, E., & Lousberg, H. (1987). Specificity
16 of interoceptive fear to panic disorders. *Journal of Psychopathology and Behavioral*
17 *Assessment*, 9(1), 99-106. doi:10.1007/bf00961635
- 18 Van der Molen, G. M., & van den Hout, M. A. (1988). Expectancy effects on respiration
19 during lactate infusion. *Psychosomatic Medicine*, 50(4), 439–443. doi:10.1097/00006842-
20 198807000-00011
- 21 Van Diest, I. (2017). Interoceptive fear conditioning and generalization in panic disorder.
22 *Oral presentation at the Anxiety and Depression conference of the Anxiety & Depression*
23 *Association of America (ADAA), San Francisco, April 2017.*

INTEROCEPTIVE FEAR CONDITIONING

- 1 Van Diest, I., Bradley, M. M., Guerra, P., Van den Bergh, O., & Lang, P. J. (2009). Fear-
2 conditioned respiration and its association to cardiac reactivity. *Biological psychology*,
3 80(2), 212-217. doi:10.1016/j.biopsycho.2008.09.006
- 4 Van Diest, I., De Peuter, S., Piedfort, K., Bresseleers, J., Devriese, S., Van de Woestijne, K.
5 P., & Van den Bergh, O. (2006). Acquired lightheadedness in response to odors after
6 hyperventilation. *Psychosomatic medicine*, 68(2), 340-347.
7 doi:10.1097/01.psy.0000204782.49159.79
- 8 Van Diest, I., Proot, P., Van de Woestijne, K. P., Han, J. N., Devriese, S., Winters, W., &
9 Van den Bergh, O. (2001a). Critical conditions for hyperventilation responses: The role
10 of autonomic response propositions during emotional imagery. *Behavior Modification*,
11 25(4), 621-639. doi:10.1177/0145445501254008
- 12 Van Diest, I., Winters, W., Devriese, S., Vercamst, E., Han, J. N., Van de Woestijne, K. P.,
13 & Van den Bergh, O. (2001b). Hyperventilation beyond fight/flight: Respiratory
14 responses during emotional imagery. *Psychophysiology*, 38, 961-968. doi:10.1111/1469-
15 8986.3860961
- 16 Vlemincx, E., Sprenger, C., & Büchel, C. (2017). A breath of air: Nocebo-effects in
17 dyspnea. *Biological Psychology*, 129, 376. doi:10.1016/j.biopsycho.2017.08.023
- 18 Vollmer, L. L., Strawn, J. R., & Sah, R. (2015). Acid–base dysregulation and chemosensory
19 mechanisms in panic disorder: a translational update. *Translational psychiatry*, 5(5),
20 e572. doi:10.1038/tp.2015.67
- 21 von Leupoldt, A., Sommer, T., Kegat, S., Baumann, H. J., Klose, H., Dahme, B., & Büchel,
22 C. (2008). The unpleasantness of perceived dyspnea is processed in the anterior insula
23 and amygdala. *American Journal of Respiratory and Critical Care Medicine*, 177(9),
24 1026-1032. doi:10.1164/rccm.200712-1821oc

INTEROCEPTIVE FEAR CONDITIONING

- 1 Wan, L., Van Diest, I., De Peuter, S., Bogaerts, K., Oyen, N., Hombroux, N., ... & Van den
2 Bergh, O. (2008). Repeated experiences of air hunger and ventilatory behavior in
3 response to hypercapnia in the standardized rebreathing test: effects of anxiety. *Biological*
4 *psychology*, 77(2), 223-232. doi:10.1016/j.biopsycho.2007.10.013
- 5 Wendt, J., Neubert, J., Koenig, J., Thayer, J. F., & Hamm, A. O. (2015). Resting heart rate
6 variability is associated with inhibition of conditioned fear. *Psychophysiology*, 52(9),
7 1161-1166. doi:10.1111/psyp.12456
- 8 Wiertelak, E. P., Smith, K. P., Furness, L., Mooney-Heiberger, K., Mayr, T., Maier, S. F., &
9 Watkins, L. R. (1994). Acute and conditioned hyperalgesic responses to illness. *Pain*,
10 56(2), 227-234. doi:10.1016/0304-3959(94)90098-1
- 11 Wirth, T., Ober, K., Prager, G., Vogelsang, M., Benson, S., Witzke, O., ... & Schedlowski,
12 M. (2011). Repeated recall of learned immunosuppression: evidence from rats and men.
13 *Brain, behavior, and immunity*, 25(7), 1444-1451. doi:10.1016/j.bbi.2011.05.011
- 14 Wolpe, J., & Rowan, V. C. (1988). Panic disorder: A product of classical conditioning.
15 *Behaviour research and therapy*, 26(6), 441-450. doi:10.1016/0005-7967(88)90138-6
- 16 Woods, S. W., Charney, D. S., Loke, J., Goodman, W., Redmond, D. E., & Heninger, G. R.
17 (1986). Carbon dioxide sensitivity in panic anxiety. *Archives of General Psychiatry*,
18 43(9), 900. doi:10.1001/archpsyc.1986.01800090090013
- 19 Woods, S. C., & Ramsay, D. S. (2000). Pavlovian influences over food and drug intake.
20 *Behavioural brain research*, 110(1), 175-182. doi:10.1016/s0166-4328(99)00194-1
- 21 Zaman, J., Van den Bergh, O., Fannes, S., & Van Diest, I. (2014). Learning to breathe?
22 Feedforward regulation of the inspiratory motor drive. *Respiratory physiology &*
23 *neurobiology*, 201, 1-6. doi:10.1016/j.resp.2014.05.009
- 24 Zaman, J., Vlaeyen, J. W., Van Oudenhove, L., Wiech, K., & Van Diest, I. (2015).
25 Associative fear learning and perceptual discrimination: a perceptual pathway in the

INTEROCEPTIVE FEAR CONDITIONING

development of chronic pain. *Neuroscience & Biobehavioral Reviews*, 51, 118-125.

doi:10.1016/j.neubiorev.2015.01.009

Zaman, J., Weltens, N., Ly, H. G., Struyf, D., Vlaeyen, J. W., Van den Bergh, O., ... & Van

Diest, I. (2016). Influence of interoceptive fear learning on visceral perception.

Psychosomatic medicine, 78(2), 248-258. doi:10.1097/psy.0000000000000257

Zandbergen, J., & Bright, M. (1991). Higher lifetime prevalence of respiratory diseases in

panic disorder?. *The American journal of psychiatry*, 148(11), 1583-1585.

doi:10.1176/ajp.148.11.1583

Ziemann, A. E., Allen, J. E., Dahdaleh, N. S., Drebot, I. I., Coryell, M. W., Wunsch, A. M.,

... & Wemmie, J. A. (2009). The amygdala is a chemosensor that detects carbon dioxide

and acidosis to elicit fear behavior. *Cell*, 139(5), 1012-1021.

doi:10.1016/j.cell.2009.10.029

Zinbarg, R., Brown, T., Barlow, D., & Rapee, R. (2001). Fear responses to hyperventilation

and 5.5% carbon dioxide enriched air and anxiety sensitivity: A reanalysis of Rapee,

Brown, Antony, & Barlow (1992). *Journal of Abnormal Psychology*, 10, 372-377.

Zucker, N., Mauro, C., Craske, M., Wagner, H. R., Datta, N., Hopkins, H., ... & Mayer, E.

(2017). Acceptance-based interoceptive exposure for young children with functional

abdominal pain. *Behaviour research and therapy*, 97, 200-212.

doi:10.1016/j.brat.2017.07.009

Zvolensky, M. J. (2001). Evaluating differential predictions of emotional reactivity during

repeated 20% carbon dioxide-enriched air challenge. *Cognition and Emotion*, 15(6), 767-

786. doi:10.1080/02699930143000284

INTEROCEPTIVE FEAR CONDITIONING

1 **Footnotes**

2

3 (1) Findings indicated that only the group with a low intensity load as the CS+ and a high intensity

4 as CS- showed the typical pattern of differential fear responding and a intensity-based

5 generalization gradient for US-expectancy ratings. In contrast, participants who received the high

6 intensity CS+ and low intensity CS- displayed impaired differential learning and complete

7 generalization of fear (Schroijen et al., 2015).

8

INTEROCEPTIVE FEAR CONDITIONING

Table 1. Fear conditioning studies with exteroceptive CSs and CO₂-inhalation as US

Authors & publication year	Study design	Phases	N	US CO ₂	CS	CR self-report	CR physiol	Effects
Forsyth & Eiffert (1996)	differential	hab (#4)	63	20%	4 video clips	SUDs*	HR*	stronger acq, slower ext with fear-relevant CSs
	W: Fear relevance CS	acq (#8)	48% men	20 s	25s		SCR*	
	B: body relevance of CS (2 cells)	ext (#10)					frontalis EMG	
Forsyth & Eiffert (1998)	non-differential	hab (#4)	96	13% 20%	3 video clips	SUDs*	HR*	stronger acq, slower ext in HR and SCR for fear-relevant CSs
	B: Fear relevance CS, US intensity (2x3cells)	acq (#8)	50% men	20 s	25 s		SCR*	
	no control group	ext (#10)					frontalis EMG	
Forsyth, Palav, & Duff (1999)	same study as Forsyth & Eiffert (1998)					SUDs*	SCR*	AS associated to SUDs*, but not to SCR*
Stegen, De Bruyne, Rasschaert, Van de Woestijne, & Van den Bergh (1999)	differential	acq (#3)	56	5.5%	4 imagery scripts	anxiety* panic-like symptoms*	resp*, ETCO ₂ *, HR	only with claustrophobic script as CS+
	B: Fear relevance (2 cells)	ext (#2)	0% men	120 s	2 min			
Fannes, Van Diest, Meulders, De Peuter, Vansteenwegen,	differential	acq (#3)	55	20%	2 unpleasant odors	US-exp* anxiety*	resp*	conditioned avoidance

INTEROCEPTIVE FEAR CONDITIONING

& Van den Bergh (2008)		ext (#3)	50% men	30 s			EtCO ₂ *	in resp
	B: odor serving as CS+/- (2 cells)				40 s			
Finlay & Forsyth (2009)	B: Context change (3 cells: control, A-B-A, A-A-B)	hab (#3)	61	20%	2 geometric shapes	SUDs;*	SCR*	ren only in A-B-A group
		acq (#6)	54% men	20 s		arousal* control*		
		ext (#8)			25 s	fearfulness*		
		ren (#4)				panic*		
De Cort, Griez, Büchler, Schruers (2012)	B: Script content (3 cells)	acq (#3)	33	35%	3 imagery scripts	fear* panic symptoms	HR resp* EtCO ₂	only with suffocation script as CS+
		ext (#2)	42% men	1 inhalation	1 min			
Valdivia-Salas, Forsyth, Berghoff, & Ritzert (2014)	differential	hab (#2)	27	20%	nonwords	SUDs*	SCR*	attentional bias for CS+
		acq (#7)	48% men	20 s	26 s	intensity of bodily reaction* fear* panic*		relative to CS-

1 *Note.* *physiol:* physiological; CS: conditional stimulus; US: unconditional stimulus; CR: conditioned reaction; B: between subject factor; W: within subject factor; *
2 means that the authors concluded that conditioning effects were observed for this outcome; hab: habituation; acq: acquisition; ext: extinction; ren: renewal; SUDS:
3 Subjective Units of Distress Scale; US-exp: US expectancy rating; HR: heart rate; SCR: skin conductance response; resp: respiration; EtCO₂: end-tidal carbon dioxide
4 level; AS: Anxiety Sensitivity

INTEROCEPTIVE FEAR CONDITIONING

Table 2. Fear conditioning studies with interoceptive CSs and CO₂-inhalation as US

Authors & publication year	Study design	Phases	N	US CO ₂	CS	CR self-report	CR physio I	Effects
Acheson, Forsyth, Prenoveau, & Bouton (2007)	non-differential	hab (#1)	42	20%	20% CO ₂	SUDs*	SCR*	unpaired >
	B: P / UP / CS-only	acq (#6)	50% men	15 s	5 s	fear*		paired >
		ext (#6)						CS only
Acheson, Forsyth, & Moses (2012)	non-differential	hab (#4)	104	20%	20% CO ₂	SUDs*	SCR*	paired >
	B: P / UP / UP° / CS-only	acq (#6)	50% men	15 s	5 s	fear*		unpaired°
		ext (#10)						
De Cort et al. (2017)	differential	hab (#1)	44	35%	infusion	US-exp*	SCR	only with
	B: pentagas-trine / saline as CS+	acq (#3)	39% men	1 inhal- ation	penta- gastrine /saline	anxiety		pentagas- trine as CS+
		ext (#3)						

Note. physiol: physiological; CS: conditional stimulus; US: unconditional stimulus; CR: conditioned reaction; B: between subject factor; P: CS and US are paired; UP: CS and US are unpaired; ° means that an exteroceptive discrimination cue was presented concurrently with the CS; * means that the authors concluded that conditioning effects were observed for this outcome; hab: habituation; acq: acquisition; ext: extinction; SUDS: Subjective Units of Distress Scale; US-exp: US expectancy rating; SCR: skin conductance response

INTEROCEPTIVE FEAR CONDITIONING

1

Table 3. Fear conditioning studies with exteroceptive CSs and limitations to breathe as US								
Authors & publication year	study design	phases	N	US	CS	CR self-report	CR physiol	Effects
Campbell et al. (1964)	non-differential	hab	9	respiratory	70 dB	free report*	fluct SC*	no ext
	B: P / UP / CS-only	acq (#1)	alcohol-depent persons	paralysis by scoline infusion	tone		resp*	
		ext (#100: #30 at t0 #30 at t1 #40 at t2)	100% men	90-103 s	50 s		muscle tension* HR*	
Pappens et al. (2013)	non-differential	hab (#2)	42	flow resistor	neutral picture	fear*	resp*	acq for all outcomes
	B: P / UP	acq (#6)	19% men	40 cmH ₂ O/l/s	8 s		startle*	no ext for physiol outcomes ratings
		ext (#6)		20 s				
Pappens et al. (2015a)	differential	acq (#4)	39	occlusion 40% of BHT	6 geo-metrical shapes	US-exp*	SCR*	acq for all outcomes
		ext (#8)	23% men				resp*	no ext for resp
					8 s		startle*	

2

3

4

Note. physiol: physiological; CS: conditional stimulus; US: unconditional stimulus; CR: conditioned reaction; B: between subject factor; P: CS and US are paired; UP: CS and US are unpaired; BHT: breath holding time; * means that the authors concluded that conditioning effects were observed for this outcome; hab: habituation; acq: acquisition; ext: extinction; fluct SC: spontaneous fluctuations in skin conductance; SCR: skin conductance response; resp: respiration

INTEROCEPTIVE FEAR CONDITIONING

1
2

Table 4 . Fear conditioning studies with interoceptive CSs and limitations to breathe as US

Authors and publication year	study design	phases	N	US	CS interoceptive	CR self-report	CR physiol	Effects
Pappens et al. (2012c)	non-differential B: P / UP	hab (#1) acq (#6) ext (#6)	40 25% men	occlusion 40% of BHT	flow resistor 10 cmH ₂ O/l/s 8 s	US-exp*	SCR* startle*	acq for all outcomes no ext for startle
Pappens et al. (2014)	<i>same design, partially same data as Pappens et al. (2012)</i>		57** 18% men			US-exp*	SCR* startle*	association of resting HRV with ext learning
Pappens et al. (2013)	non-differential B: P / UP	hab (#2) acq (#6) ext (#6)	74 22% men	flow resistor 40 cmH ₂ O/l/s 20 s	flow resistor 10 cm H ₂ O/l/s 8 s	fear rating	resp* startle*	acq + ext only for resp and startle
Pappens et al. (2015b); <i>replication of Pappens et al. (2013)</i>	non-differential B: P / UP	hab (#3) acq (#6) ext (#6)	56 9% men	flow resistor 40 cmH ₂ O/l/s 20 s	flow resistor 10 cm H ₂ O/l/s 8 s	US-exp	SCR resp* startle *	acq + ext only for resp and startle
Benke et al. (2018)	W: P / UP / COnly	acq (#8)	34 44% men	flow resistor maximally tolerable 20 s	flow resistor slight dyspnea 15 s	Valence Arousal	startle *	acq only for startle and for persons with high FoS

INTEROCEPTIVE FEAR CONDITIONING

Schroijen et al. (2015)	differential	hab (#3)	43	occlusion 40% of BHT	2 PTLs 6 cmH ₂ O 25 cmH ₂ O	US-exp*	SCR	acq only for US-exp and for low intensity load as CS+
		acq (#8)	33% men				resp	
		gen (#2)					startle	
Van Diest (2017)	non-differential	hab (#3)	55	occlusion 40% of BHT	flow resistor 15 cmH ₂ O/l/s	US-exp*	SCR*	acq for all outcomes
		acq (#8)	(15 PD / 40 healthy)				startle *	
		gen (#3)	33% men					

Note. *physiol:* physiological; *CS:* conditional stimulus; *US:* unconditional stimulus; *CR:* conditioned reaction; *B:* between subject factor; *P:* CS and US are paired; *UP:* CS and US are unpaired; *BHT:* breath holding time; * means that the authors concluded that conditioning effects were observed for this outcome; ** To increase the power to study interindividual differences in safety learning and , the original dataset of Pappens et al. (2012) was extended with another 17 participants (all women); *hab:* habituation; *acq:* acquisition; *ext:* extinction; *gen:* generalization; *HRV:* heart rate variability; *US-exp:* US expectancy rating; *SCR:* skin conductance response; *resp:* respiration; *FoS:* Fear of Suffocation; *PTLs:* pressure threshold loads; *PD:* persons with panic disorder; *IA:* interoceptive accuracy

INTEROCEPTIVE FEAR CONDITIONING

1 Figure Caption

2

3

4

5 Figure 1. Figure 1A (left panel) depicts the load-load paradigm. Credit: Irina Strelnikova/Shutterstock.com

6

7 Figure 1B (right panel) depicts the load-occlusion paradigm. Credit: Good_Stock/Shutterstock.com

8

9

10

Figure 1A (left panel).
Load-load paradigm. Credit: Irina Strelnikova/Shutterstock.com

Figure 1B (right panel).
Load-occlusion paradigm. Credit: Good_Stock/Shutterstock.com

CS = conditioned stimulus; US = unconditional stimulus; CR= conditioned response; UR = unconditioned response; BHT% = percentage of an individual's post-expiratory breath holding time; SCR: skin conductance response