Synthesis and Conformation of Pentopyranoside Nucleoside Phosphonates

Xiaochen Li, Shrinivas Dumbre, Eveline Lescrinier, Elisabetta Groaz, and Piet Herdewijn*
Medicinal Chemistry, Rega Institute for Medical Research, KU Leuven, Herestraat 49, Bus 1041, 3000 Leuven, Belgium

GRAPHICAL ABSTRACT

ABSTRACT

In contrast to natural nucleosides, where the nucleobase is positioned at the anomeric center, we report the synthesis of pentopyranoside nucleosides with a phosphonate functionality at the 1'-anomeric oxygen. Starting from L-arabinose, key functionalized *L-glycero*- and *L-erythro*-pentopyranose carbohydrate synthons were prepared and further elaborated into the final six-membered ring nucleosides via nucleobase incorporation and phosphonomethylation reactions. NMR analysis demonstrated that these nucleoside phosphonates exist in solution as conformers predominantly adopting a chair structure in which the base moiety is equatorially positioned. Such conformation prevents unfavorable 1,3-diaxial steric and electronic interactions. Notably, the stereochemical outcome of the Vorbrüggen glycosylation step utilized en route to the thymine analogue clearly suggests the absence of anchimeric assistance, as opposed to what is usually observed during nucleoside synthesis using protected furanose precursors. The finding that the diphosphates of the compounds developed in this study are recognized by DNA polymerases is important in view of the future selection of artificial genetic systems and dedicated polymerases as well as applications in therapy.

INTRODUCTION

Advances in the application of nucleosides and oligonucleotides as antiviral/anticancer therapeutics as well as gene silencing and editing tools largely depend on the systematic exploration of related chemical space through nucleobase and/or backbone modifications.¹⁻⁴ In DNA and RNA, the hydrophobic nitrogenous nucleobases encode information and ensure genetic fidelity through the formation of Watson-Crick pairs, while the furanose sugar and phosphate backbone moieties define the structure and stability of the genetic material. Chemical variations at the sugar moiety have been frequently exploited to modulate the biological and physicochemical properties of these molecules by inducing changes in nucleoside conformation without interfering with the accuracy of base pairing. This strategy has proven highly effective in altering enzyme recognition and activation by kinases and polymerases as well as binding affinity towards RNA or DNA targets.

Our group previously prepared nucleoside analogues comprising a hexopyranose sugar mimic, i.e., 1,5-anhydro-2,3-dideoxy- β -D-arabinohexitol, linked to the nucleobase at C2' rather than the anomeric position.⁵ In contrast to pyranosyl nucleosides, anhydrohexitol derivatives were found to preferentially adopt a slightly distorted chair conformation with an axial orientation of the nucleobase [**1a**, Figure 1(A)] instead of the usual equatorial position [**1a'**, Figure 1(A)].⁵⁻⁶ This is presumably a consequence of the sterically unfavorable 1,3-diaxial repulsive interactions in the six-membered chair. Such an atypical sugar conformation resulted in both distinguishing biological activities and physicochemical properties at the monomer and oligonucleotide level, respectively. For example, hexitol nucleosides proved to be potent and highly selective inhibitors of herpes simplex virus type 1 and 2 (HSV-1 and HSV-2).⁵ With regard to the corresponding oligomers, hexitol nucleic acid (HNA) sequences demonstrated the ability to fold into helical structures⁷ as well as form stable hybrids with natural counterparts,⁸ by reproducing the A-type form of dsRNA.⁹ HNA has been also proven to act as xenobiotic genetic material (XNA) capable of evolution in vitro,¹⁰⁻¹¹ while directing DNA synthesis in *E. coli*.¹²

However, while HNA oligonucleotides are resistant against cellular nucleases,⁷ their phosphorylated nucleoside precursors might be dephosphorylated by phosphatases in a cellular environment. On the other hand, HNA nucleosides are phosphorylated only in herpes virus infected cells by the viral thymidine kinase and not in non-infected cells. One important approach to obtain nucleoside phosphate mimics is to replace the natural P-O-C bond with a phosphonate (P-C-O) linkage.¹³⁻¹⁴ The phosphonate moiety is a bioisostere of the phosphate group with similar polar and electronic properties, which may function just like its natural counterpart within an artificial information system. The improved stability of phosphonate mimics may also lead to prolonged intracellular half-lives.

Figure 1. (A) Conformational preference of a hexitol nucleoside; (B) Structural diversity between hexitol nucleoside monophosphates (**1**) and target *D-threo*-pentopyranoside nucleoside phosphonates **2a** and **2b**.

A number of methods are available in the literature for the preparation of five-membered cyclic as well as acyclic nucleoside phosphonates,¹³⁻¹⁵ however in the case of six-membered ring analogues the synthesis is complicated by the greater conformational diversity arising from the pyranosyl moiety. A general method to generate these derivatives has not yet been described and reported properties of the corresponding monomers/oligomers have therefore been limited.¹⁶⁻²⁰ In this context, we set out to investigate plausible synthetic pathways that would enable the construction of a suitably functionalized pentopyranoside scaffold as key intermediate for further nucleobase incorporation and glycosylation reactions.

Herein, we report the results of our synthetic studies that ultimately led to phosphonomethyl *D-threo* pentopyranoside nucleoside analogues **2a** and **2b** [Figure 1(B)] bearing a 4'-nucleobase, which differ from hexitol nucleoside monophosphates [**1**, Figure 1(B)] by the presence of an O-C-P bond at the anomeric position, along with their conformational analysis and preliminary biological data.

RESULTS AND DISCUSSION

Nucleoside phosphonates are usually synthesized either by initial insertion of the nucleobase on the glycone moiety followed by phosphonomethylation (S_N2 or glycosylation reaction) or *vice versa*. The key disconnections in our retrosynthetic analysis of target *D-threo*-pentopyranoside nucleoside phosphonates **2a,b** are shown in Figure 2. For both proposed routes A and B, it was envisaged that the synthesis of a suitably functionalized *L-glycero*-pentopyranose (**II**) or *L-erythro*-pentopyranose (**IV**) synthon could begin from a five carbon L-sugar; thus, *L*-arabinose **3** was chosen as common starting material.

Figure 2. Retrosynthetic analysis of planned *D-threo*-pentopyranoside nucleoside phosphonates **2a** and **2b**.

Since the glycosidic linkage between the anomeric OH and the phosphonate functionality might be susceptible to epimerization during other synthetic manipulations that need to be performed along the overall synthetic path, route A in Figure 2, which entails the introduction of this group towards the end of the synthetic sequence, was initially pursued in order to minimize this possibility. Thus, benzyl isopropylidene arabinoside **4** was readily synthesized from L-arabinose **3** in four steps and used without need of chromatographic purification (Scheme 1).²¹ The key step entailed the inversion of configuration of the 2-hydroxyl group via TEMPO-mediated oxidation, followed by reduction of the resultant 2-ketone. Given that the glycosylation step for the introduction of the phosphonomethyl moiety usually requires the anchimeric assistance from a neighboring 2-*O*-acyl group, the 2-hydroxyl of compound **4** was protected as an acetate, affording **5** in excellent yield. While initial attempts to hydrolyze the isopropylidene acetal of **5** in the presence of *p*-toluenesulfonic acid resulted in the concurrent cleavage of the 2-*O*-acetyl group, the use of 80% aq. acetic acid was found to allow for selective cleavage of the acetal functionality to afford diol **6**. The next step in the synthesis entailed the regioselective protection of the equatorially oriented 4-hydroxy group, however only an inseparable mixture of products was obtained in the presence of TBDMSCl due to concurrent silylation at different positions of the sugar ring. To minimize 3-*O*-silylation, a bulkier reagent such as TBDPSCl was employed, which successfully led to the highly predominant formation of the desired compound **7**. Subsequently, 3-hydroxy glycoside **7** was subjected to Barton-McCombie deoxygenation, however no desired product (**8**) was formed upon reaction with a variety of sulfur reagents such as CS₂ MeI, 1,1'-dithiocarbonyldiimidazole (TCDI), and *O*-phenyl chlorothionoformate. This result may be attributed to the axial orientation as well as the increased steric hindrance of the bulky 4-OTBDPS moiety, which might prevent the formation of the corresponding *O*-thiocarbonyl intermediate **8**. Alternatively, compound **7** was converted to 3-*O*-triflate derivative **9**, which underwent successive S_N2 iodination and radical reduction to furnish fully protected sugar synthon **11**. Later, desilylation at the 4-

position afforded coupling partner **12** required for the construction of 1'-*O*-phosphonomethyl-D-*threo*-pentapyranosides carrying a nucleobase at the 4'-position.

Scheme 1. Initial Routes for the Preparation of Functionalized L-Glycero-pentopyranoside Scaffolds **12, **14a-c**, and **16**.^a**

^aReagents and conditions: (a) BnOH, AcCl, 75%; (b) DMP, CSA, DMF, 96%; (c) TEMPO, NaClO, KBr, DCM, H₂O, 90%; (d) NaBH₄, CH₃OH, 90%; (e) Ac₂O, Et₃N, DMAP, DCM, 93%; (f) 80% aq. AcOH, 83%; (g) TBDPSCl, imidazole, DCM, 71%; (h) Tf₂O, pyridine, DCM, 95%; (i) NaI, DMF, 96%; (j) Bu₃SnH, AIBN, toluene, 94%; (k) TBAF, THF, 87%; (l) thymine or *N*³-benzoylthymine, PPH₃, DIAD or DEAD, THF or dioxane, rt; (m) MsCl or TsCl, Et₃N/pyridine, DCM; (n) TBAF, THF, 95%.

However, all attempts to introduce thymine or *N*³-benzoyl protected thymine as nucleobase by reacting **12** under Mitsunobu conditions⁵ resulted in the formation of E2 elimination product **13** rather than the desired nucleoside, most likely due to the axial orientation of the 4-hydroxyl group (for detailed conditions, see Experimental Section). Further efforts to achieve the nucleophilic substitution of 4-OTs and 4-OMs substituted analogues **14a,b** by thymine in the presence of different bases such as DBU, NaH, Cs₂CO₃, and K₂CO₃ alone or in combination with a variety of crown ethers were also unsuccessful. Moreover, the conversion of the 4-OH of **12** into a better leaving group using triflic anhydride did not provide the expected activated product, but rather led to a 3,4-elimination reaction (**13**) and nucleophilic substitution with pyridine at the 4-position. Therefore, another option for the introduction of the nucleobase was investigated focusing on the preparation of a 3,4-epoxy sugar, which could in principle undergo a ring-opening reaction upon nucleophilic attack of a heterocyclic nitrogen. Initially, the envisaged epoxide **16** was directly obtained in one step from 3-iodo derivative **10** upon removal of the 4-*O*-silyl protecting group (Scheme 1). It was found that the basic conditions due to the deprotecting reagent TBAF favored efficient epoxide formation once the 4-hydroxyl group was unmasked. However, when epoxide **16** was reacted with thymine under basic (DBU) and high temperature (110 °C) conditions, conversion to the desired nucleoside occurred

concomitantly with the migration and hydrolysis of the 2'-acetyl group leading to a complex mixture of products.

This problem was addressed by adopting an alternative synthetic strategy that would allow to generate a differently substituted key sugar intermediate with higher stability under base-sugar coupling conditions (Scheme 2). In particular, a Corey-Winter olefination was employed to install the alkene at the 3,4-position to obtain the L-*glycero*-pentopyranose ring. Treatment of **6** with TCDI afforded cyclic thionocarbonate **17**, which subsequently underwent a *syn*-elimination in the presence of triethyl phosphite to furnish 3,4-unsaturated pentose **13** in good yield (84% over two steps). It should be noted that in this case the standard use of trimethyl phosphite was ineffective to promote the desulfurization step. Epoxidation of **13** using either *m*-CPBA or H₂O₂ did not proceed stereoselectively, while resulting in a mixture of diastereomeric epoxides. Hydrolysis of the 2-*O*-acetyl group of **13** prior to epoxidation was therefore necessary, in order for the corresponding free hydroxyl moiety to establish a hydrogen bonding interaction with the oxidizing agent that allowed for the reaction to selectively take place from the same side of the 2-hydroxy group. Thus, epoxide **19** was obtained in good yield, and successively subjected to MEM protection, epoxide ring opening with thymine, and Barton deoxygenation to yield 4'-thymidylated sugar **22**. The MEM protecting group was then removed under acidic conditions affording a 15:1 mixture of α - and β -anomers **23a** and **23b**, which were characterized after column chromatography separation, and further benzoylated at the 2' position to furnish **24a** and **24b** in an overall 92% yield. While the Pd/C-catalyzed hydrogenolysis of the anomeric OBn functionality proceeded sluggishly in either methanol or acetic acid (3 days), a significant enhancement of the reaction rate was observed when the debenzoylation was conducted in THF as solvent (8 h). Later, we found that treatment of a mixture of **24a** and **24b** under transfer hydrogenation conditions (cyclohexene in ethanol) produced the desired glycone **25** in 88% yield.

For the final glycosylation step, glycone **25** was first activated at the anomeric position either as a *N*-phenyltrifluoroacetimidate (PTFAI, **26a**), trichloroacetimidate (TCAI, **26b**), or acetate (**26c**), while diisopropyl phosphonmethanol was used as glycosyl acceptor. A variety of standard conditions for the formation of the O-C-P bond were screened and are summarized in Table 1. Glycosylated product **27** could not be isolated as a pure compound by silica gel column chromatography at this stage due to diisopropylphosphonmethanol contamination, thus it was used as such in the following debenzoylation step, and product characterization was carried out at the level of compound **28**. Surprisingly, glycosyl donors **26a** and **26b** yielded either starting material **25** (entries 1 and 3, Table 1) or a product with undesired stereochemistry at the anomeric center (β -**28**, entries 2 and 4, Table 1). The formation of β -**28** suggests the non-participation of the 2'-*O*-acyl moiety, with the aglycone acceptor approaching the oxocarbenium intermediate from the β -face to form the 1,4-*anti* conformer. In contrast, 1'-acetylated nucleoside donor **26c** furnished the desired α -**28** isomer as major compound (α -**28**: β -**28** = 2:1) in the presence of a strong Lewis acid (TMSOTf) at room temperature. It can be postulated that at low temperature (-78 °C) the presence of aglycone moieties with an increased leaving group ability as those in **26a** and **26b** support the formation of product β -**28** as a result of the absence of neighboring group assistance, whereas in the case of **26c** bearing a poorer leaving group the reaction favors isomer α -**28**. Final removal of the diisopropyl ester

functionalities under standard conditions afforded nucleoside phosphonate **2a** in 40% yield (0.98% overall yield).

Scheme 2. Synthesis of Thymine α -D-Threo-pentopyranoside Nucleoside Phosphonate **2a.^a**

^aReagents and conditions: (a) TCDI, DCM, 83%; (b) triethylphosphite, 140 °C, 85%; (c) MeONa, methanol, 91%; (d) *m*-CPBA, DCM, 89%; (e) 2-Methoxyethoxymethyl chloride (MEMCl), DCM, 91%; (f) thymine, DBU, DMF, 85%; (g) TCDI, DMAP, DCM; (h) Bu₃SnH, AIBN, toluene, 83% over two steps; (i) 1M HCl in dioxane, 75% for α -**25** and 5% for β -**25**; (j) BzCl, pyridine, 0 °C, 92%; (k) Pd/C, cyclohexene, ethanol, 88%; (l) Ac₂O, pyridine, 85%; (m) TMSOTf, hydroxymethyl phosphonic acid diisopropyl ester, CH₂Cl₂; (n) 7 M NH₃ in methanol, 20% over two steps; (o) TMSBr, 2,6-lutidine, acetonitrile, 40%.

Table 1. Optimization of the Phosphonylation Step

entry	glycosyl donor	Lewis acid and T (°C)	product (yield%)
1	26a^b	TMSOTf, -78 °C	25
2		BF ₃ .Et ₂ O, -78 °C	<i>β</i>-28 (30%) ^a
3	26b^b	TMSOTf, -78 °C	25
4		BF ₃ .Et ₂ O, -78 °C	<i>β</i>-28 (40%) ^a
5	26c	TMSOTf, rt	<i>α</i>-28/<i>β</i>-28 (20/10%) ^a
6		BF ₃ .Et ₂ O, rt	<i>β</i>-28 (traces) ^a

***α*-28** ***β*-28**

^aYields refer to the glycosylated product isolated after the following debenzoylation step.

^bExperimental procedures for the phosphonylation of **26a** and **26b** are detailed in the Supporting Information (S192-S193).

By analogy to early work towards thymine D-*threo*-pentopyranoside phosphonate **2a**, we initially investigated the nucleophilic substitution of 4-sulfonate activated sugars **14a-c** as well as ring opening reactions of epoxide **20** with 6-chloropurine and *N*⁶-benzoyl protected adenine (Scheme S1, in Supporting Information). However, all attempts to directly introduce the purine ring at the 4-position by changing the base or using various additives failed.

Therefore, we opted for the construction of the purine ring via a linear approach. As shown in Scheme 3, radical deoxygenation of previously synthesized thionocarbonate derivative **17** afforded 3-deoxy compound **12** in 60% yield along with small amounts of the undesired 4-deoxy product (10%), under high dilution reaction conditions (0.02 mol/L). It should be noted that when the reaction was conducted at a concentration of 0.10 mol/L, the reduction of the C=S bond to a methylene moiety (-CH₂) took place instead of the desired deoxygenation reaction at the 3-position. Mesylation and subsequent azidation of the 4-hydroxyl group occurred smoothly to afford 4-azido compound **29** in good yield. After deacetylation at the 2-position, compound **30** was reacted under Staudinger conditions to give 4-amino-D-*threo*-pentopyranoside **31**. Next, a literature procedure was employed for the construction of the purine ring starting from precursor **31** and 4,6-dichloro-5-formamidopyrimidine.²² The initially formed pyrimidine nucleoside intermediate underwent in situ cyclization under basic conditions to smoothly provide 6-chloropurine pyranoside **32**. Subsequent amination at the 6-position of purine in **32** yielded adenine analogue **33** in excellent yield (95%), whose further reaction with an excess of benzoyl chloride furnished

fully protected derivative **34**. Following attempts to achieve the removal of the 1'-OBn group along with mono-debenzoylation at the primary amino group of **34** under different hydrogenation conditions in the presence of a range of Pd catalysts (10% Pd/C, 20% Pd(OH)₂/C, Pd(OAc)₂, and Pd-black) did not deliver chemoselectively glycone **35**, due to the contamination of the desired compound with other fully and partially *N*-debenzoylated by-products. However, the use of a Lewis-acid such as boron trichloride enabled benzyl removal with the exclusive formation of the desired anomeric hydroxyl glycone **35** in good yield. After acetylation of the anomeric 1'-OH, the resulting glycosyl donor **36** was reacted with diisopropyl phosphonmethanol under conditions similar to those developed earlier for the phosphomethylation of thymine analogue **2a** (Table 1, entry 5). To our dismay, none of these conditions provided the desired target phosphonate glycoside **37** and similar attempts to prepare a 6-chloropurine containing nucleoside starting from **32** through a similar sequence of steps also met with failure.

Scheme 3. Building of the Purine Ring on 4-Amino-D-Threo-pentopyranoside Sugar **31.**

^aReagents and conditions: (a) Bu₃SnH, AIBN, toluene; (b) MsCl, TEA, DMAP, DCM, 54% over two steps; (c) NaN₃, DMF, 91%; (d) MeONa, MeOH, 95%; (e) triphenyl phosphine, THF then water, 92%; (f) 4,6-dichloro-5-formamidopyrimidine, DIPEA, BuOH, 75%; (g) NH₃ in ethanol, 95%; (h) BzCl, pyridine, 88%; (i) 1 M BCl₃, DCM, -78 to 0 °C, 80%; (j) Ac₂O, pyridine, 78%; (k) TMSOTf, hydroxymethyl phosphonic acid diisopropyl ester, CH₂Cl₂.

In view of these results, we therefore decided to turn to route B in our retrosynthetic plan (Figure 2). As illustrated in Scheme 4, the amino moiety of sugar scaffold **31** was first protected with a trifluoroacetyl group to yield **38**, whose benzyl group was later cleaved by Pd/C catalytic hydrogenation. Diol **39** was then transformed into 1,2-diacetyl glycosyl donor **40**, which underwent smooth glycosylation with diisopropyl

phosphonomethanol at the 1-position affording **41**.²³ Subsequent trifluoroacetyl deprotection under basic conditions provided key amino sugar intermediate **42**, which served as synthon for the stepwise construction of the purine nucleobase. The final adenine nucleoside phosphonate **2b** was successfully obtained in 40% yield (0.78%, overall yield) following the cleavage of the diisopropyl esters with TMSBr in the presence of lutidine. It is worth mentioning that 1-*O*-diisopropylphosphonomethyl-3,4-dideoxy-4-amino- α -D-*threo*-pentopyranoside **42** might constitute a versatile intermediate for the construction of a variety of purine or pyrimidine nucleobases.²²

Scheme 4. Synthesis of Adenine α -D-*threo*-pentopyranoside Nucleoside Phosphonate **2b.**

^aReagents and conditions: (a) ethyl trifluoroacetate, TEA, methanol, 91%; (b) Pd/C, H₂, methanol, 82%; (c) Ac₂O, pyridine, 80%; (d) hydroxymethyl phosphonic acid diisopropyl ester, TMSOTf, DCM, rt; (e) 7 M NH₃ in methanol, 40% over two steps; (f) 4,6-dichloro-5-formamidopyrimidine, DIPEA, BuOH, 75%; (g) NH₃ in ethanol, 92%; (h) TMSBr, acetonitrile, 40%.

A detailed solution conformational analysis of the final phosphonate products **2a** and **2b** was carried out using NMR spectroscopy. A preference for a 1,4-*syn* substitution pattern (Figure 3) was established for both pyrimidine and purine derivatives **2a** and **2b** following complete assignment using 1D and 2D NMR experiments and based on the values of the coupling constants observed in the ¹H NMR spectra. Table 2 summarizes the proton NMR spectroscopic data acquired for thymine derivative **2a**. The corresponding data obtained for adenine containing nucleoside phosphonate **2b** can be found in the Supporting Information (Table S2) together with all relevant spectra.

Figure 3. Schematic representation of the solution phase preferential conformation of 1'-phosphonate D-threo-pentopyranose nucleoside bearing thymine as base (**2a**) as determined by NMR studies. Dashed lines indicate the observed NOE interaction between the H6 of thymine and the axially positioned H3' and H5'.

As shown in the ^1H NMR spectrum of **2a** (see Supporting Information), H1' (δ 4.68) and H2' (δ 4.04) appear as a doublet and triplet of doublets, respectively, with a small $^3J_{1,2'}$ diequatorial coupling constant, which is consistent with a chair conformation with both phosphonomethyl group and 2'-OH occupying an axial orientation. The td pattern of H2' with the largest splitting equal to 3.6 Hz is characteristic of an equatorial position of this proton. Furthermore, the H4' signal at δ 4.79 consists of a triplet of triplets arising from the coupling with the protons at C3' and C5'; notably, the observed $^3J_{4',3'}$ and $^3J_{4',5'}$ values of 12.1 and 10.8 Hz between H4' and H3'a and H5'a, respectively, support an equatorial orientation of the nucleobase at the 4'-position. This chair conformation of the pyranose ring was further confirmed by the presence of a clear NOE correlation between the H6 of thymine and the axially positioned H3' and H5'.

Such 1,4-*syn*-relationship is similar to that previously reported between the heterocyclic base and either the 5'-hydroxymethyl to be phosphorylated in 1,5-anhydrohexitol nucleosides⁵ or the 4'-*O*-phosphonomethyl moiety in 2',3'-dideoxy-2',3'-didehydro-pentopyranosyl¹⁶ nucleoside phosphonates. However, nucleoside phosphonates **2a,b** differ in their conformational behavior from **1** by the orientation of the nucleobase, which is equatorial rather than axial, with both the hydroxymethyl and secondary OH groups at the 1'- and 2'-positions axially positioned. Thus, it can be assumed that the preferred conformation in **2a,b** is sterically enforced in order to avoid the 1,3-diaxial steric interactions among bulky substituents and/or between the lone pairs of the ring oxygen and the anomeric oxygen, along with the opposite direction of the dipole moment.

For comparison, the β -anomer of compound **2a** (**β -S2a**) was obtained upon deprotection of **β -28** and fully characterized by 1D and 2D NMR experiments (for details, see Table S3 and Figure S3 in the Supporting Information). Accordingly, the analysis of these NMR data indicated that the preferred chair conformation of the pyranose ring in **β -S2a** adjusts to a 1,4-*trans*-diaxial orientation of the bulky substituents.

It is noteworthy that the conformation of such nucleoside analogues is expected to easily adapt to external conditions such as those required by enzyme binding²⁴ or insertion within a nucleic acid chain,²⁵ thus influencing oligonucleotides properties in a variable way.

Table 2. ¹H NMR Analysis of the Sugar Skeleton of Thymine D-Threo-pentopyranoside Nucleoside Phosphonate **2a(600 MHz, D₂O).**

bond	orientation ^a	δ (ppm)	multiplicity	<i>J</i> (Hz) (D ₂ O)	³ <i>J</i> connection
C-H1'	e	4.68	d	2.2	H1'e-H2'e
C-H2'	e	4.04	td	2.2–3.0 3.6	H2'e-H1'e H2'e-H3'e H2'e-H3'a
C-H3'	a	2.38	ddd	3.0 12.1 13.8	H3'a-H2'e H3'a-H4'a H3'a-H3'e
C-H3''	e	1.93	dtd	1.2 3.9 13.1	H3'e-H2'e H3'e-H4'a H3'e-H3'a
C-H4'	a	4.79	tt	10.8 4.8	H4'a-H3'a H4'a-H5'a H4'a-H3'e H4'a-H5'e
C-H5'	a	3.91	t	10.7	H5'a-H5'e H5'a-H4'a
C-H5''	e	3.73	ddd	1.4 4.7 11.2	H5'e-H3'e ^b H5'e-H4'a H5'e-H5'a

^ae stands for equatorial, a for axial. ^bW proton-proton coupling (⁴*J*_{HH}).

In order to define whether D-*threo*-pentopyranoside nucleoside phosphonates are still recognized by enzymes involved in nucleoside and nucleotide metabolism, compounds **2a,b** were at first evaluated for their antiviral activity against different herpesviruses (varicella zoster virus (VZV), HSV-1, and HSV-2). Contrary to the results obtained for **1**⁵ as well as known furanose nucleoside phosphonates,²⁶ these studies revealed that **2a,b** exerted poor inhibitory activity of DNA viral replication. This could be due to a lack of cell penetration (the negatively charged phosphonate moiety is very polar) as well as inefficient intracellular enzymatic phosphorylation. Specifically, the very high EC₅₀ values determined for **2a,b** (Table S4, in the Supporting Information) suggest that these issues cannot be overcome by synthesizing the corresponding prodrugs.

Furthermore, aiming to gain knowledge about the potential utility of these synthetic analogues for the selection of artificial information systems, the adenine containing nucleoside phosphonate **2b** was converted to its corresponding diphosphate **45** using a known phosphorylation procedure (Scheme 5).²⁷ Compound **45** was assessed for its ability to be accepted as substrate by DNA polymerases in an enzymatic primer extension assay where a primer-template duplex with a seven 5'-dT overhang was used. Enzymes lacking 3',5'-proofreading activity such as Klenow fragment (*exo*-) and Vent (*exo*-) were tested in the presence or absence of Mn²⁺ ions and shown to be able to use **45** as substrate, although only the formation

of P+1 or P+2 products was detected, as exemplified in Figure 4 (for a complete account of used conditions and results, see Supporting information, Figure S1 and S2).

Scheme 5. Phosphorylation of Adenine α -D-*Threo*-pentopyranoside Phosphonate **2b.^a**

^aReagents and conditions: (a) (i) 1,1'-carbonyldiimidazole, DMF; (ii) (HNBu₃)₂H₂P₂O₇, (ii) TEAB buffer, 30%.

Figure 4. Primer extension of compound **45** in the presence of Vent polymerase (*exo*-) and Mn²⁺ ions. The positions **45** has to be incorporated opposite the template oligonucleotide are in bold. dA stands for 2'-deoxyadenosine triphosphate (positive control), while Neg for negative control.

CONCLUSION

In summary, D-*threo*-pentopyranoside nucleoside analogues bearing a phosphonate functionality rather than a nucleobase (i.e., in natural nucleosides) at the anomeric center have been synthesized starting from L-arabinose. Specifically, stereocontrolled routes to suitably functionalized key carbohydrate scaffolds have been established and optimized. Such intermediates were further assembled and elaborated into the final six-membered ring nucleosides using two main strategies entailing the initial insertion of the nucleobase on the glycone moiety followed by phosphonomethylation or by inverting this reaction sequence. Notably, when a thymine containing pentopyranose glycosyl donor was subjected to Vorbrüggen glycosylation conditions, the reaction proceeded without neighboring group participation, in contrast to the usual behavior of protected furanose nucleoside precursors. The synthesis of the adenine analogue relied on a linear approach for building the purine base on a phosphonometylated 4-amino pentopyranoside synthon. This method allowed for an improved stereoselectivity and could potentially be extended to the stepwise construction of other purine as well as pyrimidine bases. According to NMR studies, the solution

conformation of such pentopyranoside nucleoside phosphonates prefers an equatorial orientation of the nucleobase that circumvents unfavorable 1,3-diaxial interactions, in contrast to HNA nucleosides. Preliminary primer incorporation reactions using a mesophilic and thermophilic polymerase revealed that the compounds synthesized by these pathways constitute potentially useful monomers for medicinal and biotechnological applications.

EXPERIMENTAL SECTION

General Information. All reagents and solvents were purchased from commercial sources and used as obtained. Moisture sensitive reactions were performed using oven-dried glassware under a nitrogen or argon atmosphere. NMR spectra were recorded on a Bruker Advance 300 MHz (^1H NMR, 300 MHz; ^{13}C NMR, 75 MHz; ^{31}P NMR, 121 MHz), 500 MHz (^1H NMR, 500 MHz; ^{13}C NMR, 125 MHz; ^{31}P NMR, 202 MHz), or 600 MHz (^1H NMR, 600 MHz; ^{13}C NMR, 150 MHz) spectrometer with tetramethylsilane as internal standard or referenced to the residual solvent signal, and 85% H_3PO_4 for ^{31}P NMR. All intermediates and final compounds were characterized by using 2D NMR (^1H -COSY, HSQC, NOESY, and HMBC) spectroscopic techniques. For NMR assignment of sugar protons and carbons, prime numbering is used. High-resolution mass spectra [HRMS (ESI)] were obtained on a quadrupole orthogonal acceleration time-of-flight mass spectrometer (Synapt G2 HDMS, Waters, Milford, MA). Samples were infused at 3 $\mu\text{L}/\text{min}$ and spectra were obtained in positive (or in negative) ionization mode with a resolution of 15000 (fwhm) using leucine enkephalin as lock mass. Precoated aluminum sheets (254 nm) were used for TLC. Products were purified by column chromatography on silica gel (60 $\text{\AA}$, 0.035–0.070 mm, Acros Organics). Preparative RP-HPLC purifications were carried out on a Phenomenex Gemini 110A column (C18, 10 μm , 21.2 mm $\times$ 250 mm) using $\text{CH}_3\text{CN}/0.05 \text{ M TEAB}$ buffer or $\text{H}_2\text{O}/\text{CH}_3\text{CN}$ as eluent gradient.

Benzyl 2-*O*-acetyl-3,4-*O*-isopropylidene- β -L-ribopyranoside (5). Benzyl 3,4-*O*-isopropylidene- β -L-ribopyranoside **4** was prepared in 4 steps from L-arabinose **3** following a literature procedure that did not require column chromatography purification.²¹ Next, to a stirred solution of **4** (34.7 g, 123.8 mmol), DMAP (0.76 g, 6 mmol) and triethylamine (69.0 mL, 495.2 mmol) in DCM (500 mL), acetic anhydride (23.4 mL, 247.6 mmol) was added dropwise at 0 $^\circ\text{C}$. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. After completion of the reaction, the mixture was cooled to 0 $^\circ\text{C}$ and quenched with saturated aq. NaHCO_3 . The organic layer was washed with brine (200 mL), dried over Na_2SO_4 , and concentrated in vacuo. The resulting residue was purified by column chromatography on silica gel (10:1, hexane/EtOAc, R_f = 0.3) to afford **5** (37.3 g, 93% yield) as a white solid. ^1H NMR (300 MHz, CDCl_3) δ 7.38–7.26 (m, 5H, Ph), 5.00 (dd, J = 6.1, 3.3 Hz, 1H, H-2), 4.93 (d, J = 6.2 Hz, 1H, H-1), 4.80 (d, J = 12.3 Hz, 1H, CH_2Ph), 4.80 (d, J = 12.3 Hz, 1H, CH_2Ph), 4.57 (dd, J = 7.0, 3.2 Hz, 1H, H-3), 4.56 (d, J = 12.2 Hz, 1H, CH_2Ph), 4.32 (dt, J = 7.0, 1.9 Hz, 1H, H-4), 3.87 (dd, J = 13.1, 2.5 Hz, 1H, H-5), 3.71 (dd, J = 13.1, 1.6 Hz, 1H, H-5'), 2.13 (s, 3H, CH_3CO), 1.53 (s, 3H, $[\text{C}(\text{CH}_3)_2]$), 1.33 (s, 3H, $[\text{C}(\text{CH}_3)_2]$); ^{13}C NMR (75 MHz, CDCl_3) δ 170.3 (CH_3CO), 137.6, 128.5, 127.8, 127.7 (Ph), 110.4 $[\text{C}(\text{CH}_3)_2]$, 96.9 (C-1), 73.6 (C-2), 71.8 (CH_2Ph), 70.3 (C-3), 69.5 (C-4), 62.5 (C-5), 26.5 $[\text{C}(\text{CH}_3)_2]$, 25.3 $[\text{C}(\text{CH}_3)_2]$, 21.2 (CH_3CO); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{17}\text{H}_{22}\text{O}_6\text{Na}$ 345.1309; Found 345.1308.

Benzyl 2-*O*-acetyl- β -L-ribose (6). A solution of **5** (20.0 g, 62.04 mmol) in 80% aq. acetic acid (400 mL) was stirred at 60 °C for 2 h. After removal of all the volatiles under reduced pressure, the residue was partitioned between water (200 mL) and EtOAc (300 mL). The water layer was extracted with EtOAc (1 $\times$ 200 mL). The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The resulting residue was purified by column chromatography on silica gel (100:5 DCM/methanol, R_f = 0.25) to afford **6** (14.6 g, 83%) as a white solid. ¹H NMR (500 MHz, DMSO-d₆) δ 7.41–7.18 (m, 5H, Ph), 5.10 (d, J = 4.7 Hz, 1H, OH-3), 4.76 (d, J = 7.0 Hz, 1H, H-1), 4.73 (d, J = 12.4 Hz, 1H, CH₂Ph), 4.67 (d, J = 7.0 Hz, 1H, OH-4), 4.55 (dd, J = 6.8, 2.9 Hz, 1H, H-2), 4.54 (d, J = 12.3 Hz, 1H, CH₂Ph), 3.96 (dt, J = 4.3, 2.6 Hz, 1H, H-3), 3.69–3.55 (m, 3H, H-4, H-5 and H-5'), 2.11 (s, 3H, CH₃CO); ¹³C NMR (125 MHz, DMSO-d₆) δ 169.6 (CH₃CO), 137.8, 128.3, 127.5, 127.4 (Ph), 97.1 (C-1), 72.0 (C-2), 69.5 (CH₂Ph), 67.8 (C-3), 66.7 (C-4), 63.5 (C-5), 20.9 (CH₃CO); HRMS (ESI-TOF) m/z : [M+Na]⁺ calcd for C₁₄H₁₈O₆Na 305.0996; Found 305.0998.

Benzyl 2-*O*-acetyl-4-*O*-*tert*-butyldiphenylsilyl- β -L-ribose (7). To a stirred solution of **6** (10.0 g, 35.41 mmol) and imidazole (6.03 g, 88.5 mmol) in dry acetonitrile (300 mL) at 0 °C, a solution of *tert*-butyldiphenylchlorosilane (10.1 mL, 38.9 mmol) in anhydrous acetonitrile (50 mL) was added dropwise. The reaction mixture was stirred at room temperature for 8 h. After removal of all the volatiles under reduced pressure, the resulting residue was partitioned between water (100 mL) and EtOAc (200 mL). The water layer was extracted with EtOAc (1 $\times$ 200 mL). The combined organic layers were washed with brine (150 mL), dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by column chromatography on silica gel (10:1 hexane/EtOAc, R_f = 0.3) to afford **7** (13.0 g, 71% yield) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.75–7.62 (m, 4H, Ph), 7.47–7.36 (m, 6H, Ph), 7.32–7.22 (m, 5H, Ph), 4.92 (d, J = 5.2 Hz, 1H, H-1), 4.82 (dd, J = 5.1, 3.6 Hz, 1H, H-2), 4.71 (d, J = 12.0 Hz, 1H, CH₂Ph), 4.50 (d, J = 12.0 Hz, 1H, CH₂Ph), 4.10 (t, J = 3.3 Hz, 1H, H-3), 3.92 (dt, J = 5.9, 3.9 Hz, 1H, H-4), 3.63–3.49 (m, 2H, H-5 and H-5'), 2.12 (s, 3H, CH₃CO), 1.09 (s, 9H, [C(CH₃)₃]); ¹³C NMR (75 MHz, CDCl₃) δ 170.4 (CH₃CO), 137.3, 135.8, 135.75, 133.2, 132.5, 130.3, 130.2, 128.5, 128.1, 128.0, 127.8, 127.6 (Ph), 97.3 (C-1), 71.3 (C-2), 70.1 (CH₂Ph), 69.1 (C-4), 67.6 (C-3), 63.1 (C-5), 27.0 [C(CH₃)₃], 21.2 (CH₃CO), 19.4 [C(CH₃)₃]; HRMS (ESI-TOF) m/z : [M+Na]⁺ calcd for C₃₀H₃₆O₆SiNa 543.2174; Found 543.2167.

Benzyl 2-*O*-acetyl-4-*O*-*tert*-butyldiphenylsilyl-3-*O*-trifluoromethanesulfonyl- β -L-ribose (9). To a stirred solution of **7** (10.0 g, 19.22 mmol) and pyridine (4.8 mL, 57.7 mmol) in dry DCM (300 mL) at 0 °C, triflic anhydride (6.5 mL, 38.44 mmol) was added dropwise. The reaction mixture was stirred at room temperature for 1 h. After completion of the reaction, the mixture was cooled to 0 °C and quenched with saturated aq. NaHCO₃. The reaction mixture was further washed with saturated aq. NaHCO₃ (100 mL) and brine (100 mL). The organic layer was dried over Na₂SO₄ and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography (silica gel was presoaked with 0.5% of triethylamine in hexane; chromatography was performed using 15:1 hexane/EtOAc, R_f = 0.2) to afford **9** (11.9 g, 95% yield) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.79–7.58 (m, 4H, Ph), 7.54–7.15 (m, 11H, Ph), 5.31 (dd, J = 2.4, 2.2 Hz, 1H, H-2), 4.82 (dd, J = 7.3, 2.4 Hz, 1H, H-3), 4.74 (d, J = 12.0 Hz, 1H, CH₂Ph), 4.72 (d, J = 7.2 Hz, 1H, H-1), 4.50 (d, J = 12.0 Hz, 1H, CH₂Ph), 3.94 (ddd, J = 9.4, 5.8, 2.3 Hz, 1H, H-4), 3.56 (dd, J = 11.6, 9.5 Hz, 1H, H-5), 3.39 (dd, J = 11.6, 4.8 Hz, 1H, H-5'), 2.08 (s, 3H, CH₃CO), 1.08 (s, 9H, [C(CH₃)₃]);

^{13}C NMR (75 MHz, CDCl_3) δ 169.6 (CH_3CO), 135.9, 135.85, 133.0, 130.6, 130.5, 128.6, 128.3, 128.1, 127.7 (Ph), 120.8 (CF_3), 97.2 (C-1), 86.0 (C-3), 70.9 (C-2), 68.7 (CH_2Ph), 67.3 (C-4), 63.7 (C-5), 26.8 [$\text{C}(\text{CH}_3)_3$], 20.7 (CH_3CO), 19.2 [$\text{C}(\text{CH}_3)_3$]; HRMS (ESI-TOF) m/z : $[\text{M}+\text{NH}_4]^+$ calcd for $\text{C}_{31}\text{H}_{39}\text{NF}_3\text{O}_8\text{SSi}$ 670.2118; Found 670.2119.

Benzyl 2-*O*-acetyl-4-*O*-*tert*-butyldiphenylsilyl-3-deoxy-3-iodo- β -L-xylopyranoside (10). A suspension of **9** (4.00 g, 6.12 mmol) and NaI (2.76 g, 18.38 mmol) in THF (70 mL) was stirred at room temperature for 12 h. After removal of all the volatiles under reduced pressure, the resulting residue was separated between EtOAc (100 mL) and saturated aq. NaHCO_3 (50 mL). The water layer was extracted with EtOAc (50 mL) once. The combined organic layers were washed with brine (50 mL), dried over Na_2SO_4 , and concentrated in vacuo. The residue was purified by column chromatography on silica gel (15:1 hexane/EtOAc) to afford **10** (3.71 g, 96% yield) as a colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.94–7.62 (m, 4H, Ph), 7.56–7.10 (m, 11H, Ph), 5.06 (dd, J = 10.5, 7.2 Hz, 1H, H-2), 4.76 (d, J = 12.4 Hz, 1H, CH_2Ph), 4.49 (d, J = 12.4 Hz, 1H, CH_2Ph), 4.29 (d, J = 7.2 Hz, 1H, H-1), 4.14–3.87 (m, 2H, H-3 and H-4), 3.64 (dd, J = 11.6, 4.4 Hz, 1H, H-5), 3.28–3.03 (m, 1H, , H-5'), 2.07 (s, 3H, CH_3CO), 1.09 (s, 9H, [$\text{C}(\text{CH}_3)_3$]); ^{13}C NMR (75 MHz, CDCl_3) δ 169.0 (CH_3CO), 136.3, 135.9, 130.3, 130.1, 128.5, 128.0, 127.9, 127.8, 127.7 (Ph), 100.4 (C-1), 73.7 (C-2), 73.1 (C-4), 70.4 (CH_2Ph), 68.2 (C-5), 33.5 (C-3), 27.2 [$\text{C}(\text{CH}_3)_3$], 21.0 (CH_3CO), 19.7 [$\text{C}(\text{CH}_3)_3$]; HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{30}\text{H}_{35}\text{IO}_5\text{SiNa}$ 653.1193; Found 653.1196.

Benzyl 2-*O*-acetyl-4-*O*-*tert*-butyldiphenylsilyl-3-deoxy- β -L-erythro-pentopyranoside (11). To a stirred solution of **10** (2.00 g, 3.17 mmol) and AIBN (0.26 g, 1.59 mmol) in toluene (100 mL), tributyltin hydride (1.7 mL, 6.34 mmol) was added dropwise. The reaction mixture was heated at 110 °C for 1 h. After removal of all the volatiles under reduced pressure, the resulting crude residue was purified by column chromatography on silica gel (15:1 hexane/EtOAc, R_f = 0.2) to afford **11** (1.5 g, 94% yield) as a colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.70–7.60 (m, 4H, Ph), 7.45–7.34 (m, 6H, Ph), 7.33–7.25 (m, 5H, Ph), 4.75 (d, J = 12.2 Hz, 1H, CH_2Ph), 4.72–4.65 (m, 1H, H-2), 4.56 (d, J = 5.3 Hz, 1H, H-1), 4.52 (d, J = 12.2 Hz, 1H, CH_2Ph), 3.88–3.78 (m, 1H, H-4), 3.74 (ddd, J = 11.4, 3.7, 1.2 Hz, 1H, H-5a), 3.32 (dd, J = 11.4, 6.3 Hz, 1H, H-5e), 2.21 (dtd, J = 13.2, 4.6, 1.2 Hz, 1H, H-3e), 2.04 (s, 3H, CH_3CO), 1.77 (dt, J = 13.2, 7.6 Hz, 1H, H-3a), 1.06 (s, 9H, [$\text{C}(\text{CH}_3)_3$]); ^{13}C NMR (75 MHz, CDCl_3) δ 170.2 (CH_3CO), 137.6, 135.8, 133.9, 133.6, 123.0, 128.5, 127.9, 127.8 (Ph), 99.2 (C-1), 69.7 (CH_2Ph), 68.6 (C-2), 67.4 (C-5), 65.5 (C-4), 34.3 (C-3), 27.0 [$\text{C}(\text{CH}_3)_3$], 21.3 (CH_3CO), 19.3 [$\text{C}(\text{CH}_3)_3$]; HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{30}\text{H}_{36}\text{O}_5\text{SiNa}$ 527.2224; Found 527.2216.

Benzyl 2-*O*-acetyl-3-deoxy- β -L-erythro-pentopyranoside (12). To a stirred solution of **11** (1.00 g, 1.98 mmol) in THF (40 mL) at 0 °C, TBAF (1 M in THF, 3.96 mL, 3.96 mmol) was added dropwise. The reaction mixture was slowly warmed to room temperature and stirred for 5 h. After completion of the reaction, all volatiles were removed in vacuo. The resulting residue was separated between EtOAc (40 mL) and saturated aq. NaHCO_3 (20 mL). The water layer was extracted with EtOAc (30 mL) once. The combined organic layers were washed with brine (20 mL), dried over Na_2SO_4 , and concentrated in vacuo. The residue was purified by column chromatography on silica gel (gradient hexane/EtOAc, 5:1, v/v; 1:1, v/v) affording **12** (0.46 g, 87%) as a colorless oil. ^1H NMR (600 MHz, CDCl_3) δ 7.41–7.30 (m, 5H, Ph), 4.90 (dt, J = 3.2,

2.3 Hz, 1H, H-2), 4.78 (bs, 1H, H-1e), 4.77 (d, $J = 11.6$ Hz, 1H, CH_2Ph), 4.56 (d, $J = 11.6$ Hz, 1H, CH_2Ph), 4.01 (dd, $J = 12.0$, 1.8 Hz, 1H, H-5a), 3.75 (dq, $J = 2.7$, 2.4 Hz, 1H, H-4e), 3.67 (dt, $J = 11.9$, 2.3 Hz, 1H, H-5e), 2.22 (td, $J = 15.0$, 3.7 Hz, 1H, H-3a), 2.1 (s, 3H, CH_3CO), 1.96 (ddtd, $J = 15.0$, 3.2, 2.4, 0.93 Hz, 1H, H-3e); ^{13}C NMR (151 MHz, CDCl_3) δ 169.5 (CH_3CO), 137.0, 128.5, 127.9, 127.8 (Ph), 95.8 (C-1), 69.3 (CH_2Ph), 68.5 (C-2), 64.5 (C-5), 64.2 (C-4), 29.3 (C-3), 21.2 (CH_3CO); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{14}\text{H}_{19}\text{O}_5\text{Na}$ 289.1046; Found 289.1045.

Benzyl 2-*O*-acetyl- β -L-glycero-pent-3-enopyranoside (13). This compound was first obtained as a side product during the Mitsunobu reaction (see Scheme 1), and it was later synthesized by using the triethyl phosphite method.

Mitsunobu conditions: To a stirred suspension of **12** (1 eq), thymine or Bz-thymine (2 eq), and triphenylphosphine (2.2 eq) in anhydrous dioxane or THF, a solution of DEAD or DIAD in 2 mL of THF or dioxane was slowly added. The mixture was stirred until disappearance of starting material **12**. After removal of all the volatiles under reduced pressure, the resulting residue was separated between EtOAc and saturated aq. NaHCO_3 . The water layer was extracted with EtOAc. The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel using 15:1(v/v) hexane/ EtOAc. The reaction was conducted at different temperatures (-78, -20, 0 °C, and rt), however only the elimination product **13** was formed instead of the desired nucleoside.

Triethyl phosphite conditions: A stirring solution of **17** (8.00 g, 24.7 mmol) in triethylphosphite (250 mL) was maintained at 140 °C for 14 h and the reaction was monitored by TLC for completion. After removal of all the volatiles under reduced pressure, the crude residue was subjected to column chromatography on silica gel (15:1 hexane/EtOAc) to give **13** (5.20 g, 85% yield) as a colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.40–7.27 (m, 5H, Ph), 6.08 (td, $J = 10.3$, 2.4 Hz, 1H, H-3), 5.95–5.71 (m, 1H, H-4), 5.11–4.94 (m, 1H, H-2), 4.90 (bs, 1H, H-1), 4.81 (d, $J = 12.0$ Hz, 1H, CH_2Ph), 4.62 (d, $J = 12.0$ Hz, 1H, CH_2Ph), 4.37–4.09 (m, 2H, H-5 and H-5'), 2.07 (s, 3H, CH_3CO); ^{13}C NMR (75 MHz, CDCl_3) δ 170.4 (CH_3CO), 137.4, 131.3, 128.6, 128.0 (Ph), 127.9 (C-3), 120.4 (C-4), 96.8 (C-1), 69.9 (C-2), 66.1 (CH_2Ph), 59.7 (C-5), 21.2 (CH_3CO); HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{14}\text{H}_{17}\text{O}_4$ 249.1121; Found 249.1123.

General procedure for the synthesis of benzyl 2-*O*-acetyl-3-deoxy-4-sulfonate- β -L-erythro-pentopyranosides 14a-c. To a stirred solution of **12** (0.2 g, 0.75 mmol) and pyridine (0.18 mL, 2.25 mmol) in DCM (10 mL) at 0 °C, a solution of a sulfonyl chloride in DCM (2 mL) was added dropwise. The progress of the reaction was monitored by TLC, and 0.5 mL of saturated aq. NaHCO_3 was added to quench the reaction at 0 °C. The reaction mixture was washed with saturated aq. NaHCO_3 (5 mL) and brine (5 mL). The organic layer was dried over Na_2SO_4 and concentrated in vacuo. The crude mesylate or tosylate was purified by column chromatography on silica gel to afford the desired sulfonate, after prewashing the column with 0.1% of triethylamine in hexane.

Mesylate (14a). Following the general procedure in the presence of mesyl chloride (0.12 mL, 1.5 mmol), a crude product was obtained, which was purified by column chromatography using 5:1 hexane/EtOAc as eluent to afford **14a** (0.23 g, 87% yield) as a colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.39–7.27 (m, 5H, Ph), 4.84–4.70 (m, H-1, H-2, H-4 and CH_2Ph), 4.54 (d, $J = 12.1$ Hz, 1H, CH_2Ph), 4.03 (dd, $J = 13.0$, 1.8

Hz, 1H, H-5), 3.82 (d, $J = 13.0$ Hz, 1H, H-5'), 3.01 (s, 3H, $\text{CH}_3 \text{SO}_2$), 2.29–2.21 (bs, 2H, H-3 and H-3'), 2.07 (s, 3H, CH_3CO); ^{13}C NMR (75 MHz, CDCl_3) δ 179.9 (CH_3CO), 136.7, 128.3, 127.8, 127.7 (Ph), 95.9 (C-1), 72.5 (C-2), 69.2 (CH_2Ph), 66.2 (C-4), 61.5 (C-5), 38.4 ($\text{CH}_3 \text{Ms}$), 27.8 (C-3), 20.8 (CH_3CO); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{15}\text{H}_{20}\text{O}_7\text{S}_1\text{Na}$ 367.0822; Found 367.0816.

Tosylate (14b). Following the general procedure in the presence of tosyl chloride (0.29 g, 1.5 mmol), a crude product was obtained, which was purified by column chromatography using 10:1 hexane/EtOAc to afford **14b** (0.267 g, 85% yield) as a colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.79 (d, $J = 8.1$ Hz, 2H, Ph), 7.45–7.18 (m, 7H, Ph), 4.84–4.66 (m, 3H, H-1, H-2 and CH_2Ph), 4.61–4.56 (m, 2H, H-4 and CH_2Ph), 3.95 (d, $J = 12.8$ Hz, 1H, H-5), 3.68 (d, $J = 12.8$ Hz, 1H, H-5'), 2.45 (s, 3H, $\text{CH}_3 \text{Ts}$), 2.24–2.06 (m, 2H, H-3 and H-3'), 2.04 (s, 3H, CH_3CO); ^{13}C NMR (75 MHz, CDCl_3) δ 170.3 (CH_3CO), 145.0, 137.0, 134.2, 130.0, 128.6, 128.1, 128.0, 127.8 (Ph), 96.3 (C-1), 73.0 (C-2), 69.6 (CH_2Ph), 66.6 (C-4), 61.9 (C-5), 28.0 (C-3), 21.7 ($\text{CH}_3 \text{Ts}$), 21.2 (CH_3CO); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{21}\text{H}_{24}\text{O}_7\text{S}_1\text{Na}$ 443.1135; Found 443.1135.

Benzyl 2-*O*-acetyl-3,4-anhydro- β -L-ribofuranoside (16). Following a similar procedure as that used for the synthesis of **12**, epoxide **16** was obtained starting from **10** (0.57 g, 0.32 mmol), TBAF (0.63 mL, 0.63 mmol, 1 M in THF) in THF (10 mL). The crude residue was purified by column chromatography using hexane/EtOAc (8:1, v/v; 3:1 v/v) to give **16** as a white solid (0.2 g, 95%). ^1H NMR (300 MHz, CDCl_3) δ 7.51–7.24 (m, 5H, Ph), 4.95 (dd, $J = 4.7, 3.2$ Hz, 1H, H-2), 4.74 (d, $J = 12.0$ Hz, 1H, CH_2Ph), 4.65 (d, $J = 3.0$ Hz, 1H, H-1), 4.52 (d, $J = 12.0$ Hz, 1H, CH_2Ph), 4.08 (dd, $J = 13.3, 1.6$ Hz, 1H, H-5), 3.98 (d, $J = 13.3$ Hz, 1H, H-5'), 3.59 (appt, $J = 4.0$ Hz, 1H, H-3), 3.34 (dd, $J = 3.9, 1.2$ Hz, 1H, H-4), 2.14 (s, 3H, CH_3CO); ^{13}C NMR (75 MHz, CDCl_3) δ 170.3 (CH_3CO), 137.0, 128.6, 128.1, 127.9 (Ph), 95.8 (C-1), 69.9 (C-2), 68.3 (CH_2Ph), 58.7 (C-5), 51.2 (C-3), 49.8 (C-4), 20.9 (CH_3CO); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{14}\text{H}_{16}\text{O}_5\text{Na}$ 287.0890; Found 287.0894.

Benzyl 2-*O*-acetyl-3,4-*O*-thiocarbonyl- β -L-ribofuranoside (17). A solution of **6** (10.0 g, 35.42 mmol) and TCDI (12.6 g, 70.9 mmol) in anhydrous DCM (500 mL) was stirred at room temperature for 10 h. The reaction mixture was washed with saturated aq. NaHCO_3 (200 mL). The organic layer was separated, and the aqueous layer was extracted with DCM. The combined organic layers were washed with brine and dried over Na_2SO_4 . After removal of all the volatiles under reduced pressure, the crude residue was purified by column chromatography (3:1 DCM/hexane) to afford **17** (9.60 g, 83% yield) as a white solid. ^1H NMR (300 MHz, CDCl_3) δ 7.43–7.30 (m, 5H, Ph), 5.22 (dd, $J = 7.7, 4.0$ Hz, 1H, H-4), 5.11 (appt, $J = 4.2$ Hz, 1H, H-2), 5.00 (d, $J = 4.6$ Hz, 1H, H-1), 4.98 (dd, $J = 7.8, 1.2$ Hz, 1H, H-3), 4.78 (d, $J = 11.9$ Hz, 1H, CH_2Ph), 4.60 (d, $J = 11.9$ Hz, 1H, CH_2Ph), 4.10 (dd, $J = 14.1, 0.7$ Hz, 1H, H-5), 4.01 (dd, $J = 14.1, 2.0$ Hz, 1H, H-5'), 2.15 (s, 3H, CH_3CO); ^{13}C NMR (75 MHz, CDCl_3) δ 191.2 (CS), 169.9 (CH_3CO), 136.5, 128.7, 128.4, 128.1 (Ph), 96.1 (C-1), 78.4 (C-4), 75.6 (C-3), 70.1 (CH_2Ph), 66.7 (C-2), 58.6 (C-5), 20.8 (CH_3CO); HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{15}\text{H}_{17}\text{O}_6\text{S}_1$ 325.0740; Found 325.0740.

Benzyl β -L-glycero-pent-3-enopyranoside (18). To a stirred solution of **13** (3.75 g, 15.1 mmol) in methanol (150 mL) at 0 °C, a 30% NaOMe solution in MeOH (5.59 mL, 30.2 mmol) was added dropwise. The reaction mixture was stirred at room temperature for 2 h. After completion of the reaction, all the volatiles were removed in vacuo. The remaining residue was partitioned between EtOAc (100 mL) and

saturated aq. NaHCO₃ (100 mL). The water layer was extracted with EtOAc (1 × 50 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, and concentrated in vacuo. The crude residue was purified by column chromatography on silica gel (4:1 hexane/EtOAc) to give **18** (2.86 g, 91% yield) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.56–7.24 (m, 5H, Ph), 6.08–5.78 (m, 2H, H-3 and H-4), 4.80 (d, *J* = 11.8 Hz, 1H, CH₂Ph), 4.77 (d, *J* = 2.6 Hz, 1H, H-1), 4.58 (d, *J* = 11.8 Hz, 1H, CH₂Ph), 4.27–4.03 (m, 2H, H-5 and H-5'), 3.89 (bs, 1H, H-2), 2.61 (s, 1H, OH); ¹³C NMR (75 MHz, CDCl₃) δ 137.4, 128.7, 128.5, 128.0 (Ph), 127.9 (C-3), 124.6 (C-4), 99.9 (C-1), 70.0 (CH₂Ph), 64.8 (C-2), 60.8 (C-5); HRMS (ESI-TOF) *m/z*: [M+H]⁺ calcd for C₁₂H₁₅O₃ 229.08353; Found 229.0842.

Benzyl 3,4-anhydro-β-L-ribose (19). A solution of *m*-chloroperbenzoic acid (*m*CPBA, 77%, 4.12 g, 9.46 mmol) in DCM (5 mL) was added dropwise to a stirring solution of **18** (2.60 g, 12.6 mmol) in dry DCM (120 mL) at 0 °C. After the addition was completed, the reaction mixture was stirred at 8–9 °C for 24 h. It was then cooled to 0 °C and 20% aq. Na₂S₂O₃ (25 mL) was added, and the stirring was continued for 2 h. The organic layer was separated and successively washed with saturated aq. NaHCO₃ and brine, dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by silica gel column chromatography (3:1 hexane/EtOAc) to afford **19** (2.50 g, 89% yield) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 7.46–7.22 (m, 5H, Ph), 4.72 (d, *J* = 11.8 Hz, 1H, CH₂Ph), 4.58 (d, *J* = 2.3 Hz, 1H, H-1), 4.49 (d, *J* = 11.8 Hz, 1H, CH₂Ph), 4.01 (dd, *J* = 13.4, 1.5 Hz, 1H, H-5), 3.93 (d, *J* = 13.4 Hz, 1H, H-5'), 3.82 (dd, *J* = 4.5, 2.4 Hz, 1H, H-2), 3.50 (appt, *J* = 4.5 Hz, 1H, H-3), 3.33 (ddd, *J* = 4.2, 1.5, 0.6 Hz, 1H, H-4); ¹³C NMR (75 MHz, CDCl₃) δ 137.0, 128.6, 128.1, 128.0 (Ph), 97.9 (C-1), 69.7 (CH₂Ph), 64.7 (C-2), 58.0 (C-5), 51.8 (C-3), 51.4 (C-4); HRMS (ESI-TOF) *m/z*: [M+Na]⁺ calcd for C₁₂H₁₄O₄Na 245.0785; Found 245.0791.

Benzyl 3,4-anhydro-2-O-methoxyethoxymethyl-β-L-ribose (20). To a stirred solution of **19** (2.10 g, 9.45 mmol) and DIPEA (4.90 mL, 28.4 mmol) in dry DCM (50 mL) at 0 °C was added MEMCl (2.16 mL, 18.9 mmol). After the addition was completed, the reaction mixture was stirred at room temperature for 1 h, and later at 40 °C for 8 h. The reaction mixture was washed with saturated aq. NaHCO₃ and brine. The organic layer was dried over Na₂SO₄ and concentrated in vacuo. The remaining residue was purified by column chromatography on silica gel (15:1 hexane/EtOAc) to afford **20** (2.68 g, 91% yield) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.39–7.20 (m, 5H, Ph), 4.89 (s, 2H, OCH₂O), 4.75 (d, *J* = 11.9 Hz, 1H, CH₂Ph), 4.61 (d, *J* = 4.0 Hz, 1H, H-1), 4.50 (d, *J* = 11.8 Hz, 1H, CH₂Ph), 4.07 (dd, *J* = 13.4, 2.3 Hz, 1H, H-5), 3.92 (d, *J* = 13.4 Hz, 1H, H-5'), 3.92–3.79 (m, 2H, H-2 and OCH₂CH₂O), 3.76–3.63 (m, 1H, OCH₂CH₂O), 3.56–3.47 (m, 3H, H-3 and OCH₂CH₂O), 3.39–3.31 (m, 1H, H-4), 3.35 (s, 3H, OCH₃); ¹³C NMR (75 MHz, CDCl₃) δ 137.1, 128.3, 127.7, 127.7 (Ph), 97.6 (OCH₂O), 95.1 (C-1), 71.7 (C-2), 71.6 (OCH₂CH₂O), 69.9 (CH₂Ph), 67.1 (OCH₂CH₂O), 59.9 (C-5), 58.8 (OCH₃), 51.9 (C-3), 51.7 (C-4); [M+Na]⁺ calcd for C₁₆H₂₂O₆Na 333.1309; Found 333.1317.

Benzyl 4-deoxy-2-O-(2-methoxyethoxymethyl)-4-(thymid-1-yl)-α-D-lyxopyranoside (21). To a stirred suspension of **20** (1.90 g, 6.12 mmol) and thymine (2.32 g, 18.4 mmol) in dry DMF (50 mL) at 0 °C was slowly added DBU (3 mL, 18.87 mmol). The suspension was stirred at room temperature for 30 min until all the solid dissolved and then heated at 80 °C for 8 h. After removal of all the volatiles under reduced pressure, the remaining syrup was dissolved in EtOAc and washed with saturated aq. NaHCO₃. The organic phase was collected, dried over Na₂SO₄ and concentrated in vacuo. The product was purified by silica gel

column chromatography (100:2 DCM/Methanol) to afford **21** (2.27 g, 85% yield) as a colorless oil. ^1H NMR (500 MHz, CDCl_3) δ 9.74 (s, 1H, NH), 7.35 (m, 5H, Ph), 7.03 (s, 1H, H-6), 5.02 (d, J = 1.5 Hz, 1H, H-1'), 4.87 (d, J = 7.2 Hz, 1H, OCH_2O), 4.80 (d, J = 7.2 Hz, 1H, OCH_2O), 4.76 (d, J = 11.7 Hz, 1H, CH_2Ph), 4.54 (d, J = 11.7 Hz, 1H, CH_2Ph), 4.28 (appt, J = 7.9 Hz, 1H, H-3'), 3.98 (appt, J = 2.4 Hz, 1H, H-2'), 3.89–3.66 (m, 4H, H-5', H-5' and $\text{OCH}_2\text{CH}_2\text{O}$), 3.51 (s, 2H, $\text{OCH}_2\text{CH}_2\text{O}$), 3.34 (s, 3H, OCH_3), 1.88 (s, 3H, T CH_3); ^{13}C NMR (126 MHz, CDCl_3) δ 163.9 (C-4), 151.8 (C-2), 137.1 (C-6), 137.0, 128.6, 128.1, 127.9 (Ph), 111.0 (C-5), 98.3 (C-1'), 96.7 (OCH_2O), 78.9 (C-2'), 71.5 ($\text{OCH}_2\text{CH}_2\text{O}$), 69.4 (CH_2Ph), 67.8 ($\text{OCH}_2\text{CH}_2\text{O}$), 65.9 (C-3'), 60.2 (C-5'), 58.9 (OCH_3), 54.7 (C-4'), 12.5 (T CH_3); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_8\text{Na}$ 459.1738; Found 459.1731.

Benzyl 3,4-dideoxy-2-O-(2-methoxyethoxymethyl)-4-(thymid-1-yl)- α -D-threo-pentopyranoside (22).

To a solution of **21** (1.40 g, 2.6 mmol) and DMAP (100 mg, 0.8 mmol) in anhydrous DCM (30 mL) was added TCDI (940 mg, 5.20 mmol) at room temperature. The reaction mixture was stirred at 40 °C overnight. It was then washed with brine, dried over Na_2SO_4 , and concentrated under reduced pressure. The resulting residue was dissolved in toluene (50 mL), and AIBN (170 mg, 1.10 mmol) was added followed by tributyltin hydride (1.70 mL, 4.20 mmol). The reaction mixture was refluxed for 1 h. After removal of all the volatiles under reduced pressure, the crude residue was purified by column chromatography on silica gel (gradient $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 100:0, v/v; 100:2, v/v; 30:1, v/v) to afford **22** (1.10 g, 83% yield over two steps) as a colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 9.95 (s, 1H, NH), 7.47–7.19 (m, 5H, Ph), 7.06 (s, 1H, H-6), 5.10–4.89 (m, 1H, H-4'), 4.80 (m, 4H, OCH_2O , CH_2Ph and H-1'), 4.59 (d, J = 11.1 Hz, 1H, CH_2Ph), 3.93 (bs, 1H, H-2'), 3.85–3.64 (m, 4H, H-5', H-5', $\text{OCH}_2\text{CH}_2\text{O}$), 3.51 (dd, J = 5.4, 3.6 Hz, 2H, $\text{OCH}_2\text{CH}_2\text{O}$), 3.34 (s, 3H, OCH_3), 2.30 (td, J = 12.7, 2.8 Hz, 1H, H-3'), 2.03 (dt, J = 12.6, 3.1 Hz, 1H, H-3''), 1.90 (s, 3H, T CH_3); ^{13}C NMR (75 MHz, CDCl_3) δ 164.3 (C-4), 151.4 (C-2), 137.5 (C-6), 137.0, 128.8, 128.20, 128.25 (Ph), 111.2 (C-5), 96.7 (OCH_2O), 95.2 (C-1'), 73.0 (C-2'), 71.9 ($\text{OCH}_2\text{CH}_2\text{O}$), 69.5 (CH_2Ph), 67.5 ($\text{OCH}_2\text{CH}_2\text{O}$), 61.3 (C-5'), 59.2 (OCH_3), 47.6 (C-4'), 29.2 (C-3'), 12.8 (T CH_3); HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{21}\text{H}_{29}\text{N}_2\text{O}_7$ 421.1969; Found 421.1964.

Benzyl 3,4-dideoxy-4-(thymid-1-yl)- α/β -D-threo-pentopyranoside (23). Hydrochloric acid in 1,4-dioxane (4 M, 8 mL) was added dropwise to a solution of **22** (1.20 g, 2.86 mmol) in 1,4-dioxane (24 mL), and the reaction mixture was stirred at room temperature for 8 h. After removal of all the volatiles under reduced pressure, the remaining syrup was co-evaporated with triethylamine (5 mL) once and then subjected to column chromatography on silica gel to give α/β -**23**. Data for the α -**23**: (0.72 g, 75% yield); ^1H NMR (600 MHz, CDCl_3) δ 9.96 (s, 1H, NH), 7.46–7.29 (m, 5H, Ph), 7.12 (d, J = 1.1 Hz, 1H, H-6), 5.13–4.97 (m, 1H, H-4'), 4.80 (d, J = 11.9 Hz, 1H, CH_2Ph), 4.71 (d, J = 2.4 Hz, 1H, H-1'), 4.57 (d, J = 11.9 Hz, 1H, CH_2Ph), 3.96 (bs, 1H, H-2'), 3.82 (m, 2H, H-5' and H-5''), 3.67 (d, J = 6.1 Hz, 1H, OH), 2.27 (appt d, J = 13.0, 3.3 Hz, 1H, H-3'), 2.01 (d appt, J = 13.3, 4.3 Hz, 1H, H-3'), 1.88 (d, J = 1.1 Hz, 3H, T CH_3); ^{13}C NMR (151 MHz, CDCl_3) δ 163.8 (C-4), 151.3 (C-2), 137.0 (C-6), 136.6, 128.5, 128.0, 128.0 (Ph), 111.0 (C-5), 98.3 (C-1'), 69.5 (CH_2Ph), 67.0 (C-2'), 61.4 (C-5'), 46.8 (C-4'), 30.8 (C-3'), 12.5 (T CH_3); HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{21}\text{N}_2\text{O}_5$ 333.1445; Found 333.1443.

Data for the β -**23**: (48 mg, 5% yield); ^1H NMR (500 MHz, CDCl_3) δ 9.52 (s, 1H, NH), 7.61 (d, J = 1.2 Hz, 1H, H-6), 7.42–7.29 (m, 5H, Ph), 4.92 (d, J = 2.9 Hz, 1H, H-1'), 4.84 (d, J = 11.9 Hz, 1H, CH_2Ph), 4.76–

4.52 (m, 2H, CH_2Ph , H-4'), 4.13 (dd, $J = 13.0, 3.9$ Hz, 1H, H-5'), 3.94–3.73 (m, 2H, H-5', H-2'), 2.48 (d, $J = 8.3$ Hz, 1H, OH), 2.21–2.17 (dT, $J = 12.9, 3.8$ Hz, 1H, H-3'), 2.15–2.09 (m, 1H, H-3''), 1.95–1.86 (d, $J = 1.2$ Hz, 3H, T CH_3); ^{13}C NMR (126 MHz, CDCl_3) δ 164.0 (C-4), 150.9 (C-2), 138.2, 136.4 (C-6), 128.5, 128.1, 128.0 (Ph), 110.6 (C-5), 97.3 (C-1'), 69.9 (CH_2Ph), 64.4 (C-2'), 61.1 (C-5'), 51.2 (C-4'), 31.1 (C-3'), 12.5 (T CH_3).

Benzyl 2-*O*-benzoyl-3,4-dideoxy-4-(thymid-1-yl)- α/β -D-threo-pentopyranoside (24). To a solution of α/β -**23** (400 mg, 1.20 mmol) in pyridine (15 mL) at 0 °C, benzoyl chloride (0.24 mL, 1.81 mmol) was added dropwise. The reaction mixture was stirred at 0 °C for 2 h. After removal of all the volatiles under reduced pressure, the remaining residue was partitioned between DCM (30 mL) and saturated aq. NaHCO_3 (20 mL). The aqueous layer was extracted again with DCM (20 mL). The combined organic layers were washed with saturated aq. NaHCO_3 and brine, dried over Na_2SO_4 , and concentrated in vacuo. The residue was purified by silica gel column chromatography (3:1 hexane/EtOAc) to afford α/β -**24** (483 mg, 92% yield) as a colorless oil.

Data for α -**24**: ^1H NMR (600 MHz, CDCl_3) δ 9.17 (s, 1H, NH), 8.09 (m, 2H, Ph), 7.64–7.30 (m, 8H, Ph), 7.04 (d, $J = 1.2$ Hz, 1H, H-6), 5.31 (td, $J = 3.5, 3.2$ Hz, 1H, H-2'), 5.09 (tt, $J = 11.7, 5.0$ Hz, 1H, H-4'), 4.92 (d, $J = 1.6$ Hz, 1H, H-1'), 4.82 (d, $J = 11.9$ Hz, 1H, CH_2Ph), 4.62 (d, $J = 11.9$ Hz, 1H, CH_2Ph), 3.87 (appt, $J = 10.8$ Hz, 1H, H-5'), 3.81 (ddd, $J = 10.7, 5.0, 1.9$ Hz, 1H, H-5'), 2.49 (td, $J = 13.2, 3.2$ Hz, 1H, H-3'), 2.26–2.04 (dt, $J = 13.8, 3.6$ Hz, 1H, H-3''), 1.92 (d, $J = 1.2$ Hz, 3H, T CH_3); ^{13}C NMR (151 MHz, CDCl_3) δ 165.5 (PhCO), 163.5 (C-4), 150.8 (C-2), 136.8, 136.1 (C-6), 133.4, 129.8, 129.4 (Ar-C), 128.6, 128.5, 128.3, 128.1, 127.9 (Ph), 111.3 (C-5), 95.0 (C-1'), 69.5 (CH_2Ph), 69.1 (C-2'), 60.7 (C-5'), 47.0 (C-4'), 28.2 (C-3'), 12.5 (T CH_3); HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{24}\text{H}_{25}\text{N}_2\text{O}_6$ 437.1707; Found 437.1700.

Data for β -**24**: ^1H NMR (300 MHz, CDCl_3) δ 9.36 (s, 1H, NH), 8.18–7.72 (m, 2H, Ph), 7.46 (m, 3H, Ph), 7.19 (m, 5H, Ph), 5.29 (dd, $J = 6.9, 9.9$ Hz, 1H, H-2'), 5.01 (td, $J = 5.9, 10.9$ Hz, 1H, H-4'), 4.85 (d, $J = 12.4$ Hz, 1H, CH_2Ph), 4.63 (d, $J = 12.4$ Hz, 1H, CH_2Ph), 4.58 (d, $J = 7.4$ Hz, 1H, H-1'), 4.11 (dd, $J = 3.7, 11.1$ Hz, 1H, H-5'), 3.60 (appt, $J = 11.0$ Hz, 1H, H-5''), 2.18–1.74 (m, 5H, , H-3' , H-3'' and T CH_3); ^{13}C NMR (75 MHz, CDCl_3) δ 165.4 (PhCO), 163.6 (C-4), 151.4 (C-2), 136.8 , 136.0 (C-6), 133.5, 130.0, 129.0, 128.5, 128.4, 127.9, 127.9 (Ph), 111.7 (C-5), 100.6 (C-1'), 71.1 (C-2'), 70.3 (CH_2Ph), 62.6 (C-5'), 54.0 (C-4'), 30.8 (C-3'), 12.7 (T CH_3).

2-*O*-Benzoyl-3,4-dideoxy-4-(thymid-1-yl)- α -D-threo-pentopyranose (25). To a solution of α/β -**24** (300 mg, 0.69 mmol) in ethanol (10 mL), Pd/C (366 mg, 0.34 mmol) and cyclohexene (1.39 mL, 13.8 mmol) were added. The reaction mixture was stirred at 80 °C for 6 h. It was then cooled and filtered through a pad of Celite to give **25** (209 mg, 88%) as a white solid. ^1H NMR (600 MHz, DMSO-d_6) δ 11.3 (s, 1H, NH), 8.01 (m, 2H, Ph), 7.72–7.68 (m, 1H, Ph), 7.65 (d, $J = 1.2$ Hz, 1H, H-6), 7.59–7.56 (m, 2H, Ph), 7.00 (d, $J = 4.7$ Hz, 1H, OH), 5.04 (td, $J = 4.7, 3.6$ Hz, 1H, H-2'), 5.01 (dd, $J = 4.7, 1.4$ Hz, 1H, H-1'), 4.79 (tt, $J = 10.7, 4.3$ Hz, 1H, H-4'), 4.03 (appt, $J = 10.8$ Hz, 1H, H-5'), 3.59 (ddd, $J = 10.5, 4.4, 1.7$ Hz, 1H, H-5''), 2.59 (td, $J = 13.4, 3.1$ Hz, 1H, H-3'), 2.01 (dt, $J = 13.4, 4.5$ Hz, 1H, H-3'), 1.79 (d, $J = 0.9$ Hz, 3H, T CH_3); ^{13}C NMR (151 MHz, DMSO-d_6) δ 165.0 (PhCO), 163.8 (C-4), 151.0 (C-2), 138.0, 133.7 (C-6), 129.7, 129.4, 129.3, 128.9, 128.8 (Ph), 109.2 (C-5), 89.7 (C-1'), 70.9 (C-2'), 59.8 (C-5'), 47.3 (C-4'), 27.3 (C-3'), 12.1 (T CH_3); HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{19}\text{N}_2\text{O}_6$ 347.1238; Found 347.1241.

General procedure for the synthesis of trichloroacetimidate and *N*-phenyltrifluoroacetimidate glycosylation donors **26a and **26b**.** To a stirred solution of **25** (100 mg, 0.29 mmol) and K₂CO₃ (44 mg, 0.32 mmol) in acetone (3 mL) at 0 °C, *N*-aryltrifluoroacetimidoyl chloride (120 mg, 0.58 mmol) or trichloroacetonitrile (0.06 mL, 0.58 mmol) was added dropwise. The reaction was monitored by TLC for completion. The reaction mixture was then filtered through a pad of Celite and concentrated under reduce pressure to afford the corresponding crude donor.

Data for **26a**: ¹H NMR (300 MHz, (CD₃)₂CO) δ 9.27 (s, 1H, NH), 8.00–7.97 (m, 2H, Ph), 7.59–7.50 (m, 1H, Ph), 7.42 (m, 2H, Ph), 7.31 (s, 1H, H-6), 6.21 (bs, 1H, H-1'), 5.31 (bs, 1H, H-2'), 4.89 (m, 1H, H-4'), 4.14 (t, *J* = 10.8 Hz, 1H, H-5'), 3.88 (m, 1H, H-5''), 2.76 (t, *J* = 13.3 Hz, 1H, H-3'), 2.29 (d, *J* = 13.6 Hz, 1H, H-3'), 1.68 (s, 3H, T CH₃), ¹³C NMR (75 MHz, (CD₃)₂CO) δ 165.8 (C-4), 151.8 (C-2), 138.6, 134.4 (C-6), 130.5, 129.5 (Ph), 110.9 (C-5), 93.9 (C-1'), 69.4 (C-2'), 63.1 (C-5'), 49.2 (C-4'), 28.5 (C-3'), 12.4 (T CH₃).

Data for **26b**: ¹H NMR (300 MHz, CDCl₃) δ 8.18–8.00 (m, 2H, Ph), 7.61 (m, 1H, Ph), 7.48 (m, Ph), 7.32 (m, 2H, Ph), 7.13 (m, 1H, Ph), 6.96 (bs, 1H, H-6), 6.88 (d, *J* = 7.7 Hz, 2H, Ph), 6.27 (bs, 1H, H-1'), 5.46 (bs, 1H, H-2'), 5.12 (m, 1H, H-4'), 3.95 (bs, 2H, H-5', H-5'), 2.54 (m, 1H, H-3'), 2.31 (m, 1H, H-3'), 1.96 (d, *J* = 1.0 Hz, 3H, T CH₃); ¹³C NMR (75 MHz, CDCl₃) δ 165.4, 163.1 (C-4), 150.7 (C-2), 143.2, 135.9, 133.9 (C-3), 130.1, 129.2, 129.0, 128.8, 124.9 (Ph), 119.7 (CF₃), 111.9 (C-5), 92.4 (C-1'), 67.9 (C-2'), 62.4 (C-5'), 47.0 (C-4'), 28.3 (C-3'), 12.7 (T CH₃); [M+Na]⁺ calcd for C₂₅H₂₂F₃N₃O₆Na 540.1353; Found 540.1362.

Acetyl 2-*O*-benzoyl-3,4-dideoxy-4-(thymid-1-yl)- α -D-threo-pentopyranoside (26c**).** To a solution of **25** (150 mg, 0.43 mmol) in pyridine (5 mL) at 0 °C, acetic anhydride (0.082 mL, 0.87 mmol) was added dropwise. The reaction mixture was slowly warmed to room temperature and left stirring for 2 h. After removal of all the volatiles under reduced pressure, the remaining residue was taken up with DCM (10 mL), and 10 mL of saturated aq. NaHCO₃ was added. The aqueous layer was then extracted with DCM (2 × 10 mL). The combined organic layers were washed with saturated aq. NaHCO₃ (5 mL) and brine (5 mL), dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by silica gel column chromatography (2:1 hexane/EtOAc) to give **26c** (142.8 mg, 85%) as a colorless oil. ¹H NMR (600 MHz, CDCl₃) δ 9.26 (s, 1H, NH), 8.17–8.01 (m, 2H, Ph), 7.63–7.54 (m, 1H, Ph), 7.53–7.40 (m, 2H, Ph), 7.08 (d, *J* = 1.2 Hz, 1H, H-6), 6.10 (d, *J* = 1.9 Hz, 1H, H-1'), 5.28 (td, *J* = 3.4, 1.9 Hz, 1H, H-2'), 4.93 (tt, *J* = 10.4, 4.3 Hz, 1H, H-4'), 4.02 (appt, *J* = 10.8 Hz, 1H, H-5'), 3.92 (ddd, *J* = 11.0, 4.7, 1.8 Hz, 1H, H-5''), 2.61 (ddd, *J* = 13.6, 12.3, 3.3 Hz, 1H, H-3'), 2.27 (dt, *J* = 13.6, 3.7 Hz, 1H, H-3''), 2.21 (s, 3H, CH₃CO), 1.95 (d, *J* = 1.1 Hz, 3H, T CH₃); ¹³C NMR (151 MHz, CDCl₃) δ 168.8 (CH₃CO), 165.4 (PhCO), 163.7 (C-4), 150.9 (C-2), 136.9 (C-6), 133.8, 130.0, 129.2, 128.7 (Ph), 111.6 (C-5), 89.6 (C-1'), 68.3 (C-2'), 62.5 (C-5'), 48.7 (C-4'), 28.5 (C-3'), 21.1 (CH₃CO), 12.7 (T CH₃); HRMS (ESI-TOF) *m/z*: [M+Na]⁺ calcd for C₁₉H₂₀N₂O₇Na 411.1163; Found 411.1157.

Diisopropylphosphonomethyl 2-*O*-benzoyl-3,4-dideoxy-4-(thymid-1-yl)- α -D-threo-pentopyranoside (27**).** To a stirred solution of **26c** (120 mg, 0.31 mmol) and diisopropylphosphonomethanol (121 mg, 0.62 mmol) in dry DCM (3 mL) was added 4 Å molecular sieves, and the mixture was stirred for 30 min at room temperature. The reaction mixture was cooled to 0 °C and TMSOTf (0.073 mL, 0.40 mmol) was added. The

mixture was warmed to room temperature and stirred for 10 h. The reaction was quenched by addition of saturated aq. NaHCO₃ at 0 °C. It was then diluted with DCM (10 mL) and saturated aq. NaHCO₃ (10 mL). The aqueous layer was extracted with DCM (2 × 10 mL). The combined organic layers were washed with saturated aq. NaHCO₃ and brine, dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by silica gel column chromatography to afford **27** as a mixture of α and β isomers, along with residual diisopropylphosphonomethanol. This material was used in the next step without further purification. HRMS (ESI-TOF) m/z: [M+Na]⁺ calcd for C₂₄H₃₃N₂O₉P₁Na 547.1816; Found 547.1801.

Diisopropylphosphonomethyl 3,4-dideoxy-4-(thymid-1-yl)- α/β -D-threo-pentopyranoside (28). A solution of crude **27** in 7 N NH₃ in MeOH (5 mL) was stirred at room temperature for 2 h. The mixture was concentrated under reduced pressure and the resultant residue was purified by silica gel column chromatography (20:1 CH₂Cl₂/MeOH) to give α/β -**28** as a white foam.

Data for α -**28**: 25 mg, 20% over two steps; ¹H NMR (500 MHz, CDCl₃) δ 9.69 (s, 1H, NH), 7.24 (s, 1H, H-6), 5.01 (tt, J = 10.3, 4.0 Hz, 1H, H-4'), 4.84–4.73 (m, 2H, [CH(CH₃)₂]), 4.68 (d, J = 2.6 Hz, 1H, H-1'), 4.05–3.93 (m, 2H, PCH₂ and H-2'), 3.86 (td, J = 10.8, 4.2 Hz, 1H, H-5'), 3.80–3.70 (m, 2H, H-5'' and PCH₂), 2.27 (ddd, J = 12.8, 3.7, 1.2 Hz, 1H, H-3'), 2.05–1.97 (dt, J = 13.2, 4.3 Hz, 1H, H-3''), 1.91 (d, J = 1.1 Hz, 3H, T CH₃), 1.41–1.29 (m, 12H, [CH(CH₃)₂]); ¹³C NMR (126 MHz, CDCl₃) δ 163.9 (C-4), 151.3 (C-2), 137.0 (C-6), 111.2 (C-5), 100.4 (d, ³J_{P,C} = 11.1 Hz, C-1'), 71.6 [CH(CH₃)₂], 66.5 (C-2'), 61.9 (C-5'), 61.8 (d, ¹J_{P,C} = 171.8 Hz, PCH₂), 47.0 (C-4'), 31.0 (C-3'), 24.2 [CH(CH₃)₂], 12.6 (T CH₃); ³¹P NMR (121 MHz, CDCl₃) δ 20.0; HRMS (ESI-TOF) m/z: [M+H]⁺ calcd for C₁₇H₃₀N₂O₈P₁ 421.1734; Found 421.1735.

Data for β -**28**: 13 mg, 10% over two steps; ¹H NMR (500 MHz, CDCl₃) δ 9.24 (s, 1H, NH), 7.64 (d, J = 1.2 Hz, 1H, H-6), 4.94 (d, J = 2.6 Hz, 1H, H-1'), 4.79 (tdd, J = 12.4, 6.2, 1.3 Hz, 2H, [CH(CH₃)₂]), 4.62 (tt, J = 4.4, 3.7 Hz, 1H, H-4'), 4.17 (dd, J = 13.2, 3.9 Hz, 1H, H-5'), 4.03 (dd, J = 13.8, 9.4 Hz, 1H, PCH₂), 3.90–3.67 (m, 3H, H-5'', PCH₂ and H-2'), 2.19–2.16 (dt, J = 13.8, 4.9 Hz, 1H, H-3'), 2.19–2.06 (ddd, J = 13.8, 9.8, 3.9 Hz, 1H, H-3''), 1.93 (d, J = 1.1 Hz, 3H, T CH₃), 1.34 (m, 12H, [CH(CH₃)₂]); ¹³C NMR (126 MHz, CDCl₃) δ 163.9 (C-4), 151.0 (C-2), 138.2 (C-6), 110.8 (C-5), 99.8 (d, ³J_{P,C} = 9.9 Hz, C-1'), 71.8, 71.7 (d, ²J_{P,C} = 6.6 Hz, [CH(CH₃)₂]), 64.3 (C-2'), 62.6 (d, ¹J_{P,C} = 171.0 Hz, PCH₂), 61.2 (C-5'), 51.2 (C-4'), 30.9 (C-3'), 24.2 [CH(CH₃)₂], 12.8 (T CH₃); ³¹P NMR (121 MHz, CDCl₃) δ 20.0.

Phosphonomethyl 3,4-dideoxy-4-(thymid-1-yl)- α -D-threo-pentopyranoside (2a). To a solution of α -**28** (20 mg, 0.048 mmol) and 2,6-lutidine (0.04 mL, 0.38 mmol) in dry CH₃CN (2 mL) was added bromotrimethylsilane (0.05 mL, 0.38 mmol) at 0 °C. The reaction mixture was stirred at room temperature overnight and quenched with 1.0 M aq. TEAB solution (1 mL). After removal of all the volatiles under reduced pressure, the remaining residue was partitioned between water and EtOAc/ether (1:1) and the water layer was lyophilized. The crude residue was first purified by silica gel column chromatography (10:1:0 to 10:5:1, CH₂Cl₂/MeOH/1 M aq. TEAB) and then by preparative reverse phase HPLC with a gradient of CH₃CN in 0.05 M aq. TEAB ranging from 2 to 30% to give **2a** (**2a**·0.67 Et₃N salt, 8.3 mg, 40%) as a white foam. $\epsilon_{267} = 9.7 \times 10^3$ (M⁻¹ cm⁻¹) in water; ¹H NMR (600 MHz, D₂O) δ 7.66 (s, 1H, H-6), 4.79 (tt, J = 10.8, 4.8 Hz, 1H, H-4'), 4.68 (d, J = 2.2 Hz, 1H, H-1'), 4.04 (td, J = 3.0, 3.6 Hz, 1H, H-2'), 3.91 (appt t, J = 10.7 Hz, 1H, H-5'), 3.87 (dd, J = 13.1, 9.3 Hz, 1H, PCH₂), 3.73 (ddd, J = 11.2, 4.7, 1.4 Hz, 1H, H-5''), 3.63 (dd, J = 13.2, 9.3 Hz, 1H, PCH₂), 3.18 (q, J = 7.4 Hz, 4H, CH₂CH₃), 2.38 (ddd, J = 13.8, 12.1, 3.0 Hz, 1H, H-

3'), 1.93 (dtd, $J = 13.1, 3.9, 1.2$ Hz, 1H, H-3'), 1.88 (s, 3H, T CH₃), 1.26 (t, $J = 7.3$ Hz, 6H, CH₂CH₃); ¹³C NMR (151 MHz, D₂O) δ 166.1 (C-4), 151.9 (C-2), 139.1 (C-6), 110.7 (C-5), 99.3 (d, $^3J_{\text{PC}} = 12.0$ Hz, C-1'), 99.3 (C-1'), 65.7 (C-2'), 62.9 (d, $^1J_{\text{PC}} = 156.6$ Hz, PCH₂), 60.4 (C-5'), 46.7 (C-4'), 29.2 (C-3'), 11.0 (T CH₃); ³¹P NMR (121 MHz, D₂O) δ 15.7; HRMS (ESI-): [M-H]⁻ calcd for C₁₁H₁₆N₂O₈P₁ 335.0650; Found 335.0654.

Benzyl 2-*O*-acetyl-3-deoxy- β -L-erythro-pentopyranoside (12). A solution of **17** (2.00 g, 6.20 mmol) and AIBN (306 mg, 1.86 mmol) in toluene (310 mL) was heated at 100 °C until complete dissolution of the reagents. To this solution, Bu₃SnH (3.34 mL, 12.4 mmol) was added and the mixture was stirred for 30 min at 100 °C under an argon atmosphere. After removal of all the volatiles under reduced pressure, the remaining residue was directly subjected to silica gel column chromatography to afford a mixture of 3'-deoxy (60%) and 4'-deoxy (10%) products, which were separated after the next mesylation step.

Benzyl 2-*O*-acetyl-4-azido-3,4-dideoxy- α -D-threo-pentopyranoside (29). A suspension of **14a** (5.60 g, 16.3 mmol) and sodium azide (2.11 g, 32.52 mmol) in DMF (150 mL) was stirred at 90 °C for 8 h. After removal of all the volatiles under reduced pressure, the resulting residue was taken up with DCM (150 mL), and 100 mL of saturated aq. NaHCO₃ was added. The aqueous layer was extracted with DCM (2 $\times$ 75 mL). The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The crude residue was purified by silica gel column chromatography (15:1 hexane/EtOAc) to give **29** (4.30 g, 91%) as a colorless oil. ¹H NMR (500 MHz, CDCl₃) δ 7.44–7.21 (m, 5H, Ph), 4.94 (dt, $J = 3.3, 2.9$ Hz, 1H, H-2), 4.72 (d, $J = 11.9$ Hz, 1H, CH₂Ph), 4.69 (d, $J = 1.6$ Hz, 1H, H-1), 4.49 (d, $J = 11.9$ Hz, 1H, CH₂Ph), 3.77 (tt, $J = 11.1, 5.5$ Hz, 1H, H-4), 3.70 (ddd, $J = 10.7, 4.9, 1.9$ Hz, 1H, H-5), 3.60 (appt, $J = 10.8$ Hz, 1H, H-5'), 2.21–1.90 (m, 5H, H-3, H-3', CH₃CO); ¹³C NMR (126 MHz, CDCl₃) δ 169.7 (CH₃CO), 136.9, 128.4, 127.8, 127.7 (Ph), 94.8 (C-1), 69.0 (CH₂Ph), 68.6 (C-2), 61.4 (C-5), 52.1 (C-4), 29.1 (C-3), 20.9 (CH₃CO); HRMS (ESI-TOF) m/z : [M+Na]⁺ calcd for C₁₄H₁₇N₃O₄Na 314.1111; Found 314.1113.

Note: Concentrating azide-containing reaction mixtures and products through rotary evaporation have caused documented explosions. NaN₃ is known to decompose violently causing exploding at 275 °C

Benzyl 4-azido-3,4-dideoxy- α -D-threo-pentopyranoside (30). A methanolic solution of NaOMe (5.4 M, 5.09 mL, 27.5 mmol) was added dropwise to a solution of **29** (4.00 g, 13.73 mmol) in methanol (150 mL). The reaction was stirred at room temperature for 3 h. After removal of all the volatiles under reduced pressure, the resulting residue was taken up with DCM (100 mL), and 50 mL of saturated aq. NaHCO₃ was added. The aqueous layer was extracted again with DCM (50 mL). The combined organic layers were washed with saturated aq. NaHCO₃ and brine, dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by silica gel column chromatography (10:1 hexane/EtOAc) give **30** (3.25 g, 95%) as a colorless oil. ¹H NMR (500 MHz, CDCl₃) δ 7.41–7.25 (m, 5H, Ph), 4.79 (d, $J = 11.8$ Hz, 1H, CH₂Ph), 4.50 (d, $J = 11.9$ Hz, 1H, CH₂Ph), 4.47 (d, $J = 4.2$ Hz, 1H, H-1), 3.80 (td, $J = 5.3, 2.0$ Hz, 1H, H-2), 3.76 (m, 1H, H-4), 3.74–3.69 (dd, $J = 7.3, 11.2$ Hz, 1H, H-5), 3.63 (ddd, $J = 11.0, 3.8, 1.2$ Hz, 1H, H-5'), 2.68 (d, $J = 3.0$ Hz, 1H, OH), 2.06–1.97 (dddd, $J = 13.5, 8.9, 3.7, 1.0$ Hz, 1H, H-3), 1.90 (dddd, $J = 13.5, 6.3, 4.2, 1.1$ Hz, 1H, H-3'); ¹³C NMR (126 MHz, CDCl₃) δ 137.0, 128.4, 128.3, 127.9 (Ph), 99.6 (C-1), 69.5 (CH₂Ph), 66.5 (C-2), 63.4 (C-5), 53.4 (C-4), 32.1 (C-3); HRMS (ESI-TOF) m/z : [M+Na]⁺ calcd for C₁₂H₁₅N₃O₃Na 272.1006; Found 272.1003.

Benzyl 4-amino-3,4-dideoxy- α -D-*threo*-pentopyranoside (31). To a solution of **30** (2.00 g, 8.02 mmol) in dry THF (80 mL) was added Ph_3P (0.46 g, 1.75 mmol). The mixture was stirred at room temperature for 1 h. Water (2 mL) was added and the mixture was stirred at 50 °C for 6 h. It was then evaporated, and the resulting residue was purified by silica gel column chromatography (100/15 DCM/MeOH) to give **31** (1.65 g, 92%) as a colorless oil. ^1H NMR (500 MHz, CDCl_3) δ 7.46–7.28 (m, 5H, Ph), 4.80 (d, J = 11.8 Hz, 1H, CH_2Ph), 4.52 (d, J = 11.8 Hz, 1H, CH_2Ph), 4.50 (d, J = 3.3 Hz, 1H, H-1), 3.80 (dt, J = 5.9, 3.5 Hz, 1H, H-2), 3.60 (ddd, J = 11.0, 4.1, 1.3 Hz, 1H, H-5), 3.50 (dd, J = 11.0, 8.0 Hz, 1H, H-5), 3.15 (tt, J = 12.7, 4.2 Hz, 1H, H-4), 2.05–1.53 (m, 5H, H-3, H-3, OH, NH_2); ^{13}C NMR (126 MHz, CDCl_3) δ 137.3, 128.4, 127.9, 127.8 (Ph), 99.7 (C-1), 69.4 (CH_2Ph), 67.6 (C-5), 66.9 (C-2), 43.6 (C-4), 36.4 (C-3); HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{12}\text{H}_{18}\text{N}_1\text{O}_3$ 224.1281; Found 224.1283.

Benzyl 4-(6-chloropurine-9-yl)-3,4-dideoxy- α -D-*threo*-pentopyranoside (32). 4,6-Dichloro-5-formamidopyrimidine (860 mg, 4.48 mmol) and DIPEA (1.17 mL, 6.72 mmol) were added to a solution of pentopyranoside **31** (500 mg, 2.24 mmol) in *n*-butanol (25 mL). The reaction mixture was stirred for 2 h at 100 °C, the temperature was then raised to 140 °C and maintained at this temperature for 10 h. After removal of all the volatiles under reduced pressure, the remaining residue was purified by silica gel column chromatography (gradient $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 100:0, v/v; 99:1, v/v; 20:1, v/v) to give **32** (605 mg, 75%) as a colorless oil. ^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ 8.25 (s, 1H, H-8), 8.16 (s, 1H, H-2), 7.45–7.25 (m, 5H, Ph), 5.34 (d, J = 3.9 Hz, 1H, OH), 4.92 (tt, J = 11.1, 4.2 Hz, 1H, H-4'), 4.76 (d, J = 12.2 Hz, 1H, CH_2Ph), 4.66 (d, J = 1.8 Hz, 1H, H-1'), 4.54 (d, J = 12.2 Hz, 1H, CH_2Ph), 4.10 (appt, J = 10.6 Hz, 1H, H-5'), 3.82 (dt, J = 6.0, 3.3 Hz, 1H, H-2'), 3.76 (ddd, J = 10.6, 4.4, 1.6 Hz, 1H, H-5'), 2.71 (td, J = 12.6, 3.0 Hz, 1H, H-3'), 1.99 (dt, J = 12.8, 3.8 Hz, 1H, H-3'); ^{13}C NMR (150 MHz, $\text{DMSO}-d_6$) δ 156.2 (C-6), 152.4 (C-2), 149.6 (C-4), 139.6 (C-8), 137.9, 128.4, 127.9, 127.7 (Ph), 119.1 (C-5), 98.1 (C-1'), 68.2 (CH_2Ph), 65.7 (C-2'), 61.7 (C-5'), 46.7 (C-4'), 31.6 (C-3'); HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{18}\text{Cl}_1\text{N}_4\text{O}_3$ 361.1061; Found 361.1056.

Benzyl 4-(adenine-9-yl)-3,4-dideoxy- α -D-*threo*-pentopyranoside (33). A solution of **32** (300 mg, 0.83 mmol) in saturated ammonia in ethanol (10 mL) was stirred in a sealed flask at 60 °C for 8 h. After removal of all the volatiles under reduced pressure, the resulting residue was purified by silica gel column chromatography ($\text{CH}_2\text{Cl}_2/\text{MeOH}$, 15:1) to give **33** (269 mg, 95%) as a colorless oil. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.82 (s, 1H, H-8), 8.79 (s, 1H, H-2), 7.46–7.29 (m, 5H, Ph), 5.38 (d, J = 4.0 Hz, 1H, OH), 5.07 (tt, J = 11.5, 4.3 Hz, 1H, H-4'), 4.76 (d, J = 12.2 Hz, 1H, CH_2Ph), 4.67 (d, J = 1.8 Hz, 1H, H-1'), 4.55 (d, J = 12.2 Hz, 1H, CH_2Ph), 4.16 (appt, J = 10.6 Hz, 1H, H-5'), 3.99–3.74 (m, 2H, H-5' and H-2'), 2.76 (td, J = 12.5, 3.0 Hz, 1H, H-3'), 2.06 (ddd, J = 12.7, 5.2, 2.8 Hz, 1H, H-3''); ^{13}C NMR (126 MHz, $\text{DMSO}-d_6$) δ 152.0 (C-4), 151.4 (C-2), 149.2 (C-6), 146.4 (C-8), 137.8, 131.2 (C-5), 128.4, 127.8, 127.6 (Ph), 98.0 (C-1'), 68.3 (CH_2Ph), 65.6 (C-2'), 61.2 (C-5'), 47.9 (C-4'), 31.4 (C-3'); HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{20}\text{N}_5\text{O}_3$ 342.1561; Found 342.1566.

Benzyl 2-*O*-benzoyl-4-(N^6,N^6 -dibenzoyladenine-9-yl)-3,4-dideoxy- α -D-*threo*-pentopyranoside (34). To a solution of **33** (200 mg, 0.59 mmol) in dry pyridine (6 mL) at 0 °C was added benzoyl chloride (0.46 mL, 3.52 mmol), and the reaction mixture was warmed to room temperature and stirred for 3 h. After removal of all the volatiles under reduced pressure, the resulting residue was partitioned between saturated aq.

NaHCO₃ (10 mL) and DCM (20 mL). The water layer was further extracted with DCM (1 × 10 mL). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, and concentrated in vacuo. The crude residue was purified by silica gel column chromatography (gradient hexane/EtOAc, 10:1, v/v; 3:1) to give **34** (282 mg, 88%) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ 8.67 (s, 1H, H-2), 8.21–8.04 (m, 3H, H-8, Ph), 8.00–7.74 (m, 4H, Ph), 7.65–7.53 (m, 1H, Ph), 7.51–7.27 (m, 14H, Ph), 5.36 (td, *J* = 4.2, 1.5 Hz, 1H, H-2'), 5.15 (tt, *J* = 11.7, 4.3 Hz, 1H, H-4'), 5.01 (d, *J* = 1.3 Hz, 1H, H-1'), 4.86 (d, *J* = 11.9 Hz, 1H, CH₂Ph), 4.64 (d, *J* = 11.9 Hz, 1H, CH₂Ph), 4.24 (appt, *J* = 10.9 Hz, 1H, H-5'), 4.03 (ddd, *J* = 10.4, 5.1, 1.8 Hz 1H, H-5'), 3.01 (td, *J* = 13.3, 3.1 Hz, 1H, H-3'), 2.44 (dt, *J* = 13.2, 5.2 Hz, 1H, H-3'); ¹³C NMR (126 MHz, CDCl₃) δ 172.2 (PhCO), 165.4 (PhCO), 153.1 (C-4), 152.0 (C-2), 152.0 (C-6), 142.8 (C-8), 136.8, 134.1, 133.5, 132.9, 129.8, 129.4, 128.7, 128.6, 128.5, 128.1, 127.9(Ph), 127.4 (C-5), 95.1 (C-1'), 69.5 (CH₂Ph), 68.9 (C-2'), 61.5 (C-5'), 47.8 (C-4'), 29.0 (C-3'); HRMS (ESI-TOF) *m/z*: [M+H]⁺ calcd for C₃₈H₃₂N₅O₆ 654.2346; Found 654.2358.

2-*O*-Benzoyl-4-(*N*⁶-benzoyladenine-9-yl)-3,4-dideoxy- α -D-*threo*-pentopyranose (35). To a stirred solution of **34** (140 mg, 0.25 mmol) in anhydrous CH₂Cl₂ (4 mL) was added 1 M BCl₃ (0.76 mL, 0.76 mmol) at -78 °C. Then, the reaction mixture was slowly warmed to 0 °C over 2.5 h and stirred at this temperature for 30 min. After completion of the reaction, the mixture was cooled to -20 °C and quenched by dropwise addition of saturated aq. NaHCO₃ (10 mL). The aqueous layer was extracted with CH₂Cl₂ (3 × 5 mL). The combined organic layers were washed with brine (10 mL) and dried over Na₂SO₄, filtered, and concentrated in vacuo. The resulting residue was filtered through a pad of silica gel to give crude **35** as a colorless oil, which was used as such in the following step. HRMS (ESI-TOF) *m/z*: [M+H]⁺ calcd for C₂₄H₂₂N₅O₅ 460.1615; Found 460.1617.

Acetyl 2-*O*-benzoyl-4-(*N*⁶-benzoyladenine-9-yl)-3,4-dideoxy- α -D-*threo*-pentopyranoside (36). Compound **36** was prepared using a similar procedure as that described for **26c** starting from **35** (117 mg, 0.25 mmol), pyridine (5 mL), and acetic anhydride (0.048 mL, 0.5 mmol), and obtained after silica gel column chromatography (gradient hexane/EtOAc, 8:1, v/v; 3:1, v/v;) as a colorless sticky oil (99 mg, 78%). ¹H NMR (500 MHz, CDCl₃) δ 8.76 (s, 1H, H-2), 8.19–8.05 (m, 3H, H-8, Ph), 8.02 (d, *J* = 7.4 Hz, 2H, Ph), 7.63–7.53 (m, 2H, Ph), 7.49 (td, *J* = 7.9, 2.1 Hz, 4H, Ph), 6.19 (d, *J* = 1.8 Hz, 1H, H-1'), 5.33 (td, *J* = 5.3, 1.8 Hz, 1H, H-2'), 5.14 (tt, *J* = 11.0, 4.3 Hz, 1H, H-4'), 4.34 (appt, *J* = 10.8 Hz, 1H, H-5'), 4.12 (ddd, *J* = 11.1, 4.7, 1.8 Hz, 1H, H-5'), 3.04 (ddd, *J* = 14.3, 12.1, 3.2 Hz, 1H, H-3'), 2.51 (dt, *J* = 14.2, 4.7 Hz, 1H, H-3'), 2.24 (s, 3H, CH₃CO); ¹³C NMR (126 MHz, CDCl₃) δ 168.8 (CH₃CO), 165.3 (PhCO), 152.6 (C-2), 152.2 (C-4), 150.0 (C-6), 141.3(C-8), 133.8, 133.6, 132.9, 123.0, 129.2, 128.9, 128.7, 128.1 (Ph), 123.5 (C-5), 89.6 (C-1'), 68.0 (C-2'), 63.3 (C-5'), 47.9 (C-4'), 29.3 (C-3'), 21.1 (CH₃CO); HRMS (ESI-TOF) *m/z*: [M+H]⁺ calcd for C₂₆H₂₄N₅O₆ 502.1721; Found 502.1723.

Benzyl 3,4-dideoxy-4-trifluoroacetyl-amino- α -D-*threo*-pentopyranoside (38). To a mixture of **31** (1 g, 4.48 mmol) and triethylamine (1.87 mL, 13.44mmol) in methanol (50 mL), ethyl trifluoroacetate (1.07 mL, 8.96 mmol) was added dropwise. The mixture was stirred at room temperature for 2 h. After removal of all the volatiles under reduced pressure, the resulting residue was purified by silica gel column chromatography (hexane/EtOAc, 3:1) to give **38** (1.3g, 91%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.55–7.16 (m, 5H, Ph), 6.91 (d, *J* = 7.9 Hz, 1H, NH), 4.80 (d, *J* = 11.6 Hz, 1H, CH₂Ph), 4.53 (d, *J* = 11.6 Hz, 1H, CH₂Ph),

4.45 (d, $J = 4.8$ Hz, 1H, H-1), 4.27 (m, 1H, H-2), 3.80–3.61 (m, 3H, H-5, H-5' and OH), 3.01 (d, $J = 3.8$ Hz, 1H, H-4), 2.17–1.98 (ddd, $J = 13.5, 6.7, 4.0$ Hz, 1H, H-3), 1.88–1.69 (ddd, $J = 13.5, 7.4, 4.1$ Hz, 1H, H-3'); ^{13}C NMR (75 MHz, CDCl_3) δ 157.0 (q, $^2J_{\text{FC}} = 37.8$ Hz, COCF_3), 137.0, 128.64, 128.20 (Ph), 115.8 (q, $^1J_{\text{FC}} = 288.9$ Hz, CF_3), 101.3 (C-1), 70.3 (C-2), 66.6 (CH_2Ph), 64.6 (C-5), 44.4 (C-4), 32.8 (C-3); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{14}\text{H}_{16}\text{F}_3\text{N}_1\text{O}_4\text{Na}$ 342.0924; Found 342.0916.

3,4-dideoxy-4-trifluoroacetyl-amino- α/β -D-threo-pentopyranose (39). To a solution of **38** (1.00 g, 3.13 mmol) in methanol (50 mL) was added 10% Pd/C (1 g, 0.94 mmol), and evacuation-replacement cycles were carried out using a hydrogen balloon (3 $\times$). The reaction mixture was stirred at room temperature for 20 h under an atmospheric pressure of hydrogen. After completion of the reaction, the catalyst was removed by filtration through a pad of Celite and washed with methanol. The filtrate was concentrated under reduced pressure to afford **39** (590 mg, 82%) as a colorless foam without further purification. ^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ 9.32 (d, $J = 6.7$ Hz, 0.16 H, $\text{NH}\beta$), 9.26 (d, $J = 7.9$ Hz, 1H, $\text{NH}\alpha$), 6.34 (d, $J = 4.0$ Hz, 1H, $\text{OH}-1\alpha$), 6.29 (d, $J = 4.3$ Hz, 0.16 H, $\text{OH}-1\beta$), 4.94 (bs, 1H, $\text{OH}-2\alpha$), 4.85–4.77 (m, 0.22H, $\text{OH}-2\beta$), 4.69 (dd, $J = 4.8, 2.0$ Hz, 0.18 H, H-1 β), 4.68 (dd, $J = 3.3, 2.2$ Hz, 1H, H-1 α), 4.17–4.08 (m, 1H, H-4 α), 4.07–4.00 (m, 0.2 H, H-4 β), 3.80 (ddd, $J = 11.5, 3.6, 0.9$ Hz, 0.18 H, H-5 β), 3.66 (t, $J = 10.2$ Hz, 1H, H-5 α), 3.56 (bs, 1.14 H, H-2 α), 3.43 (ddd, $J = 10.6, 4.6, 1.6$ Hz, 1 H, H-5' α and H-2 β), 3.30 (dd, $J = 11.1, 6.6$ Hz, 0.3 H, H-5' β), 1.96 (ddd, $J = 13.1, 10.9, 3.2$ Hz, 1H, H-3 α), 1.88 (dddd, $J = 13.3, 7.3, 4.3, 0.9$ Hz, 0.2 H, H-3 β), 1.65 (dtd, $J = 13.1, 4.8, 1.4$ Hz, 1H, H-3' α), 1.42 (ddd, $J = 13.3, 7.8, 3.7$ Hz, 0.14 H, H-3' β); ^{13}C NMR (151 MHz, $\text{DMSO}-d_6$) δ 156.2 (q, $^2J_{\text{FC}} = 38$ Hz, COCF_3), 115.9 (q, $^1J_{\text{FC}} = 289$ Hz, CF_3), 93.7 (C-1 β), 93.4 (C-1 α), 68.1 (C-2 β), 66.4 (C-2 α), 62.8 (C-5 β), 60.8 (C-5 α), 44.3 (C-4 β), 42.6 (C-4 α), 32.2 (C-3 β), 31.3 (C-3 α); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_7\text{H}_{10}\text{F}_3\text{N}_1\text{O}_4\text{Na}$ 252.0454; Found 252.0448.

1,2-Di-O-acetyl-3,4-dideoxy-4-trifluoroacetyl-amino- α -D-threo-pentopyranoside (40). 1,2-O-Diacetylated compound **40** was prepared using a similar procedure as that described for **26c** starting from **39** (500 mg, 2.18 mmol), Ac_2O (0.62 mL, 6.55 mmol), and pyridine (25 mL), and obtained after silica gel column chromatography (gradient hexane/EtOAc, 6:1, v/v; 2:1, v/v) as a colorless sticky oil (550 mg, 80%), ^1H NMR (600 MHz, CDCl_3) δ 6.78 (d, $J = 7.9$ Hz, 1H, NH), 5.88 (d, $J = 2.8$ Hz, 1H, H-1), 4.93 (td, $J = 6.1, 3.8$ Hz, 1H, H-2), 4.37 (dtt, $J = 17.6, 7.9, 4.6$ Hz, 1H, H-4), 3.88 (ddd, $J = 11.4, 4.2, 1.5$ Hz, 1H, H-5), 3.66 (dd, $J = 11.0, 9.7$ Hz, 1H, H-5'), 2.17–2.09 (m, 8H, CH_3CO , CH_3CO , H-3, H-3); ^{13}C NMR (151 MHz, CDCl_3) δ 170.0 (CH_3CO), 168.9 (CH_3CO), 157.0 (q, $^2J_{\text{FC}} = 38$ Hz, COCF_3), 115.5 (q, $^1J_{\text{FC}} = 289$ Hz, CF_3), 89.8 (C-1), 67.1 (C-2), 63.4 (C-5), 42.5 (C-4), 29.5 (C-3), 21.0 (CH_3CO), 20.9 (CH_3CO); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{11}\text{H}_{14}\text{F}_3\text{N}_1\text{O}_6\text{Na}$ 336.0666; Found 336.0660.

Diisopropylphosphonomethyl 2-O-acetyl-3,4-dideoxy-4-trifluoroacetyl-amino- α -D-threo-pentopyranoside (41). Phosphonomethylated compound **41** was prepared using a similar procedure as that described for **27** starting from **40** (300 mg, 0.96 mmol), TMSOTf (0.35 mL, 1.92 mmol), diisopropylphosphonomethanol (564 mg, 2.87 mmol) in dry DCM (15 mL). Purification by silica gel column chromatography (gradient $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 99:1, v/v; 49:1, v/v; 24:1, v/v) afforded **41** as a colorless sticky oil, which was still contaminated with diisopropylphosphonomethanol. ^1H NMR (600 MHz, CDCl_3) δ 7.45 (d, $J = 8.3$ Hz, 1H, NH), 4.91 (td, $J = 4.7, 2.2$ Hz, 1H, H-2), 4.83–4.69 (m, 2H, $[\text{CH}(\text{CH}_3)_2]$), 4.69 (d, $J = 1.8$ Hz, 1H, H-1), 4.45–4.34 (m, 1H, H-4), 3.92 (dd, $J = 13.2, 10.0$ Hz, 1H, PCH_2), 3.79–3.72 (ddd,

$J = 10.9, 5.1, 1.7$ Hz, 1H, H-5), 3.72–3.66 (m, 1H, PCH_2), 3.64 (appt, $J = 10.7$ Hz, 1H, H-5'), 2.12 (s, 3H, CH_3CO), 2.11–2.08 (m, 1H, H-3), 2.07–1.99 (dt, $J = 13.7, 4.2$ Hz, 1H, H-3'), 1.38–1.30 (m, 12H, $[CH(CH_3)_2]$); ^{13}C NMR (151 MHz, $CDCl_3$) δ 170.2 (CH_3CO), 157.1 (q, $^2J_{FC} = 37.0$ Hz, $COCF_3$), 115.8 (q, $^1J_{FC} = 288.9$ Hz, CF_3), 96.4 (d, $^3J_{PC} = 11.1$ Hz, C-1), 71.8, 71.7 (d, $^2J_{PC} = 6.2$ Hz, $[CH(CH_3)_2]$), 68.0 (C-2), 60.9 (d, $^1J_{PC} = 171.7$ Hz, PCH_2), 61.4 (C-5), 41.8 (C-4), 29.0 (C-3), 24.0 $[CH(CH_3)_2]$; ^{31}P NMR (121 MHz, $CDCl_3$) δ 19.0; HRMS (ESI-TOF) m/z : $[M+H]^+$ calcd for $C_{16}H_{28}F_3N_1O_8P_1$ 450.1499; Found 450.1497.

Diisopropylphosphonomethyl 4-amino-3,4-dideoxy- α -D-threo-pentopyranoside (42). Compound **42** was prepared using a similar procedure as that described for **28** starting from crude **41** and 7 M NH_3 in MeOH (15 mL), and obtained after silica gel column chromatography (gradient $CH_2Cl_2/MeOH$, 50:1, v/v; 25:1, v/v; 6:1, v/v) as a colorless sticky oil (121 mg, 40% over two steps), 1H NMR (600 MHz, $CDCl_3$) δ 4.81–4.67 (m, 2H, $[CH(CH_3)_2]$), 4.54 (d, $J = 3.0$ Hz, 1H, H-1), 3.99 (dd, $J = 13.6, 9.1$ Hz, 1H, PCH_2), 3.91 (m, 1H, H-2), 3.76–3.68 (m, 2H, H-5 and PCH_2), 3.65 (appt, $J = 9.5$ Hz, 1H, H-5'), 3.49–3.42 (m, 1H, H-4), 2.06–2.01 (m, 1H, H-3), 1.96 (td, $J = 11.2, 3.2$ Hz, 1H, H-3'), 1.37–1.29 (m, 12H, $[CH(CH_3)_2]$); ^{13}C NMR (151 MHz, $CDCl_3$) δ 101.2 (d, $^3J_{PC} = 10.4$ Hz, C-1'), 71.7, 71.5 (d, $^2J_{PC} = 5.6$ Hz, $[CH(CH_3)_2]$), 65.7 (C-2), 64.7 (C-5), 61.6 (d, $^1J_{PC} = 170.9$ Hz, PCH_2), 43.6 (C-4), 33.6 (C-3), 24.0 $[CH(CH_3)_2]$; ^{31}P NMR (121 MHz, $CDCl_3$) δ 19.3; HRMS (ESI-TOF) m/z : $[M+H]^+$ calcd for $C_{12}H_{27}N_1O_6P_1$ 312.1570; Found 312.1570.

Diisopropylphosphonomethyl 4-(6-chloropurine-9-yl)-3,4-dideoxy- α -D-threo-pentopyranoside (43). Compound **43** was prepared using a similar procedure as that described for **32** starting from **42** (120 mg, 0.39 mmol), 4,6-dichloro-5-formamidopyrimidine (96 mg, 0.5 mmol), and DIPEA (0.2 mL, 1.16 mmol) in *n*-butanol (5 mL), and obtained after silica gel column chromatography (gradient $CH_2Cl_2/MeOH$, 99:1, v/v; 49:1, v/v; 24:1, v/v) as a colorless sticky oil (130 mg, 75%). 1H NMR (600 MHz, $CDCl_3$) δ 8.75 (s, 1H, H-2), 8.37 (s, 1H, H-8), 5.14 (tt, $J = 12.6, 4.1$ Hz, 1H, H-4'), 4.80 (tt, $J = 12.4, 6.2$ Hz, 2H, $[CH(CH_3)_2]$), 4.74 (d, $J = 3.8$ Hz, 1H, H-1'), 4.29 (dd, $J = 11.5, 7.7$ Hz, 1H, H-5'), 4.09 (dd, $J = 13.8, 9.4$ Hz, 1H, PCH_2), 3.98 (dd, $J = 11.7, 3.8$ Hz, 1H, H-5'), 3.98 (td, $J = 7.0, 3.8$ Hz, 1H, H-2'), 3.84 (dd, $J = 13.8, 8.3$ Hz, 1H, PCH_2), 2.71 (ddd, $J = 13.0, 9.1, 3.6$ Hz, 1H, H-3'), 2.20 (dt, $J = 13.2, 4.8$ Hz, 1H, H-3'), 1.40–1.32 (m, 12H, $[CH(CH_3)_2]$); ^{13}C NMR (151 MHz, $CDCl_3$) δ 152.0 (C-2), 151.9 (C-4), 151.4 (C-6), 144.0 (C-8), 131.8 (C-5), 102.2 (d, $^3J_{PC} = 11.1$ Hz, C-1'), 71.9, 71.7 (d, $^2J_{PC} = 6.2$ Hz, $[CH(CH_3)_2]$), 66.0 (C-2'), 63.8 (C-5'), 62.1 (d, $^1J_{PC} = 171.1$ Hz, PCH_2), 48.8 (C-4'), 32.7 (C-3'), 24.2 $[CH(CH_3)_2]$; ^{31}P NMR (121 MHz, $CDCl_3$) δ 19.6; HRMS (ESI-TOF) m/z : $[M+H]^+$ calcd for $C_{17}H_{27}Cl_1N_4O_6P_1$ 449.1351; Found 449.1349.

Diisopropylphosphonomethyl 4-(adenine-9-yl)-3,4-dideoxy- α -D-threo-pentopyranoside (44). Compound **44** was prepared using a similar procedure as that described for **37** starting from **43** (100 mg, 0.22 mmol) in a mixture of aq. NH_3 (25%)/1,4-dioxane = 1:1 (10 mL), and obtained after silica gel column chromatography (gradient $CH_2Cl_2/MeOH$, 50:1, v/v; 25:1, v/v; 15:1, v/v) as a colorless sticky oil (88 mg, 92%). 1H NMR (600 MHz, methanol- d_4) δ 8.23 (s, 1H, H-8), 8.19 (s, 1H, H-2), 5.08–4.97 (tt, $J = 10.7, 3.7$ Hz, 1H, H-4'), 4.82–4.75 (m, 2H, $[CH(CH_3)_2]$), 4.74 (d, $J = 2.3$ Hz, 1H, H-1'), 4.14 (appt, $J = 10.5$ Hz, 1H, H-5'), 4.10–4.05 (dd, $J = 14.0, 9.2$ Hz, 1H, PCH_2), 3.95–3.89 (m, 2H, H-2' and H-5'), 3.88 (dd, $J = 14.0, 9.2$ Hz, 1H, PCH_2), 2.70 (ddd, $J = 14.6, 11.1, 3.0$ Hz, 1H, H-3'), 2.19–2.14 (m, 1H, H-3'), 1.40–1.34 (dd, $J = 6.2, 1.9$ Hz, 12H, $[CH(CH_3)_2]$); ^{13}C NMR (151 MHz, methanol- d_4) δ 157.3 (C-6), 153.7 (C-2), 150.8 (C-4), 140.9 (C-8), 120.1 (C-5), 101.7 (d, $^3J_{PC} = 11.6$ Hz, C-1'), 73.4, 73.3 $[CH(CH_3)_2]$, 67.2 (C-2'), 63.6 (C-

5'), 62.2 (d, $^1J_{\text{PC}} = 170.9$ Hz, PCH_2), 48.5 (C-4'), 32.9 (C-3'), 24.3 [$\text{CH}(\text{CH}_3)_2$]; ^{31}P NMR (121 MHz, methanol- d_4) δ 20.0; HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{29}\text{N}_5\text{O}_6\text{P}_1$ 430.1850; Found 430.1848.

Phosphonomethyl 4-(adenine-9-yl)-3,4-dideoxy- α -D-threo-pentopyranoside (2b). Compound **2b** was prepared using a similar procedure as that described for **2a** starting from **44** (70 mg, 0.16 mmol), 2,6-lutidine (0.15 mL, 1.3 mmol), and TMSBr (0.17 mL, 0.16 mmol) in dry acetonitrile (2 mL) and obtained as a colorless sticky oil (**2b**·1Et₃N salt, 22 mg, 40%). $\epsilon_{259} = 16.9 \times 10^3$ ($\text{M}^{-1}\text{cm}^{-1}$) in water; ^1H NMR (600 MHz, D₂O) δ 8.22 (s, 1H, H-8), 8.05 (s, 1H, H-2), 4.72 (m, 1H, H-4'), 4.71 (d, $J = 3.3$ Hz, 1H, H-1'), 4.09 (dd, $J = 11.4, 8.7$ Hz, 1H, H-5'), 3.97 (td, $J = 4.8, 3.3$ Hz, 1H, H-2'), 3.92 (dd, $J = 11.4, 4.0$ Hz, 1H, H-5'), 3.90 (dd, $J = 13.2, 9.3$ Hz, 1H, PCH_2), 3.64 (dd, $J = 13.0, 9.5$ Hz, 1H, PCH_2), 3.16 (q, $J = 7.4$ Hz, 6H, CH_2CH_3), 2.56 (ddd, $J = 13.4, 10.1, 3.4$ Hz, 1H, H-3'), 2.15 (dt, $J = 13.3, 4.3$ Hz, 1H, H-3''), 1.23 (t, $J = 7.3$ Hz, 9H, CH_2CH_3); ^{13}C NMR (151 MHz, D₂O) δ 156.5(C-6), 153.3(C-2), 149.8 (C-4), 141.7 (C-8), 119.4 (C-5), 102.4 (d, $^3J_{\text{PC}} = 11.8$ Hz, C-1'), 67.1 (C-2'), 65.3 (d, $^1J_{\text{PC}} = 158.5$ Hz, PCH_2), 64.1 (C-5'), 48.7 (C-4'), 32.6 (C-3'); ^{31}P NMR (121 MHz, D₂O) δ 15.2; HRMS (ESI-TOF) m/z : $[\text{M}-\text{H}]^+$ calcd for $\text{C}_{11}\text{H}_{15}\text{N}_5\text{O}_6\text{P}_1$ 344.0765; Found 344.0760.

Diphosphorylphosphonomethyl 4-(adenine-9-yl)-3,4-dideoxy- α -D-threo-pentopyranoside (45).

Adenine α -D-threo-pentopyranoside nucleoside phosphonic acid **2b** (30 mg, 0.087 mmol) was dissolved in anhydrous DMF (1 mL) and carbonyldiimidazole (70.4 mg, 0.434 mmol) was added. The mixture was stirred for 30 min at room temperature, and then a solution of pyrophosphate (333 mg, 0.608 mmol) in dry DMF (2 mL) was added and the stirring was continued overnight. An excess of 25% aq. ammonia (1 mL) was added and the mixture was concentrated in vacuo. The resulting residue was purified by chromatography on a DEAE-cellulose column (gradient: 0.1 to 1 M TEAB (v/v); 1/0, 1/0.2, 1/0.5, 1/1). Further purification was performed by RP-HPLC to afford diphosphate(phosphonomethyl) 3-deoxy-4-deoxy-4-(adenine-9-yl)- α -D-threo-pentopyranoside triethylammonium salt **45** as a white solid (30% yield). ^1H NMR (600 MHz, D₂O): δ 8.38 (s, 1H, H-2), 8.23 (s, 1H, H-8), 4.91 (tt, $J = 9.3, 4.1$ Hz, 1H, H-4'), 4.82 (d, $J = 3.2$ Hz, 1H, H-1'), 4.19 (dd, $J = 12.0, 8.6$ Hz, 1H, H-5'), 4.08–4.03 (m, 2H, PCH_2 and H-2'), 3.99 (dd, $J = 12.0, 4.1$ Hz, 1H, H-5''), 4.19 (dd, $J = 13.2, 9.8$ Hz, 1H, PCH_2), 2.63 (ddd, $J = 13.5, 9.9, 3.1$ Hz, 1H, H-3'), 2.15 (dt, $J = 13.2, 5.1$ Hz, 1H, H-3''); ^{13}C NMR (150 MHz, D₂O): δ 157.0 (C-6), 153.7 (C-2), 150.3 (C-4), 142.1 (C-8), 119.8 (C-5), 102.6 (d, $^3J_{\text{PC}} = 9.8$ Hz, C-1'), 67.0 (C-2'), 65.2 (d, $^1J_{\text{PC}} = 167.1$ Hz, PCH_2), 60.3 (C-5'), 48.7 (C-4'), 32.6 (C-3'); ^{31}P NMR (121 MHz, D₂O): δ 7.6 (d, $J = 23.1$ Hz, $\text{P}\alpha$), -6.4 (d, $J = 18.9$ Hz, $\text{P}\gamma$), -22.7 (t, $J = 21.4$ Hz, $\text{P}\beta$); HRMS (ESI-TOF) m/z : $[\text{M}-\text{H}]^-$ calcd for $\text{C}_{11}\text{H}_{17}\text{N}_5\text{O}_{12}\text{P}_3$, 504.0092; found 504.0096.

ASSOCIATED CONTENT

Supporting Information

NMR spectra, supplementary NMR data, procedures, scheme, and figures.

AUTHOR INFORMATION

Corresponding author

[*piet.herdewijn@kuleuven.be](mailto:piet.herdewijn@kuleuven.be)

ORCID

Piet Herdewijn: 0000-0003-3589-8503

Elisabetta Groaz: 0000-0002-8866-2265

Eveline Lescrinier: 0000-0001-7066-4329

Shrinivas Dumbre: 0000-0002-1237-9586

ACKNOWLEDGEMENTS

X.L. acknowledges the China Scholarship Council (CSC) for funding (No. 201307040022). We express our gratitude to Hoai Nguyen and Luc Baudemprez for excellent technical assistance. We wish to thank Prof. Graciela Andrei for carrying out the antiviral screening, Prof. Jef Rozenski for conducting mass spectrometry measurements, and Dr. Mikhail Abramov for diphosphate synthesis.

REFERENCES

1. Jordheim, L. P.; Durantel, D.; Zoulim, F.; Dumontet, C., Advances in the Development of Nucleoside and Nucleotide Analogues for Cancer and Viral Diseases. *Nat. Rev. Drug Discov.* **2013**, *12* (6), 447-464.
2. Khvorova, A.; Watts, J. K., The Chemical Evolution of Oligonucleotide Therapies of Clinical Utility. *Nature Biotechnol.* **2017**, *35* (3), 238-248.
3. Wan, W. B.; Seth, P. P., The Medicinal Chemistry of Therapeutic Oligonucleotides. *J. Med. Chem.* **2016**, *59* (21), 9645-9667.
4. Ryan, D. E.; Taussig, D.; Steinfeld, I.; Phadnis, S. M.; Lunstad, B. D.; Singh, M.; Vuong, X.; Okochi, K. D.; McCaffrey, R.; Olesiak, M.; Roy, S.; Yung, C. W.; Curry, B.; Sampson, J. R.; Bruhn, L.; Dellinger, D. J., Improving CRISPR-Cas Specificity with Chemical Modifications in Single-Guide RNAs. *Nucleic Acids Res.* **2018**, *46* (2), 792-803.
5. Verheggen, I.; Van Aerschot, A.; Toppet, S.; Snoeck, R.; Janssen, G.; Balzarini, J.; Declercq, E.; Herdewijn, P., Synthesis and Antiherpes Virus Activity of 1,5-Anhydrohexitol Nucleosides. *J. Med. Chem.* **1993**, *36* (14), 2033-2040.
6. Declercq, R.; Herdewijn, P.; VanMeervelt, L., 1,5-Anhydro-2,3-Dideoxy-2-(Guanin-9-yl)-D-Arabin-Hexitol. *Acta Crystallogr. C* **1996**, *52*, 1213-1215.
7. Van Aerschot, A.; Verheggen, I.; Hendrix, C.; Herdewijn, P., 1,5-Anhydrohexitol Nucleic-Acids, a New Promising Antisense Construct. *Angew. Chem. Int. Ed. Eng.* **1995**, *34* (12), 1338-1339.
8. Hendrix, C.; Rosemeyer, H.; Verheggen, I.; Seela, F.; VanAerschot, A.; Herdewijn, P., 1',5'-Anhydrohexitol Oligonucleotides: Synthesis, Base Pairing and Recognition by Regular Oligodeoxyribonucleotides and Oligoribonucleotides. *Chem. Eur. J.* **1997**, *3* (1), 110-120.
9. Lescrinier, E.; Esnouf, R.; Schraml, J.; Busson, R.; Heus, H. A.; Hilbers, C. W.; Herdewijn, P., Solution Structure of a HNA-RNA Hybrid. *Chem. Biol.* **2000**, *7* (9), 719-731.
10. Pinheiro, V. B.; Taylor, A. I.; Cozens, C.; Abramov, M.; Renders, M.; Zhang, S.; Chaput, J. C.; Wengel, J.; Peak-Chew, S. Y.; McLaughlin, S. H.; Herdewijn, P.; Holliger, P., Synthetic Genetic Polymers Capable of Heredity and Evolution. *Science (Washington, DC, US)* **2012**, *336* (6079), 341-344.

11. Taylor, A. I.; Pinheiro, V. B.; Smola, M. J.; Morgunov, A. S.; Peak-Chew, S.; Cozens, C.; Weeks, K. M.; Herdewijn, P.; Holliger, P., Catalysts from Synthetic Genetic Polymers. *Nature* **2015**, *518* (7539), 427-430.
12. Pezo, V.; Liu, F. W.; Abramov, M.; Froeyen, M.; Herdewijn, P.; Marliere, P., Binary Genetic Cassettes for Selecting XNA-Templated DNA Synthesis In Vivo. *Angew. Chem. Int. Ed.* **2013**, *52* (31), 8139-8143.
13. Piperno, A.; Chiacchio, M. A.; Iannazzo, D.; Romeo, R., Synthesis and Biological Activity of Phosphonated Nucleosides: Part 1. Furanose, Carbocyclic and Heterocyclic Analogues. *Curr. Med. Chem.* **2006**, *13* (30), 3675-3695.
14. De Clercq, E.; Holy, A., Acyclic Nucleoside Phosphonates: A Key Class of Antiviral Drugs. *Nat. Rev. Drug Discov.* **2005**, *4* (11), 928-940.
15. Pradere, U.; Garnier-Amblard, E. C.; Coats, S. J.; Amblard, F.; Schinazi, R. F., Synthesis of Nucleoside Phosphate and Phosphonate Prodrugs. *Chem. Rev.* **2014**, *114* (18), 9154-9218.
16. Perezperez, M. J.; Rozenski, J.; Herdewijn, P., Stereospecific Synthesis of a Pentopyranosyl Analog of D4t Monophosphate. *Bioorg. Med. Chem. Lett.* **1994**, *4* (10), 1199-1202.
17. Perezperez, M. J.; Balzarini, J.; Rozenski, J.; Declercq, E.; Herdewijn, P., Synthesis and Antiviral Activity of Phosphonate Derivatives of Enantiomeric Dihydro-2h-Pyranyl Nucleosides. *Bioorg. Med. Chem. Lett.* **1995**, *5* (11), 1115-1118.
18. Perezperez, M. J.; Doboszewski, B.; Declercq, E.; Herdewijn, P., Phosphonates Derivatives of 2',3'-Dideoxy-2',3'-Didehydro-Pentopyranosyl Nucleosides. *Nucleosides Nucleotides* **1995**, *14* (3-5), 707-710.
19. Perezperez, M. J.; Doboszewski, B.; Rozenski, J.; Herdewijn, P., Stereocontrolled Synthesis of Phosphonate Derivatives of Tetrahydro-2h-Pyranyl and Dihydro-2h-Pyranyl Nucleosides - the Selectivity of the Ferrier Rearrangement. *Tetrahedron: Asymmetry* **1995**, *6* (4), 973-984.
20. Perezperez, M. J.; Rozenski, J.; Busson, R.; Herdewijn, P., Application of the Mitsunobu-Type Condensation Reaction to the Synthesis of Phosphonate Derivatives of Cyclohexenyl and Cyclohexanyl Nucleosides. *J. Org. Chem.* **1995**, *60* (6), 1531-1537.
21. Du, J. F.; Choi, Y.; Lee, K.; Chun, B. K.; Hong, J. H.; Chu, C. K., A Practical Synthesis of L-FMAU from L-Arabinose. *Nucleosides Nucleotides* **1999**, *18* (2), 187-195.
22. Pav, O.; Zbornikova, E.; Budesinsky, M.; Rosenberg, I., A New Class of Phosphanucleosides Containing a 3-Hydroxy-1-Hydroxymethylphospholane 1-Oxide Ring. *Tetrahedron* **2013**, *69* (43), 9120-9129.
23. The formation of the β isomer was also observed, however isolation by column chromatography provided only trace amounts of material insufficiently pure for complete characterization.
24. Ostrowski, T.; Wroblowski, B.; Busson, R.; Rozenski, J.; De Clercq, E.; Bennett, M. S.; Champness, J. N.; Summers, W. C.; Sanderson, M. R.; Herdewijn, P., 5-Substituted Pyrimidines with a 1,5-Anhydro-2,3-Dideoxy-D-Arabino-Hexitol Moiety at N-1: Synthesis, Antiviral Activity, Conformational Analysis, and Interaction with Viral Thymidine Kinase. *J. Med. Chem.* **1998**, *41* (22), 4343-4353.
25. Robeyns, K. H., P.; Van Meervelt, L., Direct Observation of Two Cyclohexenyl (CeNA) Ring Conformations in Duplex DNA. *Artif DNA: PNA XNA* **2010**, *1*, 2-8.
26. Kim, C. U.; Luh, B. Y.; Martin, J. C., Regiospecific and Highly Stereoselective Electrophilic Addition to Furanoid Glycals - Synthesis of Phosphonate Nucleotide Analogs with Potent Activity against HIV. *J. Org. Chem.* **1991**, *56* (8), 2642-2647.
27. Koh, Y. H.; Shim, J. H.; Wu, J. Z.; Zhong, W. D.; Hong, Z.; Girardet, J. L., Design, Synthesis, and Antiviral Activity of Adenosine 5'-Phosphonate Analogues as Chain Terminators against Hepatitis C Virus. *J. Med. Chem.* **2005**, *48* (8), 2867-2875.