Underlying structure of collective bands and self-organization in quantum systems

T. Otsuka*,^{1,2,3} Y. Tsunoda,⁴ T. Abe,⁴ N. Shimizu,⁴ and P. Van Duppen³

¹*Department of Physics, The University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan*

²*RIKEN Nishina Center, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan*

³*KU Leuven, Instituut voor Kern- en Stralingsfysica, 3000 Leuven, Belgium*

⁴*Center for Nuclear Study, The University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan*

(Dated: October 22, 2019)

The underlying structure of low-lying collective bands of atomic nuclei is discussed from a novel perspective on the interplay between single-particle and collective degrees of freedom, by utilizing state-of-the-art configuration interaction calculations on heavy nuclei. Besides the multipole components of the nucleon-nucleon interaction that drive collective modes forming those bands, the monopole component is shown to control the resistance against such modes. The calculated structure of ^{154}Sm corresponds to coexistence between prolate and triaxial shapes, while that of ^{166}Er exhibits a deformed shape with a strong triaxial instability. Both findings differ from traditional views based on β/γ vibrations. The formation of collective bands is shown to be facilitated from a self-organization mechanism.

The structure of atomic nuclei exhibits single-particle as well as collective-mode aspects created by the protons and neutrons (nucleons). The former has been characterized by the shell structure shown initially by Mayer [1] and Jensen [2], while the latter has presented a variety of nuclear shapes following Rainwater [3], and Bohr and Mottelson [4–6]. The two aspects lead to “the problem of reconciling the simultaneous occurrence of single-particle and collective degrees of freedom ...” [7]. This is one of the most important basic questions in nuclear structure research, and it remains open. For instance, G.E. Brown has addressed the question, “how single particle states can coexist with collective modes”, throughout his life [8]. We discuss in this Letter this problem from a novel perspective.

Nucleons in an atomic nucleus occupy single-particle orbits in various configurations. The effective nucleon-nucleon (NN) interaction in nuclei induces multi-nucleon correlations by mixing such configurations. This mixing occurs, in many cases, basically for “valence” nucleons in single-particle orbits on top of the appropriate closed proton and neutron shell (inert core). The ellipsoidal shapes correspond to such correlations having the nature of quadrupole surface deformation from a sphere, driven by the quadrupole component of the NN interaction [9–14]. This gives rise to an interplay between collective mode and single-particle states (SPS’s). If SPS’s relevant to these correlations are separated by large gaps, the mixing between them and the resulting correlations are reduced. Thus, the SPS’s can act as a “resistance” to (the formation of) collective modes. In this Letter, we first present how ellipsoidal shapes emerge from multi-nucleon systems by using the state-of-the-art Configuration Interaction (CI) simulations, called Monte Carlo Shell Model (MCSM)[15, 16]. This allowed us to uncover a novel mechanism: the monopole component of the NN interaction shifts the single-particle energies (SPEs) effectively, weakening the “resistance” against deformation, and thus

FIG. 1. Systematic changes of the 2_1^+ and 4_1^+ levels in Sm isotopes, as functions of N . **a** Energy levels[25], **b** $B(E2; 2_1^+ \rightarrow 0_1^+)$ and $B(E2; 4_1^+ \rightarrow 2_1^+)$ values[25], **c**. spectroscopic electric quadrupole moment of the 2_1^+ state[26].

enlarging effects of the quadrupole interaction. For textbook examples of strongly deformed nuclei, such as ^{154}Sm and ^{166}Er , the obtained properties agree with experiments, but the underlying structures are shown to differ from the traditional interpretation [4]. This mechanism can be interpreted as a quantum-mechanical self-organization[17].

We performed CI calculations on the samarium isotopes (proton number, $Z=62$) with even numbers of neutrons, $N=82-92$, and the ^{166}Er ($Z=68$, $N=98$) nucleus, without any assumptions of collective modes or shapes. A many-body Schrödinger equation is solved for the input NN interaction, which remains the same for all calculations. The valence proton (neutron) orbits are all orbits in the $sdg(pfh)$ -shell and the lower half of the next shell, implying the one-and-half harmonic oscillator shell on top of a ^{110}Zr inert core. This large model space is essential for the present study. The CI calculations need the SPEs with respect to the inert core, the appropriate values of which are taken and kept; if available, based on known ones at ^{132}Sn . The effective NN interaction is taken from the V_{MU} interaction for the proton-neutron channel [18] with a factor of 0.94 to its $T=0$ (T :isospin) central part. The proton-proton and neutron-neutron channels are taken from [19]. For the cases where this recipe is not possible, the V_{MU} interaction is used. Note that the V_{MU} interaction was determined as

FIG. 2. Levels of ^{154}Sm . **a** experimental levels[25], **b,c,d** calculated levels of the original and monopole-frozen Hamiltonians.

a simple modeling to the microscopic shell-model interactions [18], and that it was used in earlier studies on Zr, Sn and Hg isotopes[20–24]. The dimension of the many-body Hilbert space reaches beyond 10^{31} , which is formidably larger than the current limit ($\sim 10^{11}$) of the conventional CI method.

Figure 1a exhibits the experimental levels for Sm isotopes[25] as a function of N in comparison to the presently calculated ones. The $E_x(2_1^+)$ is high for $N=82$, a magic number, but becomes lower steadily as N increases. The 2_1^+ states around $N=86$ imply a quadrupole surface oscillation on top of a spherical ground state. With larger N , the deformation becomes static out to an ellipsoid, which generates a rotational band, a Nambu-Goldstone effect [27–29]. The $J^\pi=4_1^+$ levels exhibit a vibrational two-phonon pattern with $E_x(4_1^+)/E_x(2_1^+) \sim 2$ at $N=86$, while it evolves to the rotational value ($10/3$) for $N=92$, both in experiment and calculation[30].

Figure 1b,c display the $B(E2; 2_1^+ \rightarrow 0_1^+)$ and $B(E2; 4_1^+ \rightarrow 2_1^+)$ values and the spectroscopic electric quadrupole moment of the 2_1^+ state. The standard effective charges $1.5e$ ($0.5e$) for protons (neutrons) are taken [4]. An overall agreement between calculation and experiment suggests the validity of the present work, which is actually the first CI calculation for the shape evolution of samarium isotopes. Previously, this topic was studied in different frameworks (see *e.g.* [31–35]).

We now focus on ^{154}Sm ($N=92$). Figure 2a,b show its observed and calculated energy levels classified in four rotational bands[25]. A good agreement between experiment and calculation is observed. The band built on the ground state is prolate deformed, as confirmed by the large quadrupole moment and $B(E2)$ values shown in Fig. 1.

Figure 3 shows a deeper insight in the calculation performed for ^{154}Sm . It includes Potential Energy Surface (PES) obtained by Hartree-Fock calculation using the same Hamiltonian with, as constraints, the quadrupole moments corresponding to specific values of β_2 and γ (their relations are explained in [23, 24, 36]). The latter are shape variables

of the ellipsoidal shape being the magnitude and the proportion of the ellipsoid axes, respectively (see Fig. 3a)[4]. Figure 3 includes panels of three-dimensional PES, relative to the lowest energy. Figure 3b exhibits the lowest values of such PES values for a given value of γ .

We can now “visualize” the eigenstates obtained for ^{154}Sm by using the so-called *T-plot*[37, 38]. An eigenstate is expanded by MCSM basis vectors. One can assign partial coordinates to each MCSM basis vector by its β_2 and γ values, and can plot it on the PES. In this plot, the importance of each basis vector for this particular eigenstate is expressed by the size of the plotted circle. Figure 3c indicates that the T-plot circles for the ground state are concentrated around $\beta_2 = 0.28$ and $\gamma = 0^\circ$, a prolate shape. The T-plots of 2^+-8^+ members of the band, independently obtained, exhibit very similar patterns. This is interpreted as a strong signature of belonging to the same band. Figure 3d shows T-plot circles belonging to the 0_2^+ state at 1.1 MeV. The circles are concentrated in a local minimum at $\gamma \sim 20^\circ$, a triaxial shape[39, 40]. The 0_2^+ , $2_{2,3}^+$, 3_1^+ and $4_{2,3}^+$ states show almost identical T-plots. Thus the present calculation indicates a coexistence between prolate and triaxial shapes, in a stark contrast to the conventional picture of the β/γ vibrations (see *e.g.* [30, 41, 42]).

In order to clarify the underlying mechanisms giving rise to this picture, we decompose NN interaction into two components: monopole and multipole interactions (see reviews *e.g.* [38, 43–45]). The effect of the monopole interaction between protons in the orbit j_p and neutrons in the orbit j_n is expressed as

$$v_{pn}^m(j_p, j_n) n_p(j_p) n_n(j_n) \quad (1)$$

where v_{pn}^m is a coefficient called *monopole matrix element*, $n_p(j_p)$ denotes the number of protons in the orbit j_p , and $n_n(j_n)$ means likewise for neutrons. We first discuss how this interaction works. For the state being considered (*e.g.* the 0_1^+ state), $n_n(j_n)$ takes integers, 0, 1, 2, ... with the corresponding probabilities. Once one of these integers, k , substitutes $n_n(j_n)$, this interaction energy becomes $k v_{pn}^m(j_p, j_n) n_p(j_p)$, which represents a shift of the SPE of the orbit j_p by $k v_{pn}^m(j_p, j_n)$. This has two important aspects, (i) the SPE is modified, (ii) the effect is proportional to the number of neutrons in j_n . So, this varies effectively the SPE of the proton j_p orbit depending which neutron orbits are occupied. By including contributions from all neutron orbits and also those due to the proton-proton interaction, we define the effective SPE (ESPE) of the proton orbit j_p . Its expectation value is depicted in Fig. 3g for the $0_{1,2}^+$ states. The expectation value of $n_p(j_p)$ is shown also. The same quantities for neutrons are displayed in panel h.

If the monopole matrix elements were identical with respect to j_p and j_n in eq. (1), the energy would be moved by the same amount for all states of a given nucleus. This is, however, not the case with the realistic NN interaction: the finite range property of central force and the spin-isospin-dependence of tensor force produce significant variations with respect to j_p and j_n in eq. (1) [18, 45, 46].

FIG. 3. Properties of the $0_{1,2}^+$ states of ^{154}Sm . **a** Deformation parameters and shapes. **b** lowest value of PES for a given γ value for the original case and the prolate and spherical monopole-frozen cases. **c, d, e, f** 3-dimensional PES in the original or spherical monopole-frozen case (lower parts only in **d** and **f**). **g, h** ESPE (vertical position) and occupation number (horizontal width).

The multipole interaction contains various pieces, but the part most relevant to this work is expressed as the coupling between the proton and neutron quadrupole moments (operators) as $v_{pn}^q Q_p \cdot Q_n$, where v_{pn}^q is the interaction strength, and Q_p and Q_n denote, respectively, proton and neutron quadrupole moments coupled by a scalar product. With negative v_{pn}^q , quadrupole moments produce binding energy through this interaction, which can result in strongly deformed states. Namely, the quadrupole interaction thus defined is the driving force of the collective mode of quadrupole deformation.

The quadrupole moment can become quite large, if several relevant single-particle orbits mix coherently, as a realization of the Jahn-Teller effect [47]. If those orbits are far away from each other in energy, they cannot mix much. However, thanks to the monopole interaction and possible variations of the occupation pattern, the ESPEs can be shifted so that a more coherent mixing occurs giving more binding energy to the nucleus. This ESPE optimization does occur, and arises differently for each collective mode

or shape, as visualized by the average ESPEs. Figure 3g,h exhibit them for the 0_1^+ and 0_2^+ states of ^{154}Sm , which are of prolate and triaxial shapes, respectively. Such differences are due to different occupation patterns (see horizontal levels in Fig. 3g,h) and the afore-mentioned orbital dependences of the monopole interaction.

In order to shed further light on this effect, we performed CI calculations without the monopole interaction and instead the average ESPEs are adopted as input SPEs (*i.e.* constants), as denoted ‘monopole-frozen’ analysis. Figure 2c depicts levels by adopting the average ESPEs of the 0_1^+ state. The properties of ground-band members are rather unchanged from the original calculation naturally. However, because the ESPE optimization for the 0_2^+ state is completely ignored, the 0_2^+ energy is raised by 50%, as well as the 2_2^+ and 4_2^+ levels, as shown in Fig. 2c. This is consistent with the PES: Fig. 3b indicates that the local minimum around $\gamma=15^\circ$ on the ‘prolate’ line is higher by about 500 keV than the corresponding one of the ‘original’ line. Another monopole-frozen analysis was made by taking average ESPEs calculated at the spherical limit of the PES. Figure 3e, f display the PES and T-plot, indicating that the 0_1^+ state is no longer prolate but triaxial, whereas the 0_2^+ state becomes prolate. Figure 2d verifies this in terms of energy levels with the 0_1^+ state being triaxial (green level) and the 0_2^+ state being prolate (red level). These two examples of the monopole-frozen analysis demonstrate the crucial roles of the monopole interaction.

The energy gain of the prolate minimum measured from the spherical limit is ~ 8 MeV in the original calculation (see Fig. 3c). While it is only ~ 5 MeV in the monopole-frozen analysis with the spherical limit (see Fig. 3e). Thus, the ESPE optimization is shown to lower the energy of the prolate state by ~ 3 MeV. The same analysis for the triaxial states indicates gain of about 1 MeV. The ESPE optimization thus yields varying energy gains for the different shapes, lowering the prolate bands more.

Finally we note that also the presence of the negative-parity band built on the 1_1^- level has been reproduced (see Fig. 2a), reinforcing the validity of the present calculation, particularly the shell gaps.

The second example is ^{166}Er , which has shape characteristics different from those of ^{154}Sm . The Hamiltonian is the same as that for the Sm isotopes except for a minor change for a better description: proton $1d_{3/2}$ ($0g_{7/2}$) SPE shifted by 0.5 (-0.5) MeV. Figure 4a displays the lowest energy levels, calculated and measured [25]. The lowest side band starts from the $J^\pi=2_2^+$ level, called the $K^\pi=2^+$ band, which has been considered to be a γ vibration, where two short axes of the elongated ellipsoid oscillate keeping the volume constant [4, 25, 48]. One of the crucial quantities for this picture has been a relatively large value of $B(E2; 2_2^+ \rightarrow 0_1^+)$, which is 5.17 (21) W.u. experimentally[25]. This value is reproduced by the MCSM calculation to be 5.3 W.u. The electric spectroscopic quadrupole moments of the $2_{1,2}^+$ states are reproduced well[26]. Thus, we have a salient de-

FIG. 4. Experimental and calculated properties of the lowest states of ^{166}Er . **a** Energy levels and electromagnetic transitions (W.u.) [25] as well as spectroscopic electric quadrupole moments (e.b.) [26]. **b** Three-dimensional PES and its cut surface for $\beta_2=0.3$. **c, d, e** T-plots for the 0_1^+ and 2_2^+ states, and for the monopole-frozen 0_1^+ state at spherical shape.

scription of the 0_1^+ and $2_{1,2}^+$ states of ^{166}Er as well as the other states (see Fig. 4a).

Figure 4b displays the PES for ^{166}Er , where the energy minimum is stretched in the γ direction. Figure 4c, d display T-plots for the 0_1^+ and 2_2^+ state, which appear to be almost identical. We thus see γ -softness with γ between 5 - 15° . This does not seem to be compatible with the traditional picture of the prolate shape for the 0_1^+ state and the γ -vibrational excitation for the 2_2^+ state [4]. Rather, the T-plots indicate that these states stem from the same minimum in the triaxial plane. The quadrupole invariant analysis [49] made with the calculated quadrupole matrix elements of the states shown in Fig. 4a results in $\gamma=9^\circ$ for the electric matrix elements and $\gamma=10^\circ$ for the mass ones, respectively, consistently with the above range of γ . In case of rigid triaxial deformation with $\gamma=9^\circ$, the 2_2^+ level is expected to be much higher in energy [40], indicating that the present γ -softness lowers the band built on this state. The monopole-frozen analysis using the spherical ESPE produces a completely different T-plot for the ground state with a concentration into the region of $\gamma < 4^\circ$ (see Figure 4e), being consistent with a prolate shape. Thus, the present ESPE optimization is crucial also for the γ -softness being discussed.

The traditional interpretation of the “band” structures in ^{154}Sm and ^{166}Er is not supported by the MCSM calculations presented here. The underlying structure of ^{154}Sm appears much more like a shape coexistence [44-47] between a prolate minimum and a triaxial minimum. While the possibility of another equilibrium was already mentioned by Bohr and Mottelson [4], it was not further investigated in detail (see also a recent review paper on the experimental findings [55]). The present calculation indicated that by adding six protons and six neutrons, the two minima of ^{154}Sm are moved closer and merged by reaching ^{166}Er , suggesting that nuclear forces can produce a wide diversity of structures.

Finally, we put the quadrupole-monopole interplay in the context of self-organization in atomic nuclei [17]. In self-organization, a system is initially disordered, which corresponds to the SPEs without the optimization of the ESPE. In the present work, order implies that the ESPEs are tailored to a specific shape. Generally, some order may arise due to the self-organization in response to a change in external conditions. Atomic nuclei are, to a large extent, isolated quantum systems. However, if there are two kinds of ingredients, like protons and neutrons, one can behave like the source of an external force on the other. By activating the present monopole mechanism, the ESPEs are organized so that more binding energies are gained, as compared to the SPEs without the optimization (“disordered” system). While these SPEs can be a resistance to the collective mode (as stated earlier), the monopole interaction can act as a resistance-control force; in the present context, it can bring a certain order to the system. While the final solution is determined self-consistently by including all components of the NN interaction, a “positive feedback” can occur in this process especially between the quadrupole and monopole effects. As the present optimization effect varies for different shapes, it may appear to act purposely for a particular shape, for instance, a prolate one, although the monopole interaction as such has no connection to the deformation. This “to act purposely” is one of the features of self-organization.

In summary, CI calculations with a largest scale have clarified the structure of various bands in heavy nuclei. The interplay between the quadrupole and monopole interaction produces significant effects, as a quantum version of the self-organization. Namely, in a simplified view based on Landau’s Fermi liquid theory [54], nucleons are like free particles (quasiparticles) in the mean potential with fixed SPEs, and interact only weakly through a (residual) NN interaction. However, the actual structure seems to lie beyond this scope, because of the richness of the nuclear forces. The resistance-control force, *i.e.* the monopole interaction, does not promote the collective mode by itself, but can change the “environment”: the ESPEs are optimized to different collective modes giving rise to a large diversity of phenomena [20, 21, 23, 24, 37, 38, 56, 57]. Although the CI calculations can be refined, the overall agree-

ment to experiments suggests the validity of the underlying picture, shifting our basic understanding from the Liquid Drop model including its quantized forms to a more explicit multi-nucleon description with aspects such as shape coexistence, triaxial instability and shell evolution. This mechanism is expected to be more important in heavier nuclei including the superheavy ones, because more valence orbits and more nucleons imply more degrees of freedom for optimizing the SPEs. The relevance to superdeformation and clustering is of interest also, as particular configurations can be important. In heavy nuclei like uranium, new local minima or widening of minima in the γ direction may appear at larger deformations on the PES, providing a new scope for the fission process, where the time evolution proceeds through possible tunneling paths linking those minima. It will be interesting also to explore similar self-organization mechanisms in other quantum systems as they can give rise to unexpected features.

-
- [1] Goeppert Mayer, M. Phys. Rev. **75**, 1969 (1949).
- [2] Haxel, O., Jensen, J. H. D. & Suess, H. E. Phys. Rev. **75**, 1766 (1949).
- [3] Rainwater, J. Phys. Rev. **79**, 432 (1950).
- [4] Bohr, A. & Mottelson, B.R. *Nuclear Structure* (Benjamin, New York, 1975), Vol. II.
- [5] Bohr, A. Mat. Fys. Medd. Dan. Vid. Selsk. **26** (1952).
- [6] Bohr, A. & Mottelson, B. R. Mat. Fys. Medd. Dan. Vid. Selsk. **27** (1952).
- [7] Bohr, A. & Mottelson, B.R. *Nuclear Structure* (Benjamin, New York, 1969), Vol. I.
- [8] Schäfer, T Nucl. Phys. A **928**, 180 (2014).
- [9] Nilsson, S. G. Dan. Mat. Fys. Medd. **29**, No. 16 (1955).
- [10] Elliott, J.P. Proc. Roy. Soc. (London) A **245**, 128 (1958).
- [11] Elliott, J.P. Proc. Roy. Soc. (London) A **245**, 562 (1958).
- [12] Bes, D. R. & Sorensen, R. A. in *Advances in Nuclear Physics* (Plenum, New York, 1969; Springer, Berlin), Ch. 3.
- [13] Dufour, M. & Zuker, A. P. Phys. Rev. C **54**, 1641 (1996).
- [14] Kaneko, K., Sun, Y., Mizusaki, T. & Hasegawa, M. Phys. Rev. C **83**, 014320 (2011).
- [15] Otsuka, T. Honma, M. Mizusaki, T. Shimizu, N. & Utsuno, Y. Prog. Part. Nucl. Phys. **47**, 319 (2001).
- [16] Shimizu, N. *et al.* Prog. Theor. Exp. Phys. **2012**, 01A205 (2012).
- [17] Wikipedia, <https://en.wikipedia.org/wiki/Self-organization>.
- [18] Otsuka, T. Suzuki, T. Honma, M. Utsuno, Y. Tsunoda, N. Tsukiyama, K. and Hjorth-Jensen, M. Phys. Rev. Lett. **104**, 012501 (2010).
- [19] Brown, B. A. Phys. Rev. Lett. **85**, 5300 (2000).
- [20] Togashi, T., Tsunoda, Y., Otsuka, T., Shimizu, N. Phys. Rev. Lett., **117**, 172502 (2016).
- [21] Kremer, C. *et al.* Phys. Rev. Lett., **117**, 172503 (2016).
- [22] Togashi, T., Tsunoda, Y., Otsuka, T., Shimizu, N. and Honma, M Phys. Rev. Lett., **121**, 062501, (2018).
- [23] Marsh, B. A. *et al.* Nature Physics **14**, 1163-1167 (2018).
- [24] Sels, S. *et al.* Phys. Rev. C **99**, 044306 (2019).
- [25] National Nuclear Data Center. Evaluated Nuclear Structure Data File. <http://www.nndc.bnl.gov/ensdf/>.
- [26] Stone, N. J. At. Data and Nucl. Data Tables, **111-112**, 1, (2016).
- [27] Nambu, Y. Phys. Rev. Lett. **4**, 380 (1960).
- [28] Goldstone, J. Il Nuovo Cimento **19**, 154 (1961).
- [29] Goldstone, J. Salam, A. and Weinberg, S. Phys. Rev. **127**, 965 (1962).
- [30] Casten, R. F. *Nuclear structure from a simple perspective*, (Oxford University Press, New York, 2000).
- [31] Scholten, O. Iachello, F. Arima, A. Ann. Phys. **115**, 325 (1978).
- [32] Nomura, K. Shimizu, N. and Otsuka, T. Phys. Rev. Lett. **101**, 142501 (2008).
- [33] Robledo, L. M. Rodríguez-Guzmán, R. R. and Sarri-guren, P. Phys. Rev. C **78**, 034314 (2008).
- [34] Li, Z. P. Nikšić, T. Vretenar, D. Meng, J. Lalazissis, G. A. and Ring, P. Phys. Rev. C **79**, 054301 (2009).
- [35] Delaroche, J. -P. Girod, M. Libert, J. Goutte, H. Hilaire, S. Péru, S. Pillet, N. and Bertsch, G. F. Phys. Rev. C **81**, 014303 (2010).
- [36] Utsuno, Y. *et al.*, Phys. Rev. Lett., **114**, 032501, (2015).
- [37] Tsunoda, Y., Otsuka, T., Shimizu, N., Honma, M. & Utsuno, Y. Phys. Rev. C **89**, 031301(R) (2014).
- [38] Otsuka, T. and Tsunoda, Y. J. Phys. G, **43**, 024009 (2016).
- [39] Davydov, A. S. and Filippov, G. F. Nucl. Phys. **8**, 237 (1958).
- [40] Davydov, A. S. and Rostovsky, V. S. Nucl. Phys. **12**, 58 (1959).
- [41] Krücken, R. Barton, C. J. Beausang, C. W. Casten, R. F. Cata-Danil, G. Cooper, J. R. Novak, J. Yang, L. Wilhelm, M. Zamfir, N. V. and Zilges, A. Phys. Lett. B **454**, 15 (1999).
- [42] Möller, T. *et al.* Phys. Rev. C **86**, 031305 (2012).
- [43] Poves, A. & Zuker, A. Phys. Rep. **70**, 235 (1981).
- [44] Caurier, E., Martínez-Pinedo, G., Nowacki, F., Poves, A. & Zuker, A.P. Rev. Mod. Phys. **77**, 427-488 (2005).
- [45] Otsuka, T. Gade, A. Sorlin, O. Suzuki, T. and Utsuno, Y. Preprint at <http://arxiv.org/1805.06501> (2018).
- [46] Otsuka, T., Suzuki, T., Fujimoto, R., Grawe, H. & Akaishi, Y. Phys. Rev. Lett. **95**, 232502 (2005).
- [47] Jahn, H. A. and Teller, E. Proc. R. Soc. A, **161**, 220, (1937).
- [48] Bohr, A. N. *Nobel Lectures, Physics 1971-1980*, Editor Stig Lundqvist (World Scientific, Singapore, 1992); <https://www.nobelprize.org/prizes/physics/1975/bohr/facts/>.
- [49] Kumar, K., Phys. Rev. Lett. **28**, 249 (1971).
- [50] Morinaga, H. Phys. Rev. **101**, 254 (1956).
- [51] Wood, J. L. Heyde, K. Nazarewicz, W. Huyse, M. and Van Duppen, P. Phys. Rep. **215**, 101 (1992).
- [52] Andreyev, A. N. *et al.* Nature **405**, 430 (2000).
- [53] Heyde, K. and Wood, J. L. Rev. Mod. Phys. **83**, 1467 (2011).
- [54] Landau, L. Sov. Phys. JETP, **3**, 920 (1957).
- [55] Sharprey-Schafer, J. F. Bark, R. A. Bvumbi, S. P. Dinoko, T. R. S. and Majola, S. N. T. Eur. Phys. J. A. **55**, 15 (2019).
- [56] Leoni, S. *et al.* Phys. Rev. Lett. **118**, 162502 (2017).
- [57] Morales, A.I. *et al.* Phys. Lett. B **765**, 328-333 (2017).

ACKNOWLEDGEMENTS

T.O., Y.T., T.A. and N.S. acknowledge the support from MEXT as "Priority Issue on post-K computer" (Elucidation of the Fundamental Laws and Evolution of the Universe) (hp160211, hp170230, hp180179, hp190160) and JICFuS. T.O. acknowledges Prof. N. Nagaosa for valu-

able comments. T.O. and Y.T. acknowledge Prof. B. A. Brown for helping in handling interaction files. This work has received funding from the Research Foundation Flanders (FWO, Belgium), the Excellence of Science program (EOS, FWO-FNRS, Belgium) and the GOA/2015/010 (BOF KU Leuven).