1 Seismically-enhanced hydrothermal plume advection through the process zone of the
2 Compione extensional Fault, Northern Apennines, Italy

3 **Alessio Lucca¹, Fabrizio Storti¹, Giancarlo Molli^{1,2}, Philippe Muchez³, Andrea Schito⁴, Andrea**
4 **Artoni⁵, Fabrizio Balsamo¹, Sveva Corrado⁴, and Emma Salvioli Mariani⁵**

5 *¹Department of Chemistry, Life Sciences and Environmental Sustainability, Natural and*
6 *Experimental Tectonics research group, University of Parma, I-43124 Parma, Italy*

7 *²Department of Earth Sciences, University of Pisa, via S. Maria, 53, I-56126 Pisa, Italy*

8 *³Katholieke Universiteit Leuven, Department of Earth and Environmental Sciences, Celestijnenlaan*
9 *200E, B-3001 Heverlee, Belgium*

10 *⁴Department of Sciences, Geology Section, Roma Tre University, L.go S. Leonardo Murialdo 1, I-*
11 *00146, Roma, Italy*

12 *⁵Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, I-*
13 *43124 Parma, Italy*

14

15 *Corresponding Author Details

16 E-mail: alessio.lucca@studenti.unipr.it

17 Telephone number: +39 393 1323420

18

19

20 **ABSTRACT**

21 **Reconstructing the paleofluid evolution in mature fault zones, which typically have complex**
22 **structural architectures, is a challenging task because re-activation of pre-existing deformation**
23 **structures and dissolution-reprecipitation processes are very abundant. Understanding why**
24 **specific structural elements are preferentially mineralized and what are the factors leading to**
25 **rapid fluid migration and accumulation, bears geological and economic implications, especially**
26 **in seismically active fault zones. We studied the Compione Fault on the Tyrrhenian Sea side of**
27 **the Northern Apennines orogenic wedge, Italy, which is a segment of the 30 km long Northern**
28 **Lunigiana high-angle extensional fault system still active today. The Compione Fault**
29 **propagated from the metamorphic basement and accumulated about 1.5 km of displacement.**

30 We used structural, petrographic, isotopic, microthermometric, compositional and organic
31 matter analyses to constrain fluid and host rock properties during fault zone evolution. This
32 approach allowed us to quantify the thermal anomaly in the fault zone and to infer the processes
33 responsible for such a disequilibrium. Specifically, we show that in the fault process zone ahead
34 of the upper fault tip, which is twice as wide as the damage zone, seismic pumping caused
35 suprahydrostatic fluid pressures and that local dilation promoted the nucleation of a highly
36 permeable mesh of conjugate extensional shear fractures hosting calc-silicate mineralization.
37 The thermal difference between hydrothermal minerals in the conjugate fracture mesh and the
38 host rock is 60-90 °C. The mineralizing fluids were deeply-sourced from metamorphic
39 reactions. Propagation of the upper fault tip caused process zone folding and incorporation into
40 the fault damage zones. As the upper fault tip breached through shallower structural levels, it
41 favored mixing between deep and meteoric fluids.

42

43 1. INTRODUCTION

44 Fluid-rock interactions have been widely studied to better understand fluid migration and
45 accumulation and their effects on compositional, petrophysical and rheological rock modifications
46 during deformation (Nesbitt and Muehlenbachs, 1989; Evans and Battles, 1999, Roure et al., 2005;
47 Vilasi et al., 2009; Vandeginste et al., 2012). Textural, geochemical and microthermometric analyses
48 of syntectonic vein cements allow constraint of paleofluid properties, such as their origin, migration
49 pathways, temperature and pressure of crystallization and the local state of stress during deformation
50 (Mullis, 1979, 1987, 1988; Carter and Dworkin, 1990; Fisher et al., 1995; Muchez et al., 1995;
51 Milliken et al., 1998; Montomoli et al., 2001; Montomoli, 2002; Clemenzi et al., 2014; Honlet et al.,
52 2017). In particular, when deformation is thick-skinned, regional-scale fault systems with kilometric
53 offsets are characterized by highly-connected fracture networks in their damage zones, which are
54 preferential sites for fluid migration and mixing from the metamorphic basement up to surficial

55 aquifers (Gratier et al., 2002; Beaudoin et al., 2011; Doglioni et al., 2014; Mamadou et al., 2016;
56 Laurent et al., 2017; Wüstefeld et al., 2017). Cements hosted in fault-related fractures commonly
57 record a multi-stage paleofluid and deformational evolution characterized by dissolution-
58 reprecipitation processes and repeated, episodic fracturing (Phillips, 1972; Ramsay, 1980; Parry and
59 Bruhn, 1987; Fisher et al., 1995; Parry, 1998). The latter can be related to the earthquake cycle that,
60 depending on whether seismic pumping or fault-valve occurred (Sibson et al., 1975, 1988), causes
61 high fluid pressures in fault zones after or before earthquake ruptures, respectively (McCaig, 1988;
62 Boullier and Robert, 1992; Cox, 1995; Robert et al., 1995; Cox, 1999). Therefore, cement patterns in
63 seismically active fault damage zones can provide important exhumed analogues to better understand
64 the properties of fluids involved in seismic ruptures at depth. The permeability of fault zones major
65 components, which can act as either conduits, barriers or mixed conduits-barriers systems, exerts a
66 first-order control on the generation of fluid pressures and fluid flow. Fault zones with kilometric
67 displacement affecting sandstones are typically characterized by low-permeability cores and highly-
68 permeable damage zones (Caine et al., 1996; Faulkner and Rutter, 2001; Faulkner et al., 2010). The
69 latter, however, are heterogeneous rock volumes, which can be subdivided in wall damage zones, tip
70 damage zones and linking damage zones (*sensu* Kim et al., 2004) according to their structural
71 position. Tip damage zones, or process zones (*sensu* Cowie and Shipton, 1998) and linking damage
72 zones, at the tips of propagating faults, have higher permeability due to extensive fracture nucleation
73 with multiple orientations and are preferential sites for fluid flow compared to wall damage zones
74 (Curewitz and Karson, 1997). Accordingly, process zones and linking damage zones can enhance
75 deep fluid advection, producing positive temperature anomalies and chemical and barometric
76 disequilibrium between fluids and host rocks. However, fluid sources and pathways and,
77 consequently, the scales of fault-controlled fluid flow deserve further investigations, as well as the
78 relationships between vein infilling phases and damage zone evolution during fault growth.

79 In this contribution we present the results of a study of the structural architecture and paleofluid
80 evolution recorded in fault-related veins of the Compione fault zone, a segment of the about 30 km

81 long Northern Lunigiana high-angle basin-boundary extensional fault system cross-cutting the whole
82 nappe pile in the inner portion of the Northern Apennines (Fig.1). The Northern Lunigiana Fault
83 started developing since Early Pliocene times during the uplift and exhumation of the Apenninic
84 tectonic wedge and is still active today (Boncio et al., 2000; Eva et al., 2014; Bonini et al., 2013). The
85 Compione Fault accumulated about 1.5 km of displacement (Bernini and Lasagna, 1988; Bernini,
86 1991; Bernini and Papani, 2002) and offers suitable exposure conditions for performing a
87 representative structural transect across the fault zone. A multidisciplinary dataset was collected,
88 including independent geothermometers to evaluate thermal anomalies, and geochemical analysis to
89 constrain the fluid sources. Structural analysis was combined with vein cement petrography and
90 optical cold cathodoluminescence, SEM-EDS analysis, carbon and oxygen stable isotope analysis
91 and microthermometry. Moreover, maximum paleotemperatures recorded in the Macigno Sandstones
92 Formation were constrained by vitrinite reflectance of organic matter and thermal modeling.

93 Our results show that hydrothermal calc-silicate mineralization occurred in the process zone fracture
94 mesh during upward propagation of the Compione Fault, from fluids at temperatures higher than at
95 the burial peak in the host rock and synchronously with methane migration. Crystallization
96 temperatures of hydrothermal minerals in the footwall damage zone record the progressive
97 cooling/exhumation and folding of the process zone fracture network by extensional fault-
98 propagation folding (Withjack et al., 1990; Schlische, 1995; Hardy and McClay, 1999; Ferrill et al.,
99 2004a; Jin and Groshong, 2006), causing shear reactivation of favorably oriented hydrothermal
100 fractures. We propose a model where mineralization preferentially forms in process zones ahead of
101 propagating fault-tips and fluids responsible for fault-related mineralization are supplied by the
102 metamorphic basement and by the different stratigraphic horizons cut by the fault zone through a
103 seismic pumping mechanism.

104

105 2. GEOLOGIC SETTING

106 The Apennines segments of the Alpine-Himalayan orogenic belt were growing in Cenozoic times due
107 to the southwestward subduction and related slab retreat of the Adria micro-plate below the overriding
108 European plate (Principi and Treves, 1984; Malinverno and Ryan, 1986; Royden, 1988; Dewey et al.,
109 1989; Patacca et al., 1990; Doglioni, 1991; Jolivet et al., 1998). Two major paleogeographic domains
110 are telescoped in the Apennines: the ocean-derived Jurassic to Paleogene Ligurian succession and the
111 Adria passive margin domain (Boccaletti et al., 1971; Elter and Pertusati, 1973, Elter et al., 1975).
112 The Subligurian domain was originally located in the ocean-continent transition (Plesi, 1975;
113 Montanari and Rossi, 1982; Vescovi, 1998). Building up of the Northern Apennines thrust wedge
114 included underthrusting up to 15-20 km depth and greenschist facies metamorphism (Apuan Alps in
115 Fig. 1; Di Pisa et al., 1985; Franceschelli et al., 1986; Jolivet et al., 1998; Molli et al., 2000a, 2000b,
116 2002); while the non-metamorphic successions (Tuscan, Subligurian, Ligurian, Epiligurian
117 successions in Fig. 1) were affected by synorogenic extension episodes (Carmignani and Kligfield,
118 1990; Carmignani et al., 1994; Decandia et al., 1998; Jolivet et al., 1998; Molli, 2008; Clemenzi et
119 al., 2014) and out-of-sequence thrusting (Storti, 1995; Argnani, 2002; Boccaletti et al., 2011; Bonini
120 et al., 2013; Clemenzi et al., 2014). Extensional faulting and magmatism took place since upper
121 Miocene to Pleistocene times and migrated northeastward behind the advancing thrust fronts (Elter
122 et al., 1975; Serri et al., 1993; Bartole, 1995; Carmignani et al., 1995; Barchi et al., 1998; Martini et
123 al., 2001).

124 Conditions of deformation in the non-metamorphic, dominantly carbonatic thrust sheets of the
125 Northern Apennines have been determined through different methodologies: vitrinite reflectance
126 (Reutter et al., 1981; Corrado et al., 2010; Carlini et al., 2013), illite crystallinity (Cerrina Feroni et
127 al., 1983; Carosi et al., 2003; Carlini et al., 2013), stable isotopes (Carter and Dworkin, 1990; Milliken
128 et al., 1998; Mazzarini et al., 2010; Clemenzi et al., 2014), calcite-dolomite geothermometry (Carosi
129 et al., 2003), fluid inclusion microthermometry (Montomoli et al., 2001; Montomoli, 2002; Mazzarini
130 et al., 2010; Clemenzi et al., 2014) and apatite and zircon fission tracks (Abbate et al., 1994; Zattin et
131 al., 2002; Balestrieri et al., 2003; Bernet et al., 2004; Fellin et al., 2007; Corrado et al., 2010; Thomson

132 et al., 2010; Carlini et al., 2013). The estimated maximum burial is about 7 km at temperatures ranging
133 between 200 and 250 °C. Geothermal gradients calculated with different methodologies span a wide
134 range from 18 °C/km to 41 °C/km with a mean value of 31 ± 4 °C/km (Molli et al., 2011).

135

136 **2.1 The Northern Lunigiana Basin**

137 The Northern Lunigiana Extensional Basin (NLB) is about 25 km long and 10 to 15 km wide. It is
138 the northwesternmost onshore extensional basin in the Northern Apennines and is located to the
139 northwest of the Alpi Apuane metamorphic complex, on the Tyrrhenian Sea side of the orogenic
140 wedge (Fig. 1). The Northern Lunigiana Basin is separated from the Southern Lunigiana Basin by the
141 Secchia transversal line (Fig. 1B; Ghelardoni, 1965). The Northern Lunigiana Basin developed in the
142 hanging wall of a regional-scale out-of-sequence thrust (Storti, 1995; Vescovi, 2005; Clemenzi et al.,
143 2014; Molli et al., 2018) that caused duplexing of the Northern Apennines nappe pile, i.e. subligurian
144 rocks overthrusting the Adria-derived Tuscan succession and overlain by Ligurian thrust sheets (Fig.
145 1D). As a result of such multiple thrusting events, the basin-boundary fault system juxtaposes the
146 Cretaceous Ligurian Ottone Flysch Formation in the hanging wall, against the Late Oligocene-Early
147 Miocene Macigno Sandstones Formation in the footwall (Figs. 1B and 1D; Elter and Schwab, 1959;
148 Giglia, 1974; Elter et al., 1975; Bernini and Lasagna, 1988). The Campanian to early Maastrichtian
149 Ottone Flysch Formation has a maximum thickness of 300-400 m and is composed of marly-
150 calcareous turbiditic and calcarenitic to lithoarenitic strata. The Chattian to Aquitanian Macigno
151 Sandstones Formation is in the studied area about 2300 m thick and is composed of massive sandstone
152 strata in the lower and upper part of the formation, separated by clay- and silt-rich strata (Ghibaudo,
153 1980). Massive sandstone beds are up to 5 m thick, while the silty and clayey facies beds are a
154 centimetre up to maximum 1.5 m thick. Macigno Sandstones are arkoses and contain abundant quartz,
155 feldspars, biotite, muscovite and chlorite.

156 The onset of extension in the NLB has been dated using palynology and mammal fauna (Azzaroli,
157 1950, 1977; Federici, 1978, 1981; Raggi, 1985; Bertoldi, 1988, 1995) in the fluvial-lacustrine deposits
158 that rest on the Ottone Flysch Formation in the depressed central area. These data indicate that two
159 sub-basins developed in response to extensional tectonics: Aulla-Olivola in the SE and Pontremoli in
160 the NW, starting from Early Pliocene and Early Pleistocene times respectively. The deposits follow
161 a regressive trend from lacustrine to alluvial-fan at the top. In the SE they are characterized by two
162 unconformities, the first Late Pliocene in age and the second dated at Middle-Pleistocene times, which
163 occurs also in the Pontremoli depocenter (Boccaletti et al., 1992; Boccaletti and Sani, 1998; Bernini
164 and Papani, 2002). Extension is interpreted to be active during a general uplifting phase of the inner
165 Apenninic belt (Bartolini et al., 1982; Cerrina Feroni et al., 1983; Bernini et al., 1990; Di Naccio et
166 al., 2013).

167 Both the Northern and the Southern Lunigiana extensional basins are tectonically active, as indicated
168 by extensional earthquakes occurring at depths typically shallower than 15 km, as well as few
169 contractional ones having their hypocenters at about 50 km (Bossolasco et al., 1973; Bossolasco et
170 al., 1974; Augliera et al., 1990; Frepoli and Amato, 1997; Boncio et al., 2000; Eva et al., 2014; Bonini
171 et al., 2013). This is further supported by morphotectonic evidence (Di Naccio et al., 2013). The NLB
172 formed above a northeastward-dipping low-angle detachment fault (Artoni et al., 1992; Camurri et
173 al., 2001; Argnani et al., 2003; Di Naccio et al., 2013) and it attains a half-graben geometry produced
174 by the activity of the NW-SE striking, SW dipping Northern Lunigiana basin-boundary extensional
175 fault system (Bernini, 1988; Bernini and Lasagna, 1988; Bernini, 1991; Bernini and Papani, 2002).
176 The asymmetric topography, with the Apenninic watershed to the NE (Fig. 1D), the northeastward
177 tilt of syn-extensional deposits (Bernini, 1988) and morphotectonic evidence (Di Naccio et al., 2013)
178 support the importance of the Northern Lunigiana extensional fault system in controlling the
179 development of the NLB. In the study area, the NLB is bounded to the NE by the Compione Fault
180 (Figs. 1B, 1D, 2 and Fig. DR1¹) which can be considered the SE prosecution of the Groppodalsio
181 fault (Fig. 3). Subsurface information provided by a seismic line located in the study area (Figs. 1B,

2) is quite scarce due to the poorly imaged geological complexity of the region (Fig. 2). The proposed geometry shows the Compione Fault cutting through the thrust sheet pile and penetrating into the seismic basement, as previously mapped (Argnani et al.; 2003; Camurri et al., 2001), at around 6 to 7 km depth with an almost planar geometry (Fig. 2), confirming its role in shaping the NLB. At the surface, the Compione Fault is located at the forelimb-crest transition of the anticline associated with the regional-scale out-of-sequence thrust (Figs. 1D; 3A and 3B; Vescovi, 2005; Clemenzi et al., 2014; Molli et al., 2018). Synthetic fault zones occur in the footwall, whereas several synthetic and antithetic fault zones dissect the hanging wall (Bernini, 1991; Bernini and Papani, 2002).

190

191 **3. METHODS**

192 **3.1 Structural Analysis**

About 600 structural data were collected at six structural sites, five of which located in the fault damage zones and one in the footwall host-rock, for comparative purposes. Structural data are reported according to the right-hand rule (strike/dip), and stereographic projections (lower hemisphere of Schmidt net) are plotted with the Daisy3 software (Salvini, 2017).

197

198 **3.2 Petrography and Cathodoluminescence**

Fifty veins were sampled in fault damage zones and in the hanging wall host rocks. Moreover, three more sampling sites were identified along strike of the footwall damage zone, to the west of the study transect, at a distance of 0.7, 1.75 and 15 km from it (Figs. 3A and 3C). Each sample was cut in two slabs, one of which was stained with Alizarin Red S and potassium ferricyanide to discriminate the different carbonates such as calcite and dolomite and their iron-rich equivalents (Dickson, 1966). Thin sections and wafers were obtained from the other slab. Petrographic and microstructural analyses were carried out on fifty 30 µm thick thin sections, through standard optical and cold CL

206 microscopy. A Zeiss Axioplan 2 microscope was used for optical petrography. A Technosyn 8200
207 Mark II cold CL stage, mounted on a LEICA DM2700P optical microscope, was used at 15 kV and
208 220-250 μ A gun current to perform CL analysis. Compositional analyses were carried out on
209 prehnites with a Jeol 6400 SEM equipped with an Oxford EDS. Operating conditions were 15 kV and
210 1.2 nA, an electron beam with a diameter of about 1 μ m and a counting time of 100 s. Errors are ± 2 –
211 5% for major elements and ± 5 –10% for minor components. Standards used to calibrate the EDS
212 include pure elements, simple oxides and simple silicate compositions (cobalt, anorthoclase, apatite,
213 augite, microcline and olivine).

214

215 **3.3 Carbon and Oxygen Stable Isotopes**

216 Stable isotope analyses for oxygen and carbon were performed on host rocks carbonate components
217 and on calcite cements previously identified by petrography and CL. Carbon isotope results are
218 expressed in Vienna Pee Dee Belemnite (V-PDB) while oxygen isotope results are expressed both in
219 V-PDB and in Vienna Standard Mean Ocean Water (V-SMOW) for convention purposes. $\delta^{18}\text{O}$ values
220 were converted using the equation (Friedman and O'Neil, 1977):

$$221 \quad \delta^{18}\text{O SMOW} = 1.03086 \cdot \delta^{18}\text{O VPDB} + 30.86 \quad (1)$$

222 Stable isotope analyses for oxygen and carbon (cf. paragraph 5.2) were carried out on 98 sub-samples
223 of veins and host rocks. Sub-samples were drilled directly from 33 thin sections using an ESI New
224 Wave Research Micromill with a 6.7X to 40X optical zoom, 3.3 mm to 24.5 mm field-of-view,
225 automated 50 mm travel in X, Y and Z directions stage with sub-micron step resolution and a milling
226 chuck speed ranging from 1200 rpm to 35000 rpm. 100-150 μ g of pure carbonate powder for each
227 sub-sample were loaded into a GasBenchII autosampler interspaced with three isotopically different
228 kinds of reference materials (NBS18, NBS19 and MAB99). After flushing the vessels with ultrapure
229 helium (5.5 grade) in order to replace the air, powders were reacted with 100% orthophosphoric acid

230 at 25 °C for 12 h (McCrea, 1950). Resulting gases were analyzed automatically using a Thermo
231 Finnigan Delta V+ mass spectrometer. For each sample, four reference gas peaks were measured and
232 the sample gas was introduced ten times into the mass spectrometer. Each sample was analyzed at
233 least in double so the uncertainty on the samples value may be considered $\pm 0.10\text{‰}$ and $\pm 0.23\text{‰}$ for
234 $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ respectively. Isotope fractionation curves were calculated according to O'Neil et al.
235 (1969).

236

237 **3.4 Micro-Raman**

238 Micro-Raman measurements were performed using a Jobin-Yvon Horiba LabRam spectrometer
239 equipped with a He-Ne laser (emission line 632.8 nm) and motorized XY stage. The spectral
240 resolution is about 2 cm^{-1} . The confocal hole was adjusted in order to obtain a spatial resolution
241 (lateral and depth) of 1-2 μm . Spectra were obtained using a 50X objective (0.75 N.A. [numerical
242 aperture]). The calibration was made using the 520.7 cm^{-1} Raman line of silicon. The scanned spectral
243 range spans from 1100 to 3300 cm^{-1} . Acquisition time was 120 s. The power on the sample surface
244 is around 1 mW but the power on the analysed inclusions has to be considered lower due light
245 reflection and scattering.

246

247 **3.5 Fluid Inclusion Microthermometry**

248 Fluid inclusions were studied in 44 wafers, i.e. doubly polished sections with a thickness of about
249 $100\text{ }\mu\text{m}$. Fluid inclusion petrography was carried out on a standard petrographic microscope in order
250 to distinguish assemblages, trails and isolated inclusions. Fluid inclusion assemblages (FIAs) are
251 groups of inclusions that occur along growth zones or randomly in the minerals and represent the
252 fluid conditions during precipitation or recrystallization (Goldstein and Reynolds, 1994). FIAs were
253 systematically measured. Fluid inclusion trails (FITs) are related to fracturing events after

crystallization of at least part of the host mineral and were not considered during microthermometric analysis. Isolated inclusions were measured and were considered reliable if they showed comparable temperatures to the inclusions organized in FIAs in the same sample. After fluid inclusion petrography, the doubly polished wafers were broken in smaller pieces (chips) and analyzed in a Linkam THMSG600 heating-freezing stage. The instrument was calibrated weekly using SynFlinC synthetic standards. Calibration lines were calculated from melting temperature of CO₂, final melting temperature of clathrate, homogenization temperature of CO₂, NaCl eutectic temperature, final ice melting temperature of pure H₂O and the critical homogenization temperature of pure H₂O. Three temperatures were acquired in aqueous biphasic inclusions: 1) homogenization into the liquid phase (Th_{tot}) indicating the minimum temperature of fluid entrapment; 2) first melting temperature (T_{fm}), related to the fluid composition; 3) ice melting temperature (T_{m_{ice}}), which is inversely proportional to the amount of solutes in the liquid phase, i.e. the salinity of the fluid. Heating was always performed before freezing runs to avoid artificial stretching of the inclusions during freezing. Monophase gaseous inclusions at room temperature develop a liquid meniscus during cooling in the heating-freezing stage. In these gaseous inclusions, only Th_{tot} was measured and homogenization occurred into the vapour phase. Monophase aqueous inclusions were kept in a freezer for 2 weeks at -20 °C to nucleate bubbles in metastable inclusions. Inclusions that did not nucleate the vapour phase, were cyclically heated and frozen to induce artificial stretching. In this way, it was possible to measure T_{m_{ice}} in monophase inclusions. Homogenization temperatures are evidently not measured in artificially stretched aqueous inclusions. Accuracy of measurements is ± 2 °C for homogenization temperatures (Th_{tot}) and ± 0.2 °C for first melting (T_{fm}) and ice melting temperatures (T_{m_{ice}}).

275

276 **3.6 Organic Matter Optical Analysis and Thermal Modelling**

277 Ten organic-rich laminae were sampled in the footwall damage zone (Sites 4 and 5) and in the
278 footwall host rock (Site 14) to constrain the thermal maturity of the Macigno Sandstones Formation

279 through vitrinite reflectance analyses. Vitrinite is the product of thermal maturation of terrestrial plant
280 remnants included in sediments and it is one of the most reliable indicator of peak temperatures at
281 maximum burial depth because it is very sensitive to temperature increase and not affected by
282 retrograde processes (Tissot and Welte, 1984; Teichmüller, 1987). Samples for vitrinite reflectance
283 analyses were crushed and then mounted in epoxy resin and polished, according to the method of
284 Bustin et al. (1990). Samples were analysed in reflected, non-polarized, monochromatic light ($\lambda =$
285 546 nm) under oil immersion ($v = 1.518$) using a Zeiss Axioplan MPM400 microscope equipped with
286 a J&M Analytik Tidas S 800 spectrometer and calibrated with CRAIC vitrinite reflectance standards.
287 Up to 40 Ro% (randomly oriented vitrinite reflectance in oil immersion) measurements per sample
288 were acquired. Thermal modeling was carried out using the BasinMod2 software, based on the
289 Easy%Ro kinetic model of Burnham and Sweeney (1989). The software requires organic maturity
290 indicators, sedimentary successions lithologies, thicknesses and ages along with paleo-heat flow or
291 paleo-geothermal gradient as basic input data. Sample stratigraphic locations were modelled as a
292 pseudo-well according to Oncken, 1982; Di Paolo et al., 2012; Caricchi et al., 2015; Schito et al.,
293 2018. The main assumptions for the modeling are: (1) decompaction of the burial curves is corrected
294 according to the method of Sclater and Christie (1980); (2) sea-level changes are neglected, as the
295 thermal evolution is influenced more by sediment thickness than water depth (Butler, 1992); (3)
296 thrusting is considered instantaneous when compared with the duration of sedimentation, as generally
297 suggested in theoretical models (Endignoux and Wolf, 1990); (4) geothermal gradient (25-30°C/km)
298 is calculated from the correlation of vitrinite reflectance data based on the kinetic model of vitrinite
299 maturation of Burnham and Sweeney (1989).

300

301 **4. STRUCTURAL ARCHITECTURE OF THE COMPIONE FAULT**

302 The area near Compione village was mapped in detail to describe the structural architecture of the
303 Compione Fault, which strikes ~NW-SE and dips to the SW (Fig. 3C). The two different lithologies

304 observed determine different deformation patterns in the footwall and in the hanging wall,
305 respectively (Figs. 4A-C). In the footwall, Macigno Sandstones Formation strata dip 5° - 10° towards
306 the NE in the footwall far from the fault zone. Approaching the footwall damage zone, bedding is
307 progressively tilted towards the SW to become horizontal at around 0.7 km distance from the fault
308 core, then SW-dipping of $\sim 40^{\circ}$ at around 0.5 km, and up to 55° at the damage zone-fault core
309 transition. Similarly, the Ottone Flysch Formation in the hangingwall is characterized by strata that
310 dip $\sim 60^{\circ}$ SW in proximity of the fault core and become subhorizontal outside the hanging wall
311 damage zone (Figs. 3B and 3C). A major E-W striking, S dipping fault zone occurs to the ESE of
312 Compione village (Fig. 4C), abutting the Compione Fault, as well as subsidiary fault segments, to the
313 south (Fig. 3C). The Compione fault zone has a core of up to 50 meters wide, consisting of deformed
314 shear lenses of mostly Macigno Sandstones Formation, bounded by anastomosed shear bands of
315 comminuted, incoherent sand and gouge (Bernini and Lasagna, 1988). In the northwestern corner of
316 the map, the Compione Fault affects a Pleistocene conglomerate consisting of boulders and pebbles
317 of Macigno Sandstones Formation, named Iera Conglomerate (Fig. 3C, Bernini and Papani, 2002)

318

319 **4.1 Footwall Damage Zone**

320 The most abundant deformation structures in the footwall damage zone are conjugate shear vein pairs
321 inclined at $\sim 60^{\circ}$ to bedding in both dipping directions and with bisector lines perpendicular to bedding
322 regardless of bedding dip (Figs. 5A and 5B). Outside the damage zone, 900 meters to the fault core,
323 veins occur only associated to subsidiary extensional faults with metric offset. In the outer boundary
324 area, where strata are subhorizontal, vein orientation is $N306^{\circ}$, 60° for the antithetic set and $\sim N120^{\circ}$ -
325 $N140^{\circ}$, 60° - 80° for the synthetic (Figs. 4A and 5B). Conjugate shear vein arrays are passively rotated
326 with bedding at decreasing distance from the fault core (Figs. 5C and 5D). Such a rotation and the
327 activity of late subsidiary fault zones causes significant dispersion of vein strike, which varies from
328 NNW-SSE to E-W. This is evident when shear vein data collected in subsidiary fault-bounded blocks

329 to the north and south of the major E-W striking fault zone are rotated to restore local bedding attitude
330 to horizontal (Fig. 6). The dip direction in non-rotated data varies from north- to eastward (Fig. 6A).
331 Rotation systematically produces conjugate arrays striking either NW-SE or E-W, respectively (Fig.
332 6B). In addition to fault-parallel conjugate shear vein arrays, cross-fault veins striking almost
333 perpendicular to the Compione Fault are also abundant (Fig. 4).
334 Most fault-related veins are reactivated as strike-slip (cross-fault veins) or conjugate extensional
335 faults (fault-parallel veins; Fig. 5E). Strike-slip faulting in the area is related both to lateral
336 propagation of the Compione fault zone and to the far-field stress of the left-lateral Secchia transversal
337 line, 4 km to the E of the study area. These fault-related shear veins and faults (extensional and strike-
338 slip) are well developed especially in coarse strata, have offsets ranging from centimetres to meters,
339 and are characterized by millimetre to centimetres slickenfibers. Strike-slip and extensional faults
340 mutually crosscut and their slip surfaces frequently show evidence for variable directions of
341 movement (Fig. 5F). In sites 4 and 5, conjugate extensional faults have a more complex pattern that
342 includes two synthetic trends, oriented 255° , 70° and 290° , 56° and two antithetic ones, oriented 350° ,
343 10° and 240° , 30° , respectively. It is important to note that, after bedding restoration to the horizontal,
344 the synthetic sets are 127° , 75° and 75° , 77° and the antithetic sets are 308° , 60° and 257° , 74° ,
345 respectively, and that ~E-W striking conjugate veins show re-activation as strike-slip to oblique-slip
346 subsidiary faults. Reverse kinematics are apparent, indicating that normal to oblique faulting, with
347 top-to-the-WNW shearing, was ongoing during passive bedding rotation in the damage zone sector
348 comprised between the Compione fault core and major E-W footwall splay fault (Fig. 6B). As a result,
349 S and SW dipping extensional faults in Fig. 6B were re-activated antithetically as soon as they
350 attained a NE dipping attitude due to bedding rotation. Late-stage faults strike from $N110^{\circ}$ to $N130^{\circ}$,
351 dip $60-80^{\circ}$ either to the NE or SW (Fig. 4) and have cores made of cataclastic loose sand and clay
352 smears, bounded by thin slip zones including extremely comminuted material. These late-stage faults
353 occur up to 400 meters from the fault core.

355 **4.2 Hanging Wall Damage Zone**

356 Subsidiary fault zones mainly arranged in synthetic anastomosed arrays are abundant in the hanging
357 wall damage zone near the fault core, isolating extensional shear lenses affected by intense pressure
358 solution as indicated by abundant shallow-dipping stylolites (Fig. 4). Moving away from the fault
359 core, both synthetic and antithetic subsidiary faults and fault zones occur, frequently with a ramp-flat
360 geometry. Here very thick beds are cut at high angle and the marly and clayish interlayers are
361 exploited as flat segments, (e.g. site 2 in Fig. 4) producing significant block tilting about horizontal
362 axes. Eventually, at about 300-400 m away from the fault core of the Compione Fault, the deformation
363 pattern is dominated by high-angle antithetic faults and fault zones (e.g. site 1 in Fig. 4). Overall,
364 subsidiary faults and fault zones in the hanging wall strike NW-SE, i.e. parallel to the master fault
365 strike, with a subordinate population striking ~E-W and dipping S (e.g. site 2 in Fig. 4). Abundant
366 veins nearly perpendicular to bedding occur in the hanging wall and crosscut pre-existing bedding-
367 parallel veins. Both populations grew by crack-seal opening (cf. Ramsay, 1980) before the onset of
368 extensional fault-related deformation, during thrusting and stacking of the Ligurian successions.
369 Subvertical veins are passively rotated in the hanging wall blocks of subsidiary fault zones and are
370 commonly re-activated as low-displacement shear surfaces showing slickensides coherent with
371 extensional, almost dip-slip, kinematics.

372

373 **5. VEIN CEMENT ANALYSIS**

374 **5.1 Petrographic Description**

375 Fault-related veins in the footwall damage zone have an aperture varying from 2 to 27 mm and show
376 evidence for antitaxial multiple opening (cf. Durney and Ramsay, 1973; Ramsay and Huber, 1983;
377 Passchier and Trouw, 2005; Bons et al., 2012). Vein cement consists of prehnite, quartz and three
378 different calcite generations, labelled as MC1A, MC1B and MC2, respectively (Fig. 7A). Prehnite
379 crystals are generally euhedral to subhedral and their dimension can reach 1500 µm parallel to the c

axis and 750 μm orthogonal to it. Prehnite was the first mineral phase to crystallize in columnar-radiating structures that are overgrown by or inter-grown with euhedral to subhedral quartz (Figs. 7A and 7B). When analysed in cold CL, prehnite luminescence color switches abruptly from “olive green” to “lime green”, from the cores towards the rims (Fig. 7B; Huber et al., 2007). Compositional analyses were carried out by SEM-EDS on three prehnite crystals sampled from site 5. Ten areas were analysed both along the long and the short axes of the crystals to investigate compositional variations during crystallization. Sixty spectra were acquired and eight of them did evidence calcite contamination, which is often found as isomorphous replacement of prehnite (Figs. 7C and 7D). Analyses results did not evidence any trend in amounts of Fe^{3+} - Al^{3+} along crystal axes from the cores to the rims of crystals. Al_2O_3 ranges between 24.5 and 25.5 wt%, with a mean value of 25.07 ± 0.19 wt%. The mean abundance of SiO_2 is 45.04 ± 0.20 wt%, with values ranging between 44.6 and 45.5 wt%. CaO is comprised between 27.3 and 29 wt% and has a mean value of 27.76 ± 0.23 , in agreement with the prehnite formula $\text{Ca}_2\text{Al}_2\text{Si}_3\text{O}_{10}(\text{OH})_2$. FeO was detected in twenty-four spectra and was always below 1 wt%. Quartz sometimes displays bridge structures, growth competition textures and dissolution embayments (Fig. 7C; Hilgers and Urai, 2002; Bons, 2001; Okamoto and Sekine, 2011). MC1 calcite grew both as isomorphous replacement of prehnite crystals and in the remaining open sites, mostly at the centre of fractures. The latter are characterized by poikilotopic rhomboedric crystals up to 1 cm in dimension with abundant Type I and Type II twinning (cf. Burkhard, 1993; Ferrill et al., 2004b) and sweeping extinction (Figs. 7A, 7C, and 7G). Stained MC1 is pink, (Fig. 7E) indicating that calcite is non-ferroan (Dickson, 1966). MC1 calcite crystals can be subdivided in MC1A and MC1B: B is mostly characterized by twinning Type I and a “clearer texture” compared to A, which displays abundant twinning Type II. Moreover, MC1A has dark red color in CL while MC1B is red to orange (Fig. 7B and 7D). Multiple antitaxial opening events (cf. Holland and Urai, 2010; Virgo et al., 2014) caused the cyclic repetition of prehnite + quartz + MC1 calcites bands infilling the veins (Figs. 7E-G). Shearing of veins caused quartz dynamic recrystallization and formation of straight, micrometric wide mirror surfaces, which truncate vein crystals (Figs. 7E, 7F

406 and 7G). MC2 is the last cement and is composed of microsparitic to blocky calcite up to maximum
407 500 μm crystal size, sometimes displaying Type I twinning (cf. Burkhard, 1993). MC2 calcite has
408 bright orange to yellow colors under CL and is hosted in sub-millimetric fracture networks that exploit
409 cleavage surfaces orthogonal and parallel to the c axis of prehnite (Fig. 7A), twinning planes in MC1
410 calcite and re-opened shear surfaces. Even MC2 calcite, in places, occurs as prehnite isomorphous
411 replacement (Figs. 7H). Veins nearest to the fault core are more intensely affected by shearing, quartz
412 recrystallization, isomorphous replacement of prehnite by MC1 crystals and by MC2 calcite
413 cementation (Figs. 7G and 7I). In the fault core, in fact, the host rock is disaggregated and cemented
414 by MC2 calcite, which displays an interparticle cement texture (Fig. 7J). Footwall fault-related veins
415 in silt and clay beds have MC1 and MC2 calcite cements (labelled as Macigno host rock calcite
416 cement), but they show a microsparitic texture, whereas prehnite and quartz are absent.

417 Hanging wall veins, hosted in the Ottone Flysch Formation, are composed of two generations of
418 calcite, labelled as OC1 and OC2, and contain traces of quartz. OC1 calcite veins are considered for
419 comparative purposes only because they formed before the onset of extensional deformation in the
420 studied area. OC2 calcite veins formed by multiple antitaxial fracturing-precipitation events (cf.
421 Holland and Urai, 2010) and crosscut or run parallel to OC1 calcite (Figs. 7K and 7L). OC2 calcites
422 have a blocky texture with a “clear” appearance and dimensions up to 1 mm (Fig. 7K), showing rarely
423 Type I twinning (cf. Burkhard, 1993). Their color under CL is red, slightly brighter than OC1 calcites,
424 but it changes depending on structural position (Fig. 7L). In particular, samples collected less than 50
425 meters from the fault core have OC2 calcites with the same dull luminescence as the OC1 calcites.
426 Quartz crystals smaller than 100 μm sometimes occur in association with OC2 calcites (Figs. 7M and
427 7N). A crack-seal bedding parallel vein in the hanging wall damage zone, in proximity of the fault
428 core, shows quartz associated with OC2 calcite, which preserve the prismatic habit of prehnite,
429 similarly to the footwall damage zone cement textures (Fig. 7N). Moreover, close to the fault core,

430 OC2 calcite cement shows a proto-breccia texture in extensional S-C (S—schistosity, C—
431 cisaillement [French for shear]) lithons (Fig. 7O).

432

433 **5.2 Stable Isotopes**

434 Results of carbon and oxygen stable isotopes analyses are summarized in Table 1. MC1 calcites have
435 the most depleted $\delta^{18}\text{O}$ values, from +11.8‰ to +14.0 ‰ V-SMOW while MC2 calcites range from
436 +13.4‰ to +17.9‰ V-SMOW. Mean $\delta^{18}\text{O}$ is +13.1‰ and +15.8‰ V-SMOW for MC1 and MC2
437 calcites respectively (Fig. 8A). $\delta^{13}\text{C}$ in footwall MC1 and MC2 calcites ranges from -5.8‰ to -0.9‰
438 and from -7.1‰ to -0.4‰ V-PDB respectively. Plotting $\delta^{18}\text{O}$ vs. distance from the Compione fault
439 core for MC1 and MC2 calcites, a slight trend appears (Fig. 8B). In particular, $\delta^{18}\text{O}$ values become
440 enriched approaching the fault core. The same plot for $\delta^{13}\text{C}$ shows that MC1 and MC2 calcites are
441 characterized by different values depending on structural position (Fig. 8C). Site 5 is located in a
442 different footwall block, compared to other structural sites, and shows depleted $\delta^{13}\text{C}$, ranging from -
443 7.1‰ to -3.8‰ V-PDB, compared to other sampling sites whose $\delta^{13}\text{C}$ is comprised between -2.8‰
444 and -0.4‰ V-PDB (Fig. 8C). MC1 and MC2 calcites from a sample collected near the NW tip of the
445 Compione Fault (Site 13) have $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values that are enriched compared to the bulk of the
446 samples that are located along the studied cross-section of the fault zone (Fig. 8A). The footwall host
447 rock carbonate component is composed of microsparitic calcite crystals, present in silt and clay-sized
448 intervals. It shows $\delta^{18}\text{O}$ values between +13.9‰ and +16.0‰ V-SMOW and $\delta^{13}\text{C}$ from -3.2‰ to -
449 0.7‰ V-PDB.

450 Hanging wall OC1 and OC2 calcites present enriched and more clustered $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values
451 compared to footwall MC1 and MC2 calcite veins. $\delta^{13}\text{C}$ values are comparable for hanging wall host
452 rock and both the hanging wall OC1 and OC2 calcite generations. $\delta^{13}\text{C}$ in hanging wall carbonates
453 ranges mostly between +1.2‰ and +2.4‰ V-PDB, excluding outliers. Mean $\delta^{13}\text{C}$ is +1.9‰ V-PDB

454 for hanging wall host rock and OC1 calcites, and +1.8‰ V-PDB for OC2 calcites. $\delta^{18}\text{O}$ values,
455 however, display large variability between hanging wall host rock, OC1, and OC2 calcites, defining
456 a clear horizontally elongated trend in Fig. 8A. Host rock $\delta^{18}\text{O}$ is between +26.9‰ and +27.5‰ V-
457 SMOW; OC1 calcites $\delta^{18}\text{O}$ are similar to the host rock, ranging from +23.5‰ to +26.9‰ V-SMOW,
458 while OC2 calcites show depleted $\delta^{18}\text{O}$ values, comprised between +13.1‰ and +19.7‰ V-SMOW
459 (Fig. 8A). Mean $\delta^{18}\text{O}$ are +25.7‰ and +16.5‰ V-SMOW for OC1 and OC2 calcites, respectively.
460 The OC2 outlier has a $\delta^{13}\text{C}$ value of -0.46‰ V-PDB, comparable to footwall fault-related veins and
461 corresponds to the replacive calcite, associated with quartz, shown in Fig. 7N.

462

463 5.3 Fluid Inclusions

464 Microthermometry measurements were done on 11 footwall and on 6 hanging wall veins and their
465 results are summarized in Table 2. Quartz in footwall veins shows abundant and clearly visible trails
466 of decrepitated and leaked inclusions. Two types of inclusions have been recognized: biphasic aqueous
467 (Q1) and monophasic gaseous (Q2). Both of them have a maximum length ranging from 5 to 15 μm
468 and rounded shapes (Figs. 9A and 9C). Only one Q1 primary FIA and two isolated inclusions were
469 large enough to be measured. Vapour to liquid phase ratios are constant and the vapour bubble is less
470 than 15% of the total inclusion volume. Homogenization, always into the liquid phase, occurs
471 between 127 °C and 215 °C (Fig. 10B) with a mean value of 157 °C. First melting temperatures are
472 in the range of -50 °C to -45 °C, suggesting a NaCl-CaCl₂-H₂O system (Roedder, 1984), and are
473 followed by the hydrohalite melting temperature between -32 °C and -28 °C. Ice melting temperatures
474 range from -4.8 °C to -11.4 °C (Fig. 10C) with a mean value of -7.2 °C, which corresponds to salinities
475 of 7.6, 15.4 and 10.7 wt% NaCl eq., respectively (Bodnar, 1993). Q2 monophasic gaseous inclusions
476 have dimensions similar to Q1 inclusions and a rounded shape. They were measured in three FIAs.
477 Homogenization was into the vapour phase at temperatures ranging from -89 °C to -83 °C (Fig. 10A).
478 Mean homogenization temperature is -88 °C. No solid was formed in these gaseous inclusions. Q2

479 inclusions were analysed through micro-Raman spectroscopy to check their composition. Results of
480 Raman analysis are briefly illustrated in Fig. 9B and show a peak at 2914 cm^{-1} , which corresponds to
481 the Raman shift of CH_4 . Homogenization temperatures, in agreement with Raman results, indicate
482 that Q2 are monophasic CH_4 inclusions.

483 Fluid inclusions in footwall MC1A calcite are biphasic aqueous, with a size from $3\text{ }\mu\text{m}$ to $10\text{--}15\text{ }\mu\text{m}$.
484 They have a negative crystal shape and constant vapour/liquid ratio (Fig. 9D). The vapour phase fills
485 10 to 15% of the inclusion's volume. Inclusions in MC1A calcite are organized in FIAs and in fluid
486 inclusion (FI) trails that follow twinning planes. FI trails and inclusions near twinning planes were
487 not taken into account during microthermometric measurements. Some inclusions leaked at their
488 homogenization temperatures and some bubbles did not reappear upon cooling. These inclusions were
489 also discarded. A total amount of twenty-two MC1A inclusions were measured, distributed in six
490 FIAs and three isolated inclusions. MC1A inclusions show homogenization temperatures ranging
491 from $178\text{ }^{\circ}\text{C}$ to $198\text{ }^{\circ}\text{C}$. The mean T_{tot} is $189\text{ }^{\circ}\text{C}$ and the modal peak is between 190 and $195\text{ }^{\circ}\text{C}$ (Fig.
492 10B). First melting in MC1A inclusions occurred at temperatures between $-17\text{ }^{\circ}\text{C}$ and $-21\text{ }^{\circ}\text{C}$,
493 indicating a $\text{H}_2\text{O}\text{--NaCl}$ composition. The MC1A ice melting temperatures range from $-0.8\text{ }^{\circ}\text{C}$ to -6.2
494 $^{\circ}\text{C}$, corresponding to salinities between 1.4 and 9.3 wt% NaCl eq. Mean T_{mice} is $-2.7\text{ }^{\circ}\text{C}$, which
495 corresponds to 4.5 wt% NaCl eq. (Bodnar, 1993; Fig. 10C).

496 Fluid inclusions hosted in impurity-poor MC1B calcite crystals are slightly smaller than MC1A
497 inclusions, ranging from 2 to $10\text{ }\mu\text{m}$. They have rounded to negative crystal shape and constant
498 vapour/liquid ratio where vapour fills around 10% of the inclusion's volume. Nineteen MC1B
499 inclusions were measured, distributed in five FIAs. They show homogenization temperatures between
500 $140\text{ }^{\circ}\text{C}$ and $161\text{ }^{\circ}\text{C}$ with a mean value of $151\text{ }^{\circ}\text{C}$ and a modal peak at $150\text{--}155\text{ }^{\circ}\text{C}$ (Fig. 10B). First
501 melting temperatures of MC1B inclusions range between $-8\text{ }^{\circ}\text{C}$ and $-20\text{ }^{\circ}\text{C}$ in three FIAs while they
502 are $-31.5\text{ }^{\circ}\text{C}$ in another FIA. Ice melting temperatures for MC1B inclusions range between $-0.8\text{ }^{\circ}\text{C}$
503 and $-17.2\text{ }^{\circ}\text{C}$, which corresponds to salinities ranging from 1.4 to 20.3 wt% NaCl eq. The mean T_{mice}

504 is -5 °C, corresponding to 7.9 wt% NaCl eq. (Bodnar, 1993; Fig. 10C). Only the four MC1B
505 inclusions belonging to the FIA characterized by first melting temperatures of -31.5 °C showed T_{mice}
506 lower than -6.2 °C, i.e. ranging from -10.1 °C to -17.7 °C.

507 Footwall MC2 calcite contains biphasic aqueous inclusions with the same petrographic characteristics
508 as MC1 inclusions but they were not found aligned in FI trails. Six MC2 FIAs and two isolated
509 inclusions were measured, representing 20 inclusions. They have rounded shapes and dimensions
510 ranging from 2-3 up to 10 µm. Homogenization temperatures range from 69 °C to 115 °C with a
511 mean value of 88 °C (Fig. 10B). Ice melting temperatures range from -0.7 °C to -20.7 °C,
512 corresponding to salinities from 1.2 to 22.9 wt% NaCl eq. The mean T_{mice} is -8.7 °C and the mean
513 salinity 12.5 wt% NaCl eq. (Bodnar, 1993) even if the distribution is formed by two clusters, with
514 T_{mice} comprised between -16.7 °C and -20.7 °C, and -0.7 °C and -5.8 °C, respectively. The modal
515 peak is comprised between -1 °C and -2 °C (Fig. 10C). MC2 inclusions belonging to the first cluster
516 froze at temperatures lower than -70 °C and showed melting evidence at temperatures comprised
517 between -34 °C and -27 °C as Q1 inclusions. The first melting temperatures were difficult to evaluate
518 but their similar behaviour with Q1 inclusions (this paragraph) suggests a NaCl-CaCl₂-H₂O system.
519 Only a first melting temperature of -17.8 °C, indicating a NaCl-H₂O system, was measurable in MC2
520 inclusions belonging to the second cluster.

521 Hanging wall OC2 calcite is characterized by fluid inclusions that are monophasic and biphasic
522 aqueous with rounded shapes and smaller than 10 µm (Fig. 9F). The vapour phase in biphasic
523 inclusions fills 5 to 10% of their volume. Small monophasic OC2 inclusions did not nucleate bubbles
524 after two weeks at -20 °C. Two FIAs and an isolated inclusion for a total of sixteen were measured.
525 Homogenization temperatures range from 60 °C to 75 °C in the first FIA and from 110 °C to 115 °C
526 in the second FIA (Fig. 10D). First melting temperature was never observed. Ice melting temperatures
527 range from -1.3 °C to -6.9 °C, with a mean value of -4.4 °C (Fig. 10E) and they correspond

528 respectively to salinities of 2.2, 10.4 and 7 wt% NaCl eq. (Bodnar, 1993). OC2 inclusions have higher
529 salinities at lower homogenization temperatures.

530 Homogenization vs. ice melting temperatures of measured inclusions are plotted in Fig. 11. Fluid
531 inclusions from quartz in footwall veins show a rather constant salinity, around 10-15 wt% NaCl eq.,
532 at different homogenization temperatures. Calcites MC1A, MC1B and MC2 in footwall veins are
533 characterized by a different temperature-salinity trend, generally displaying less than 5 wt% NaCl eq.
534 at 180-200 °C (subgroup MC1A) and an increasingly wider range of salinities at decreasing
535 temperatures. Subgroup MC1B with homogenization temperatures between 140-160 °C, can have
536 salinities up to 20 wt% NaCl eq. At lower temperatures, between 70 and 110 °C, MC2 inclusions
537 have salinities up to 23 wt% NaCl eq. The data belonging to hanging wall fluid inclusions OC2 show
538 a trend similar to footwall calcites, characterized by higher salinities at lower homogenization
539 temperatures.

540

541 **6. VITRINITE REFLECTANCE AND THERMAL MODELLING**

542 Results of vitrinite reflectance measurements acquired in footwall Macigno Sandstones Formation
543 are summarized in Table 3. Ten samples were analyzed from sites 14, 4 and 5. Samples from sites 4-
544 5 are located in the footwall damage zone while site 14 is located outside the damage zone. Mean
545 values range between 0.42 and 0.70 Ro% and standard deviations are generally below 0.1 Ro%. More
546 in detail, vitrinite reflectance measured on samples in the footwall damage zone (sites 4 and 5) show
547 very dispersed average values ranging from 0.42 to 0.61 Ro% (Table 3), which were based on the in
548 situ vitrinite. The lowest values were measured on very small and fractured fragments with dark
549 oxidized rims along the irregular micro-fissure in the vitrinite particle, making the reflectance
550 assessment scarcely reliable, while only few sufficiently large and unfractured fragments were found
551 showing generally slightly higher values around 0.65-0.70 Ro%. A further family of vitrinite

552 fragments, excluded from the aforementioned average, was found showing values between 1.0 and
553 1.30 %. These fragments were interpreted as reworked vitrinite.

554 On the other hand, samples from site 14, far from the damage zone, indicate average values between
555 0.66 and 0.70 Ro% (Table 3). In situ vitrinite is here represented by large and unfractured fragments
556 even if averages are based on a small number of measurements (between 8 and 19). Also in these
557 samples reworked vitrinite showing higher values (between 1.10 and 1.80 Ro%) was found.
558 Nevertheless, the lowest mean values between 0.42 and 0.61 Ro% are found systematically in the
559 footwall damage zone. These values are always associated with oxidized and pervasively fractured
560 vitrinite fragments and thus were not considered in the calibration of the thermal model. Samples in
561 the footwall sandstones outside the Compione fault damage zone, on the other hand, show consistent
562 Ro% values between 0.66 and 0.70.

563 Assumptions about the burial/exhumation history and heat flow in the Macigno Sandstones
564 Formation have to be made, as they are input data necessary for the thermal modelling: (1)
565 stratigraphic location of the samples is comprised between 600 m and 1100 m from the top of the
566 Macigno Sandstones Formation, which is 2300 m thick (Ghibaudo, 1980). In detail, samples from
567 site 14 are located between 600-800 m and samples from sites 4-5 between 900-1100 m; (2) the
568 Macigno Sandstones Formation was rapidly buried below the 4000-6000-m-thick allochthonous
569 Ligurian and Subligurian units (Carlini et al., 2013); (3) syncontractional exhumation took place up
570 to Late Messinian times through low angle normal faulting in the inner part of the orogenic wedge
571 (Fellin et al., 2007; Carlini et al., 2013; Molli et al., 2018). Moreover, heat flow in the Northern
572 Apennines foredeep has values lower than 30°C/km due to rapid burial and thrusting of foredeep
573 units, whose thermal regime is far from equilibrium (Mongelli et al., 1991; Della Vedova et al., 1995).

574 Ro% values at maximum burial depth could only be fitted using 3000 m of Ligurian and Subligurian
575 units overburden, lower than the minimum thickness estimates of allochthonous units (according to
576 Carlini et al., 2013) and adopting a geothermal gradient of 25-30°C/Km. Thinning of the overburden

577 thickness moving towards the foreland is logical considering the wedge-shaped overall geometry of
578 fold and thrust belts (Davis et al., 1983). The best fit between calculated maturity profile and
579 measured vitrinite reflectance was attained using a geothermal gradient of 25°C/Km up to the end of
580 Miocene that gradually increased up to a present-day value of 30°C/Km. As a consequence, the
581 Macigno Sandstones Formation stratigraphic sector, hosted in the Compione fault footwall damage
582 zone, at 5 km depth, experienced peak temperatures between 140-150°C (Fig. 12).

583

584 **7. DISCUSSION**

585 **7.1 Process Zone Width**

586 An outstanding feature of the Compione Fault is the abundance in the footwall damage zone of veins
587 that, when bedding is rotated to the horizontal, restore to a pattern of conjugate shear fractures with
588 a vertical bisector, i.e. the typical geometry produced in an extensional tectonic regime (Anderson,
589 1951; Sibson, 1996). This evidence supports vein formation in an early evolutionary stage of the
590 Compione Fault, when bedding in the process zone was still horizontal, before extensional folding
591 and shear localization. Accordingly, the area affected by such extensional shear veins can be
592 interpreted as the process zone sector preserved in the footwall of the Compione Fault, which formed
593 ahead of the upward-propagating master shear zone (cf. Lockner et al., 1992; Reches and Lockner,
594 1994; Cowie and Shipton, 1998; Vermilye and Scholz, 1998). In the studied across-fault transect, the
595 width of the footwall process zone is about 900 m. The present-day tectonic juxtaposition of the
596 Ottone Flysch Formation in the hanging wall prevents any possibility to quantify the total width of
597 the process zone in the Macigno Sandstones Formation. However, by assuming that the Upper
598 Triassic Burano Evaporites provided a thick plastic layer suitable to enhance initial extensional
599 folding in the overburden (Schlische, 1995; Withjack and Callaway, 2000), it is possible to
600 hypothesize that the tip of the master shear zone was temporarily arrested at the base of the Burano
601 Evaporites while their top, now at around 6 to 7 km depth, provided the apical point of the extensional

602 fault-propagation fold (Fig. 13). By applying the geometric construction of Jin and Groshong (2006)
603 and assuming a linear velocity field in the trishear zone ahead of the propagating upper fault tip
604 (Hardy and Ford, 1997) it is possible to infer a total process zone width of about 1800 m and an apical
605 angle of $\sim 10^\circ$ (Fig. 13). Taking into account that the total displacement of the Compione Fault is
606 around 1.5 km (Bernini and Lasagna, 1988), our estimate of the process zone width is out-of-scale
607 compared to displacement (D) to damage zone (DZ) ratios reported in the literature, even though
608 D/DZ is strongly dependent on the criteria used to define damage zone thickness and on lithological
609 properties (Knott et al., 1996; Fossen et al., 2007; Childs et al., 2009; Fossen, 2010; Torabi and Berg,
610 2011; Solum and Huisman, 2016). In fact, assuming that late-stage extensional fault zones are the
611 structural elements that define the damage zone width, then the D/DZ ratio of the Compione Fault
612 would be comparable to those reported in published datasets.

613

614 **7.2 Cyclical Vein Development and Earthquake Cycle**

615 Footwall damage zone fault-related veins show multiple subparallel fracturing-sealing events
616 indicating that fracturing, cementation, dissolution and shearing were cyclic (Figs. 14A-C; Ramsay,
617 1980; Boullier and Robert, 1992; Boullier et al., 1994; Sibson, 1996; Renard et al., 2000; Sibson,
618 2004). The majority of veins is intersected by localized shear-reactivation, forming abundant mirror
619 surfaces and, in cases displaying straight micrometric wide slip surfaces truncating crystals, which
620 are interpreted as evidence for coseismic slip (Smith et al., 2011; Fondriest et al., 2013; Smeraglia et
621 al., 2017). Fracturing-sealing cycles caused strain-softening promoting localization of younger
622 fracturing and shearing events at vein-host rock interfaces or in between different openings (Jessell
623 et al., 1994; Virgo et al., 2014). Prehnite exhibits euhedral crystals, organized in columnar-radiating
624 aggregates while quartz shows euhedral to subhedral crystals, in cases also displaying growth
625 competition (Fig. 14A). Both textures require fractures to remain fluid-filled and open during crystal
626 growth (Fisher et al., 1995; Koehn and Passchier, 2000; Oliver and Bons, 2001; Bons et al., 2012),

627 thus implying fluid pressures higher than the local σ_3 for fault-parallel extensional veins. Conversely,
628 blocky rhombohedral calcite texture (Figs. 14A-C), which occludes completely the remaining
629 fracture space, could be caused by different processes: a) supersaturation in response to a pressure
630 drop; b) texture obliteration due to repeated fracturing; c) fast crystal nucleation caused by a sudden
631 arrest of an ascending fluid (Oliver and Bons, 2001). Moreover, prehnite dissolution can promote
632 permeability enhancement and, consequently, pressure reduction (Figs. 14B and 14C; Boullier et al.,
633 1994). In the literature models such as the fault-valve and seismic pumping have been proposed to
634 relate cyclical fracturing-sealing events to the seismic cycle (Sibson et al., 1975, 1988; McCaig, 1988;
635 Boullier and Robert, 1992; Robert et al., 1995; Cox, 1995, 1999).

636 Our microstructural data support a model of extensional faulting in the upper crust, triggered by
637 shortening and thrusting in a seismically active metamorphic basement (McCaig, 1988). At shallow
638 crustal levels, as in this specific case, fluid pressure is governed by seismic-pumping (Sibson et al.,
639 1975), while the fault-valve mechanism explains supralithostatic pressures at deeper crustal levels
640 (Sibson et al., 1988). In the hypothesis that precipitation of the described prehnite-quartz-calcite
641 assemblages was triggered by seismic activity during upward fault propagation from the basement,
642 silicates (prehnite and quartz) may have crystallized at suprahydrostatic fluid pressure conditions (P_f)
643 after seismically-induced fracturing, i.e. in the post-seismic stages (Figs. 14D and 14E).
644 Suprahydrostatic P_f in the extensional process zone may have been generated by ascending fluids that
645 breached a low-permeability layer at depth (Sibson et al., 1988; McCaig, 1988), reasonably provided
646 by the thick evaporitic sequences at the top of the metamorphic basement, which is deformed in a
647 regional scale antiformal stack structure (e.g. Molli et al., 2018). The decrease of P_f to hydrostatic
648 values led to supersaturation of calcite, which precipitated in the remaining voids thus completing
649 vein infilling and favouring a new cycle of pore fluid pressure increase (Figs. 14D and 14E). The
650 causal link between seismic activity and precipitation of the mineralogical assemblage in rotated shear
651 veins exposed in the footwall damage zone of the Compione Fault is tentatively proposed as a working

652 hypothesis that deserves further studies specifically designed for acquiring a comprehensive dataset
653 suitable to either support or reject this possibility.

654

655 **7.3 Process Zone Temperature Anomaly**

656 Paleothermal data obtained from the host Macigno Sandstones Formation and from the shear vein
657 network allow estimating the thermal disequilibrium associated with the upward migrating fluids that
658 infiltrated the process zone in the early stages of faulting and related extensional folding.

659 An anomalous feature is the maturity difference between sites 4 and 5 with respect to site 14, which
660 cannot be explained by different burial since low maturity samples have a lower stratigraphic position.
661 Actually, lower Ro% values from sites 4 and 5 locate into the footwall damage zone and were
662 obtained from very small and fractured fragments with oxidized rims around fractures. This suggests
663 that anomalously low reflectance values are due to oxidation from mixed meteoric and deep fluid
664 weathering occurring probably during the last pulse of fluid circulation around the fault. The increase
665 in permeability during the last stage of the fault's activity created conditions that favoured oxidation
666 of the surface of the organic matter which is subsequently degraded during weathering (Petsch et al.,
667 2000).

668 Accordingly, the thermal model was calibrated using data from site 14. Thermal modelling of vitrinite
669 reflectance data provided peak temperatures in the footwall damage zone ranging between 140-150
670 °C at maximum depths of about 5 km and geothermal gradients between 25-30°C/km (Fig. 12). In
671 addition, published data from apatite fission tracks in the Macigno Sandstones Formation sampled in
672 the study area show complete annealing (Thomson et al., 2010; Carlini et al., 2013), which generally
673 indicates temperatures higher than 110 °C (Ketcham et al., 1999). This supports results from our
674 model.

675 Microthermometric data from quartz Q1 and calcite MC1A cements show homogenization
676 temperatures of 155 °C and 180 °C, respectively. Accordingly, MC1A calcite in the footwall damage
677 zone fracture network shows homogenization temperatures at least 30 °C higher than the surrounding
678 Macigno Sandstones Formation. If we correct data by pressure, assuming that MC1A crystallized
679 crystallized at about 5 km depth in hydrostatic conditions at the onset of exhumation, real fluid
680 trapping temperatures are predicted to be around 210-230 °C (Fig. 15A). Moreover, prehnite in the
681 studied veins, despite the impossibility to provide direct constraints on paleotemperatures may
682 suggest minimum temperature values of about 230 °C based on occurrences in hydrothermal and in
683 geothermal systems in Tuscany, Italy, and other areas showing calc-silicate mineralization (Browne,
684 1978; Arnason et al., 1994;). It is worth noting that uncertainty is associated with this inference
685 because crystallization conditions might depend on fluid chemistry. Hydrothermalism is abundant
686 and ongoing on the Tyrrhenian Sea side of the Apennines and is related to high-temperature-low-
687 pressure contact metamorphism due to the intrusion of igneous bodies into the crust associated with
688 the Tyrrhenian extension (Cavarretta et al., 1982; Boccaletti et al., 1997; Gianelli et al., 1997; Dini et
689 al., 2005; Boiron et al., 2007 and many others). If our assumptions are correct, then the difference in
690 temperature between hydrothermal fluids that circulated through the fracture network in the process
691 zone and the surrounding host Macigno Sandstones Formation was between 60 and 90 °C. Such a
692 high thermal disequilibrium is supported both by the geometry of the process zone, which displays a
693 low apical angle (Fig. 13), and by supra-hydrostatic fluid pressures that promote fast advection of hot
694 fluids from the basement in a channelized, highly-fractured and narrow deformation zone (Sibson et
695 al., 1988, 1996, 2000; Renard et al., 2000; Gratier et al., 2002; Beaudoin et al., 2011). This result
696 highlights the importance of combining different methodologies to constrain host rock and fault-
697 related fluid paleotemperatures (Mamadou et al., 2016; Honlet et al., 2017; Laurent et al., 2017;
698 Wustefeld et al., 2017).

700 **7.4 Fluid Sources and Migration Pathways**

701 Assuming that Q1 and Q2 inclusions are cogenetic, this would imply that the source fluid underwent
702 fluid immiscibility before entrapment. Cogenetic aqueous biphasic and CH₄ gaseous monophasic
703 inclusions have already been documented in authigenic quartz in sandstones of the Northern
704 Apennines (Mullis, 1979, 1987, 1988; Montomoli et al., 2001; Montomoli, 2002; Mazzarini et al.,
705 2010). This fluid immiscibility is generally caused by decreasing fluid pressure during upward fluid
706 migration in the fault damage zone, (Parry and Bruhn, 1987, 1990; Sibson et al., 1975, 1988, 2000).
707 MC1B inclusions show the same composition and salinity as MC1A inclusions except for a FIA in
708 Site 5, with higher salinity and lower first melting temperatures, indicating a NaCl-CaCl₂-H₂O
709 composition as Q1 inclusions.

710 Assuming equilibrium precipitation, the calculated $\delta^{18}\text{O}_{\text{fluid}}$ for MC1 calcites is shown in Fig. 15B
711 (Friedman and O'Neil, 1977), and is between +2‰ and +4‰ V-SMOW for MC1A calcite and
712 between 0‰ and +2‰ V-SMOW for MC1B calcite indicating ¹⁸O enriched waters due to different
713 degrees of water-rock interaction (e.g. Muchez et al., 1995). Taking into account the different
714 lithologies cut by the Compione Fault (Figs. 1C and 2), $\delta^{13}\text{C}$ values of MC1 calcites may indicate
715 different degrees of mixing between methane fluid originating from the thermal maturation of organic
716 matter inside the Macigno Sandstones Formation, inorganic marine carbon-rich fluids from the
717 underlying Mesozoic carbonates and, also, a contribution of metamorphic fluids coming from the
718 basement (cf. Hoefs, 1997; Milliken et al., 1998; Mazzarini et al., 2010; Boschetti et al., 2017). The
719 latter is supported by the occurrence of prehnite crystallization in stage 1, which is a typical mineral
720 of anchizone metamorphism (Merriman and Frey, 1999). Moreover, fluid trapped in MC1A
721 inclusions shows low-salinity and high temperature, which can be ascribed to devolatilization
722 reactions in the underlying metamorphic basement (Walther and Orville, 1982; Oliver, 1996;
723 Connolly, 2010; Ingebritsen and Manning, 2010), while locally higher salinities in MC1B reflect a
724 decreasing contribution of metamorphic fluids.

725 Hanging wall calcite OC2 is interpreted to be associated with stage 2 on the basis of structural,
726 microstructural and isotopic observations: a) OC2 cements in microfractures that display, locally,
727 quartz crystals along rims; b) OC2 in Fig. 7N occurs, as in the footwall, as isomorphous replacement
728 of prehnite; c) isotopic analysis results of OC2 calcite in the same sample shows a $\delta^{13}\text{C}$ shift towards
729 MC1 calcites values indicating local mixing between hanging wall and footwall fluids at stage 2 (Fig.
730 8). Isotopic results of OC2 calcite show that $\delta^{13}\text{C}$ values, except the outlier in Fig. 7N, are similar to
731 OC1 ones, indicating no external source of carbon in the Ottone Flysch Formation. $\delta^{18}\text{O}_{\text{fluid}}$ from
732 which OC2 precipitated is characterized by a wide variability (Fig. 15B), ranging from around -6‰
733 to 0‰ V-SMOW. This range could be interpreted as a fluid mix between Ottone Flysch formational
734 waters and meteoric waters ($\delta^{18}\text{O}$ around -8‰ to -6‰ V-SMOW, cf. Longinelli and Selmo, 2003;
735 Giustini et al., 2016) even this is difficult to prove since we do not have data on the composition of
736 the fluid.

737 MC2 calcite crystallized in micro-fractures in the internal footwall damage zone during stage 3. As
738 in stage 1 and 2, even MC2 locally replaces prehnite crystals. The wide range of homogenization
739 temperatures of MC2 inclusions could indicate they were stretched. We, therefore, assume the lower
740 temperatures ranging from 70 to 90 °C as representative for these inclusions. MC2 inclusions with
741 low-salinity (< 10 wt% NaCl eq.) have a NaCl-H₂O composition while those with high-salinity (up
742 to 22.9 wt% NaCl eq.) have a NaCl-CaCl₂-H₂O composition. The high salinity could indicate the
743 dissolution of salts in the subsurface (Goldstein and Reynolds, 1994; Boschetti et al., 2017). Carbon
744 forming MC2 calcites was derived from the same sources discussed for MC1 while $\delta^{18}\text{O}_{\text{Fluid}}$ values,
745 comprised between -6‰ and -2‰ V-SMOW, could indicate mixing between MC1 source fluid and
746 meteoric waters (Fig. 15B).

747 Phase diagrams of $\log(a\text{Ca}^{2+}/a^2\text{H}^+)$ vs. $\log a\text{SiO}_2(\text{aq})$ in Figs. 16A and 16B have been plotted for,
748 respectively, temperatures of 200 and 150 °C show the stability fields of the mineral assemblages in
749 stage 1 and stage 2. In this framework, at the beginning of stage 1, the upward migrating fault-related

750 fluid is initially recorded by crystallization of prehnite and quartz, which at 200 °C (Fig.16A) are
751 stable at lower $\log (a \text{ Ca}^{2+} / a^2 \text{ H}^+)$ and $\log a \text{ SiO}_2 (\text{aq})$ compared to 150 °C (Fig.16B; cf. Bird and
752 Helgeson, 1980, 1981; Cavarretta et al., 1982). Then it evolved, at lower temperatures, causing
753 prehnite crystals dissolution and further precipitation of quartz and calcite (MC1 A and B), both in
754 stages 1 and 2. Replacive calcite indicate a dissolution-reprecipitation process, lowering CO₂ content
755 and increasing H₂O, silica and alumina activity in the fluid. Quartz cementation is coeval with upward
756 methane migration which, in contrast, is not recorded in calcite. It can be inferred that quartz
757 crystallization was inhibited when methane was no longer in the fluid and was oxidized to CO₂,
758 necessary to precipitate calcite. Silicates precipitation from an upward migrating and cooling hot fluid
759 is easily explained by decreasing silica solubility at lower temperatures and pressures, while
760 precipitation of calcite, which solubility increases with decreasing temperature and pressure, is
761 promoted by decreasing $f\text{CO}_2$ (cf. Bird and Helgeson, 1980, 1981; Cavarretta et al., 1982).

762 MC1 and MC2 calcite cements in Site 5, located in the footwall block between the Compione fault
763 core and the E-W footwall splay, show lighter carbon values compared to other sites (Fig. 8C). This
764 $\delta^{13}\text{C}$ variation indicates that the sectors of the process zone were characterized by different quantities
765 of organic matter maturation-derived carbon during upward migration. Stable isotope values of MC1
766 and MC2 in the sample from Groppodalsio (Fault NW tip area, Site 13) are slightly enriched in ¹⁸O
767 (triangles in Fig. 8A) compared to Compione, while minimum temperatures of entrapment were the
768 same. Therefore, the $\delta^{18}\text{O}_{\text{Fluid}}$ composition in the NW were slightly heavier compared to the
769 Compione area.

770

771 **7.5 Evolutionary Model**

772 The structural fabric preserved in the footwall of the Compione Fault indicates that fault activity in
773 the Macigno Sandstones Formation started with the formation of a km-scale network of shear
774 fractures in conjugate arrays with vertical σ_1 bisector, as expected in extensional Andersonian faulting

775 (Anderson, 1951; Sibson, 1996). The strike of the fractures was mainly parallel to the main trend of
776 the Compione Fault, but also E-W, i.e. parallel to the major footwall fault splay occurring in the study
777 area. This suggests that linkage within and among the major footwall segments constituting the
778 Northern Lunigiana fault system (Fig. 3) occurred in the very early stages of extensional deformation.
779 Such an early fracture network formed the process zone (Lockner et al., 1992; Reches and Lockner,
780 1994; Cowie and Shipton, 1998; Vermilye and Scholz, 1998) of the Compione Fault in the Macigno
781 Sandstones Formation, ahead of the upward-propagating master fault surface. The process zone was
782 a preferential site for effective fluid circulation and advection of a hydrothermal plume at minimum
783 temperatures of 210-230 °C, i.e. the trapping temperature of MC1A inclusions assuming hydrostatic
784 conditions at 5 km depth, and possibly exceeding ~230 °C, as suggested by fracture cementation with
785 prehnite (Fig. 17A). Results from microthermometry and stable isotope geochemistry indicate an
786 open system circulation with upward directed and channelized high-temperature and low-salinity
787 fluids coming from the metamorphic basement (Mazzarini et al., 2010; Boschetti et al., 2017) which
788 mixed with carbon derived from the maturation of organic matter in the Macigno host rock.
789 Petrographic evidence supports cross-formational fluid flow up to the base of the Ottone Flysch
790 Formation in this early stage.

791 With increasing extension, the vein network in the process zone was progressively tilted by
792 extensional fault-propagation folding (Withjack et al., 1990; Schlische, 1995; Hardy and McClay,
793 1999; Ferrill et al., 2004a; Jin and Groshong, 2006) and bedding attained a synthetic dip attitude.
794 During folding, the fault-parallel extensional conjugate vein arrays was preferentially reactivated by
795 antithetic shearing, accompanied by precipitation of quartz and MC1B calcite (Fig. 17B).
796 Mineralization occurred at minimum temperatures between 140 and 160 °C and increasing fluid
797 salinities indicating mixing with a fluid characterized by lower temperature and higher salinity. At
798 this stage there was still stratigraphic continuity across the Compione fault zone, which near surface
799 expression was likely a flexure in the crestal region of the regional-scale anticline deforming the
800 previously stacked thrust sheets of Tuscan and Ligurian rocks. The presence of the clay-rich

801 Subligurian succession at the top of the Macigno Sandstones Formation and the change from
802 siliciclastic to carbonate composition helps explain why conjugate fracture arrays comparable to the
803 underlying ones did not develop in the Ottone Flysch Formation. In the latter, deformation was
804 accommodated through a network of fault segments that partially exploited the pre-existing structural
805 inheritance, accompanied by precipitation of OC2 calcite and rare quartz, at a temperature lower than
806 110 °C (Fig. 17B).

807 When bedding dip in the extensional monocline reached values exceeding ~50°, fault propagation
808 and breakthrough was accompanied by formation of a footwall damage zone with a width of about
809 half that of the corresponding process zone. Many rotated shear veins and subsidiary faults were re-
810 activated and together with newly-formed ones, produced a network of high-angle faults with
811 cataclastic cores of disaggregated and gouge layers. MC2 calcite precipitated in fractures of the
812 damage zone, from fluids at temperature between 70° and 90° C, resulting from mixing between
813 deeply-sourced fluids and meteoric waters which likely interacted, at different degrees, with the
814 Burano Evaporites in the subsurface (Fig. 17C). This would explain the high salinity of a low
815 temperature fluid with a stable isotopic composition typical of meteoric fluids. The damage zone in
816 the hanging wall had a comparable width as the footwall but was affected by less intense fracturing,
817 partly replaced by dissolution and discrete subsidiary faulting.

818

819 **8. CONCLUSION**

820 The Compione Fault is part of the Northern Lunigiana regional-scale extensional fault system,
821 exposed for around 30 km along strike, which accumulated about 1.5 km offset since Early Pliocene
822 times. It is located in the inner portion of the Northern Apennines, at the forelimb-crest transition of
823 a major out-of-sequence thrust-related anticline, that deformed the previously stacked thrust sheet
824 pile. The Compione Fault can be traced in seismic reflection profile down to the seismic basement
825 top at about 6 to 7 km depth and dissects the previously produced contractional architecture. This

826 fault zone offers the possibility to study the interaction between deformation, fluid flow and fracture
827 cementation that progressed from a depth of ~5 km up to near surface conditions. The following
828 major points can be drawn from this multidisciplinary study of the Compione Fault cross-sectional
829 architecture, resulting from the combination of structural, petrographical, geochemical,
830 microthermometric, and paleothermal analyses.

- 831 - The footwall damage zone, affecting thick sandstone strata, is characterized by a network of
832 shear veins with bisectors perpendicular to bedding, which was passively rotated by
833 extensional fault-propagation folding during upward fault growth. Such a fracture mesh
834 testifies for the presence of a wide process zone ahead of the fault tip in the early evolutionary
835 stages. A comparable deformation pattern does not occur in the hanging wall damage zone
836 because of either the carbonate composition of the Ottone Flysch Formation that favored an
837 important role of dissolution, or the presence of clay-rich sediments tectonically juxtaposed
838 at its base, which provided a strong mechanical discontinuity, or both.
- 839 - Mineralization of the process zone fracture network by a prehnite-quartz-calcite assemblage
840 from a fluid at a minimum temperature of 210 °C, possibly exceeding 230 °C indicates that:
841 (i) the process zone provided a well-connected fracture mesh that significantly improved
842 porosity and favored effective circulation and upward fluid migration; (ii) such a deep fluid
843 volume constituted a hydrothermal plume in strong thermal disequilibrium with the host rock
844 outside the fault zone, which experienced maximum temperatures of less than 140-150 °C.
- 845 - Shear vein cementation in the process zone was cyclic and episodic, indicating fluid pressure
846 variations that might relate to the earthquake cycle. According to this hypothesis, seismic
847 pumping may have promoted fast channelized fluid migration from the metamorphic
848 basement, along the fault zone and up to the process zone.
- 849 - Synthetic rotation about a horizontal axis of the process zone caused antithetic re-activation
850 of pre-existing shear veins as subsidiary faults and formation of new, non-mineralized high-

angle extensional faults. This event of deformation localization represents the formation of the fault damage zones *sensu-stricto*.

- The structural and paleofluid framework exposed in the thick sandstone beds at the footwall of the Compione Fault highlights the importance of the process zone for both fault scaling properties and hydrology. Process zone width is twice that of the damage zone *sensu-stricto*, produced by deformation localization during fault slip. This means that the total volume of footwall fractured rocks, typically included into the damage zone as a whole, is much thicker than what can be expected from statistical scaling laws and has a structural fabric mainly imprinted at the process zone stage. Furthermore, it developed diachronously during fault evolution, with maximum permeability and fluid advection ahead of the upward propagating fault tip, followed by fracture cementation, deformation localization, and porosity and permeability reduction in more mature stages.
- Development of a process zone at the onset of extensional faulting can significantly contribute creating economically valuable fractured reservoirs ahead of fault tips. Extensional fault-propagation folding favors migration and accumulation of fluids from deeper stratigraphic horizons and the metamorphic basement into the extensional process zone, where high fluid pressures may likely occur. Eventually, fault breakthrough causes reservoir compartmentalization and sealing, preserving favorable conditions for fluid storage in the footwall damage zone and the corresponding process zone sector.
- Depending on the first-order mechanical stratigraphy, development of crustal-scale extensional fault systems can create a strong vertical variability of fractured rock volumes resulting from the interplay among several factors, including: (i) the dominant deformation mechanisms, (ii) fault-propagation versus slip rates, (iii) possible development and width of a vertically-compartmentalized process zone ahead of the upward-migrating fault tip, (iv) competition between faulting and folding, etc. Such a large variability and vertical compartmentalization of the cross-sectional damage zone width strongly impacts fluid storage

potential and partitioning in rift-related fault-bounded blocks, as well as the seismic behavior of fault zones.

ACKNOWLEDGMENTS

Discussions with Mahtab Mozafari and Alessandra Montanini on vein cement analysis are gratefully acknowledged. We are indebted to Andrea Comelli for thin section preparation, Enrico Selmo for providing isotopic data, Luca Barchi for support with scanning electron microscopy and Zita Kelemen and Steven Bouillon for help with micromilling equipment. Anna Laura Cazzola, Alessandro Fattorini and Claudio Cattaneo of ENI S.p.A. are thanked for providing reflection seismic data and authorizing their publication and Midland Valley for providing the software Move in the frame of the Academic Software Initiative. Alessio Lucca and Fabrizio Storti ideated the study, collected most of the data and wrote the manuscript, Giancarlo Molli and Fabrizio Balsamo participated to the fieldwork, to data discussion and tectonic interpretations, Philippe Muchez participated to the analysis and interpretation of vein cement petrographic, microthermometric and geochemical data, Andrea Schito and Sveva Corrado provided vitrinite reflectance data and their thermal modelling, and discussed their meaning in the framework of this research, Andrea Artoni depth converted and interpreted reflection seismic data, and Emma Salvioli Mariani helped with Raman spectroscopy and fluid inclusion analysis.

REFERENCES

- Abbate, E., Balestrieri, M.L., Bigazzi, G., Norelli, P., and Quercioli, C., 1994, Fission-track dating and recent rapid denudation in Northern Apennines, Italy: *Mem. Soc. Geol. It.*, v. 48, p. 579–585.
- Anderson, E.M., 1951, *The Dynamics of Faulting and Dyke Formation with Application to Britain* (2nd edn): Edinburgh, Scotland, Oliver and Boyd, p. 206.

901 Argnani, A., 2002, The Northern Apennines and the kinematics of Europe-Africa convergence:
 902 Bollettino della Società Geologica Italiana, v. 1, p. 47–60.

903 Argnani, A., Barbacini, G., Bernini, M., Camurri, F., Ghielmi, M., Papani G., Rizzini F., Rogledi S.,
 904 and Torelli, L., 2003, Gravity tectonics driven by Quaternary uplift in the Northern Apennines:
 905 insights from the La Spezia-Reggio Emilia geo-transect: Quaternary International, v. 101-102, p. 13-
 906 26.

907 Arnason, J.G., Bird, D.K., and Liou, J.G., 1993, Variables Controlling Epidote Composition in
 908 Hydrothermal and Low-Pressure Regional Metamorphic Rocks: Abhandlungen der Geologischen
 909 Bundesanstalt, v. 49, pp. 17-25.

910 Artoni, A., Bernini, M., Papani, G., Vescovi, P., and Zanzucchi, G., 1992, Sezione geologica
 911 schematica Bonassola (SP)-Felino (PR): Studi Geologici Camerti, Volume Speciale 1992/2, CROP
 912 01-1A, p. 61-63.

913 Augliera, P., Pastore, S., and Tomaselli, A., 1990, Sismicità della Lunigiana-Garfagnana: Primi
 914 risultati da una rete mobile: Proc. 9, Gruppo Naz. Geofis. Della Terra Solida, v. 1, p. 221-232.

915 Azzaroli, A., 1950, Osservazioni sulla formazione villafranchiana di Olivola in Val di Magra: Atti
 916 della Società Toscana di Scienze Naturali Memorie, v. 57, p. 104-111.

917 Azzaroli, A., 1977, The Villafranchian stage in Italy and the Plio-Pleistocene boundary: Giornale di
 918 Geologia, v. 41, p. 61-79.

919 Balestrieri, M.L., Bernet, M., Brandon, M.T., Picotti, V., Reiners, P., and Zattin, M., 2003, Pliocene
 920 and Pleistocene exhumation and uplift of two key areas of the Northern Apennines: Quaternary
 921 International, v. 101– 102, p. 67–73.

922 Barchi, M.R., Minelli, G., and Pialli, G., 1998, The CROP 03 Profile: A synthesis of results on deep
 923 structures of the Northern Apennines: Memorie della Società Geologica Italiana, v. 52, p. 383–400.

924 Bartole, R., 1995, The North Tyrrhenian–Northern Apennines post-collisional system: constraints for
 925 a geodynamic model: Terra Nova, v. 7, p. 7-30.

926 Bartolini, C., Bernini, M., Carloni, G.C., Costantini, A., Federici, P.R., Gasperi, G., Lazzarotto, A.,
 927 Marchetti, G., Mazzanti, R., Papani, G., Pranzini, G., Rau, A., Sandrelli, F., Vercesi, P.L., Castaldini,
 928 D., and Francavilla, F., 1982, Carta Neotettonica dell'Appennino Settentrionale. Note Illustrative:
 929 Bollettino della Società Geologica Italiana, v. 101, p. 355-371.

930 Beaudoin, N., Bellahsen, N., Lacombe, O., and Emmanuel, L., 2011, Fracture-controlled
931 paleohydrogeology in a basement-cored, fault-related fold: Sheep Mountain Anticline, Wyoming,
932 United States: *Geochemistry, Geophysics, Geosystems*, v. 12, n. 6, Q06011.

933 Bernet, M., Brandon, M.T., Garver, J.I., and Molitor, B.R., 2004, Fundamentals of detrital zircon
934 fission-track analysis for provenance and exhumation studies with examples from the European Alps,
935 in Bernet M., and Spiegel C., eds., *Detrital Thermochronology; Provenance Analysis, Exhumation,*
936 *and Landscape Evolution of Mountain Belts: Geological Society of America Special Paper 378*, p.
937 25–36.

938 Bernini, M., 1988, Il bacino dell'Alta Val Magra: primi dati mesostrutturali sulla tettonica distensiva:
939 *Boll. Soc. Geol. It.*, v. 107, p. 355-371.

940 Bernini, M., and Lasagna, S., 1988, Rilevamento geologico e analisi strutturale del bacino dell'Alta
941 Val Magra tra M. Orsaro e Pontremoli (Appennino Settentrionale): *Atti della Società Toscana di*
942 *Scienze Naturali Memorie, Serie A*, 95, p. 139-183.

943 Bernini, M., Boccaletti, M., Moratti, G., Papani, G., Sani, F., and Torelli, L., 1990, Episodi
944 compressivi neogenico-quadernari nell'area estensionale tirrenica nord-orientale. Dati in mare e a
945 terra: *Mem. Soc. Geol. It.*, v. 45, p. 577-589.

946 Bernini, M., 1991, Le strutture estensionali della Lunigiana (Appennino Settentrionale): proposta di
947 un modello deformativo: *Atti Ticinese di Scienze della Terra*, v. 34, p. 29-38.

948 Bernini, M., and Papani, G., 2002, La distensione della fossa tettonica della Lunigiana nord-
949 occidentale (con Carta Geologica alla scala 1:50.000): *Boll. Soc. Geol. It.*, v. 121, p. 313-341.

950 Bertoldi, R., 1988, Una sequenza palinologica di età rusciniana nei sedimenti palustri basali del
951 bacino di Aulla-Olivola (Val di Magra): *Rivista Italiana di Paleontologia e Stratigrafia*, v. 94, p. 105-
952 138.

953 Bertoldi, R., 1995, Palinostratigrafia dei bacini intramontani dell'Appennino nord-occidentali: *Il*
954 *Quaternario*, v. 8, p. 61-70.

955 Bird, D.K., and Helgeson, H.C., 1980, Chemical interaction of aqueous solutions with epidote-
956 feldspar mineral assemblages in geologic systems. I. Thermodynamic analysis of phase relations in
957 the system $\text{CaO-FeO-Fe}_2\text{O}_3\text{-Al}_2\text{O}_3\text{-SiO}_2\text{-H}_2\text{O-CO}_2$: *Am. Jour. Sci.*, v. 280, p. 907-941.

958 Bird, D.K., and Helgeson, H.C., 1981, Chemical interaction of aqueous solutions with epidote-
 959 feldspar mineral assemblages in geologic systems. II. Equilibrium constraints in
 960 metamorphic/geothermal processes: *Am. Jour. Sci.*, v. 281, p. 576-614.

961 Boccaletti, M., Elter, P., and Guazzone, G., 1971, Plate Tectonic Models for the Development of the
 962 Western Alps and Northern Apennines: *Nature*, v. 234, p. 108-111.

963 Boccaletti, M., Cerrina Ferroni, A., Martinelli, P., Moratti, G., Plesi, G., and Sani, F., 1992, Late
 964 Miocene-Quaternary compressive events in the Tyrrhenian side of the Northern Apennines: *Annales*
 965 *Tectonicae*, v.6, p. 214-230.

966 Boccaletti, M., Gianelli, G., and Sani F., 1997, Tectonic regime, granite emplacement and crustal
 967 structure in the inner zone of the Northern Apennines (Tuscany, Italy): A new hypothesis:
 968 *Tectonophysics*, v. 270, p. 127-143.

969 Boccaletti, M., and Sani, F., 1998, Cover Thrust reactivations related to internal basement
 970 involvement during Neogene-Quaternary evolution of the Northern Apennines: *Tectonics*, v. 17, p.
 971 112-130.

972 Boccaletti, M., Corti, G., and Martelli, L., 2011, Recent and active tectonics of the external zone of
 973 the Northern Apennines (Italy): *Int. J. Earth Sci.*, v. 100, p. 1331-1348.

974 Bodnar, R.J., 1993, Revised equation and table to determine the freezing point depression of H₂O-
 975 NaCl solutions: *Geochim. Cosmochim. Acta*, v. 57., pp. 683-684.

976 Boiron, M.C., Cathelineau, M., Ruggieri, G., Jeanningros, A., Gianelli, G., and Banks, A., 2007,
 977 Active contact metamorphism and CO₂-CH₄ fluid production in the Larderello geothermal field
 978 (Italy) at depths between 2.3 and 4 km: *Chemical Geology*, v. 237, p. 303–327.

979 Boncio, P., Brozzetti, F., and Lavecchia, G., 2000, Architecture and seismotectonics of a regional
 980 low-angle normal fault zone in central Italy: *Tectonics*, v. 19, p. 1038-1055.

981 Bonini, M., Moratti, G., Sani, F., and Balestrieri, M.L., 2013, Compression-to-extension record in the
 982 late Pliocene–Pleistocene Upper Valdarno Basin (Northern Apennines, Italy): Structural and
 983 thermochronological constraints: *Italian Journal of Geosciences*, v. 132, p. 54–80.

984 Bons, P.D., 2001, Development of crystal morphology during unitaxial growth in a progressively
 985 widening vein: I. The numerical model: *Journal of Structural Geology*, v. 23, p. 865-872.

986 Bons, P.D., Elburg, M.A., and Gomez-Rivas, E., 2012, A review of the formation of tectonic veins
 987 and their microstructures: *Journal of Structural Geology*, v. 43, p. 33-62.

988 Boschetti, T., Toscani, L., Barbieri, M., Mucchino, C., and Marino, T., 2017, Low enthalpy Na-
 989 chloride waters from the Lunigiana and Garfagnana grabens, Northern Apennines, Italy: Tracing fluid
 990 connections and basement interactions via chemical and isotopic compositions: *Journal of*
 991 *Volcanology and Geothermal Research*, v. 348, p. 12–25.

992 Bossolasco, M., Eva, C., and Pasquale, C., 1973, Il terremoto del 25 Ottobre in Lunigiana: *Rivista*
 993 *Italiana di Geofisica*, v. 22, p. 314-322.

994 Bossolasco, M., Eva, C., and Pasquale, C., 1974, On seismotectonics of the Alps and Northern
 995 Apennines: *Rivista Italiana di Geofisica*, v. 23(1/2), p. 57-63.

996 Boullier, A.-M., and Robert, F., 1992, Paleoseismic events recorded in Archean gold-quartz vein
 997 networks, Val d'Or, Abitibi, Quebec, Canada: *Journal of Structural Geology*, v. 14, p. 161-180.

998 Boullier, A.-M., Charoy, B., and Pollard, P.J., 1994. Fluctuation in porosity and fluid pressure during
 999 hydrothermal events: textural evidence in the Emuford District, Australia: *Journal of Structural*
 1000 *Geology*, v. 16 (10), p. 1417-1429.

1001 Browne, P.R.L., 1978, Hydrothermal alteration in active geothermal fields: *Ann. Rev. Earth Planet.*
 1002 *Sci.*, v. 6, p. 229-250.

1003 Burkhard, M., 1993, Calcite twins, their geometry, appearance and significance as stress-strain
 1004 markers and indicators of tectonic regime: a review: *Journal of Structural Geology*, v. 15, p. 351-368.

1005 Burnham, A.K., and Sweeney, J.J., 1989, A chemical kinetic model of vitrinite maturation and
 1006 reflectance: *Geochimica et Cosmochimica Acta*, v. 53, p. 2649–2657, doi: 10.1016/0016-7037
 1007 (89)90136-1.

1008 Bustin, R.M., Barnes, M.A., and Barnes, W.C., 1990, Determining levels of organic diagenesis in
 1009 sediments and fossil fuels, in McIlreath, I.A., and Morrow, D.W., eds., *Diagenesis: Geoscience*
 1010 *Canada Reprint Series 4*, p. 205–226.

1011 Butler, R.W.H., 1992, Hydrocarbon maturation, migration and tectonic loading in the western Alps,
 1012 in England, W.A., and Fleet, A.J., eds., *Petroleum Migration: Geological Society of London Special*
 1013 *Publication 59*, p. 227–244.

1014 Caine, J.S., Evans, J.P., and Forster, J.B., 1996, Fault zone architecture and permeability structure:
 1015 *Geology*, v. 24, p. 1025-1028.

1016 Camurri, F., Argnani, A., Bernini, M., Papani, G., Rogledi, S., and Torelli, L., 2001, The basement
1017 of the NW Apennines: interpretation of reflection seismics and geodynamic implications: Final
1018 Proceedings, Geoitalia 2001, 3° Forum Italiano di Scienze della Terra (Fist), Chieti, Italy, p. 50-51.

1019 Caricchi, C., Aldega, L., and Corrado, S., 2015, Reconstruction of maximum burial along the
1020 Northern Apennines thrust wedge (Italy) by indicators of thermal exposure and modeling: GSA
1021 Bulletin, 127(3-4), 428-442.

1022 Carlini, M., Artoni, A., Aldega, L., Balestrieri, M.L., Corrado, S., Vescovi, P., Bernini, M., and
1023 Torelli, L., 2013, Exhumation and reshaping of far-travelled/allochthonous tectonic units in mountain
1024 belts. New insights for the relationships between shortening and coeval extension in the western
1025 Northern Apennines (Italy): Tectonophysics, v. 608, p. 267–287.

1026 Carmignani, L., and Kligfield, R., 1990, Crustal extension in the Northern Apennines: The transition
1027 from compression to extension in the Alpi Apuane core complex: Tectonics, v. 9, p. 1275–1303.

1028 Carmignani, L., Decandia, F.A., Fantozzi, P.L., Lazzarotto, A., Liotta, D., and Meccheri, M., 1994.
1029 Tertiary extensional tectonics in Tuscany (Northern Apennines, Italy): Tectonophysics, v. 238, p.
1030 295–315.

1031 Carmignani, L., Decandia, F.A., Disperati, L., Fantozzi, P.L., Lazzarotto, A., Liotta, D., and Oggiano,
1032 G., 1995. Relationships between the Tertiary structural evolution of the Sardinia-Corsica-Provençal
1033 Domain and the Northern Apennines: Terra Nova, v. 7, no. 2, p. 128–137.

1034 Carosi, R., Leoni, L., Montomoli, C., and Sartori, F., 2003, Very low-grade metamorphism in the
1035 Tuscan Nappe, Northern Apennines, Italy. Relationships between deformation and metamorphic
1036 indicators in the La Spezia mega-fold: Swiss Bulletin of Mineralogy and Petrology, v. 83, p. 15-32.

1037 Carter, K.E., and Dworkin, S.I., 1990, Channelized fluid flow through shear zones during fluid-
1038 enhanced dynamic recrystallization, Northern Apennines, Italy: Geology, v. 18, p. 720–723.

1039 Cavarretta, G., Gianelli, G., and Puxeddu, M., 1982, Formation of authigenic minerals and their use
1040 as indicators of the physicochemical parameters of the fluid in the Larderello-Travale geothermal
1041 field: Economic Geology, v. 77, p. 1071-1084.

1042 Cerrina Feroni, A., Moratti, G., and Plesi, G., 1983, Evidenze di episodi compressivi messiniano-
1043 pliocenici alternati alla tettonica di distensione nella Toscana sud-occidentale, emerse dall'analisi
1044 mesostrutturale. In: Atti della riunione su “Meccanismi deformativi nelle catene perimediterranee:
1045 stato di avanzamento delle ricerche e problematiche emerse”, Firenze, Italy, 5 dicembre 1983, Centro
1046 Stampa Palagi Firenze, 35-42.

1047 Childs, C., Manzocchi, T., Walsh, J.J., Bonson, C.G., Nicol, A., and Schopfer, M.P.J., 2009, A
 1048 geometric model of fault zone and fault rock thickness variations: *Journal of Structural Geology*, v.
 1049 31 (2), p. 117-127.

1050 Clemenzi, L., Molli, G., Storti, F., Muchez, P., Swennen, R., and Torelli, L., 2014, Extensional
 1051 deformation structures within a convergent orogen: The Val di Lima low-angle normal fault system
 1052 (Northern Apennines, Italy): *Journal of Structural Geology*, v. 66, p. 205-222.

1053 Connolly, J.A.D., 2010, The mechanics of metamorphic fluid expulsion: *Elements*, v. 6, p. 165-172,
 1054 doi:10.2113/gselements.6.3.165.

1055 Corrado, S., Aldega, L., and Zattin, M., 2010, Sedimentary vs. tectonic burial and exhumation along
 1056 the Apennines (Italy): *Journal of the Virtual Explorer, Electronic Edition*, ISSN 1441-8142, v. 36.
 1057 Paper 15, In: Beltrando M., Peccerillo A., Mattei M., Conticelli S. and Doglioni C., *The Geology of*
 1058 *Italy*.

1059 Cowie, P.A., Shipton, Z.K., 1998, Fault tip displacement gradients and process zone dimension:
 1060 *Journal of Structural Geology*, v. 20, p. 983-997.

1061 Cox, S. F., 1995, Faulting processes at high fluid pressures: An example of fault valve behavior from
 1062 the Wattle Gully Fault, Victoria, Australia: *J. Geophys. Res.*, v. 100, p. 12841-1285.

1063 Cox, S.F., 1999, Deformational controls in the dynamics of fluid flow in mesothermal gold systems,
 1064 in McCaffrey, K., Lonergan, L., and Wilkinson, J., eds., *Fractures, Fluid Flow and* 1160
 1065 *Mineralization: Geological Society of London Special Publication*, v. 155, p. 123-140.

1066 Curewitz, D., and Karson, J.A., 1997, Structural settings of hydrothermal outflow: Fracture
 1067 permeability maintained by fault propagation and interaction: *Journal of Volcanology and*
 1068 *Geothermal research*, v. 79, p. 149-168.

1069 Davis, D.M., Suppe, J. and Dahlen, F.A., 1983, Mechanics of fold-and thrust belts and accretionaryw
 1070 edges: *J. Geophys. Res.*, v. 88, p. 1153-1172.

1071 Decandia, F.A., Lazzarotto, A., and Liotta, D., 1998, The CROP 03 traverse: insights on post-
 1072 collisional evolution of Northern Apennines: *Mem. Soc. Geol. It.*, v. 52, p. 427–440.

1073 Della Vedova, B., Lucazeau, F., Pasquale, V., Pellis, G., and Verdoya, M., 1995, Heat flow in the
 1074 tectonic provinces crossed by the southern segment of the European Geotraverse: *Tectonophysics*, v.
 1075 244, p. 57-74.

1076 Dewey, J.F., Helman, M.L., Knott, S.D., Turco, E., and Hutton, D.H.W., 1989, Kinematics of the
 1077 western Mediterranean, In: Coward M.P., Dietrich D., and Park R.G., eds., *Alpine Tectonics*:
 1078 Geological Society of London Special Publication, v. 45, p. 265–283.

1079 Di Naccio, D., Boncio, P., Brozzetti, F., Pazzaglia, F.J., and Lavecchia, G., 2013, Morphotectonic
 1080 analysis of the Lunigiana and Garfagnana grabens (northern Apennines, Italy): Implications for active
 1081 normal faulting: *Geomorphology*, v. 201, p. 293-311.

1082 Di Paolo, L., Aldega, L., Corrado, S., and Mastalerz, M., 2012, Maximum burial and unroofing of
 1083 Mt. Judica recess area in Sicily: implication for the Apenninic Maghrebian wedge dynamics:
 1084 *Tectonophysics* 530-531, 193-207.

1085 Di Pisa, A., Franceschelli, M., Leoni, L., and Meccheri, M., 1985. Regional variation of the
 1086 metamorphic temperatures across the Tuscanid 1 unit and its implications on the alpine
 1087 metamorphism (Apuane Alps, N-Tuscany): *Neues Jahrbuch für Mineralogie, Abhandlungen*, v. 151,
 1088 p. 197–211.

1089 Dickson, J.A.D., 1966, Carbonate identification and genesis as revealed by staining: *Journal of*
 1090 *Sedimentary Petrology*, v. 32, p. 491-505.

1091 Dini, A., Gianelli, G., Puxeddu, M., and Ruggieri, G., 2005, Origin and evolution of Pliocene-
 1092 Pleistocene granites from the Larderello geothermal field (Tuscan Magmatic Province, Italy): *Lithos*,
 1093 v. 81, p. 1–31.

1094 Doglioni, C., 1991, A proposal for the kinematic modelling of W-dipping subductions - Possible
 1095 applications to the Tyrrhenian-Apennines system: *Terra Nova*, v. 3, n. 4, p. 423–434.

1096 Doglioni, C., Barba, S., Carminati, E., and Riguzzi, F., 2014, Fault on–off versus coseismic fluids
 1097 Reaction: *Geosci. Front.* V. 5, p. 767–780. <http://dx.doi.org/10.1016/j.gsf.2013.08.004>.

1098 Durney, D.W., and Ramsay, J.G., 1973, Incremental strains measured by syntectonic crystal growths,
 1099 In: De Jong K.A., Scholten, K. (Eds.), *Gravity and Tectonics*, Wiley, New York, USA, p. 67-96.

1100 Elter, P., and Schwab, K., 1959, Note illustrative della Carta Geologica alla scala 1:50.000 della
 1101 regione Carro-Zeri-Pontremoli: *Boll. Soc. Geol. It.*, v. 78, p. 157-187.

1102 Elter, P., and Pertusati, P., 1973, Considerazioni sul limite Alpi-Appennino e sulle relazioni con l’arco
 1103 delle Alpi occidentali: *Mem. Soc. Geol. It.*, v. 12, p. 359-375.

1104 Elter, P., Giglia, G., Tongiorgi, M., and Trevisan, L., 1975, Tensional and compressional areas in
 1105 recent (Tortonian to present) evolution of Northern Apennines: *Boll. Geofis. Teor. Appl.*, v. 17, p. 13-
 1106 18.

1107 Endignoux, L., and Wolf, S., 1990, Thermal and kinematic evolution of thrust basins: a 2D numerical
 1108 model, In: *Petroleum and Tectonics in Mobile Belts*, J. Letouzey (Ed.) and Editions Technip, Paris,
 1109 France, 181-192.

1110 Eva, E., Solarion, S., and Boncio, P., 2014, HypoDD relocated seismicity in northern Apennines,
 1111 (Italy) preceding the 2013: seismotectonic implications for the Lunigiana-Garfagnana area: *Boll.*
 1112 *Geofis. Teor. Appl.*, v. 55, p. 739-754.

1113 Evans, M. A., and Battles, D.A., 1999, Fluid inclusion and stable isotope analyses of veins from the
 1114 central Appalachian Valley and Ridge province: Implications for regional synorogenic hydrologic
 1115 structure and fluid migration: *Geological Society of America Bulletin*, v. 111, p. 1841-1860.

1116 Faulkner, D.R., and Rutter, E.H., 2001, Can the maintenance of overpressured fluids in large strike-
 1117 slip fault zones explain their apparent weakness?: *Geology*, v. 29 (6), p. 503-506.

1118 Faulkner, D., Jackson, C., Lunn, R., Schlische, R., Shipton, Z., Wibberley, C., and Withjack, M.,
 1119 2010, A review of recent developments concerning the structure, mechanics and fluid flow properties
 1120 of fault zones: *J. Struct. Geol.* v. 32, p. 1557-1575. <http://dx.doi.org/10.1016/j.jsg.2010.06.009>.

1121 Federici, P.R., 1978, La tettonica recente dell'appennino; 2) Il bacino fluvio-lacustre di Pontremoli
 1122 (Alta Val di Magra) e sue implicazioni neotettoniche: *G.S.Q.P.*, Quaderno n. 4, p. 121-131.

1123 Federici, P.R., 1981, Nuovi resti di vertebrato nel bacino fluvio-lacustre villafranchiano di Pontremoli
 1124 (Val di Magra): *Bolletino della Museo Storia Naturale della Lunigiana*, 1 (2), p. 71-74.

1125 Fellin, M.G., Reiners, P.W., Brandon, M.T., Wüthrich, E., Balestrieri, M.L., and Molli, G., 2007,
 1126 Thermochronologic evidence for the exhumational history of the Alpi Apuane metamorphic core
 1127 complex, Northern Apennines, Italy: *Tectonics*, v. 26, TC6015, doi: 10.1029/2006TC002085.

1128 Ferrill, D.A., Morris A.P., Sims D.W., Waiting D.J., and Hasegawa S., 2004a, Development of
 1129 synthetic layer dip adjacent to normal faults, In: Sorkhabi R., Tsuji Y. (Eds.), *Faults, Fluid Flow, and*
 1130 *Petroleum Traps* American Association of Petroleum Geologists Memoir. American Association of
 1131 Petroleum Geologists, Tulsa, OK, USA, p. 125–138.

1132 Ferrill, D.A., Morris, A.P., Evans, M.A., Burkhard, M., Groshong, R.H., and Onasch, C.M., 2004b,
 1133 Calcite twin morphology: a low-temperature deformation geothermometer: *Journal of Structural*
 1134 *Geology*, v. 26, p. 1521–1529.

1135 Fisher, D.M., Brantley, S.L., Everett, M., and Dzvonik, J., 1995, Cyclic fluid-flow through a
 1136 regionally extensive fracture network within the Kodiak accretionary prism: *Journal of Geophysical*
 1137 *Research, Solid Earth*, v. 100 (B7), p. 12881-12894.

1138 Fondriest, M., Smith, S.A.F., Candela, T., Nielsen, S.B., Mair, K., and Di Toro, G., 2013, Mirror-like
 1139 faults and power dissipation during earthquakes: *Geology*, v. 41 (11), p. 1175-1178.

1140 Fossen, H., Schultz, R.A., Shipton, Z.K., and Mair, K., 2007, Deformation bands in sandstone: a
 1141 review: *Journal of the Geological Society*, v. 164, p. 755-769.

1142 Fossen, H., 2010, *Structural Geology*, 463 pp., Cambridge Univ. Press, Cambridge, U. K.

1143 Franceschelli, M., Leoni, L., Memmi, I., and Puxeddu, M., 1986, Regional distribution of Al-silicates
 1144 and metamorphic zonation in the low-grade Verrucano metasediments from the Northern Apennines,
 1145 Italy: *Journal of Metamorphic Geology*, v. 4, p. 309–321.

1146 Frepoli, A., and Amato, A., 1997, Contemporaneous extension and compression in the Northern
 1147 Apennines from earthquake fault-plane solutions: *Geophys. J. Int.*, v. 129, p. 368-388.

1148 Friedman, I., and O’Neil, J.R., 1977, Compilation of stable isotope fractionation factors of
 1149 geochemical interest, In: Fleisher, M., ed., *Data of Geochemistry* (6th ed.): U.S. Geological Survey
 1150 Professional Paper 440-KK, 117 p.

1151 Ghelardoni, R., 1965, Osservazioni sulla tettonica trasversale dell’Appennino settentrionale: *Boll.*
 1152 *Soc. Geol. It.*, 84, pp. 276-290.

1153 Ghibaudo, G., 1980, Deep-sea fan deposits in the Macigno Formation (middle-upper Oligocene) of
 1154 the Gordana Valley, northern Apennines, Italy: *Journal of Sedimentary Research*, v. 50, p. 723-741.

1155 Gianelli, G., Ruggieri, G., and Mussi, M., 1997, Isotopic and fluid inclusion study of hydrothermal
 1156 and metamorphic carbonates in the Larderello geothermal field and surrounding areas, Italy:
 1157 *Geothermics*, v. 26, p. 393-417.

1158 Giglia, G., 1974, L’insieme Corsica-Sardegna e I suoi rapporti con l’Appennino settentrionale: rassegna
 1159 di dati cronologici e strutturali, in Maxia, C., and Pomesano A., eds., *Paleogeografia del Terziario*
 1160 *sardo nell’ambito del Mediterraneo occidentale*: Cagliari, Italy, *Rendiconti Semestrali della Facoltà*
 1161 *di Scienze, Università di Cagliari*, 43, p. 245–275.

Giustini, F., Brilli, M., and Patera, A., 2016, Mapping oxygen stable isotopes of precipitation in Italy: Journal of Hydrology: Regional Studies, v. 8, p. 162-181.

Goldstein, R.H., and Reynolds, J., 1994, Systematics of Fluid Inclusions in Diagenetic Minerals: Short Course Notes: Tulsa, Oklahoma, USA, The Society for Sedimentary Geology, v. 31, <https://doi.org/10.2110/scn.94.31>.

Gratier, J.P., Favreau, P., Renard, F., and Pili, E., 2002, Fluid pressure evolution during the earthquake cycle controlled by fluid flow and pressure-solution crack sealing, Earth Planets Space, v. 54, p. 1139-1146.

Hardy, S., and Ford, M., 1997, Numerical modelling of trishear fault propagation folding: Tectonics, v. 16, n. 5, pp. 841-854.

Hardy, S., and McClay, K., 1999, Kinematic modeling of extensional fault propagation folding: Journal of Structural Geology, v. 21, p. 695–702.

Hilgers, C., and Urai, J.L., 2002, Experimental study of syntaxial vein growth during lateral fluid flow in transmitted light: first results: Journal of Structural Geology, v. 24, p. 1029-1043.

Hoefs, J., 1997, Stable Isotope Geochemistry: Springer-Verlag, Berlin, Germany, 200 p.

Holland, M., and Urai, J.L., 2010, Evolution of anastomosing crack-seal vein networks in limestones: insights from an exhumed high pressure cell, Jabal Shams, Oman Mountains: Journal of Structural Geology, v. 32, p. 1279-1290.

Honlet, R., Gasparrini, M., Jager, H., Muchez, P., and Swennen, R., 2017, Precursor and ambient rock paleothermometry to assess the thermicity of burial dolomitization in the southern Cantabrian Zone (northern Spain): International Journal of Earth Sciences, v. 106-7, p. 1-21.

Huber, M., Halas, S., and Sikorska, M., 2007, Evolution of prehnite-albite-calcite veins in metamorphic rocks from the Lapland Granulite Belt (Kandalaksha region of Kola Peninsula): Geologija, v. 57, p. 1-7.

Ingebritsen, S.E., and Manning, C.E., 2010, Permeability of the continental crust: Dynamic variations inferred from seismicity and metamorphism: Geofluids, v. 10, p. 193–205.

Jessell, M.W., Willman, C.E., and Gray, D.R., 1994, Bedding parallel veins and their relationship to folding: Journal of Structural Geology, v. 16, p. 753-767.

Jin, G., and Groshong, Jr. R.H., 2006, Trishear kinematic modelling of extensional fault-propagation folding: Journal of Structural Geology, v. 28, p. 170-183.

1192 Jolivet, L., Faccenna, C., Goffé, B., Mattei, M., Rossetti, F., Brunet, C., Storti, F., Funicello, R.,
1193 Cadet, J.P., D'Agostino, N., and Parra, T., 1998, Midcrustal shear zones in postorogenic extension:
1194 Example from the northern Tyrrhenian Sea: *Journal of Geophysical Research*, v. 103, p. 12,123–
1195 12,160.

1196 Ketcham, R.A., Donelick, R.A., and Carlson, W.D., 1999, Variability of apatite fission-track
1197 annealing kinetics: III. Extrapolation to geological time scales: *American mineralogist*, v. 84, p. 1235–
1198 1255.

1199 Kim, Y.S., Peacock, D.C.P., and Sanderson, D.J., 2004, Fault damage zones: *Journal of Structural*
1200 *Geology*, v. 26 (3), p. 503-517.

1201 Knott, S.D., Beach, A., Brockbank, P.J., Brown, J.L., McCallum, J.E., and Welbon, A.I., 1996,
1202 Spatial and mechanical controls on normal fault population: *Journal of Structural Geology*, v. 18, p.
1203 359- 372.

1204 Koehn, D., and Passchier, C.W., 2000, Shear sense indicators in striped bedding-veins: *Journal of*
1205 *Structural Geology*, v. 22, p. 1141-1151.

1206 Laurent, D., Lopez, M., Chauvet, A., Sauvage, A.C., Buatier, M., and Spangenberg, J.E., 2017,
1207 Sedimentary fluids/fault interaction during syn-rift burial of the Lodève Permian Basin (Hérault,
1208 France): An example of seismic-valve mechanism in active extensional faults: *Marine and Petroleum*
1209 *Geology*, v. 88, p. 303-328.

1210 Lockner, D.A., Byerlee, J.D., Kuksenk, V., Ponomarev, A., and Sidorin, A., 1992, Observation of
1211 quasistatic fault growth from acoustic emissions, In: *Fault mechanics and transport properties of*
1212 *rocks*, edited by Evans B. and Wong T.F., p. 3-32, Academic, San Diego, California.

1213 Longinelli, A., Selmo, E., 2003, Isotopic composition of precipitation in Italy: a first overall map:
1214 *Journal of Hydrology*, v. 270, p. 75–88.

1215 Malinverno, A., and Ryan, W.B.F., 1986, Extension in the Tyrrhenian Sea and shortening in the
1216 Apennines as result of arc migration driven by sinking of the lithosphere: *Tectonics*, v. 5, p. 227-245.

1217 Mamadou, M.M., Chatelineau, M., Bourdelle, F., Boiron, M-C., Elmaleh, A., and Brouand, M., 2016,
1218 Hot fluid flows around a major fault identified by paleothermometric studies (Tim Mersoï Basin,
1219 Niger): *Journal of Sedimentary Research*, v. 86, p. 914-928.

1220 Martini, I.P., Sagri, M., and Colella, A., 2001, Neogene-Quaternary basins of the inner Apennines
1221 and Calabrian arc, In: *Anatomy of an orogeny: The Apennines and adjacent Mediterranean Basins*,
1222 edited by Vai G.B. and Martini I.P., p. 375-400, Kluwer Acad., Dordrecht, Netherlands.

1223 Mazzarini, F., Isola, I., Ruggieri, G., and Boschi, C., 2010, Fluid circulation in the upper brittle crust:
1224 Thickness distribution, hydraulic transmissivity fluid inclusion and isotopic data of veins hosted in
1225 the Oligocene sandstones of the Macigno Formation in southern Tuscany, Italy: *Tectonophysics*, v.
1226 493, p. 118-138.

1227 McCaig, A.M., 1988, Deep fluid circulation in fault zones: *Geology*, v. 16, p. 867-870.

1228 McCrea, J.M., 1950, On the isotopic chemistry of carbonates and a paleotemperature scale: *J. Chem.*
1229 *Phys.*, v. 18, pp. 849-857.

1230 Merriman, R.J., and Frey, M., 1999, Patterns of very low-grade metamorphism in metapelitic rocks.
1231 In: Frey, M., Robinson, D. (Eds.), *Low-grade Metamorphism*. Blackwell Science, Oxford, pp. 61–
1232 107.

1233 Milliken, K.L., McBride, E.F., Cavazza, W., Cibin, U., Fontana, D., Picard, M.D., and Zuffa, G.G.,
1234 1998, Geochemical history of calcite precipitation in Tertiary sandstones, Northern Apennines, Italy,
1235 *in* Morad, S., ed., *Carbonate Cementation in Sandstones: International Association of*
1236 *Sedimentologists Special Publications*, v. 26, p. 213-239.

1237 Molli, G., Conti, P., Giorgetti, G., Meccheri, M., and Oesterling, N., 2000a, Microfabric study on the
1238 deformational and thermal history of the Alpi Apuane marbles (Carrara marbles), Italy: *Journal of*
1239 *Structural Geology*, v. 22, p. 1809–1825.

1240 Molli, G., Giorgetti, G., and Meccheri, M., 2000b, Structural and petrological constraints on the
1241 tectono-metamorphic evolution of the Massa unit (Alpi Apuane, NW Tuscany, Italy): *Geological*
1242 *Journal*, v. 35, p. 251–264.

1243 Molli, G., Giorgetti, G., and Meccheri M., 2002, Tectono-metamorphic evolution of the Alpi Apuane
1244 metamorphic complex. New data and constraints for geodynamic models: *Boll. Soc. Geol. It.*, Special
1245 Issue, v. 1, p. 801–817.

1246 Molli, G., 2008, Northern Apennine-Corsica orogenic system: an updated review, In: Siegesmund S.,
1247 Fugenschuh B., Froitzheim N. (Eds.), *Tectonic Aspects of the Alpine-Dinaride-Carpathian System:*
1248 *Geological Society of London Special Publication*, v. 298, p. 413–442.

1249 Molli, G., White, J.C., Kennedy, L., and Taini, V., 2011, Low-temperature deformation of limestone,
 1250 Isola Palmaria, northern Apennine, Italy—the role of primary textures, precursory veins and
 1251 intracrystalline deformation in localization: *Journal of Structural Geology*, v. 33, p. 255–270, <https://doi.org/10.1016/j.jsg.2010.11.015> .

1253 Molli, G., Carlini, M., Vescovi, P., Artoni, A., Balsamo, F., Camurri, F., Clemenzi, L., Storti, F., and
 1254 Torelli, L., 2018, Neogene 3-D structural architecture of the north-west Apennines: The role of the
 1255 low-angle normal faults and basement thrusts: *Tectonics*, <https://doi.org/10.1029/2018TC005057>,
 1256 (in press).

1257 Mongelli, F., Zito, G., Della Vedova, B., Pellis, G., Squarci, P., and Taffi, L., 1991, Geothermal
 1258 regime of Italy and surrounding areas, In: V. Cermàk and L. Rybach (Editors), *Terrestrial Heat Flow
 1259 and the Lithosphere structure*, Springer, Berlin, Germany, p. 381-394.

1260 Montanari, L., and Rossi, M., 1982, Evoluzione delle unità stratigrafico-strutturali terziarie del
 1261 nordappennino; 1. L'unità di Canetolo: *Boll. Soc. Geol. It.*, v. 101, p. 275-289.

1262 Montomoli, C., 2002, Vein development and fluid inclusion data: Insight on the evolution of the
 1263 Tuscan Nappe in the Northern Apennine: *Boll. della Soc. Geol. It.*, v. 1, Special Issue, p. 801–817.

1264 Montomoli, C., Ruggieri, G., Boiron, M.C., and Cathelineau, M., 2001, Pressure fluctuation during
 1265 uplift of the Northern Apennines (Italy): A fluid inclusions study: *Tectonophysics*, v. 341, p. 121-
 1266 139.

1267 Muchez, P., Slobodnik, M., Viaene, W.A., and Keppens, E., 1995, Geochemical constraints on the
 1268 origin and migration of palaeofluids at the northern margin of the Variscan foreland, southern
 1269 Belgium: *Sedimentary Geology*, v. 96, p. 191-200.

1270 Mullis, J., 1979, The system methane-water as a geological thermometer and barometer from the
 1271 external part of the Central Alps: *Bulletin de Mineralogie*, v. 102, p. 526–536.

1272 Mullis, J., 1987, Fluid inclusion studies during very low-grade metamorphism, In: Frey, M. (Ed.),
 1273 *Low Temperature Metamorphism*, Blackie, Glasgow, Scotland, p. 162– 199.

1274 Mullis, J., 1988, Rapid subsidence and upthrusting in the Northern Apennines, deduced by fluid
 1275 inclusion studies in quartz crystals from Porretta Terme: *Schweizerische Mineralogische und
 1276 Petrographische Mitteilungen*, v. 68, p. 157–170.

1277 Nesbitt, B.E., and Muehlenbachs, K., 1989, Origins and movement of fluids during deformation and
 1278 metamorphism in the Canadian Cordillera: *Science*, v. 245, p. 733-736.

- 1279 Okamoto, A., and Sekine, K., 2011, Textures of syntaxial quartz veins synthesized by hydrothermal
1280 experiments: *Journal of Structural Geology*, v. 33, p. 1764-1775.
- 1281 Oliver, N.H.S., 1996, Review and classification of structural controls on fluid flow during regional
1282 metamorphism: *Journal of Metamorphic Geology*, v. 14, p. 477-492.
- 1283 Oliver, N.H.S., and Bons, P.D., 2001, Mechanisms of fluid flow and fluid-rock interaction in fossil
1284 metamorphic hydrothermal systems inferred from vein-wallrock patterns, geometry and
1285 microstructure: *Geofluids* v. 1, 137-162.
- 1286 Oncken, O., 1982, Basin reconstruction by coalification studies: *Geol. Rundsch.*, 71, 579–602.
- 1287 O’Neil, J.R., Clayton, R.N., and Mayeda, T.K., 1969, Oxygen isotope fractionation in divalent metal
1288 carbonates: *The Journal of Chemical Physics*, v. 51, p. 5547–5558, [https:// doi .org /10 .1063 /1](https://doi.org/10.1063/1.1671982)
1289 [.1671982](https://doi.org/10.1063/1.1671982) .
- 1290 Parry, W.T., and Bruhn, R.L., 1987, Fluid inclusion evidence for a minimum 11 km vertical offset on
1291 the Wasatch fault, Utah: *Geology*, v. 15, p. 67-70.
- 1292 Parry, W.T., 1998, Fault-fluid compositions from fluid inclusion observations and solubilities of
1293 fracture-sealing minerals: *Tectonophysics*, v. 290, p. 1-26.
- 1294 Passchier, C.W., and Trouw, R.A.J., 2005, *Microtectonics*, second ed., Springer Verlag, Berlin,
1295 Germany, 371 p.
- 1296 Patacca, E., Sartori, R., and Scandone, P., 1990, Tyrrhenian Basin and Apenninic arc: Kinematic
1297 relations since late Tortonian times: *Mem. Soc. Geol. It.*, v. 45, p. 425–451.
- 1298 Petsch, S.T., Berner, R.A., and Eglinton, T.I., 2000, A field study of the chemical weathering of
1299 ancient sedimentary organic matter: *Organic Geochemistry*, v. 31, p. 475–487.
- 1300 Phillips, W.J., 1972, Hydraulic fracturing and mineralization: *Jour. Geol. Soc. Lond.*, v. 128, p. 337-
1301 359.
- 1302 Plesi, G., 1975, La nappe de Canetolo: *Bull. Soc. Géol. Fr.*, v. 6, p. 979-983.
- 1303 Principi, G., and Treves, B., 1984, Il sistema corso-appennino come prisma d'accrescimento. Riflessi sul
1304 problema generale del limite Alpi-Appennino: *Mem. Soc. Geol. It.*, v. 28, p. 529-576.
- 1305 Raggi, G., 1985, Neotettonica ed evoluzione paleogeografica plio-pleistocenica del Bacino de fiume
1306 Magra: *Mem. Soc. Geol. It.*, v. 30, p. 35-62.
- 1307 Ramsay, J.G., 1980, The crack-seal mechanism of rock deformation: *Nature*, v. 284, p. 135-139.

1308 Ramsay, J.G., and Huber, M.I., 1983, *The Techniques of Modern Structural Geology: Volume I:*
1309 *Strain Analysis*: London, UK, Academic Press, 307 p.

1310 Reches, Z., and Lockner, D.A., 1994, Nucleation and growth of faults in brittle rocks: *Journal of*
1311 *Geophysical Research*, v. 99, n. B9, p. 18,159-18,173.

1312 Renard, F., Gratier, J.P., and Jamtveit, B., 2000, Kinetics of crack-sealing, intergranular pressure
1313 solution, and compaction around active faults: *Journal of Structural Geology*, v. 22 (10), p. 1395-
1314 1407.

1315 Reutter, K.J., Teichmuller, M., Teichmuller, R., and Zanzucchi, G., 1981, Le ricerche sulla
1316 carbonificazione dei frustoli vegetali nelle rocce clastiche, come contributo ai problemi di
1317 paleogeotermia e tettonica nell'Appennino settentrionale: *Mem. Soc. Geol. It.*, v. 21, p. 111-126.

1318 Robert, F., Boullier, A.-M., and Firdaous, K., 1995, Gold-quartz veins in metamorphic terranes and
1319 their bearing on the role of fluids in faulting: *Journal of Geophysical Research*, v. 100, p. 12,861-
1320 12,879.

1321 Roedder, E., 1984, Fluid inclusions, *in* Ribbe, P.H., ed., *Reviews in Mineralogy*: Washington, D.C.,
1322 USA, Mineralogical Society of America, v. 12, 644 p.

1323 Roure, F., Swennen, R., Schneider, F., Faure, J.L., Ferket, H., Guilhaumou, N., Osadetz, K., Robion,
1324 P., and Vandeginste, V., 2005, Incidence and importance of tectonics and natural fluid migration on
1325 reservoir evolution in foreland fold-and-thrust belts: In: Brosse, E. (Ed.), *Oil and Gas Science and*
1326 *Technology. Revue de l'IFP* 60, pp. 67–106.

1327 Royden, L., 1988, Flexural behaviour of the continental lithosphere in Italy: Constraints imposed by
1328 gravity and deflection data: *Journal of Geophysical Research*, v. 93, p. 7747–7766.

1329 Salvini, F., 2017, Daisy 3: The Structural Data Integrated System Analyzer. Software: Rome, Italy,
1330 Roma Tre University, GeoQuTe Lab, <http://host.uniroma3.it/progetti/fralab/>.

1331 Schito, A., Andreucci, B., Aldega, L., Corrado, S., Di Paolo, L., Zattin, M., Szaniawski, R.,
1332 Jankowski, L., and Mazzoli, S., 2018, Burial and exhumation of the western border of the Ukrainian
1333 Shield (Podolia): a multi-disciplinary approach: *Basin Research*, 30, (Suppl. 1), p. 532–549.

1334 Schlische, R.W., 1995, Geometry and origin of fault-related folds in extensional settings: *American*
1335 *Association of Petroleum Geologists Bulletin*, v. 79, p. 1661–1678.

1336 Sclater, J.G., and Christie, P., 1980, Continental stretching: an explanation of the post-mid
1337 cretaceous subsidence of the central North Sea basin: *J. Geophys. Res. Solid Earth*, 85 (B7), 3711–
1338 3739.

1339 Serri, G., Innocenti, F., and Manetti, P., 1993, Geochemical and petrological evidence of the
1340 subduction of delaminated Adriatic continental lithosphere in the genesis of the Neogene-Quaternary
1341 magmatism of central Italy: *Tectonophysics*, v. 223, p. 117-147.

1342 Sibson, R.H., Moore, J.McM., and Rankin, A.H., 1975, Seismic pumping-a hydrothermal fluid
1343 transport mechanism: *Jour. Geol. Soc. Lond.*, v. 131, p. 653-659.

1344 Sibson, R.H., Robert, F., and Poulsen, K.H., 1988, High-angle reverse faults, fluid-pressure cycling,
1345 and mesothermal gold-quartz deposits: *Geology*, v. 16, p. 551-555.

1346 Sibson, R.H., 1996, Structural Permeability of fluid-driven fault-fracture meshes: *Journal of*
1347 *Structural Geology*, v. 18, p. 1031-1042.

1348 Sibson, R.H., 2000, Fluid involvement in normal faulting: *Journal of Geodynamics*, v. 29, p. 469-
1349 499.

1350 Sibson, R.H., 2004, Controls on maximum fluid overpressure defining conditions for mesozonal
1351 mineralization: *Journal of Structural Geology*, v. 26, p. 1127–1136, [https://doi.org/10.1016/j.jsg](https://doi.org/10.1016/j.jsg.2003.11.003)
1352 [.2003.11.003](https://doi.org/10.1016/j.jsg.2003.11.003).

1353 Smeraglia, L., Billi, A., Carminati, E., Cavallo, A., and Doglioni, C., 2017, Field- to nano-scale
1354 evidence for weakening mechanisms along the fault of the 2016 Amatrice and Norcia earthquakes,
1355 Italy: *Tectonophysics*, v. 712-713, p. 156-169.

1356 Smith, S.A.F., Billi, A., Di Toro, G., and Spiess, R., 2011, Principal slip zones in limestone:
1357 Microstructural characterization and implications for the seismic cycle (Tre Monti fault, Central
1358 Apennines, Italy): *Pure and Applied Geophysics*, v. 168, p. 1-27, doi:10.1007/s00024-011-0267-5.

1359 Solum, J.G., and Huisman, B.A.H., 2016, Toward the creation of models to predict static and dynamic
1360 fault-seal potential in carbonates: *Petroleum Geoscience*, v. 23 (1), p. 70-91.

1361 Storti, F., 1995, Tectonics of the Punta Bianca promontory: Insights for the evolution of the Northern
1362 Apennines-Northern Tyrrhenian Sea basin: *Tectonics*, v. 14, p. 832–847.

1363 Teichmüller, M., and Teichmüller, R., 1967, Diagenesis of coal (coalification), In: *Diagenesis in*
1364 *Sediments* by Larsen G. and Chilingar G.V. Edn., *Developments in Sedimentology*, Elsevier, v. 8, p.
1365 391-415.

1366 Teichmüller, M., 1987, Organic material and very low grade metamorphism, In: Frey M. (Ed.), Low-
1367 temperature Metamorphism, Chapman and Hall, Glasgow, Scotland, p. 114-161.

1368 Thomson, S.N., Brandon, M.T., Reiners, P.W., Zattin, M., Isaacson, P.J., and Balestrieri, M.L., 2010,
1369 Thermochronologic evidence for orogeny-parallel variability in wedge kinematics during extending
1370 convergent orogenesis of the Northern Apennines, Italy: *Geol. Soc. Am. Bull.*, v. 122, p. 1160-1179.

1371 Tissot, B.P., and Welte, D.H., 1984, From Kerogen to Petroleum, In: *Petroleum Formation and*
1372 *Occurrence*, Springer, Berlin Heidelberg, Germany, p. 160-198.

1373 Torabi, A., and Berg, S.S., 2011, Scaling of fault attributes: a review: *Marine and Petroleum Geology*,
1374 v. 28, p. 1444–1460, <http://dx.doi.org/10.1016/j.marpetgeo.2011.04.003>.

1375 Vandeginste, V., Swennen, R., Allaeys, M., Ellam, R.M., Osadetz, K., and Roure, F., 2012,
1376 Challenges of structural diagenesis in foreland fold-and-thrust belts: a case study on paleofluid flow
1377 in the Canadian Rocky Mountains west of Calgary: *Mar. Petrol. Geol.*, v. 35, p. 235-251.

1378 Veizer, J., Ala, D., Azmy, K., Bruckschen, P., Buhl, D., Bruhn, F., Carden, G.A.F., Diener, A.,
1379 Ebner, S., Godderis, Y., Jasper, T., Korte, C., Pawellek, F., Podlaha, O.G., and Strauss, H., 1999,
1380 $^{87}\text{Sr}/^{86}\text{Sr}$, $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ evolution of Phanerozoic seawater: *Chemical Geology*, v. 161, p. 59-88.

1381 Vermilye, J.M., and Scholz, C.H., 1998, The process zone: A microstructural view of fault growth:
1382 *Journal of Geophysical Research*, v. 103, n. B6, p. 12,223-12,237.

1383 Vescovi, P., 1998, Le Unità Subliguri dell'alta Val Parma (Provincia di Parma): *Atti Ticinesi di*
1384 *Scienze della Terra*, v. 40, p. 215-231.

1385 Vescovi, P., 2005, The Middle Miocene Mt. Ventasso – Mt. Cimone arcuate structure of the Emilia
1386 Apennines: *Bollettino della Società Geologica Italiana*, v. 124, p. 53-67.

1387 Vilasi, N., Malandain, J., Barrier, L., Callot, J-P., Amrouch, K., Guilhaumou, N., Lacombe, O.,
1388 Muska, K., Roure, F., and Swennen, R., 2009, From outcrop and petrographic studies to basin-scale
1389 fluid flow modelling: the use of the Albanian natural laboratory for carbonate reservoir
1390 characterisation. *Tectonophysics*, v. 474 (1-2), p. 367-392.

1391 Virgo, S., Abe, S., and Urai, J.L., 2014, The evolution of crack seal vein and fracture networks in an
1392 evolving stress field: Insights from Discrete Element Models of fracture sealing: *Journal of*
1393 *Geophysical Research, Solid Earth*, v. 119, p. 8708–8727.

1394 Walther, J.V., and Orville, P.M., 1982, Volatile production and transport in regional metamorphism:
1395 *Contrib. Mineral. Petrol.*, v. 79, p. 252-257.

1396 Withjack, M.O., Olson, J., and Peterson, E., 1990, Experimental models of extensional forced folds:
 1397 American Association of Petroleum Geologists Bulletin, v. 74, p. 1038–1054.

1398 Withjack, M.O., and Callaway, S., 2000, Active Normal Faulting Beneath a Salt Layer: An
 1399 Experimental Study of Deformation Patterns in the Cover Sequence: American Association of
 1400 Petroleum Geologists Bulletin, v. 84, n. 5, pp. 627–651.

1401 Wüstefeld, P., Hilse, U., Lüders, V., Wemmer, K., Koherer, B., and Hilgers, C., 2017, Kilometer-
 1402 scale fault-related thermal anomalies in tight gas sandstones: Marine and Petroleum Geology, v. 86,
 1403 p. 288-303.

1404 Zattin, M., Picotti, V., and Zuffa, G.G., 2002, Fission-track reconstruction of the front of the Northern
 1405 Apennine thrust wedge and overlying Ligurian unit: American Journal of Science, v. 302, p. 346–
 1406 379.

1407

1408 **FIGURE CAPTIONS**

1409 Figure 1. (A) Location of the study area, in the inner part of the Northern Apennines, Italy. (B)
 1410 Tectonic sketch map of the region where the Lunigiana extensional basin developed (modified from
 1411 Bernini, 1997); the black and white traces indicate the geologic cross-section represented in (D) and
 1412 the seismic line in Figure 2, while the black and white rectangle represents the area shown in Figure
 1413 3A. NLB—Northern Lunigiana basin; SLB—Southern Lunigiana basin. (C) Schematic column of
 1414 the inner Northern Apennines stratigraphy. (D) Geological cross-section passing through the studied
 1415 segment of the Compione extensional fault zone (after Bernini and Papani, 2002). OTT—Ottone
 1416 Flysch Formation; MAC—Macigno Sandstones Formation; SUBL.—Subligurian Succession; m
 1417 (a.s.l.)—meters above sea level.

1418 Figure 2. (A) Depth converted seismic line and (B) line drawing highlighting the geometry of the
 1419 Compione Fault, Northern Apennines, Italy, in the subsurface (segment of a seismic reflection profile
 1420 acquired from Eni S.p.A.). The seismic reflection profile is depth converted with the Move Software
 1421 (provided by Midland Valley Exploration Ltd.). By using the “User defined tables/checkshots”
 1422 method, 13 Time/Interval Velocity tables were defined for the common mid-point (CMP) gathers

1423 (CMP1–CMP13 in A). The interval velocities derive by Dix conversion of stacking velocity used for
1424 the processing sequence as provided for this seismic profile. The Dix interval velocities are in
1425 between 3130 and 7644 m/sec, they had an overall increase down to 5 sec (two-way time) in the
1426 seismic reflection profile before depth conversion. $v=h$ —vertical equals horizontal.

1427 Figure 3. (A) Simplified structural map of the Northern Lunigiana, Northern Apennines, Italy,
1428 extensional basin showing the main extensional fault zones that constitute the Northern Lunigiana
1429 extensional fault system, bounding the basin to the NE. The black and white trace A–B and the
1430 rectangle refer to (B) and (C), respectively. (B) Schematic geological cross-section showing the
1431 geometry of the extensional fault propagation fold in the footwall damage zone of the Compione
1432 extensional fault zone, overprinting the out-of-sequence (O-of-Seq) thrust-related fold geometry. (C)
1433 Structural map of the study area showing structural and sampling sites (S).

1434 Figure 4. (A) Geologic cross-section across the Compione fault damage zone, Northern Apennines,
1435 Italy, and stereographic projections of structural data (Schmidt lower hemisphere). The trace of the
1436 section is indicated in Figure 3C. Ex.F. c.i.—Extensional Faults contouring interval; S-s.F. c.i.—
1437 Strike-slip Faults contouring interval; FW non-tilted PZ c.i.—Footwall non-tilted Process Zone
1438 contouring interval; meters a.s.l.—meters above sea level. (B) Outcrop picture and line drawing of
1439 the hanging wall damage zone at Site 2, showing synthetic extensional faults in the Ottone Flysch
1440 Formation. (C) Outcrop picture and line drawing of the footwall damage zone at Site 4, showing the
1441 major E-W footwall splay extensional fault zone; red areas highlight the master slip surface.

1442 Figure 5. (A) Outcrop picture and line drawing of conjugate shear veins and extensional faults in the
1443 footwall damage zone to background transition (44°19'56.65"N 10°3'41.49"E). (B) Stereographic
1444 projection of structural data in this area; dashed lines represent bedding. (C) Outcrop picture and line
1445 drawing of Site 5; dashed white line corresponds to bedding, white lines highlight conjugate shear
1446 veins and faults, and dashed black lines indicate late cataclastic faults; white (black) arrows indicate
1447 kinematics before (after) bedding rotation. (D) Stereographic projection of structural data collected

1448 at this site. (E) Detail of (C) showing a shear vein reactivated along a synthetic subsidiary fault and
1449 later crosscut along an antithetic one. (F) Slickenfibers on a synthetic subsidiary fault showing
1450 multiple slip directions; coin for scale.

1451 Figure 6. (A) Stereographic projections (Schmidt lower hemisphere) of footwall shear veins and
1452 extensional subsidiary conjugate faults, measured at sites 4 and 5, and separated for sectors
1453 compartmentalized by late-stage large-scale cataclastic faults. In (B) data are restored to horizontal
1454 using the related bedding orientation; black (white) dots indicate normal (reverse) kinematics; reverse
1455 kinematics are apparent because of shear reactivation after bedding rotation. S—sampling sites.

1456

1457 Figure 7. Microphotographs of veins hosted in the footwall and hanging wall damage zones of the
1458 Compione extensional fault. (A) Prehnite (Prh) crystals overgrown by quartz (Qz) and MC1 calcite;
1459 MC2 calcite grew in dissolution fractures outlined by small dotted white lines (cross-polarized light).
1460 (B) Prh growing in columnar-radiating structures, characterized by an abrupt change in luminescence
1461 color and MC1A calcite crystals recrystallized in MC1B calcite along crystal fractures,
1462 cathodoluminescence (CL) image. (C) Vein composed of euhedral to subhedral Qz rims and MC1
1463 replacive (rep) and rhombohedric (rho) calcite crystals in the center (cross-polarized light). (D) CL
1464 detail of (C) showing MC1A and MC1B replacing prismatic Prh crystals. (E) Stained hand sample
1465 displaying pink MC1 calcite and composed of five fracturing-sealing increments and/or shear
1466 reactivations. (F) Detail of (E) illustrating micrometric wide slip surfaces formed by coseismic slip
1467 (cross-polarized light). (G) Shear bands composed of recrystallized Qz subgrains and replacive MC1
1468 calcite crystals (cross-polarized light). (H) Isomorphous replacement of Prh crystals by MC2 calcite
1469 (CL). (I) MC1A isomorphous replacements of prehnite crystals cut by late microfractures cemented
1470 with MC2 calcite (CL). (J) Sample near the Compione fault core showing disaggregated texture
1471 cemented by MC2 calcite. (K) Cross-polarized light image showing OC1 “dirty” and OC2 “clear”
1472 calcites textures. (L) CL image highlighting contrast in luminescence colors between OC1 and OC2

1473 calcites from dull red to bright red and pressure solution affecting both OC1 and OC2. (M) Host rock
1474 clast in OC1 and late fractures filled by OC2 calcite and quartz (qz). (N) CL image showing detail of
1475 a bedding parallel vein where OC2 calcite, associated with qz, isomorphously replaces prehnite
1476 crystals. (O) Thin section scan of a breccia-vein composed of OC2 calcite cement collected in
1477 extensional S-C (schistosity-cisaillement) structures near the Compione fault core. MC—Macigno
1478 Sandstones Formation calcite cements; OC—Ottone Flysch Formation calcite cements.

1479 Figure 8. Stable isotope data. (A) $\delta^{18}\text{O}$ vs. $\delta^{13}\text{C}$ plot of calcite from veins and host rocks; the dashed
1480 grey rectangle indicates the range of isotopic values of Cretaceous limestone (Lms.) (after Veizer et
1481 al., 1999) and the dashed black rectangle indicates those of Late Oligocene– Early Miocene Macigno
1482 Sandstones Formation (after Milliken et al., 1998). (B, C) Plots of $\delta^{18}\text{O}$ (B) and $\delta^{13}\text{C}$ (C) vs. distance
1483 (m) from the Compione fault core, Northern Apennines, Italy. FC—Fault core; VSMOW—Vienna
1484 standard mean ocean water; VPDB—Vienna Pee Dee belemnite. In the legend Ottone refers to calcite
1485 cements hosted in the Ottone Flysch Formation.

1486 Figure 9. Microphotographs in plane polarized light of fluid inclusion types, in the analyzed veins.
1487 (A) Monophase gaseous inclusions Q2 cooled at $-100\text{ }^{\circ}\text{C}$. (B) Raw spectra of Raman analysis
1488 performed on Q2 inclusions showing peaks corresponding to CH_4 . (C) Fluid inclusion assemblages
1489 (FIA) of Q1 aqueous biphasic inclusions in quartz from a footwall vein. (D) FIA of aqueous biphasic
1490 inclusions with a negative crystal shape in MC1. (E) FIA of aqueous biphasic inclusions in MC2. (F)
1491 FIA of aqueous monophase inclusions in OC2, overprinting a OC1 vein, big inclusions in OC2 are
1492 biphasic aqueous. MC—Macigno Sandstones Formation calcite cements; OC—Ottone Flysch Forma-
1493 tion calcite cements.

1494 Figure 10. Frequency distribution plots of homogenization (T_{htot}) and ice melting (T_{mice})
1495 temperatures. (A) T_{htot} of Q2 monophase gaseous inclusions in quartz. (B) T_{htot} of biphasic aqueous
1496 inclusions in footwall veins. (C) T_{mice} of biphasic aqueous inclusions in footwall veins. (D) T_{htot} of
1497 biphasic aqueous inclusions in hanging wall veins. (E) T_{mice} of monophase and biphasic aqueous

1498 inclusions in hanging wall veins. Macigno Calcite—Macigno Sandstones Formation calcite cements;
1499 OC—Ottone Flysch Formation calcite cements; Footwall veins, hosted in the Macigno Sandstones
1500 Formation, from Sites 4, 5, 7, and 13 in Figures 3A and 3C; Hanging wall veins, hosted in the Ottone
1501 Flysch Formation, from Sites 2, 3, 8, 9, and 10 in Figure 3C.

1502 Figure 11. Plot of homogenization temperatures (T_{htot}) vs. ice melting temperatures (T_{mice}) of
1503 measured fluid inclusions. Ice melting temperatures are reported along with salinity in NaCl eq. wt%,
1504 according to Bodnar (1993). Macigno Calcite—Macigno Sandstones Formation calcite cements;
1505 Ottone Calcite—Ottone Flysch Formation calcite cements; Footwall veins, hosted in the Macigno
1506 Sandstones Formation, from Sites 4, 5, 7, and 13 in Figures 3A and 3C; Hanging wall veins, hosted
1507 in the Ottone Flysch Formation, from Sites 2, 3, 8, 9, and 10 in Figure 3C.

1508 Figure 12. Burial and thermal history of the Macigno Sandstones Formation, Northern Apennines,
1509 Italy, shaded in grey, using a geothermal gradient of 25 °C/km up to the end of the Miocene and of
1510 30 °C/km since the Pliocene (P.). Striped areas represent the stratigraphic range of Site 14.

1511 Figure 13. Trishear predicted geometry of the Compione Fault, Northern Apennines, Italy, using the
1512 method of Jin and Groshong (2006). Monocline width is around twice the footwall monocline width
1513 assuming a linear velocity field in the trishear zone. Striped unit above the basement correspond to
1514 Burano Evaporites.

1515 Figure 14. Sketch illustrating the typical paragenetic evolution of conjugate veins in the footwall of
1516 the Compione Fault, Northern Apennines, Italy. (A) Pristine crystal textures of the initial infilling
1517 event. (B) Microstructural modifications after shear reactivation. (C) After late-stage reactivation.
1518 (D) Schematic table showing the cyclic events, indicating the relative chronology of fracturing,
1519 precipitation, and dissolution processes, along with their temperature range. (E) Schematic evolution
1520 of fluid pressure, reported as ratio to lithostatic pressure, and shear stress in time relative to earthquake
1521 rupturing and to shearing, fracturing, precipitation, and dissolution processes (D) in fault-related

1522 veins. Prh—prehnite, Qz—quartz; MC—Macigno Sandstones Formation calcite cements. P—
1523 pressure; tau—shear stress; EQ— seismic rupture.

1524 Figure 15. Data are from Sites 4, 5, 7, and 13 in the footwall damage zone (MC) and from Sites 2, 3,
1525 8, 9, and 10 in the hanging wall damage zone (OC); see Figures 3A and 3C. (A) Isochores are plotted
1526 for the different fluid inclusion types. Peak burial derived from vitrinite reflectance modelling is used
1527 as an independent constraint to calculate a maximum trapping temperature for MC1A (shaded area).
1528 (B) Oxygen isotope fractionation during equilibrium calcite precipitation. The fluid oxygen
1529 composition is reported as a function of calcite oxygen composition and temperature. Square symbols
1530 and error bars represent the mean value and range of footwall calcite MC1A, MC1B, and MC2. OC2
1531 calcite is represented by the white circles. Dashed lines describe the inferred fluid evolution during
1532 time. MC—Macigno calcite cements; OC—Ottone Flysch Formation calcite cements; VR (Ro%)—
1533 Maximum burial depth estimated from vitrinite reflectance measurements; VSMOW—Vienna stan-
1534 dard mean ocean water.

1535 Figure 16. Phase diagrams for calcium and alumina minerals in terms of $\text{Ca}^{2+}/(\text{H}^+)^2$ activity ratio
1536 and of aqueous SiO_2 activity at 200 °C (A) and at 150 °C (B).

1537 Figure 17. Cartoon showing the proposed evolution of the Compione Fault, Northern Apennines,
1538 Italy. (A) Onset of extensional deformations overprinting the previously stacked Sub-Ligurian and
1539 Ligurian thrust sheets onto the Macigno foredeep sandstones (MAC). A wide process zone forms,
1540 mainly consisting of conjugate shear fractures that enhance the advection of hydrothermal fluids and
1541 rapid cementation. (B) Extensional fault-propagation folding during upward fault migration, causing
1542 bending of part of the process zone closer to the prospect master shear zone and reactivation of
1543 conjugate shear veins as both synthetic and antithetic subsidiary extensional faults. (C) Shear
1544 localization, fault breakthrough and accommodation of most of the displacement in the fault core,
1545 hanging wall, and footwall subsidiary fault zones. Black arrows are formational fluids, white and
1546 black ones are high-temperature low-salinity hydrothermal fluids, white and grey ones are meteoric

1547 fluids and black and grey ones are low-temperature high-salinity fluids. OTT—Ottone Flysch
1548 Formation; Prh—prehnite, Qz—quartz; MC—Macigno calcite cement; OC—Ottone Flysch Forma-
1549 tion calcite cements.

1550 Table 1. Suummary of petrographic observations and stable isotope analyses results of the different
1551 carbonate cements.

1552 Table 2. Summary of petrographic observations and microthermometry analyses results of quartz and
1553 carbonate cements.

1554 Table 3. Summary of results of vitrinite reflectance measurements.

1555

1556 Figure DR1. Seismic reflection profile of Fig. 2 before being depth-converted.

1557 Figure DR2. (A-B) Outcrop pictures of coarse sandstones strata in footwall damage zone Site 5. White
1558 dotted lines represent bedding, white lines are conjugate extensional shear veins and fractures with
1559 σ_1 orthogonal to bedding, black dashed lines are late-stage extensional shear fractures and red dashed
1560 lines indicate strike-slip shear veins and fractures. White arrows indicate pre-bedding rotation
1561 kinematics while black arrows indicate kinematics after bedding rotation. (C) Stained hand specimen
1562 of footwall fault-related shear vein; black dotted lines separate different opening events. (D) XPL
1563 image showing, as in C, multiple openings (separated by white dotted lines), of which some interested
1564 by shearing and quartz recrystallization. (E) PPL image of vein showing both MC1A and MC2
1565 replacive calcite crystals along with quartz and rhombohedric MC1A. (F-G-H) CL images showing
1566 progressive prehnite dissolution and replacement by MC calcites.

1567 Figure DR3. Fluid inclusion assemblage in quartz of footwall damage zone fault-related shear veins
1568 showing aqueous inclusion (white), gaseous CH₄ inclusions (black) and an inclusion containing
1569 liquid H₂O and gaseous CH₄ (black and white). In (A) at 20 °C and in (B) at -100 °C.

1570

1571 ¹GSA Data Repository item 2018xxx, including structural sites and samples list, and stable isotopic,
1572 microthermometric and SEM-EDS analyses results, is available online at
1573 www.geosociety.org/pubs/ft20XX.htm, or on request from editing@geosociety.org or Documents
1574 Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.

Lucca et al.
Fig. 1

Lucca et al.
Fig. 2

Lucca et al.
Fig. 3

Lucca et al.
Fig. 4

Lucca et al.
Fig. 5

Lucca et al.
Fig. 6

Lucca et al.
Fig. 7

Lucca et al.
Fig. 8

Lucca et al.
Fig. 9

Lucca et al.
Fig. 10

Lucca et al.
Fig. 11

Lucca et al.
Fig. 12

Lucca et al.
Fig. 13

Lucca et al.
Fig. 14

Lucca et al.
Fig. 15

Lucca et al.
Fig. 16

Lucca et al.
Fig. 17

TABLE 1. SUMMARY OF PETROGRAPHIC OBSERVATIONS AND STABLE ISOTOPES ANALYSIS RESULTS OF THE DIFFERENT CARBONATE CEMENTS

Cement	Texture description	$\delta^{18}\text{O}$ ‰ (V-SMOW)			$\delta^{13}\text{C}$ ‰ (V-PDB)		
		range	mean	st. dev.	range	mean	st. dev.
Ottone host rock (n=4)	Anhedral calcite microcrystals in matrix; Non luminescent in clay-silt size beds; dull luminescent in silt-fine sands beds.	+ 26.9 to + 27.5	+ 27.2	$\pm$ 0.2	+ 1.7 to + 2.0	+ 1.9	$\pm$ 0.1
Ottone calcite 1 (n=26)	Well developed, rhomboedric “dirty” crystals with abundant twinning Type I, undulose extinction, intercrystalline slip and dissolution surfaces; non to dull luminescence.	+ 23.5 to + 26.9	+ 25.7	$\pm$ 0.7	+ 1.6 to + 2.4	+ 1.9	$\pm$ 0.2
Ottone calcite 2 (n=16)	Clean translucent crystals in late fractures with rare twinning Type I; non to dull luminescence.	+ 13.1 to + 19.7	+ 16.5	$\pm$ 1.6	- 0.5 to + 2.2	+ 1.8	$\pm$ 0.7
Macigno host rock (n=3)	Anhedral calcite microcrystals in matrix of fine clay-silt beds; red to orange luminescence.	+ 13.9 to + 16.0	+ 14.7	$\pm$ 1.1	- 3.2 to - 0.7	- 2.1	$\pm$ 1.3
Macigno calcite 1 (n=21)	Crystals characterized by twinning Type I and rare Type II, displaying rhomboedric and isomorphously replacing Prh; red (MC1A) to orange (MC1B) luminescence.	+ 11.8 to + 14.0	+ 13.1	$\pm$ 0.6	- 5.8 to - 0.9	- 3.3	$\pm$ 1.8
Macigno calcite 2 (n=18)	Crystals in late fractures crosscutting Prh, Qz, MC1 and locally replacing Prh; orange to yellow bright luminescence.	+ 13.4 to + 17.9	+ 15.8	$\pm$ 1.2	- 7.1 to - 0.4	- 2.7	$\pm$ 2.0

**TABLE 2. SUMMARY OF PETROGRAPHIC OBSERVATIONS AND MICROTERMOMETRY ANALYSIS RESULTS
OF QUARTZ AND CARBONATE CEMENTS**

Cement	FIA origin & type	Homogenization temperature (°C) range/mean	Ice melting temperature (°C) range/mean	Salinity (eq. wt% NaCl) range/mean
Macigno quartz 1	Primary/Pseudosecondary two- phase aqueous	127 to 212 / 157	- 4.8 to - 11.4 / - 7.2	7.6 to 15.4 / 10.7
Macigno quartz 2	Primary Monophase gaseous CH ₄	- 83 to -89 / -88	N.D.	N.D.
Macigno calcite 1A	Primary/Pseudosecondary two- phase aqueous	178 to 198 / 189	- 0.8 to - 6.2 / -2.7	1.4 to 9.3 / 4.5
Macigno calcite 1B	Primary/Pseudosecondary two- phase aqueous	140 to 161 / 151	- 0.8 to - 17.2 / - 5	1.4 to 20.3 / 7.9
Macigno calcite 2	Primary two-phase aqueous	69 to 115 / 88	- 0.7 to - 20.7 / -8.7	1.2 to 22.9 / 12.5
Ottone calcite 1	Reequilibrated two- and one-phase aqueous	< 50 to 113	0 to - 1.6 / - 0.8	0 to 2.7 / 1.4
Ottone calcite 2	Primary two- and one-phase aqueous	< 50 to 115	- 1.3 to - 6.9 / - 4.4	2.2 to 10.4 / 7

TABLE 3. SUMMARY OF RESULTS OF VITRINITE REFLECTANCE MEASUREMENTS

Site n.	Sample n.	R ₀ mean	R ₀ S.D.	Data n.
14	1	0.70	0.08	11
14	2	0.66	0.12	8
14	3	0.69	0.10	19
14	4	0.67	0.12	7
4	1	0.61	0.04	8
4	3	0.42	0.06	40
4	4	0.42	0.10	12
5	1	0.46	0.07	29
5	2	0.55	0.05	12
5	3	0.49	0.06	11