Accepted Manuscript

Title: ER Ca²⁺ release and store-operated Ca²⁺ entry - partners in crime or independent actors in oncogenic transformation?

Authors: Cristina Pierro, Flore Sneyers, Geert Bultynck, H. Llewelyn Roderick

PII: S0143-4160(19)30128-9
DOI: <https://doi.org/10.1016/j.ceca.2019.102061>
Article Number: 102061

Reference: YCECA 102061

To appear in: *Cell Calcium*

Received date: 3 May 2019
Revised date: 21 June 2019
Accepted date: 26 June 2019

Please cite this article as: { <https://doi.org/>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ER Ca^{2+} release and store-operated Ca^{2+} entry - partners in crime or independent actors in oncogenic transformation?

Cristina Pierro^{#,1}, Flore Sneyers^{#,2}, Geert Bultynck², H. Llewelyn Roderick^{1,*}

¹KU Leuven, Laboratory of Experimental Cardiology, Department of Cardiovascular Sciences, Campus Gasthuisberg O/N-I bus 802, Herestraat 49, 3000 Leuven, Belgium

²KU Leuven, Laboratory of Molecular & Cellular Signaling, Department of Cellular & Molecular Medicine, Campus Gasthuisberg O/N-I bus 802, Herestraat 49, 3000 Leuven, Belgium

*For correspondence: Llewelyn.roderick@kuleuven.be

#shared first author

Graphical abstract

Graphical Abstract Legend.

The Graphical Abstract indicates the Hallmarks of Cancer and their relationship with a cell signalling module involving store operated Ca^{2+} entry (SOCE) based on STIM/Orai and InsP_3 dependent Ca^{2+} release from the ER (via InsP_3Rs ; IICR). We propose that SOCE/IICR is dynamically modulated to meet the needs of the cell as it transitions through the various stages of oncogenic transformation. In turn, SOCE/IICR influences the physiology of the cell according to the transformation stage.

Highlights:

- Ca^{2+} signalling pathways control cellular life and death decisions
- Ca^{2+} signalling is remodelled during oncogenic transformation
- Ca^{2+} signalling contributes to oncogenic transformation
- Ca^{2+} release via ER InsP_3R channels is remodelled in transformed cells
- STIM/ORAI-associated store operated Ca^{2+} entry is altered in cancer
- ER Ca^{2+} release and store operated Ca^{2+} entry synergise to determine cell fate

Abstract

Ca^{2+} is a pleiotropic messenger that controls life and death decisions from fertilisation until death. Cellular Ca^{2+} handling mechanisms show plasticity and are remodelled throughout life to meet the changing needs of the cell. In turn, as the demands on a cell alter, for example through a change in its niche environment, Ca^{2+} handling systems may be targeted to sustain the remodelled cellular state. Nowhere is this more apparent than in cancer. Oncogenic transformation is a multi-stage process during which normal cells become progressively differentiated towards a cancerous state that is principally associated with enhanced proliferation and avoidance of death. Ca^{2+} signalling is intimately involved in almost all aspects of the life of a transformed cell and alterations in Ca^{2+} handling have been observed in cancer. Moreover, this remodelling of Ca^{2+} signalling pathways is also required in some cases to sustain the transformed phenotype. As such, Ca^{2+} handling is hijacked by oncogenic processes to deliver and maintain the transformed phenotype. Central to generation of intracellular Ca^{2+} signals is release of Ca^{2+} from the endoplasmic reticulum intracellular (ER) Ca^{2+} store via inositol 1,4,5-trisphosphate receptors (InsP_3Rs). Upon depletion of the ER of Ca^{2+} , store-operated Ca^{2+} entry (SOCE) across the plasma membrane occurs via Stim-gated Orai channels. SOCE serves to both replenish stores but also sustain Ca^{2+} signalling events. Here, we will discuss the role and regulation of these two signalling pathways and their interplay in oncogenic transformation.

Introduction

Ca^{2+} is a multifunctional second messenger that participates in almost all aspects of cellular physiology, intervening in processes including fertilisation, cell division, muscle contraction, synaptic transmission, hormone release and immune responses [1]. To perform this signalling function, variations in its levels in the cytosol in the form of complex patterns that vary in space, time and location are generated. These changes in intracellular Ca^{2+} are brought about and shaped through the concerted action of a toolkit of Ca^{2+} handling proteins comprising Ca^{2+} channels, binding proteins, pumps and exchangers. Specifically, through the gating of Ca^{2+} channels on the plasma membrane and on intracellular Ca^{2+} stores, Ca^{2+} flows down its concentration gradient, entering the cytosol where its concentration is increased. Ca^{2+} signals are then terminated by Ca^{2+} pumps and exchangers, which extrude Ca^{2+} from the cell or sequester it back into the internal stores.

Ca^{2+} is primarily stored in the endoplasmic reticulum (ER) and to a lesser extent in the Golgi apparatus and lysosomes. Ca^{2+} is released from ER stores predominantly via inositol 1,4,5-trisphosphate (InsP_3) receptor (InsP_3R) and ryanodine receptor (RyR) channels [2]. InsP_3Rs open in response to InsP_3 , produced following hydrolysis of phosphatidyl inositol 4,5-bisphosphate (PIP_2) in response to activation of phospholipase C by extracellular agonists [3]. Three InsP_3R isoforms ($\text{InsP}_3\text{R}1$, 2 and 3) are expressed in mammalian systems that are assembled in tetramers or heterotetramers, which display subtly different sensitivities to InsP_3 and Ca^{2+} itself [4]. The relative proportion of these isoforms varies according to tissue type, cellular location and development stage, thereby allowing InsP_3 signalling to fulfil specific cellular needs. RyRs are also expressed as three isoforms, which in contrast to InsP_3Rs generally form homotetrameric channels that only require Ca^{2+} for gating. Skeletal muscle RyRs (Ryr1) can however be activated via a conformational change in their associated plasmalemmal dihydropyridine receptors (DHPRs, otherwise known as $\text{Cav}1.1$), which occurs following cellular depolarisation. The Ca^{2+} sensitivity of both InsP_3Rs and RyR is bell-shaped, with a stimulatory effect of $[\text{Ca}^{2+}]$ in the physiological range ($\sim 100 \text{ nM} - 1 \mu\text{M}$) that is

maximal at 1 μM , above which activity is inhibited. However, the $[\text{Ca}^{2+}]_{\text{cyt}}$ levels at which Ca^{2+} -dependent inhibition and activation occur show isoform-specific differences. InsP_3Rs also underlie Ca^{2+} release from other intracellular Ca^{2+} storage organelles, including lysosomes and Golgi apparatus [5]. A further pathway involved in Ca^{2+} release from intracellular stores involves the second messenger NAADP, which stimulates Ca^{2+} release from lysosomes *via* two-pore channels (TPCs) [6]. In addition, members of the transient receptor potential family of channels (TRP), mediate Ca^{2+} release from the lysosomes, e.g. the mucolipins (TRPML), which are gated by PIP_2 [7]. Plasma membrane (PM) Ca^{2+} channels contributing to Ca^{2+} signals (the so-called 'ON-reactions') include voltage-gated Ca^{2+} channels (VGCCs; e.g. P, Q, L and N-type voltage Ca^{2+} channels primarily expressed in excitable cells), store-operated channels (SOC; ORAI1,2,3), TRP channels and ligand-gated Ca^{2+} channels (e.g. nACh, NMDA and purinergic receptors) [8]. Responsible for the termination of Ca^{2+} signals (the 'OFF-reactions') are plasma membrane Ca^{2+} ATPases (PMCA), sarco-endoplasmic reticulum Ca^{2+} ATPases (SERCAs) on the ER, mitochondrial Ca^{2+} uptake mechanisms via VDAC1 and MCU channels and Ca^{2+} pumps/exchangers on other organelles such as the Golgi apparatus (Secretory Pathway Calcium ATPases, SPCAs) and lysosomes [9,10]. Where rapid high capacity Ca^{2+} extrusion is required as in electrically excitable cells, Sodium Calcium Exchangers (NCX), which extrude one Ca^{2+} for three Na^+ , are expressed [11].

In contrast to the finite intracellular Ca^{2+} stores, the extracellular space represents an inexhaustible source of Ca^{2+} . The interplay between ER and PM Ca^{2+} channels is important for cytosolic Ca^{2+} dynamics. While short-lived Ca^{2+} transients in non-excitable cells primarily involve Ca^{2+} release from intracellular stores, sustained Ca^{2+} signalling events also require Ca^{2+} influx from the extracellular space. Ca^{2+} influx across the PM can be directly activated via PM-mediated mechanisms or through store-operated Ca^{2+} entry (SOCE). SOCE is mediated via the Orai-channel family (Orai1-3) of highly Ca^{2+} -selective, hexameric channels, which are gated by stromal interaction molecule 1 and 2 (STIM1 and STIM2) [12,13]. STIM proteins are ER-located Ca^{2+} sensors that detect ER Ca^{2+} content through their ER luminal N-terminal EF-hand and activate Orai channels through a physical

interaction via specialized domains located in their cytosolic C-terminal tail [14]. Under resting conditions, STIM proteins are homogeneously distributed across the ER membrane. Upon ER Ca^{2+} depletion, STIM proteins aggregate and cluster at ER-PM contact sites, where they recruit Orai channels and provoke their opening [15]. STIM2 has a lower affinity for Ca^{2+} than STIM1, enabling STIM2 to sense smaller decreases in ER Ca^{2+} content [16]. As such, STIM2 is defined as a regulator of basal $[\text{Ca}^{2+}]_{\text{ER}}$, in contrast to STIM1, which responds to more robust ER store depletion, for example following agonist stimulation [17]. Two splice variants of STIM2 have recently been described which act to promote (STIM2.2) or inhibit (STIM2.1) SOCE [18,19]. As STIM1 and 2 co-localise at punctae, SOCE can thus be finely tuned on the basis of the STIM1/STIM2 ratio and the predominant STIM2 splice variant [20,21]. Recent studies confirm the role of STIM2 as an important modulator of STIM1/Orai1 coupling [22,23].

STIM proteins play roles independent of their gating of the Orai family of channels. They have also been reported to activate SOCE *via* TRP channels, including TRPC1, 3, 4 and 5 [24]. Interestingly, Orai1 is required for TRPC1 function *via* a dynamic signalling complex involving TRPC1, STIM1 and Orai1, localised in ER-PM junctions [25,26]. At rest, TRPC1 is docked in a trafficking vesicle at the PM. Following store depletion, Orai1 associates with and is activated by STIM1, leading to Ca^{2+} entry. This causes insertion of TRPC1 into the PM, where it is subsequently gated and activated by STIM1, allowing TRPC1-mediated Ca^{2+} influx. Furthermore, Ca^{2+} currents of Orai channels and TRPC channels seem to stimulate different cellular processes. For example, Orai1-mediated Ca^{2+} influx activates NFAT through a local $[\text{Ca}^{2+}]$ increase near the Orai1 channel, which in turn activates calcineurin, triggering dephosphorylation and nuclear translocation of NFAT [27,28]. Ca^{2+} entry *via* TRPC1, however, does not affect NFAT dephosphorylation but contributes to the activation of NF κ B [29,30].

The Orai family is not only a key player in SOCE, it is an equally essential component of agonist-activated, store-independent Ca^{2+} entry pathway [31]. This pathway is mediated through the

arachidonic acid-regulated Ca^{2+} -selective (ARC) channel, which is formed by a combination of Orai1 and Orai3 subunits and is regulated by a population of STIM1 molecules localised in the PM [32].

Changes in intracellular $[\text{Ca}^{2+}]$ are transduced into a cellular response via a number of Ca^{2+} binding/sensor proteins. Ca^{2+} binds to these proteins via specialised Ca^{2+} binding motifs including EF hands and C2 domains. Through variations in affinity of these domains and localisation of the proteins, these signal transducers contribute specificity in Ca^{2+} signalling mechanisms. Notable examples amongst these Ca^{2+} signal effectors are the archetypal EF hand-containing Ca^{2+} -binding protein calmodulin (CaM) and C2 domain-containing protein kinase C (delta, PKC δ). Upon binding Ca^{2+} , CaM associates with and engages a number of downstream effectors including the protein phosphatase calcineurin and Ca^{2+} /CaM kinases (CaMK1-IV) (Fig 1). Proteins once considered as Ca^{2+} buffers including calbindin-D28k and calretinin, also function as Ca^{2+} sensors, enabling them to modulate Ca^{2+} signalling outputs [33].

Ca^{2+} regulation of cancer-associated cell processes.

Ca^{2+} plays a major role in the development, progression and maintenance of cancer [34,35]. Cells become cancerous through the progressive accumulation of mutations in their genome. Subsequent to transformation, the cell displays a series of defining traits or hallmarks [36,37]: self-sufficiency from growth signals, independency from anti-growth signals, escape from apoptosis, loss of differentiation potential, neoangiogenesis, metastasis, metabolic reprogramming, immune surveillance evasion, genome instability and exploitation of inflammatory responses. These features render the cancer cell autonomous from the physiological signals that would normally regulate its lifespan. As Ca^{2+} participates in the processes underlying nearly all of these cancer hallmarks [38–40], most notably in regulating the balance between cell proliferation and death, it is not surprising that changes in the abundance or activity of Ca^{2+} handling proteins impact oncogenesis. Indeed, a dramatic increase in research in this area over the last few years has uncovered a critical role for Ca^{2+} in key control nodes in oncogenic transformation [34,35]. Not only have changes in expression levels

of Ca^{2+} handling proteins in cancerous cells been identified, interfering in Ca^{2+} handling pathways has emerged as a promising target for therapy [41,42].

Altered expression of Ca^{2+} handling proteins has been reported in cancers from different tissue origins [34,43]. Changes in the activity of Ca^{2+} signalling pathways and regulated processes have also been observed [35,44]. A defining picture of how Ca^{2+} signalling is modified at each stage of oncogenesis and how in turn the initiating mutation and tissue of origin impacts or influences this alteration in Ca^{2+} signalling is only just emerging. Moreover, whether Ca^{2+} signalling remodelling is a driver or a passenger of oncogenesis is not clearly established [45]. Precluding this deeper understanding is that oncogenesis involves a series of phenotypic changes and that the mechanism by which oncogenes remodel Ca^{2+} signals to promote tumorigenesis are not fully resolved. Further, the same Ca^{2+} -handling proteins show both increases and decreases in expression depending on the cancer type or stage [46,47]. Here, we will attempt to summarise the involvement and interactions between Ca^{2+} signalling via InsP_3 -induced Ca^{2+} release (IICR) and SOCE in cancer. Specifically, how these Ca^{2+} signalling pathways regulate different features of transformed cells, how Ca^{2+} signals generated are altered at the different stages of oncogenic transformation, and whether modifications in Ca^{2+} handling by these pathways drive and sustain the transformed phenotype. Prior to describing specific roles of SOCE and IICR in cancer, we will also attempt to provide a comprehensive picture of how cancer cells may employ or require different Ca^{2+} signals as they progress through the stages of tumorigenesis.

Ca^{2+} regulation of cell physiology and tumorigenic stages

Ca^{2+} in Cell Proliferation. It is classically considered that Ca^{2+} oscillations sustain life and are important for regulation of cell processes, including cell proliferation and metabolism, while large elevations in $[\text{Ca}^{2+}]_i$ promote apoptosis [48]. Indeed, Ca^{2+} oscillations occur throughout the cell cycle, and play strategic roles at nodal cell cycle phases such as during the early G_1 phase and the G_1/S and G_2/M transitions, where they correspond to peaks of activity of CaM and CaMK [38,49,50]. At early

G₁, Ca²⁺ regulates the transcription of immediate-early genes, such as FOS, JUN and MYC [51]. At the G₁/S transition, mitogen dependent signalling and Ca²⁺ increases activate Ras (through GEFs and GAPs) as well as CaM. Active Ras and Ca²⁺/CaM complexes are responsible for the transcription of cyclin D1 and formation of CDK4/cyclin D1 complexes. At the same time, CaM inhibits p21 and p27 while calcineurin promotes NFATc1 dephosphorylation, which in turn translocates to the nucleus and transactivates Myc [50,52–54]. Myc and Ras activate CycE/cdk2. All these events coincide to determine the multistep phosphorylation of retinoblastoma protein (Rb), which then releases the inhibitory transcription factor E2F, allowing DNA synthesis and progression into S phase [55,56]. At S and G₂/M transition, activation of CaMKII is necessary for correct centrosome duplication [57].

The aforementioned Ca²⁺ signals that control these cell cycle checkpoints show requirements for both ER Ca²⁺ release and SOCE [50]. A particular role for SOCE in the control of the G₁/S transition has been however recently demonstrated [58–60]. In this regard, SOCE is upregulated during the G₁/S transition through mechanisms involving STIM phosphorylation and its increased trafficking to sub-plasmalemmal *punctae*, as well as through Orai1 upregulation. Subsequent to this cell cycle phase, SOCE is downregulated while InsP₃-mediated signalling is active. The mechanism of SOCE involvement in G₂/M and early G₁ is more elusive.

Cell death and metabolism.

Ca²⁺ is a key player in mitochondrial metabolism. Oscillating Ca²⁺ signals drive mitochondrial bioenergetics by boosting the activity of three key enzymes of the TCA cycle, i.e. pyruvate dehydrogenase, isocitrate dehydrogenase and alpha-ketoglutarate dehydrogenase, as well as two components of the electron transport chain (ETC), namely cytochrome c oxidase and F₀-F₁-ATPase [61].

While Ca²⁺ oscillations fuel mitochondrial bioenergetics, sustained and high amplitude intracellular Ca²⁺ transients result in mitochondrial Ca²⁺ overload and production of reactive oxygen species (ROS). ROS trigger the opening of the mitochondrial permeability transition pore (mPTP), evoking

loss of inner mitochondrial membrane potential, mitochondrial swelling and rupture of the mitochondrial membrane. The molecular identity of the mPTP has been long sought but notably has recently been shown to be established by dimers of the F₀-F₁-ATPase [62]. Ca²⁺ can directly bind to these dimers and facilitate the formation of mitochondrial megachannels upon addition of the ATP synthase inhibitor Benzodiazepine 423 [63]. An elegant experiment showed the critical role for mitochondrial matrix Ca²⁺ in the opening of mPTP in conditions of phenylarsineoxide and *p*-hydroxyphenylglyoxal, whereby mPTP opening was blunted by BAPTA, a Ca²⁺ chelator [64]. The Ca²⁺-trigger site for the mPTP was identified in the catalytic site of the F-ATP synthase β subunit, provoking a conformational change via oligomycin sensitivity-conferring protein (OSCP) and the lateral stalk to the inner membrane [65]. The actual pore-forming subunit of the permeability transition complex remains a matter of debate, however, evidence was provided that the mPTP is formed upon dissociation of the F₀-F₁-ATPase dimers, and the c subunit functions as the pore-forming subunit of the permeability transition pore complex [66]. Besides the direct impact of Ca²⁺ on the proteins forming the mPTP, mitochondrial Ca²⁺ also provokes cardiolipin dissociation from the complex II of the respiratory chain, resulting in its disassembly [67]. The catalytic subunits (SDHA and SDHB) produce excessive ROS in the mitochondrial matrix, which triggers mPTP opening. This leads to Ca²⁺-induced mitochondrial outer membrane permeabilisation (MOMP), releasing cytochrome C and other apoptotic mediators [64,68].

Although mitochondria take up Ca²⁺ irrespective of its source, efficiency of Ca²⁺ uptake is greatest at the ER-mitochondrial interface. At this location, the InsP₃R is connected with the voltage-dependent anion channel of the outer mitochondrial membrane (VDAC1) through the chaperone GRP75 [69]. In this way, microdomains, known as mitochondria-associated ER membranes (MAMs) are created, allowing quasi-synaptic Ca²⁺ transfer from the ER to the mitochondria [70,71]. These contact sites are interesting targets for modulation of Ca²⁺ signalling, either to evade or to induce cell death [68,72,73]. Modulation of InsP₃Rs at these sites are discussed below.

As cancer cells are highly dependent on anabolic pathways for the production of substrates/products critical for cell division, maintenance of mitochondrial metabolism is essential. In several tumorigenic cell types, abrogating ER-mitochondrial Ca^{2+} fluxes leads to mitotic catastrophe, while in healthy cells it results in autophagy and a slowdown of cell cycle progression. This indicates that constitutive intra-organellar Ca^{2+} fluxes are an indispensable factor for both cell growth and survival [74–76]. However, these findings have been challenged in other studies. For instance, InsP_3R inhibition in tumorigenic MCF7 breast cancer cells provoked autophagy, and HEK293 cells remain viable even in the face of genetic knockout of all three InsP_3R isoforms via CRISPR/Cas9 [77,78].

Angiogenesis, invasion and metastasis. After reaching a critical size, the growing tumour mass develops its own vasculature in order to maintain a supply of nutrients and an oxygen to the core of the tumour (tumour angiogenesis). Angiogenesis depends on, amongst other factors, a Ca^{2+} -dependent increase in expression of the hypoxia-inducible factor 1 (HIF-1) transcription factor [79]. HIF-1 subsequently drives the transcription of vascular endothelial growth factor (VEGF), that in turn promotes endothelial cell (EC) proliferation, migration and formation of new vasculature (tubulogenesis). This potent pro-angiogenic effect of VEGF is elicited in part through activation of downstream Ca^{2+} signalling pathways [39,80], including of Ca^{2+} release from intracellular stores and Ca^{2+} entry across the plasma membrane [81–86]. Depending upon VEGF levels, different patterns of Ca^{2+} signals are observed in ECs, which have specific consequences for cell function. At low VEGF, oscillatory Ca^{2+} signals are observed, which induce NFAT nuclear translocation and EC proliferation, while at higher VEGF concentrations, sustained Ca^{2+} signals of a lower amplitude that promote EC migration via MLCK have been reported [87].

Cancer cell migration and matrix invasion show a strong dependence on Ca^{2+} signalling pathways. The MCU is a key player in these processes. Indeed, MCU silencing leads to a downregulation of HIF1 α , thereby impairing transcription of HIF1 α target genes involved in tumour progression [88].

Tumour cell Invasion requires epithelial-mesenchymal transition (EMT), through which cancer cells lose cell-cell contacts, change in shape, loose the polarized organization, and downregulate epithelial markers (e.g. E-cadherin, claudins, occludins) in favour of mesenchymal markers (e.g. N-cadherin, vimentin, integrins) [89,90]. Enhanced Ca^{2+} signalling involving TRP channels and ligand-gated P2X channels is associated with EMT and plays a role in controlling the expression of certain mesenchymal markers [39,91].

The Ca^{2+} dependence of cell migration is widely reported, including in cancer. Transient Ca^{2+} microdomains, also referred to as 'calcium flickers', occur at the leading front of a migrating cell, thereby facilitating local cytoskeletal remodelling and orchestrating directional movement [92,93]. In this way, they act as a steering wheel to turn the cell in a new direction. These short elevations in local $[\text{Ca}^{2+}]$ are brought about through several mechanisms, including InsP_3R signalling, modulation of SERCA and activation of TRPM7. Importantly, the flicker activity responds to directional cues of chemoattractants without any central command. Finally, invasion (the ability of the cancer cell to degrade the extracellular matrix (ECM) in order to reach the bloodstream and colonise other tissues), is a Ca^{2+} -dependent process. Ca^{2+} oscillations and Ca^{2+} influx are necessary for the formation of invadopodia structures and the recruitment of degrading enzymes such as matrix metalloproteases (MMPs), mainly through SOCE, TRP channels and VGCCs [39].

Ca^{2+} release channels and alterations in cancer

The importance of Ca^{2+} signalling in many aspects of cell physiology including cellular processes related to tumour development, places the InsP_3Rs centre stage in cancer biology [76]. A number of observations underpin this proposed role of InsP_3Rs . For example, absence of InsP_3R -mediated Ca^{2+} signals in thymocytes leads to the development of malignancies in mice, which resemble T-cell acute lymphoblastic leukemia [94]. The maintenance of the ER-mitochondrial Ca^{2+} flux via InsP_3Rs is important to sustain mitochondrial bioenergetics, which is required by cancer cells despite their dependence on glycolysis (Warburg effect) [95–97]. It is therefore not surprising that InsP_3R

expression levels are often altered in cancer. For instance, InsP₃R3 is overexpressed in glioblastoma cells and reducing its expression *via* siRNA resulted in impaired cell migration due to inhibition of cytosolic Ca²⁺ signalling [98]. Furthermore, in MCF-7 cells, InsP₃R blockage resulted in growth inhibition and 17- β -estradiol seems to induce cell proliferation in the same cell line through elevation of InsP₃R levels [99]. Interestingly, the different InsP₃R isoforms exhibit different types of Ca²⁺ responses [100]; a feature presumably owed to different regulation by Ca²⁺, InsP₃ and cofactors. While InsP₃R1 produces regular Ca²⁺ oscillations, which in the context of cancer cells may sustain the cell cycle, InsP₃R3 produces larger transients, which are instrumental in signalling apoptosis [100–102]. Indeed, InsP₃R3 silencing (thereby increasing the contribution of InsP₃R1 to Ca²⁺ signalling) often reveals an oscillatory Ca²⁺ signalling pattern [103,104]. In line with this, we showed that InsP₃R3 is the prevalent subtype in a non-tumorigenic colon cancer cell line compared to the isogenic tumorigenic counterpart, which predominantly expresses InsP₃R1 and is protected from apoptosis [105].

The role of InsP₃Rs in cancer, particularly that of InsP₃R3, has been the subject of a number of recent studies [45,46]. It appears that the expression level of this channel is finely tuned by the cancer cells to balance proliferation and apoptosis, which can be considered during early stages of tumourigenesis. InsP₃R3 levels are dynamically regulated through ubiquitylation, leading to proteasomal degradation of the channel. Several tumour suppressors, such as BAP1 and PTEN, are known to influence the ubiquitylation status of InsP₃R3 [106]. BAP1 possesses deubiquitylation enzyme activity, preventing its proteasomal degradation. PTEN, on the other hand, competes with FBXL2 for InsP₃R3 binding, thereby preventing FBXL2-mediated ubiquitylation of the channel. While healthy cells favour deubiquitylation of the InsP₃R3, cancer cells display defects in the expression of tumour suppressors, such as BAP1 and PTEN, resulting in decreased InsP₃R3 levels. However, InsP₃R3 is upregulated in several malignant, migratory cancers and cancer cell lines (Table 1), e.g. gastric [107], small and non-small cell lung [108,109], colorectal [110], glioblastoma [98] and breast cancer [111]. Moreover, InsP₃R3 inhibition reduces cell migration [98,103,104]. Very recently, silencing of

InsP₃R3 in different cancer cell models, including colorectal cancer DLD1 cells, ovarian cancer A2780 cells, and clear cell renal cell carcinoma RCC4 cells augmented apoptosis sensitivity, whereas silencing of InsP₃R1 conferred apoptotic protection [112]. Excitingly, DLD1 cells in which the InsP₃R3 gene was knocked out displayed a reduced in vivo tumour growth. Therefore, the latter study advocates that in some cancer cells, InsP₃R3 could fulfil an anti-apoptotic role, while InsP₃R1 acts as a pro-apoptotic channel. Thus, it might be possible that early downregulation of InsP₃R3 confers a survival advantage, while increasing its expression at later stages of tumorigenesis enables the cancer cell to proliferate and migrate, as seen in breast cancer [99,103,104,113], gastric cancer [107] and glioblastoma [98]. The reported roles of InsP₃R3 as an anti- versus a pro-apoptotic Ca²⁺-release channel may also in part be related to different localisations of this channel between different cancer cell models. We can speculate that InsP₃R3 at the MAMs serves as a pro-apoptotic channel, whereas InsP₃R3 outside the MAMs serves as an anti-apoptotic channel.

On the other hand, the involvement of the InsP₃R1 and 2 in tumour development is less straightforward. For instance, hormone refractory prostate cancer cells display elevated InsP₃R1 expression levels, which is thought to increase Ca²⁺ leakage from the ER making less Ca²⁺ available for induction of apoptosis through mitochondrial Ca²⁺ overload [114]. In bladder cancer cells resistant to cisplatin, however, InsP₃R1 levels are reduced and overexpression of InsP₃R1 induced apoptosis and restored sensitivity to cisplatin [115].

In tumour cells, not only do differences in InsP₃R expression levels affect cell survival, changes in Ca²⁺ release properties can also influence crucial cellular processes. First, phosphorylation of the InsP₃R can alter its function dramatically, for example through the PKB/AKT pathway, thereby reducing pro-apoptotic Ca²⁺ transfer to the mitochondria and conferring a survival advantage [116,117].

Conversely, tumour suppressor PML [23] impacts phosphorylation of the InsP₃R3 at the MAMs to promote ER-to-mitochondria pro-apoptotic Ca²⁺ fluxes [63] Furthermore, InsP₃R activity can be greatly influenced by an array of oncogenes. For instance, cancer cells promote cell survival through activity of the Bcl-2 family at the ER. On the one hand, anti-apoptotic Bcl-2-family members are able

to prevent pro-apoptotic Ca^{2+} signalling [56]. For instance, Bcl-2 can directly interact with the modulatory domain of the InsP_3R via its BH4 domain, thus inhibiting aberrant Ca^{2+} release [57–59].

On the other hand, Bcl-2-family members, such as Bcl-2, Bcl-XL and Mcl-1, can also increase pro-survival Ca^{2+} oscillations through sensitisation of InsP_3Rs , which in turn drives mitochondrial metabolism and provides the cancer cell with ATP [60].

In several Bcl-2-dependent cancer cell models, including chronic lymphocytic leukemia (CLL) and diffuse large B-cell lymphoma (DLBCL), Bcl-2 is found in complex with InsP_3R_2 , whereby Bcl-2 displacement from the InsP_3R channel results in Ca^{2+} -driven cell death [118–120]. In DLBCL cancer cells, the high InsP_3R_2 expression in combination with constitutively active B-cell receptor signalling and thus InsP_3 signalling leads to a Bcl-2 addiction to prevent pro-apoptotic InsP_3R -mediated Ca^{2+} release events [119,120]. Peptides targeting the BH4 domain of Bcl-2 (such as BIRD-2 [121]), which is responsible for InsP_3R interaction and inhibition, disrupt InsP_3R /Bcl-2 complexes and provoke apoptosis through intracellular Ca^{2+} overload, further exacerbated through Ca^{2+} influx from the extracellular environment [120,122]. Recently, a screen (using >25000 compounds) to identify small molecules that target InsP_3R /Bcl-2 complexes in cancer has yielded a lead compound that is able to provoke Ca^{2+} increases in Jurkat and CLL cells and to provoke cell death in Bcl-2-dependent cancer cell models [123]. This compound killed Bcl-2-dependent myeloma cells with IC_{50} of $\sim 5 \mu\text{M}$ without affecting the viability of normal human lymphocytes. BH4-domain-antagonizing tools can also be combined with BH3 mimetics (such as venetoclax/ABT-199) to target cancer cells poorly responding to venetoclax/ABT-199 or to sensitize cancer cells to venetoclax/ABT-199 [124–126]. The latter observation seems to involve the Ca^{2+} dependent upregulation of Bim, a pro-apoptotic BH3-only protein, in response to low [BIRD-2]. Also, the recently identified lead compound could promote venetoclax-induced cell death [123]. As such, peptide tools/small molecules targeting Bcl-2's BH4 domain may therefore offer interesting lead compounds to tackle several diseases by promoting ER-mitochondrial Ca^{2+} fluxes [127].

Unlike that for InsP_3Rs , much less is known about the role of RyRs in cancer, possibly because these channels are mainly studied in excitable cells, where the onset of cancer itself is rarer. RyRs are known to be expressed in breast cancer [128] and prostate cancer cells, where they may have a role in regulating apoptosis [129,130]. In a breast cancer cell line, for instance, EGF-induced EMT was associated with elevated mRNA levels of several ER Ca^{2+} pumps and channels, of which RyR2 displayed the most prominent change in gene expression levels [89]. These findings might suggest that also RyRs play a significant role in tumour onset and progression.

Ca^{2+} entry channels

Increasing evidence for the role of SOCE in cancer has emerged over the last decade [131], and also Ca^{2+} entry via VGCCs and TRPs is important in the evolution of cancer [132,133]. The intimate relationship between SOCE and ER Ca^{2+} release places this pathway centre stage in the regulation of cell physiology and likely the process of oncogenic transformation. STIM1 and Orai1 are considered the main mediators of SOCE in several different cancer cell types, and their role has been particularly recognised in the metastatic stage. Both STIM1 and Orai1 are frequently upregulated in invasive, malignant and metastatic tumours or cell lines (Table 1), such as advanced colorectal cancers [134–136], hepatocellular carcinoma [137], cervical cancer [138], breast cancer [139], glioblastoma [140] and oesophageal cancer [141]. Consistently, their pharmacological or genetic inhibition reduces migration and invasion, as seen in breast cancer cells [142], glioblastoma [140], hepatocellular carcinoma [137], clear cell renal carcinoma [143], colorectal cancer [134] and melanoma [144]. A recent study suggests that, in colorectal cancer, the maintenance of high STIM1 activity is due to Heat shock protein 27 (HSP27) [145], which is involved in STIM1 stabilisation at *punctae*. To maintain their viability and allow growth, metastatic cancers also need to form new vasculature in the mass. Indeed, the low O_2 tension associated with tumour environment promotes STIM-1 expression via HIF-1, which in turn upregulates HIF-1 expression in a feed-forward fashion [146]. VEGF expression is also associated with increased expression of SOCE components (e.g. STIM1 [59], VEGF in turn stimulates Ca^{2+} signalling via amongst other pathways a complex interplay between SOCE, the PLCy-

IP₃R Ca²⁺ release, and TRPC1 [81–86]. Indeed, STIM1, Orai1 and TRPC1 are the critical mediators of SOCE in ECs [147,148]. Altered expression of SOCE components as well as some TRP channels such as TRPC6 [149], TRPA1 [150] or TRPV4 [151] in cancer leads to the proliferation, migration and tubulogenesis of tumour-derived endothelial cells (TECs), which then allow vascularisation of the tumour mass. SOCE and non-SOCE components as well as ER-Ca²⁺ release in TECs represent an important target for therapy [86]

Less is clear regarding the modulation of SOCE and its mediators at the early stages of tumorigenesis. A direct involvement of STIM1 and SOCE in the G1/S transition of the cell cycle in SiHa cervical cancer cells has been described [58]. Furthermore, proliferation is repressed through inhibition of STIM1/Orai1 or SOCE in esophageal squamous cell carcinoma [141] and clear renal cell carcinoma [143]. In malignant melanoma, enhanced SOCE was responsible for improved survival through the maintenance of PKB activity, which is known to protect from apoptosis [152]. Studies conducted in leukemic cell lines have also shown that SOCE activation can temporarily inhibit extrinsic apoptotic pathways to prevent accidental death [153,154]. On the other hand, Orai1 downregulation confers apoptotic resistance [155], consistent with the hypothesis that SOCE activation stimulates proliferation by increasing ER Ca²⁺ content and mitochondrial uptake at the same time. Physiologically, upregulation of SOCE actors is important in mediating apoptosis in a number of cell types including neurons, endothelial cells and lymphocytes (for refs. see [131]). Thus, it is possible that SOCE activation has a general pro-proliferative and pro-migratory effect but will support tumorigenesis only when apoptotic mechanisms have been hijacked in other ways (e.g. increase in PKB-dependent InsP₃R phosphorylation). Alternatively, SOCE may be reduced as a strategy to lower ER Ca²⁺ content and protect from apoptosis, as seen for example by our group [156], where SOCE was equally diminished in two different KRAS^{wt/G13D} colorectal cancer cell lines bearing a different abundance of STIM1 (and STIM2, see later). This might seem counterintuitive but the demand for Ca²⁺ signals during tumorigenic transformation presumably differs according to the stage of differentiation and functional requirements of the cell [34,49,157]. Indeed, in line with our

analysis, others have described an impairment of SOCE, even in invasive compared to non-invasive tumours, e.g. in melanoma [158]. In a comparison of primary CRC cells with metastatic cells, the expression of STIM1 and Orai1 were found not to vary however, at least at the transcript level, and blockade of these proteins did not affect both primary and metastatic CRC cell proliferation and migration, despite the fact that SOCE was inhibited [136]. In this latter study, even though Stim1 and Orai3 are upregulated in metastatic CRC cells, SOCE was larger in primary cells because ER Ca^{2+} content was lower. A similar anti-correlation between ER store load and SOCE was reported in KRAS mutated colorectal cell lines by our group [105]. To complicate this scenario, different SOCE inhibitors might exert different effects [159]. The contribution of SOCE to the onset and maintenance of the cancer phenotype is therefore incompletely defined.

An understanding of the role of SOCE in cancer is further complicated by the different patterns of expression of the other STIM and Orai isoforms (relative to STIM1 and Orai1) in cancer. Moreover, how the various STIM and Orai isoforms combine to affect SOCE remains to be fully clarified, and changes in the relative abundance of these isoforms may indeed participate in the tumorigenic process [156]. Insights into the role of STIM2 in the context of cancer and its role in tumourigenesis is now emerging. STIM2 was originally reported to be upregulated in glioblastoma [160]. In CRC however, STIM2 has been shown to be both upregulated [161] and downregulated [135], with an inverse correlation between expression and invasiveness [161]. In contrast, Orai1/STIM2 expression in melanoma directly correlates with invasiveness [144]. As mentioned, our group observed a different STIM1/2 ratio, even between CRC cell lines harbouring the same KRAS mutation, with SOCE being reduced in both [156]. It is not clear whether some of these controversies are due to the analysis of different STIM2 splice variants. In a study of 295 breast tumours, the poorer prognosis was associated with those displaying a high STIM1/STIM2 ratio [139]. More recently, different roles for STIM1 and 2 in cervical cancer cells have been proposed. While both control proliferation, only STIM1 would also regulate migration [138].

Moreover, changes in Orai2 and 3 have been described in oncogenesis. In breast cancer, a specific role for Orai3 in mediating SOCE was reported in oestrogen receptor-positive (ER+ve) but not in ER-negative (ER-ve) cells, where STIM1 and Orai1 are the main SOCE actors [162,163]. In prostate cancer cells, Orai3 is overexpressed, forms AA-gated channels and its overexpression correlates with proliferation and apoptosis resistance [164]. Orai3 and 2 were also overexpressed in CRC but their role remained elusive [135] (and Roderick group-unpublished data). Further studies are indeed needed to confirm whether Orai2 and 3 play a role in cancer and could be used as tumour biomarkers, as it has been proposed for breast cancer [139,162,163].

The contribution of excessive SOCE to cell death also seems to underlie increased cell death susceptibility of cells expressing a Bcl-2 mutant carrying a three-amino acid mutation within the Bcl-2 BH1 domain [165]. It is proposed that ER Ca^{2+} depletion caused by this mutant version of Bcl-2 leads to activation of SOCE and increased expression levels of SOCE-related molecules. As a consequence of this excessive SOCE-mediated Ca^{2+} influx, caspases are activated, and apoptosis induced. The protection from apoptosis in this model by inactivation of SOCE or chelating intra- or extracellular Ca^{2+} supports the role of SOCE.

Interdependence of ER Ca^{2+} release and SOCE.

While we and others discuss apparently distinct contributions of ER Ca^{2+} release via InsP_3R and Ca^{2+} entry via SOCE to oncogenic transformation, these two pathways are highly interdependent and cannot be readily dissociated. In its simplest terms, ER Ca^{2+} release induces SOCE, and SOCE is needed to replenish/maintain ER Ca^{2+} levels – required for further release. Thus, with greater Ca^{2+} release, Ca^{2+} entry is likely to be more highly activated and increased SOCE could modify Ca^{2+} release. How can these two pathways act independently in cell physiology and pathology? For example, and at odds with the requirement of maintenance of ER Ca^{2+} levels to allow the ER to perform its many cellular functions [138], a substantial loss of ER Ca^{2+} has been considered to be required for SOCE activation [166]. The question then arises as how physiological stimuli induce

SOCE while sustaining other ER functions related to Ca^{2+} signalling. Notably, unlike the substantial depletion in ER Ca^{2+} observed when SOCE is experimentally induced by thapsigargin, ER $[\text{Ca}^{2+}]$ remains relatively stable in cells stimulated with physiologically relevant levels of agonist [167]. This could suggest that SOCE is preferentially activated by a sub-compartment of the ER store that is more depleted by agonist than the bulk ER. The preferential activation of I_{CRAC} by agonist in type II or type III InsP_3Rs expressing DT40 B lymphocytes compared to type 1 InsP_3R expressing cells may support such a model [168]. A model for SOCE activation involving a direct role for InsP_3Rs in gating Ca^{2+} entry, which could thus contribute to selective SOCE activation at Ca^{2+} release sites, has also been considered. This model is analogous to the conformational coupling of the L-type voltage gated Ca^{2+} channel ($\text{Ca}_v1.2$ a.k.a DHPR) and RyR1 in skeletal muscle – albeit signalling from the ER to the plasmalemma rather than the reverse [169]. A proximal localisation of the InsP_3Rs and the SOCE machinery at ER-PM junctions would lend support to such a model [170]. Indeed, InsP_3 has been demonstrated to activate a PM current in excised patches that was similar to I_{CRAC} [171]. InsP_3Rs have also been shown to directly interact with Orai channels [172]. This model may however be restricted to certain cell types and is not consistently supported. The identification of STIM proteins also circumvents the need for an interaction between InsP_3Rs and Orai to bring about SOCE. Recent findings now indicate that InsP_3Rs immobilised alongside ER-PM junctions, play an important role in stimulating SOCE [170]. Activation of these InsP_3Rs provokes a substantial drop in ER $[\text{Ca}^{2+}]$ in these junctions, thereby triggering SOCE. Due to the rapid diffusion of Ca^{2+} within the ER, however, luminal $[\text{Ca}^{2+}]$ in the ER will be minimally affected. This model gives rise to the idea that every ER-PM junction forms an autonomous SOCE unit which is regulated by its own immobilised InsP_3Rs . Interestingly, only InsP_3Rs at these junctions termed ‘licensed’ InsP_3Rs , underlie the elementary Ca^{2+} puffs generated in cells stimulated with agonist or InsP_3 . This paradigm would thus allow SOCE activation without compromising other ER functions. Ca^{2+} entering at these junctions not only has local effects, regulating signalling effectors such as calcineurin/NFAT, it is also re-sequestered back into the ER and ‘tunnelled’ to the remote ER, where it may regulate other cell functions [173,174].

Additionally, by ensuring ER refilling, Ca^{2+} remains available for Ca^{2+} release and for maintenance of InsP_3R activity. This latter function may indeed contribute to the modification of Ca^{2+} oscillations to a tonic a tonic elevation in $[\text{Ca}^{2+}]_i$ brought about via Ca^{2+} influx during cell stimulation with InsP_3 [175].

A direct role of InsP_3Rs in promoting Ca^{2+} influx via SOCE, independent of their role in bringing about store depletion and subsequent activation of STIM has also been suggested. Recent findings suggest that InsP_3Rs may trigger SOCE through an additional pathway where InsP_3Rs facilitates the coupling of STIM to Orai [176]. Following their activation, InsP_3Rs translocate to STIM punctae, thereby increasing STIM oligomerisation and ultimately enhancing SOCE. Interestingly, it has been found that SOCE is strongly suppressed during mitosis through uncoupling of ER- Ca^{2+} store depletion and activation of Ca^{2+} influx [177]. STIM1 is reported to be phosphorylated in mitotic cells, thereby disabling its rearrangement into PM punctae and inhibiting SOCE activation. This latter mechanism of regulation of STIM is likely to be of greater importance in highly proliferative cells than in cancer.

The consequences of this co-regulation of Ca^{2+} release mechanism and SOCE are not fully resolved. Given that both ER Ca^{2+} release and SOCE are initially lowered, then enhanced over the course of oncogenic development, we may hypothesize that an initial downregulation of the ER Ca^{2+} content (be it through an InsP_3 isoform switch, or a change in SERCA expression or activity) may lead to impaired SOCE and impaired mitochondrial uptake, an event which would protect from apoptosis while ensuring cancer cell viability through enhanced glycolysis. Lowered ER content would then stimulate higher SOCE, progressively upregulating Ca^{2+} leak, mitochondrial uptake and activity, and even more SOCE, in a feed-forward mechanism, which sustains invasion and metastasis.

Concluding remarks and model

The remodelling of the Ca^{2+} toolkit during the stages of oncogenic transformation is complex and designed to meet the needs of the cell during these stages – for example, to promote proliferation,

invasiveness, metabolism and avoid death. Ca^{2+} signalling mechanisms show substantial co-regulation in activity and expression. As such, care should be taken in overinterpreting findings of studies in which a full detailed analysis of the Ca^{2+} signalling-ome has not been carried out. Moreover, and not covered in this review of SOCE and ER Ca^{2+} release, is the array of the growing list of regulators/protein partners of the pathways described.

The alterations in the expression of Ca^{2+} toolkit proteins occurring in cancer are summarised in Table 1. Table 2 summarises the studies discussed here in relation to early or late tumorigenic stages. As previously mentioned, we have attempted to build a general model of Ca^{2+} signalling modifications occurring in cancer (Fig. 2). In this model, a downregulation of Ca^{2+} signals and of the ER Ca^{2+} content (obtained through an InsP_3R isoform switch and/or increased phosphorylation by overactive PKB and interaction with Bcl-2 family members, a decrease in SERCA pumps expression or activity, an increase in PMCA expression and a decrease in MCU activity, together with a decrease in SOCE or a switch in the STIM1/STIM2 ratio) would initially protect from apoptosis and encourage proliferation. At later stages, on the background of compromised apoptotic mechanisms (e.g. by KO mutations in p53), augmentation of Ca^{2+} signals through upregulation of InsP_3Rs , SOCE mediators, non-SOCE Ca^{2+} entry or mitochondrial fluxes sustains the proliferating, invading and migrating cancer cell. It is envisaged that a detailed knowledge of the Ca^{2+} signalling signature of cancers at different tumorigenic stages in a patient specific manner may help in their more effective therapeutic targeting.

Acknowledgements:

Work in the authors' laboratories is supported by grants from the Research Foundation Flanders (FWO). HLR would specifically wish to acknowledge Odysseus grant (90663).

Conflict of interest.

None.

ACCEPTED MANUSCRIPT

Bibliography

- [1] M.J. Berridge, P. Lipp, M.D. Bootman, The versatility and universality of calcium signalling., *Nat. Rev. Mol. Cell Biol.* 1 (2000) 11–21. doi:10.1038/35036035.
- [2] G. Santulli, R. Nakashima, Q. Yuan, A.R. Marks, Intracellular calcium release channels: an update., *J. Physiol.* 595 (2017) 3041–3051. doi:10.1113/JP272781.
- [3] J.K. Foskett, C. White, K.-H. Cheung, D.-O.D. Mak, Inositol Trisphosphate Receptor Ca²⁺ Release Channels, *Physiol. Rev.* 87 (2007) 593–658. doi:10.1152/physrev.00035.2006.
- [4] C.L. Newton, G.A. Mignery, T.C. Südhof, Co-expression in vertebrate tissues and cell lines of multiple inositol 1,4,5-trisphosphate (InsP₃) receptors with distinct affinities for InsP₃., *J. Biol. Chem.* 269 (1994) 28613–9. <http://www.ncbi.nlm.nih.gov/pubmed/7961809>.
- [5] P. Pinton, T. Pozzan, R. Rizzuto, The Golgi apparatus is an inositol 1,4,5-trisphosphate-sensitive Ca²⁺ store, with functional properties distinct from those of the endoplasmic reticulum., *EMBO J.* 17 (1998) 5298–308. doi:10.1093/emboj/17.18.5298.
- [6] A. Jha, M. Ahuja, S. Patel, E. Brailoiu, S. Muallem, Convergent regulation of the lysosomal two-pore channel-2 by Mg²⁺, NAADP, PI(3,5)P₂ and multiple protein kinases., *EMBO J.* 33 (2014) 501–11. doi:10.1002/emboj.201387035.
- [7] K. Venkatachalam, C.-O. Wong, M.X. Zhu, The role of TRPMLs in endolysosomal trafficking and function., *Cell Calcium.* 58 (2015) 48–56. doi:10.1016/j.ceca.2014.10.008.
- [8] M.D. Bootman, Calcium signaling., *Cold Spring Harb. Perspect. Biol.* 4 (2012) a011171. doi:10.1101/cshperspect.a011171.
- [9] I. Vandecaetsbeek, P. Vangheluwe, L. Raeymaekers, F. Wuytack, J. Vanoevelen, The Ca²⁺ pumps of the endoplasmic reticulum and Golgi apparatus., *Cold Spring Harb. Perspect. Biol.* 3 (2011). doi:10.1101/cshperspect.a004184.
- [10] T. Cali, M. Brini, E. Carafoli, Regulation of Cell Calcium and Role of Plasma Membrane Calcium ATPases., *Int. Rev. Cell Mol. Biol.* 332 (2017) 259–296. doi:10.1016/bs.ircmb.2017.01.002.
- [11] M. Ottolia, S. John, Y. Xie, X. Ren, K.D. Philipson, Shedding light on the Na⁺/Ca²⁺ exchanger., *Ann. N. Y. Acad. Sci.* 1099 (2007) 78–85. doi:10.1196/annals.1387.044.
- [12] A. Gudlur, P.G. Hogan, The STIM-Orai Pathway: Orai, the Pore-Forming Subunit of the CRAC Channel., *Adv. Exp. Med. Biol.* 993 (2017) 39–57. doi:10.1007/978-3-319-57732-6_3.
- [13] R. Qiu, R.S. Lewis, Structural features of STIM and Orai underlying store-operated calcium entry., *Curr. Opin. Cell Biol.* 57 (2019) 90–98. doi:10.1016/j.ceb.2018.12.012.
- [14] Y. Zhou, X. Cai, R.M. Nwokonko, N.A. Loktionova, Y. Wang, D.L. Gill, The STIM-Orai coupling interface and gating of the Orai1 channel., *Cell Calcium.* 63 (2017) 8–13. doi:10.1016/j.ceca.2017.01.001.
- [15] J. Liou, M.L. Kim, D.H. Won, J.T. Jones, J.W. Myers, J.E. Ferrell, T. Meyer, STIM is a Ca²⁺ sensor essential for Ca²⁺-store- depletion-triggered Ca²⁺ influx, *Curr. Biol.* 15 (2005) 1235–1241. doi:10.1016/j.cub.2005.05.055.
- [16] J. Soboloff, M.A. Spassova, T. Hewavitharana, L.P. He, W. Xu, L.S. Johnstone, M.A. Dziadek, D.L. Gill, STIM2 Is an Inhibitor of STIM1-Mediated Store-Operated Ca²⁺ Entry, *Curr. Biol.* 16 (2006) 1465–1470. doi:10.1016/j.cub.2006.05.051.

- [17] O. Brandman, J. Liou, W.S. Park, T. Meyer, STIM2 is a Feedback Regulator that Stabilizes Basal Cytosolic and Endoplasmic Reticulum Ca²⁺ Levels, *Cell*. 131 (2007) 1327–1339. doi:10.1016/j.cell.2007.11.039.STIM2.
- [18] A. Rana, M. Yen, A.M. Sadaghiani, S. Malmersjö, C.Y. Park, R.E. Dolmetsch, R.S. Lewis, Alternative splicing converts STIM2 from an activator to an inhibitor of store-operated calcium channels, *J. Cell Biol.* 209 (2015) 653–670. doi:10.1083/jcb.201412060.
- [19] A.M. Miederer, D. Alansary, G. Schwär, P.H. Lee, M. Jung, V. Helms, B.A. Niemeyer, A STIM2 splice variant negatively regulates store-operated calcium entry, *Nat. Commun.* 6 (2015). doi:10.1038/ncomms7899.
- [20] A. Berna-Erro, I. Jardin, G.M. Salido, J.A. Rosado, Role of STIM2 in cell function and physiopathology, *J. Physiol.* 595 (2017) 3111–3128. doi:10.1113/JP273889.
- [21] H.A. Nelson, C.A. Leech, R.F. Kopp, M.W. Roe, Interplay between ER Ca²⁺-binding proteins, STIM1 and STIM2, is required for store-operated Ca²⁺-entry, *Int. J. Mol. Sci.* 19 (2018). doi:10.3390/ijms19051522.
- [22] K.P. Subedi, H.L. Ong, G.Y. Son, X. Liu, I.S. Ambudkar, STIM2 Induces Activated Conformation of STIM1 to Control Orai1 Function in ER-PM Junctions, *Cell Rep.* 23 (2018) 522–534. doi:10.1016/j.celrep.2018.03.065.
- [23] S. Zheng, G. Ma, L. He, T. Zhang, J. Li, X. Yuan, N.T. Nguyen, Y. Huang, X. Zhang, P. Gao, R. Nwokonko, D.L. Gill, H. Dong, Y. Zhou, Y. Wang, Identification of molecular determinants that govern distinct STIM2 activation dynamics., *PLoS Biol.* 16 (2018) e2006898. doi:10.1371/journal.pbio.2006898.
- [24] H.L. Ong, I.S. Ambudkar, STIM-TRP Pathways and Microdomain Organization: Contribution of TRPC1 in Store-Operated Ca²⁺ Entry: Impact on Ca²⁺ Signaling and Cell Function., *Adv. Exp. Med. Biol.* 993 (2017) 159–188. doi:10.1007/978-3-319-57732-6_9.
- [25] I.S. Ambudkar, L.B. de Souza, H.L. Ong, TRPC1, Orai1, and STIM1 in SOCE: Friends in tight spaces., *Cell Calcium*. 63 (2017) 33–39. doi:10.1016/j.ceca.2016.12.009.
- [26] K.T. Cheng, H.L. Ong, X. Liu, I.S. Ambudkar, Contribution of TRPC1 and Orai1 to Ca²⁺ entry activated by store depletion., *Adv. Exp. Med. Biol.* 704 (2011) 435–49. doi:10.1007/978-94-007-0265-3_24.
- [27] H. Li, M.D. Pink, J.G. Murphy, A. Stein, M.L. Dell’Acqua, P.G. Hogan, Balanced interactions of calcineurin with AKAP79 regulate Ca²⁺-calcineurin-NFAT signaling., *Nat. Struct. Mol. Biol.* 19 (2012) 337–45. doi:10.1038/nsmb.2238.
- [28] A.B. Parekh, Decoding cytosolic Ca²⁺ oscillations., *Trends Biochem. Sci.* 36 (2011) 78–87. doi:10.1016/j.tibs.2010.07.013.
- [29] K.T. Cheng, X. Liu, H.L. Ong, W. Swaim, I.S. Ambudkar, Local Ca²⁺ entry via Orai1 regulates plasma membrane recruitment of TRPC1 and controls cytosolic Ca²⁺ signals required for specific cell functions., *PLoS Biol.* 9 (2011) e1001025. doi:10.1371/journal.pbio.1001025.
- [30] H.L. Ong, S.-I. Jang, I.S. Ambudkar, Distinct contributions of Orai1 and TRPC1 to agonist-induced [Ca²⁺]_i signals determine specificity of Ca²⁺-dependent gene expression., *PLoS One*. 7 (2012) e47146. doi:10.1371/journal.pone.0047146.
- [31] X. Zhang, M. Gueguinou, M. Trebak, Store-Independent Orai Channels Regulated by STIM,

2018. <http://www.ncbi.nlm.nih.gov/pubmed/30299650>.
- [32] T.J. Shuttleworth, J.L. Thompson, O. Mignen, STIM1 and the noncapacitative ARC channels, *Cell Calcium*. 42 (2007) 183–191. doi:10.1016/j.ceca.2007.01.012.
 - [33] Schwaller, No Title, Cold Spring Harb. – Perspect. Biol. (2019).
 - [34] G.R. Monteith, N. Prevarskaya, S.J. Roberts-Thomson, The calcium–cancer signalling nexus, *Nat. Rev. Cancer*. (2017). doi:10.1038/nrc.2017.18.
 - [35] T.A. Stewart, K.T.D.S. Yapa, G.R. Monteith, Altered calcium signaling in cancer cells, *Biochim. Biophys. Acta*. 1848 (2015) 2502–2511. doi:10.1016/j.bbame.2014.08.016.
 - [36] D. Hanahan, R.A. Weinberg, Hallmarks of cancer: the next generation., *Cell*. 144 (2011) 646–74. doi:10.1016/j.cell.2011.02.013.
 - [37] D. Hanahan, R.A. Weinberg, The hallmarks of cancer., *Cell*. 100 (2000) 57–70. <http://www.ncbi.nlm.nih.gov/pubmed/10647931>.
 - [38] J. Humeau, J.M. Bravo-San Pedro, I. Vitale, L. Nuñez, C. Villalobos, G. Kroemer, L. Senovilla, Calcium signaling and cell cycle: Progression or death, *Cell Calcium*. 70 (2017) 3–15. doi:10.1016/j.ceca.2017.07.006.
 - [39] O. Iamshanova, A. Fiorio Pla, N. Prevarskaya, Molecular mechanisms of tumor invasion: regulation by calcium signals, *J. Physiol.* 10 (2017) 3063–3075. doi:10.1113/JP272844.
 - [40] M.M. Xu, A. Seas, M. Kiyani, K.S.Y. Ji, H.N. Bell, A temporal examination of calcium signaling in cancer- from tumorigenesis, to immune evasion, and metastasis, *Cell Biosci.* 8 (2018) 1–9. doi:10.1186/s13578-018-0223-5.
 - [41] M. Bonora, C. Giorgi, P. Pinton, Novel frontiers in calcium signaling: A possible target for chemotherapy., *Pharmacol. Res.* 99 (2015) 82–5. doi:10.1016/j.phrs.2015.05.008.
 - [42] M. Kerkhofs, M. Bittremieux, G. Morciano, C. Giorgi, P. Pinton, J.B. Parys, G. Bultynck, Emerging molecular mechanisms in chemotherapy: Ca²⁺ signaling at the mitochondria-associated endoplasmic reticulum membranes, *Cell Death Dis.* 9 (2018). doi:10.1038/s41419-017-0179-0.
 - [43] G.R. Monteith, D. McAndrew, H.M. Faddy, S.J. Roberts-Thomson, Calcium and cancer: targeting Ca²⁺ transport, *Nat. Rev. Cancer*. 7 (2007) 519–530. doi:10.1038/nrc2171.
 - [44] Y.-F. Chen, Y.-T. Chen, W.-T. Chiu, M.-R. Shen, Remodeling of calcium signaling in tumor progression., *J. Biomed. Sci.* 20 (2013) 23. doi:10.1186/1423-0127-20-23.
 - [45] M. Hoth, CRAC channels, calcium, and cancer in light of the driver and passenger concept, *Biochim. Biophys. Acta - Mol. Cell Res.* (2016). doi:10.1016/j.bbamcr.2015.12.009.
 - [46] M.C. Curry, A.A. Peters, P.A. Kenny, S.J. Roberts-Thomson, G.R. Monteith, Mitochondrial calcium uniporter silencing potentiates caspase-independent cell death in MDA-MB-231 breast cancer cells., *Biochem. Biophys. Res. Commun.* 434 (2013) 695–700. doi:10.1016/j.bbrc.2013.04.015.
 - [47] S. Marchi, L. Lupini, S. Patergnani, A. Rimessi, S. Missiroli, M. Bonora, A. Bononi, F. Corrà, C. Giorgi, E. De Marchi, F. Poletti, R. Gafà, G. Lanza, M. Negrini, R. Rizzuto, P. Pinton, Downregulation of the mitochondrial calcium uniporter by cancer-related miR-25, *Curr. Biol.* 23 (2013) 58–63. doi:10.1016/j.cub.2012.11.026.

- [48] P.M. Herst, M.R. Rowe, G.M. Carson, M. V Berridge, Functional Mitochondria in Health and Disease., *Front. Endocrinol. (Lausanne)*. 8 (2017) 296. doi:10.3389/fendo.2017.00296.
- [49] H.L. Roderick, S.J. Cook, Ca²⁺ signalling checkpoints in cancer: remodelling Ca²⁺ for cancer cell proliferation and survival, *Nat. Rev. Cancer*. 8 (2008) 361–375. doi:10.1038/nrc2374.
- [50] A. Arredouani, F. Yu, L. Sun, K. Machaca, Regulation of store-operated Ca²⁺ entry during the cell cycle., *J. Cell Sci.* 123 (2010) 2155–62. doi:10.1242/jcs.069690.
- [51] E. Roche, M. Prentki, Calcium regulation of immediate-early response genes, *Cell Calcium*. 16 (1994) 331–338. doi:10.1016/0143-4160(94)90097-3.
- [52] A. Köenig, T. Linhart, K. Schlengemann, K. Reutlinger, J. Wegele, G. Adler, G. Singh, L. Hofmann, S. Kunsch, T. Büch, E. Schäfer, T.M. Gress, M.E. Fernandez-Zapico, V. Ellenrieder, NFAT-induced histone acetylation relay switch promotes c-Myc-dependent growth in pancreatic cancer cells., *Gastroenterology*. 138 (2010) 1189-99.e1–2. doi:10.1053/j.gastro.2009.10.045.
- [53] M.C.X. Pinto, A.H. Kihara, V.A.M. Goulart, F.M.P. Tonelli, K.N. Gomes, H. Ulrich, R.R. Resende, Calcium signaling and cell proliferation., *Cell. Signal.* 27 (2015) 2139–49. doi:10.1016/j.cellsig.2015.08.006.
- [54] P.G. Hogan, L. Chen, J. Nardone, A. Rao, Transcriptional regulation by calcium, calcineurin, and NFAT, *Genes Dev.* 17 (2003) 2205–2232. doi:10.1101/gad.1102703.
- [55] C.D. Rasmussen, A.R. Means, Calmodulin is required for cell-cycle progression during G1 and mitosis., *EMBO J.* 8 (1989) 73–82.
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=400774&tool=pmcentrez&rendertype=abstract>.
- [56] N. Takuwa, W. Zhou, M. Kumada, Y. Takuwa, Ca²⁺-dependent stimulation of retinoblastoma gene product phosphorylation and p34cdc2 kinase activation in serum-stimulated human fibroblasts, *J. Biol. Chem.* 268 (1993) 138–145.
- [57] Y. Matsumoto, J.L. Maller, Calcium, calmodulin, and CaMKII requirement for initiation of centrosome duplication in *Xenopus* egg extracts, *Science* (80-.). 295 (2002) 499–502. doi:10.1126/science.1065693.
- [58] Y.-W. Chen, Y.-F. Chen, Y.-T. Chen, W.-T. Chiu, M.-R. Shen, The STIM1-Orai1 pathway of store-operated Ca(2+) entry controls the checkpoint in cell cycle G1/S transition., *Sci. Rep.* 6 (2016) 22142. doi:10.1038/srep22142.
- [59] Y.-F. Chen, W.-T. Chiu, Y.-T. Chen, P.-Y. Lin, H.-J. Huang, C.-Y. Chou, H.-C. Chang, M.-J. Tang, M.-R. Shen, Calcium store sensor stromal-interaction molecule 1-dependent signaling plays an important role in cervical cancer growth, migration, and angiogenesis., *Proc. Natl. Acad. Sci. U. S. A.* 108 (2011) 15225–30. doi:10.1073/pnas.1103315108.
- [60] P. Tomas-Martin, A.M. Lopez-Guerrero, V. Casas-Rua, E. Pozo-Guisado, F.J. Martin-Romero, Phospho-STIM1 is a downstream effector that mediates the signaling triggered by IGF-1 in HEK293 cells, *Cell. Signal.* 27 (2015) 545–554. doi:10.1016/j.cellsig.2014.12.017.
- [61] R.M. Denton, J.G. McCormack, P.J. Midgley, G.A. Rutter, A.P. Thomas, The role of Ca²⁺ in the hormonal control of intramitochondrial metabolism in heart, liver, and adipose tissue., *Adv. Second Messenger Phosphoprotein Res.* 21 (1988) 157–64.

- [62] P. Bernardi, A. Rasola, M. Forte, G. Lippe, The Mitochondrial Permeability Transition Pore: Channel Formation by F-ATP Synthase, Integration in Signal Transduction, and Role in Pathophysiology., *Physiol. Rev.* 95 (2015) 1111–55. doi:10.1152/physrev.00001.2015.
- [63] V. Giorgio, S. von Stockum, M. Antoniel, A. Fabbro, F. Fogolari, M. Forte, G.D. Glick, V. Petronilli, M. Zoratti, I. Szabó, G. Lippe, P. Bernardi, Dimers of mitochondrial ATP synthase form the permeability transition pore., *Proc. Natl. Acad. Sci. U. S. A.* 110 (2013) 5887–92. doi:10.1073/pnas.1217823110.
- [64] V. Giorgio, L. Guo, C. Bassot, V. Petronilli, P. Bernardi, Calcium and regulation of the mitochondrial permeability transition., *Cell Calcium.* 70 (2018) 56–63. doi:10.1016/j.ceca.2017.05.004.
- [65] V. Giorgio, V. Burchell, M. Schiavone, C. Bassot, G. Minervini, V. Petronilli, F. Argenton, M. Forte, S. Tosatto, G. Lippe, P. Bernardi, Ca²⁺ binding to F-ATP synthase β subunit triggers the mitochondrial permeability transition., *EMBO Rep.* 18 (2017) 1065–1076. doi:10.15252/embr.201643354.
- [66] M. Bonora, C. Morganti, G. Morciano, G. Pedriali, M. Lebiecinska-Arciszewska, G. Aquila, C. Giorgi, P. Rizzo, G. Campo, R. Ferrari, G. Kroemer, M.R. Wieckowski, L. Galluzzi, P. Pinton, Mitochondrial permeability transition involves dissociation of F₁FO ATP synthase dimers and C-ring conformation., *EMBO Rep.* 18 (2017) 1077–1089. doi:10.15252/embr.201643602.
- [67] M.-S. Hwang, C.T. Schwall, E. Pazarentzos, C. Datler, N.N. Alder, S. Grimm, Mitochondrial Ca(2+) influx targets cardiolipin to disintegrate respiratory chain complex II for cell death induction., *Cell Death Differ.* 21 (2014) 1733–45. doi:10.1038/cdd.2014.84.
- [68] S. Marchi, S. Patergnani, S. Missiroli, G. Morciano, A. Rimessi, M.R. Wieckowski, C. Giorgi, P. Pinton, Mitochondrial and endoplasmic reticulum calcium homeostasis and cell death., *Cell Calcium.* 69 (2018) 62–72. doi:10.1016/j.ceca.2017.05.003.
- [69] G. Szabadkai, K. Bianchi, P. Várnai, D. De Stefani, M.R. Wieckowski, D. Cavagna, A.I. Nagy, T. Balla, R. Rizzuto, Chaperone-mediated coupling of endoplasmic reticulum and mitochondrial Ca²⁺ channels., *J. Cell Biol.* 175 (2006) 901–11. doi:10.1083/jcb.200608073.
- [70] T. Vervloessem, D.I. Yule, G. Bultynck, J.B. Parys, The type 2 inositol 1,4,5-trisphosphate receptor, emerging functions for an intriguing Ca²⁺ -release channel, 15 (2014) 34–48. doi:10.1038/nrg3575.Systems.
- [71] T. Hayashi, R. Rizzuto, G. Hajnoczky, T.-P. Su, MAM: more than just a housekeeper, *Motor Control.* 27 (2009) 590–609. doi:10.1016/j.humov.2008.02.015.Changes.
- [72] S. Marchi, M. Bittremieux, S. Missiroli, C. Morganti, S. Patergnani, L. Sbano, A. Rimessi, M. Kerkhofs, J.B. Parys, G. Bultynck, C. Giorgi, P. Pinton, Endoplasmic Reticulum-Mitochondria Communication Through Ca²⁺ Signaling: The Importance of Mitochondria-Associated Membranes (MAMs)., *Adv. Exp. Med. Biol.* 997 (2017) 49–67. doi:10.1007/978-981-10-4567-7_4.
- [73] T. Gutiérrez, T. Simmen, Endoplasmic reticulum chaperones tweak the mitochondrial calcium rheostat to control metabolism and cell death., *Cell Calcium.* 70 (2018) 64–75. doi:10.1016/j.ceca.2017.05.015.
- [74] C. Cárdenas, M. Müller, A. McNeal, A. Lovy, F. Jaña, G. Bustos, F. Urra, N. Smith, J. Molgó, J.A. Diehl, T.W. Ridky, J.K. Foskett, Selective Vulnerability of Cancer Cells by Inhibition of Ca²⁺ Transfer from Endoplasmic Reticulum to Mitochondria, *Cell Rep.* 14 (2016) 2313–2324.

- doi:10.1016/j.celrep.2016.02.030.
- [75] G. Bultynck, Onco-IP3Rs Feed Cancerous Cravings for Mitochondrial Ca²⁺, *Trends Biochem. Sci.* 41 (2016) 390–393. doi:10.1016/j.tibs.2016.03.006.
 - [76] H. Ivanova, M. Kerkhofs, R.M. La Rovere, G. Bultynck, Endoplasmic Reticulum-Mitochondrial Ca²⁺ Fluxes Underlying Cancer Cell Survival., *Front. Oncol.* 7 (2017) 70. doi:10.3389/fonc.2017.00070.
 - [77] A. Singh, M. Chagtoo, S. Tiwari, N. George, B. Chakravarti, S. Khan, S. Lakshmi, M.M. Godbole, Inhibition of Inositol 1, 4, 5-Trisphosphate Receptor Induce Breast Cancer Cell Death Through Deregulated Autophagy and Cellular Bioenergetics., *J. Cell. Biochem.* 118 (2017) 2333–2346. doi:10.1002/jcb.25891.
 - [78] K.J. Alzayady, L. Wang, R. Chandrasekhar, L.E. Wagner, F. Van Petegem, D.I. Yule, Defining the stoichiometry of inositol 1,4,5-trisphosphate binding required to initiate Ca²⁺ release., *Sci. Signal.* 9 (2016) ra35. doi:10.1126/scisignal.aad6281.
 - [79] I. Azimi, The interplay between HIF-1 and calcium signalling in cancer., *Int. J. Biochem. Cell Biol.* 97 (2018) 73–77. doi:10.1016/j.biocel.2018.02.001.
 - [80] A. Fiorio Pla, K. Kondratska, N. Prevarskaya, STIM and ORAI proteins: crucial roles in hallmarks of cancer, *Am. J. Physiol. - Cell Physiol.* 310 (2016) C509–C519. doi:10.1152/ajpcell.00364.2015.
 - [81] S. Wang, X. Li, M. Parra, E. Verdin, R. Bassel-Duby, E.N. Olson, Control of endothelial cell proliferation and migration by VEGF signaling to histone deacetylase 7, *Proc. Natl. Acad. Sci.* 105 (2008) 7738–7743. doi:10.1073/pnas.0802857105.
 - [82] J. Li, R.M. Cubbon, L.A. Wilson, M.S. Amer, L. McKeown, B. Hou, Y. Majeed, S. Tumova, V.A.L. Seymour, H. Taylor, M. Stacey, D. O'Regan, R. Foster, K.E. Porter, M.T. Kearney, D.J. Beech, Orai1 and CRAC channel dependence of VEGF-activated Ca²⁺ entry and endothelial tube formation, *Circ. Res.* 108 (2011) 1190–1198. doi:10.1161/CIRCRESAHA.111.243352.
 - [83] S. Dragoni, U. Laforenza, E. Bonetti, F. Lodola, C. Bottino, R. Berra-Romani, G. Carlo Bongio, M.P. Cinelli, G. Guerra, P. Pedrazzoli, V. Rosti, F. Tanzi, F. Moccia, Vascular endothelial growth factor stimulates endothelial colony forming cells proliferation and tubulogenesis by inducing oscillations in intracellular Ca²⁺ concentration., *Stem Cells.* 29 (2011) 1898–907. doi:10.1002/stem.734.
 - [84] D. Jho, D. Mehta, G. Ahmmed, X.-P. Gao, C. Tiruppathi, M. Broman, A.B. Malik, Angiopoietin-1 opposes VEGF-induced increase in endothelial permeability by inhibiting TRPC1-dependent Ca²⁺ influx., *Circ. Res.* 96 (2005) 1282–90. doi:10.1161/01.RES.0000171894.03801.03.
 - [85] D. Mehta, G.U. Ahmmed, B.C. Paria, M. Holinstat, T. Voyno-Yasenetskaya, C. Tiruppathi, R.D. Minshall, A.B. Malik, RhoA Interaction with Inositol 1,4,5-Trisphosphate Receptor and Transient Receptor Potential Channel-1 Regulates Ca²⁺ Entry, *J. Biol. Chem.* 278 (2003) 33492–33500. doi:10.1074/jbc.m302401200.
 - [86] Scarpellino, Genova, Avanzato, Bernardini, Bianco, Petrillo, Tolosano, de Almeida Vieira, Bussolati, Fiorio Pla, Munaron, Purinergic Calcium Signals in Tumor-Derived Endothelium, *Cancers (Basel).* 11 (2019) 766. doi:10.3390/cancers11060766.
 - [87] D.P. Noren, W.H. Chou, S.H. Lee, A.A. Qutub, A. Warmflash, D.S. Wagner, A.S. Popel, A. Levchenko, Endothelial cells decode VEGF-mediated Ca²⁺ signaling patterns to produce

- distinct functional responses., *Sci. Signal.* 9 (2016) ra20. doi:10.1126/scisignal.aad3188.
- [88] A. Tosatto, R. Sommaggio, C. Kummerow, R.B. Bentham, T.S. Blacker, T. Berecz, M.R. Duchon, A. Rosato, I. Bogeski, G. Szabadkai, R. Rizzuto, C. Mammucari, The mitochondrial calcium uniporter regulates breast cancer progression via HIF-1 α , *EMBO Mol. Med.* 8 (2016) 569–585. doi:10.15252/emmm.201606255.
- [89] F.M. Davis, M.T. Parsonage, P.J. Cabot, M.-O. Parat, E.W. Thompson, S.J. Roberts-Thomson, G.R. Monteith, Assessment of gene expression of intracellular calcium channels, pumps and exchangers with epidermal growth factor-induced epithelial-mesenchymal transition in a breast cancer cell line., *Cancer Cell Int.* 13 (2013) 76. doi:10.1186/1475-2867-13-76.
- [90] I. Azimi, H. Beilby, F.M. Davis, D.L. Marcial, P.A. Kenny, E.W. Thompson, S.J. Roberts-Thomson, G.R. Monteith, Altered purinergic receptor-Ca²⁺ signaling associated with hypoxia-induced epithelial-mesenchymal transition in breast cancer cells., *Mol. Oncol.* 10 (2016) 166–78. doi:10.1016/j.molonc.2015.09.006.
- [91] F.M. Davis, I. Azimi, R.A. Faville, A.A. Peters, K. Jalink, J.W.J. Putney, G.J. Goodhill, E.W. Thompson, S.J. Roberts-Thomson, G.R. Monteith, Induction of epithelial-mesenchymal transition (EMT) in breast cancer cells is calcium signal dependent, *Oncogene.* 33 (2014) 2307–2316. doi:10.1038/onc.2013.187.Induction.
- [92] C. Wei, X. Wang, M. Chen, K. Ouyang, L.-S. Song, H. Cheng, Calcium flickers steer cell migration., *Nature.* 457 (2009) 901–5. doi:10.1038/nature07577.
- [93] J.H. Evans, J.J. Falke, Ca²⁺ influx is an essential component of the positive-feedback loop that maintains leading-edge structure and activity in macrophages, *Proc. Natl. Acad. Sci.* 104 (2007) 16176–16181. doi:10.1073/pnas.0707719104.
- [94] K. Ouyang, R. Leandro Gomez-Amaro, D.L. Stachura, H. Tang, X. Peng, X. Fang, D. Traver, S.M. Evans, J. Chen, Loss of IP3R-dependent Ca²⁺ signalling in thymocytes leads to aberrant development and acute lymphoblastic leukemia., *Nat. Commun.* 5 (2014) 4814. doi:10.1038/ncomms5814.
- [95] M.G. Vander Heiden, L.C. Cantley, C.B. Thompson, Understanding the Warburg Effect: The Metabolic Requirements of Cell Proliferation, *Science* (80-.). 324 (2009) 1029–1033. doi:10.1126/science.1160809.
- [96] A.J. de Groof, M.M. te Lindert, M.M. van Dommelen, M. Wu, M. Willemse, A.L. Smift, M. Winer, F. Oerlemans, H. Pluk, J.A. Fransen, B. Wieringa, Increased OXPHOS activity precedes rise in glycolytic rate in H-RasV12/E1A transformed fibroblasts that develop a Warburg phenotype, *Mol. Cancer.* 8 (2009) 54. doi:10.1186/1476-4598-8-54.
- [97] B. Zhivotovsky, S. Orrenius, The Warburg Effect returns to the cancer stage, *Semin. Cancer Biol.* 19 (2009) 1–3. doi:10.1016/j.semcancer.2008.12.003.
- [98] S.S. Kang, K.S. Han, B.M. Ku, Y.K. Lee, J. Hong, H.Y. Shin, A.G. Almonte, D.H. Woo, D.J. Brat, E.M. Hwang, S.H. Yoo, C.K. Chung, S.H. Park, S.H. Paek, E.J. Roh, S.J. Lee, J.Y. Park, S.F. Traynelis, C.J. Lee, Caffeine-mediated inhibition of calcium release channel inositol 1,4,5-trisphosphate receptor subtype 3 blocks glioblastoma invasion and extends survival, *Cancer Res.* 70 (2010) 1173–1183. doi:10.1158/0008-5472.CAN-09-2886.
- [99] C. Szatkowski, J.B. Parys, H. Ouadid-Ahidouch, F. Matifat, Inositol 1,4,5-trisphosphate-induced Ca²⁺ signalling is involved in estradiol-induced breast cancer epithelial cell growth., *Mol. Cancer.* 9 (2010) 156. doi:10.1186/1476-4598-9-156.

- [100] H. Ivanova, T. Vervliet, L. Missiaen, J.B. Parys, H. De Smedt, G. Bultynck, Inositol 1,4,5-trisphosphate receptor-isoform diversity in cell death and survival, *Biochim. Biophys. Acta - Mol. Cell Res.* 1843 (2014) 2164–2183. doi:10.1016/j.bbamcr.2014.03.007.
- [101] T. Miyakawa, a Maeda, T. Yamazawa, K. Hirose, T. Kurosaki, M. Iino, Encoding of Ca²⁺ signals by differential expression of IP3 receptor subtypes., *EMBO J.* 18 (1999) 1303–1308. doi:10.1093/emboj/18.5.1303.
- [102] M. Hattori, A.Z. Suzuki, T. Higo, H. Miyauchi, T. Michikawa, T. Nakamura, T. Inoue, K. Mikoshiba, Distinct Roles of Inositol 1,4,5-Trisphosphate Receptor Types 1 and 3 in Ca²⁺ Signaling, *J. Biol. Chem.* 279 (2004) 11967–11975. doi:10.1074/jbc.M311456200.
- [103] A. Mound, A. Vautrin-Glabik, A. Foulon, B. Botia, F. Hague, J.B. Parys, H. Ouadid-Ahidouch, L. Rodat-Despoix, Downregulation of type 3 inositol (1,4,5)-trisphosphate receptor decreases breast cancer cell migration through an oscillatory Ca²⁺ signal, *Oncotarget.* 8 (2017) 72324–72341. doi:10.18632/oncotarget.20327.
- [104] A. Vautrin-Glabik, B. Botia, P. Kischel, H. Ouadid-Ahidouch, L. Rodat-Despoix, IP3R3 silencing induced actin cytoskeletal reorganization through ARHGAP18/RhoA/mDia1/FAK pathway in breast cancer cell lines., *Biochim. Biophys. Acta. Mol. Cell Res.* 1865 (2018) 945–958. doi:10.1016/j.bbamcr.2018.04.002.
- [105] C. Pierro, S.J. Cook, T.C. Foets, M.D. Bootman, H.L. Roderick, Oncogenic K-Ras suppresses IP₃-dependent Ca²⁺ release through remodelling of the isoform composition of IP₃Rs and ER luminal Ca²⁺ levels in colorectal cancer cell lines, *J Cell Sci.* 127 (2014) 1607–1619. doi:10.1242/jcs.141408.
- [106] A. Bononi, C. Giorgi, S. Patergnani, D. Larson, K. Verbruggen, M. Tanji, L. Pellegrini, V. Signorato, F. Olivetto, S. Pastorino, M. Nasu, A. Napolitano, G. Gaudino, P. Morris, G. Sakamoto, L.K. Ferris, A. Danese, A. Raimondi, C. Tacchetti, S. Kuchay, H.I. Pass, E.B. Affar, H. Yang, P. Pinton, M. Carbone, BAP1 regulates IP3R3-mediated Ca²⁺ flux to mitochondria suppressing cell transformation., *Nature.* 546 (2017) 549–553. doi:10.1038/nature22798.
- [107] C. Sakakura, A. Hagiwara, K. Fukuda, K. Shimomura, T. Takagi, S. Kin, Y. Nakase, J. Fujiyama, K. Mikoshiba, Y. Okazaki, H. Yamagishi, Possible involvement of inositol 1,4,5-trisphosphate receptor type 3 (IP3R3) in the peritoneal dissemination of gastric cancers., *Anticancer Res.* 23 (2003) 3691–7. <http://www.ncbi.nlm.nih.gov/pubmed/14666665>.
- [108] A. Bergner, J. Kellner, A. Tufman, R.M. Huber, Endoplasmic reticulum Ca²⁺-homeostasis is altered in Small and non-small Cell Lung Cancer cell lines., *J. Exp. Clin. Cancer Res.* 28 (2009) 25. doi:10.1186/1756-9966-28-25.
- [109] F. Iommelli, V. De Rosa, C. Terlizzi, M. Monti, M. Panico, R. Fonti, S. Del Vecchio, Inositol Trisphosphate Receptor Type 3-mediated Enhancement of EGFR and MET Cotargeting Efficacy in Non-Small Cell Lung Cancer Detected by 18F-fluorothymidine., *Clin. Cancer Res.* 24 (2018) 3126–3136. doi:10.1158/1078-0432.CCR-17-3657.
- [110] K. Shibao, M. Fiedler, J. Nagata, N. Minagawa, K. Hirata, Y. Nakayama, Y. Iwakiri, M. Nathanson, K. Yamaguchi, The type III inositol 1,4,5-trisphosphate receptor is associated with aggressiveness of colorectal carcinoma, *Cell Calcium.* 48 (2010) 315–323. doi:10.1038/jid.2014.371.
- [111] A. Singh, R.K. Sharma, M. Chagtoo, G. Agarwal, N. George, N. Sinha, M.M. Godbole, 1H NMR metabolomics reveals association of high expression of inositol 1, 4, 5 trisphosphate receptor

- and metabolites in breast cancer patients, *PLoS One*. 12 (2017) 1–20. doi:10.1371/journal.pone.0169330.
- [112] I. Rezuchova, S. Hudecova, A. Soltysova, M. Matuskova, E. Durinikova, B. Chovancova, M. Zuzcak, M. Cihova, M. Burikova, A. Penesova, L. Lencesova, J. Breza, O. Krizanov, Type 3 inositol 1,4,5-trisphosphate receptor has antiapoptotic and proliferative role in cancer cells., *Cell Death Dis.* 10 (2019) 186. doi:10.1038/s41419-019-1433-4.
- [113] A. Mound, L. Rodat-Despoix, S. Bougarn, H. Ouadid-Ahidouch, F. Matifat, Molecular interaction and functional coupling between type 3 inositol 1,4,5-trisphosphate receptor and BKCa channel stimulate breast cancer cell proliferation., *Eur. J. Cancer*. 49 (2013) 3738–51. doi:10.1016/j.ejca.2013.07.013.
- [114] B. Boutin, N. Tajeddine, G. Monaco, J. Molgo, D. Vertommen, M. Rider, J.B. Parys, G. Bultynck, P. Gailly, Endoplasmic reticulum Ca(2+) content decrease by PKA-dependent hyperphosphorylation of type 1 IP3 receptor contributes to prostate cancer cell resistance to androgen deprivation., *Cell Calcium*. 57 (2015) 312–20. doi:10.1016/j.ceca.2015.02.004.
- [115] T. Tsunoda, H. Koga, A. Yokomizo, K. Tatsugami, M. Eto, J. Inokuchi, A. Hirata, K. Masuda, K. Okumura, S. Naito, Inositol 1,4,5-trisphosphate (IP3) receptor type1 (IP3R1) modulates the acquisition of cisplatin resistance in bladder cancer cell lines., *Oncogene*. 24 (2005) 1396–402. doi:10.1038/sj.onc.1208313.
- [116] T. Szado, V. Vanderheyden, J.B. Parys, H. De Smedt, K. Rietdorf, L. Kotelevets, E. Chastre, F. Khan, U. Landegren, O. Söderberg, M.D. Bootman, H.L. Roderick, Phosphorylation of inositol 1,4,5-trisphosphate receptors by protein kinase B/Akt inhibits Ca²⁺ release and apoptosis., *Proc. Natl. Acad. Sci. U. S. A.* 105 (2008) 2427–32. doi:10.1073/pnas.0711324105.
- [117] S. Marchi, M. Marinello, A. Bononi, M. Bonora, C. Giorgi, A. Rimessi, P. Pinton, Selective modulation of subtype III IP3R by Akt regulates ER Ca²⁺ release and apoptosis, *Cell Death Dis.* 3 (2012) e304–10. doi:10.1038/cddis.2012.45.
- [118] F. Zhong, M.W. Harr, G. Bultynck, G. Monaco, J.B. Parys, H. De Smedt, Y.-P. Rong, J.K. Molitoris, M. Lam, C. Ryder, S. Matsuyama, C.W. Distelhorst, Induction of Ca²⁺-driven apoptosis in chronic lymphocytic leukemia cells by peptide-mediated disruption of Bcl-2-IP3 receptor interaction., *Blood*. 117 (2011) 2924–34. doi:10.1182/blood-2010-09-307405.
- [119] H. Akl, G. Monaco, R. La Rovere, K. Welkenhuyzen, S. Kiviluoto, T. Vervliet, J. Molgó, C.W. Distelhorst, L. Missiaen, K. Mikoshiba, J.B. Parys, H. De Smedt, G. Bultynck, IP3R2 levels dictate the apoptotic sensitivity of diffuse large B-cell lymphoma cells to an IP3R-derived peptide targeting the BH4 domain of Bcl-2, *Cell Death Dis.* 4 (2013) 1–12. doi:10.1038/cddis.2013.140.
- [120] M. Kerkhofs, T. Vervloessem, M. Bittremieux, G. Bultynck, Recent advances in uncovering the mechanisms contributing to BIRD-2-induced cell death in B-cell cancer cells., *Cell Death Dis.* 10 (2019) 42. doi:10.1038/s41419-018-1297-z.
- [121] C.W. Distelhorst, Targeting Bcl-2-IP3 receptor interaction to treat cancer: A novel approach inspired by nearly a century treating cancer with adrenal corticosteroid hormones., *Biochim. Biophys. Acta. Mol. Cell Res.* (2018). doi:10.1016/j.bbamcr.2018.07.020.
- [122] M. Bittremieux, R.M. La Rovere, M. Schuermans, T. Luyten, K. Mikoshiba, P. Vangheluwe, J.B. Parys, G. Bultynck, Extracellular and ER-stored Ca²⁺ contribute to BIRD-2-induced cell death in diffuse large B-cell lymphoma cells., *Cell Death Discov.* 4 (2018) 101. doi:10.1038/s41420-

018-0118-6.

- [123] C.W. Distelhorst, M.D. Bootman, Creating a New Cancer Therapeutic Agent by Targeting the Interaction between Bcl-2 and IP3 Receptors., *Cold Spring Harb. Perspect. Biol.* (2019). doi:10.1101/cshperspect.a035196.
- [124] A.R. Lavik, F. Zhong, M.-J. Chang, E. Greenberg, Y. Choudhary, M.R. Smith, K.S. McColl, J. Pink, F.J. Reu, S. Matsuyama, C.W. Distelhorst, A synthetic peptide targeting the BH4 domain of Bcl-2 induces apoptosis in multiple myeloma and follicular lymphoma cells alone or in combination with agents targeting the BH3-binding pocket of Bcl-2., *Oncotarget*. 6 (2015) 27388–402. doi:10.18632/oncotarget.4489.
- [125] T. Vervloessem, H. Akl, T. Tousseyn, H. De Smedt, J.B. Parys, G. Bultynck, Reciprocal sensitivity of diffuse large B-cell lymphoma cells to Bcl-2 inhibitors BIRD-2 versus venetoclax, *Oncotarget*. 8 (2017) 111656–111671. doi:10.18632/oncotarget.22898.
- [126] E.F. Greenberg, K.S. McColl, F. Zhong, G. Wildey, A. Dowlati, C.W. Distelhorst, Synergistic killing of human small cell lung cancer cells by the Bcl-2-inositol 1,4,5-trisphosphate receptor disruptor BIRD-2 and the BH3-mimetic ABT-263., *Cell Death Dis.* 6 (2015) e2034. doi:10.1038/cddis.2015.355.
- [127] M. Kerkhofs, G. Bultynck, T. Vervliet, G. Monaco, Therapeutic implications of novel peptides targeting ER-mitochondria Ca²⁺-flux systems., *Drug Discov. Today*. (2019). doi:10.1016/j.drudis.2019.03.020.
- [128] M. Abdul, S. Ramlal, N. Hoosein, Ryanodine receptor expression correlates with tumor grade in breast cancer., *Pathol. Oncol. Res.* 14 (2008) 157–60. doi:10.1007/s12253-008-9045-9.
- [129] S.E. Kobylewski, K.A. Henderson, C.D. Eckhert, Identification of ryanodine receptor isoforms in prostate DU-145, LNCaP, and PWR-1E cells., *Biochem. Biophys. Res. Commun.* 425 (2012) 431–5. doi:10.1016/j.bbrc.2012.07.119.
- [130] P. Mariot, N. Prevarskaya, M.M. Roudbaraki, X. Le Bourhis, F. Van Coppenolle, K. Vanoverberghe, R. Skryma, Evidence of functional ryanodine receptor involved in apoptosis of prostate cancer (LNCaP) cells., *Prostate*. 43 (2000) 205–14. <http://www.ncbi.nlm.nih.gov/pubmed/10797495>.
- [131] J. Tanwar, R.K. Motiani, Role of SOCE architects STIM and Orai proteins in Cell Death, *Cell Calcium*. 69 (2018) 19–27. doi:10.1016/j.ceca.2017.06.002.
- [132] V.R. Rao, M. Perez-Neut, S. Kaja, S. Gentile, Voltage-gated ion channels in cancer cell proliferation, *Cancers (Basel)*. 7 (2015) 849–875. doi:10.3390/cancers7020813.
- [133] C. Cui, R. Merritt, L. Fu, Z. Pan, Targeting calcium signaling in cancer therapy, *Acta Pharm. Sin. B*. 7 (2017) 3–17. doi:10.1016/j.apsb.2016.11.001.
- [134] J.-Y. Wang, J. Sun, M.-Y. Huang, Y.-S. Wang, M.-F. Hou, Y. Sun, H. He, N. Krishna, S.-J. Chiu, S. Lin, S. Yang, W.-C. Chang, STIM1 overexpression promotes colorectal cancer progression, cell motility and COX-2 expression., *Oncogene*. 34 (2015) 4358–67. doi:10.1038/onc.2014.366.
- [135] D. Sobradillo, M. Hernández-Morales, D. Ubierna, M.P. Moyer, L. Núñez, C. Villalobos, A reciprocal shift in transient receptor potential channel 1 (TRPC1) and stromal interaction molecule 2 (STIM2) contributes to Ca²⁺-remodeling and cancer hallmarks in colorectal carcinoma cells, *J. Biol. Chem.* 289 (2014) 28765–28782. doi:10.1074/jbc.M114.581678.

- [136] E. Zuccolo, U. Laforenza, F. Ferulli, G. Pellavio, G. Scarpellino, M. Tanzi, I. Turin, P. Faris, A. Lucariello, M. Maestri, D.A. Kheder, G. Guerra, P. Pedrazzoli, F. Moccia, Stim and Orai mediate constitutive Ca entry and control endoplasmic reticulum Ca refilling in primary cultures of colorectal carcinoma cells, *Oncotarget*. 9 (2018) 31098–31119.
- [137] N. Yang, Y. Tang, F. Wang, H. Zhang, D. Xu, Y. Shen, S. Sun, G. Yang, Blockade of store-operated Ca²⁺ entry inhibits hepatocarcinoma cell migration and invasion by regulating focal adhesion turnover, *Cancer Lett.* 330 (2013) 163–169. doi:10.1016/j.canlet.2012.11.040.
- [138] Y.-F. Chen, L.-H. Chen, M.-R. Shen, The distinct role of STIM1 and STIM2 in the regulation of store-operated Ca²⁺ entry and cellular function., *J. Cell. Physiol.* (2018). doi:10.1002/jcp.27532.
- [139] D. McAndrew, D.M. Grice, A.A. Peters, F.M. Davis, T. Stewart, M. Rice, C.E. Smart, M.A. Brown, P.A. Kenny, S.J. Roberts-Thomson, G.R. Monteith, ORAI1-Mediated Calcium Influx in Lactation and in Breast Cancer, *Mol. Cancer Ther.* 10 (2011) 448–460. doi:10.1158/1535-7163.MCT-10-0923.
- [140] R.K. Motiani, M.C. Hyzinski-García, X. Zhang, M.M. Henkel, I.F. Abdullaev, Y.H. Kuo, K. Matrougui, A.A. Mongin, M. Trebak, STIM1 and Orai1 mediate CRAC channel activity and are essential for human glioblastoma invasion, *Pflugers Arch. Eur. J. Physiol.* 465 (2013) 1249–1260. doi:10.1007/s00424-013-1254-8.
- [141] H. Zhu, H. Zhang, F. Jin, M. Fang, M. Huang, C.S. Yang, T. Chen, L. Fu, Z. Pan, Elevated Orai1 expression mediates tumor-promoting intracellular Ca²⁺ oscillations in human esophageal squamous cell carcinoma., *Oncotarget*. 5 (2014) 3455–71. doi:10.18632/oncotarget.1903.
- [142] S. Yang, J.J. Zhang, X.-Y. Huang, Orai1 and STIM1 are critical for breast tumor cell migration and metastasis., *Cancer Cell.* 15 (2009) 124–34. doi:10.1016/j.ccr.2008.12.019.
- [143] J.-H. Kim, S. Lkhagvadorj, M.-R. Lee, K.-H. Hwang, H.C. Chung, J.H. Jung, S.-K. Cha, M. Eom, Orai1 and STIM1 are critical for cell migration and proliferation of clear cell renal cell carcinoma, *Biochem. Biophys. Res. Commun.* 448 (2014) 76–82. doi:10.1016/j.bbrc.2014.04.064.
- [144] H. Stanisiz, S. Saul, C.S.L. Müller, R. Kappl, B.A. Niemeyer, T. Vogt, M. Hoth, A. Roesch, I. Bogeski, Inverse regulation of melanoma growth and migration by Orai1/STIM2-dependent calcium entry, *Pigment Cell Melanoma Res.* 27 (2014) 442–453. doi:10.1111/pcmr.12222.
- [145] C.-Y. Huang, P.-L. Wei, W.-Y. Chen, W.-C. Chang, Y.-J. Chang, Silencing Heat Shock Protein 27 Inhibits the Progression and Metastasis of Colorectal Cancer (CRC) by Maintaining the Stability of Stromal Interaction Molecule 1 (STIM1) Proteins, *Cells.* 7 (2018) 262. doi:10.3390/cells7120262.
- [146] Y. Li, B. Guo, Q. Xie, D. Ye, D. Zhang, Y. Zhu, H. Chen, B. Zhu, STIM1 Mediates Hypoxia-Driven Hepatocarcinogenesis via Interaction with HIF-1, *Cell Rep.* 12 (2015) 388–395. doi:10.1016/j.celrep.2015.06.033.
- [147] I.F. Abdullaev, J.M. Bisailon, M. Potier, J.C. Gonzalez, R.K. Motiani, M. Trebak, Stim1 and orai1 mediate crac currents and store-operated calcium entry important for endothelial cell proliferation, *Circ. Res.* 103 (2008) 1289–1299. doi:10.1161/01.RES.0000338496.95579.56.
- [148] B.C. Paria, S.M. Vogel, G.U. Ahmmed, S. Alamgir, J. Shroff, A.B. Malik, C. Tiruppathi, Tumor necrosis factor- α -induced TRPC1 expression amplifies store-operated Ca²⁺ influx and endothelial permeability, *Am. J. Physiol. Cell. Mol. Physiol.* 287 (2004) L1303–L1313.

doi:10.1152/ajplung.00240.2004.

- [149] S. Chigurupati, R. Venkataraman, D. Barrera, A. Naganathan, M. Madan, L. Paul, J. V. Pattisapu, G.A. Kyriazis, K. Sugaya, S. Bushnev, J.D. Lathia, J.N. Rich, S.L. Chan, Receptor channel TRPC6 is a key mediator of Notch-driven glioblastoma growth and invasiveness, *Cancer Res.* 70 (2010) 418–427. doi:10.1158/0008-5472.CAN-09-2654.
- [150] J. Park, M.K. Shim, M. Jin, M.-R. Rhyu, Y. Lee, Methyl syringate, a TRPA1 agonist represses hypoxia-induced cyclooxygenase-2 in lung cancer cells., *Phytomedicine.* 23 (2016) 324–9. doi:10.1016/j.phymed.2016.01.009.
- [151] A. Fiorio Pla, H.L. Ong, K.T. Cheng, A. Brossa, B. Bussolati, T. Lockwich, B. Paria, L. Munaron, I.S. Ambudkar, TRPV4 mediates tumor-derived endothelial cell migration via arachidonic acid-activated actin remodeling., *Oncogene.* 31 (2012) 200–12. doi:10.1038/onc.2011.231.
- [152] B. Feldman, S. Fedida-Metula, J. Nita, I. Sekler, D. Fishman, Coupling of mitochondria to store-operated Ca^{2+} -signaling sustains constitutive activation of protein kinase B/Akt and augments survival of malignant melanoma cells, *Cell Calcium.* 47 (2010) 525–537. doi:10.1016/j.ceca.2010.05.002.
- [153] N. Khadra, L. Bresson-bepoldin, A. Penna, B. Chaigne-delalande, B. Ségui, CD95 triggers Orai1-mediated localized Ca^{2+} entry, regulates recruitment of protein kinase C (PKC) $\beta 2$, and prevents death-inducing signaling complex formation, *Pnas.* (2011) 2–7. doi:10.1073/pnas.1116946108/-/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1116946108.
- [154] P. Vacher, A.-M. Vacher, R. Pineau, S. Latour, I. Soubeyran, C. Pangault, K. Tarte, P. Soubeyran, T. Ducret, L. Bresson-Bepoldin, Localized Store-Operated Calcium Influx Represses CD95-Dependent Apoptotic Effects of Rituximab in Non-Hodgkin B Lymphomas, *J. Immunol.* 195 (2015) 2207–2215. doi:10.4049/jimmunol.1402942.
- [155] M. Flourakis, V. Lehen'kyi, B. Beck, M. Raphaël, M. Vandenberghe, F. V Abeele, M. Roudbaraki, G. Lepage, B. Mauroy, C. Romanin, Y. Shuba, R. Skryma, N. Prevarskeya, Orai1 contributes to the establishment of an apoptosis-resistant phenotype in prostate cancer cells, *Cell Death Dis.* 1 (2010) e75. doi:10.1038/cddis.2010.52.
- [156] C. Pierro, X. Zhang, C. Kankeu, M. Trebak, M.D. Bootman, H.L. Roderick, Oncogenic KRAS suppresses store-operated Ca^{2+} entry and ICRACthrough ERK pathway-dependent remodelling of STIM expression in colorectal cancer cell lines, *Cell Calcium.* 72 (2018) 70–80. doi:10.1016/j.ceca.2018.03.002.
- [157] F. Vanden Abeele, Y. Shuba, M. Roudbaraki, L. Lemonnier, K. Vanoverberghe, P. Mariot, R. Skryma, N. Prevarskeya, Store-operated Ca^{2+} channels in prostate cancer epithelial cells: Function, regulation, and role in carcinogenesis, *Cell Calcium.* 33 (2003) 357–373. doi:10.1016/S0143-4160(03)00049-6.
- [158] R. Hooper, X. Zhang, M. Webster, C. Go, J. Kedra, K. Marchbank, D.L. Gill, A.T. Weeraratna, M. Trebak, J. Soboloff, Novel Protein Kinase C-Mediated Control of Orai1 Function in Invasive Melanoma, *Mol. Cell. Biol.* 35 (2015) 2790–2798. doi:10.1128/MCB.01500-14.
- [159] I. Azimi, A.H. Bong, G.X.H. Poo, K. Armitage, D. Lok, S.J. Roberts-Thomson, G.R. Monteith, Pharmacological inhibition of store-operated calcium entry in MDA-MB-468 basal A breast cancer cells: consequences on calcium signalling, cell migration and proliferation., *Cell. Mol. Life Sci.* 75 (2018) 4525–4537. doi:10.1007/s00018-018-2904-y.

- [160] Y. Ruano, M. Mollejo, T. Ribalta, C. Fiaño, F.I. Camacho, E. Gómez, A.R. de Lope, J.-L. Hernández-Moneo, P. Martínez, B. Meléndez, Identification of novel candidate target genes in amplicons of Glioblastoma multiforme tumors detected by expression and CGH microarray profiling, *Mol. Cancer*. 5 (2006) 39. doi:10.1186/1476-4598-5-39.
- [161] A. Aytes, D.G. Molleví, M. Martinez-Iniesta, M. Nadal, A. Vidal, A. Morales, R. Salazar, G. Capellà, A. Villanueva, Stromal interaction molecule 2 (STIM2) is frequently overexpressed in colorectal tumors and confers a tumor cell growth suppressor phenotype., *Mol. Carcinog.* 51 (2012) 746–53. doi:10.1002/mc.20843.
- [162] R.K. Motiani, I.F. Abdullaev, M. Trebak, A novel native store-operated calcium channel encoded by Orai3: Selective requirement of Orai3 versus Orai1 in estrogen receptor-positive versus estrogen receptor-negative breast cancer cells, *J. Biol. Chem.* 285 (2010) 19173–19183. doi:10.1074/jbc.M110.102582.
- [163] M. Faouzi, F. Hague, M. Potier, A. Ahidouch, H. Sevestre, H. Ouadid-Ahidouch, Down-regulation of Orai3 arrests cell-cycle progression and induces apoptosis in breast cancer cells but not in normal breast epithelial cells, *J. Cell. Physiol.* 226 (2011) 542–551. doi:10.1002/jcp.22363.
- [164] C. Dubois, F. Vanden Abeele, V. Lehen'kyi, D. Gkika, B. Guarmit, G. Lepage, C. Slomianny, A. Borowiec, G. Bidaux, M. Benahmed, Y. Shuba, N. Prevarskaya, Remodeling of Channel-Forming ORAI Proteins Determines an Oncogenic Switch in Prostate Cancer, *Cancer Cell*. 26 (2014) 19–32. doi:10.1016/j.ccr.2014.04.025.
- [165] W.-T. Chiu, H.-A. Chang, Y.-H. Lin, Y.-S. Lin, H.-T. Chang, H.-H. Lin, S.-C. Huang, M.-J. Tang, M.-R. Shen, Bcl-2 regulates store-operated Ca²⁺ entry to modulate ER stress-induced apoptosis., *Cell Death Discov.* 4 (2018) 37. doi:10.1038/s41420-018-0039-4.
- [166] R.M. Luik, B. Wang, M. Prakriya, M.M. Wu, R.S. Lewis, Oligomerization of STIM1 couples ER calcium depletion to CRAC channel activation., *Nature*. 454 (2008) 538–42. doi:10.1038/nature07065.
- [167] M.K. Park, The endoplasmic reticulum as one continuous Ca²⁺ pool: visualization of rapid Ca²⁺ movements and equilibration, *EMBO J.* 19 (2000) 5729–5739. doi:10.1093/emboj/19.21.5729.
- [168] C. Peinelt, A. Beck, M.K. Monteilh-Zoller, R. Penner, A. Fleig, IP3 receptor subtype-dependent activation of store-operated calcium entry through ICRAC, *Cell Calcium*. 45 (2009) 326–330. doi:10.1016/j.ceca.2008.12.001.
- [169] M.J. Berridge, Capacitative calcium entry., *Biochem. J.* 312 (Pt 1 (1995) 1–11. <http://www.ncbi.nlm.nih.gov/pubmed/7492298>.
- [170] N.B. Thillaiappan, A.P. Chavda, S.C. Tovey, D.L. Prole, C.W. Taylor, Ca²⁺ signals initiate at immobile IP3 receptors adjacent to ER-plasma membrane junctions., *Nat. Commun.* 8 (2017) 1505. doi:10.1038/s41467-017-01644-8.
- [171] A.I. Zubov, E. V Kaznacheeva, A. V Nikolaev, V.A. Alexeenko, K. Kiselyov, S. Muallem, G.N. Mozhayeva, Regulation of the miniature plasma membrane Ca(2+) channel I(min) by inositol 1,4,5-trisphosphate receptors., *J. Biol. Chem.* 274 (1999) 25983–5. <http://www.ncbi.nlm.nih.gov/pubmed/10473541>.
- [172] G. Lur, M.W. Sherwood, E. Ebisui, L. Haynes, S. Feske, R. Sutton, R.D. Burgoyne, K. Mikoshiba, O.H. Petersen, A. V Tepikin, InsP₃ receptors and Orai channels in pancreatic acinar cells: co-

- localization and its consequences., *Biochem. J.* 436 (2011) 231–239. doi:10.1042/BJ20110083.
- [173] R. Courjaret, K. Machaca, Mid-range Ca^{2+} signalling mediated by functional coupling between store-operated Ca^{2+} entry and IP_3 -dependent Ca^{2+} release., *Nat. Commun.* 5 (2014) 3916. doi:10.1038/ncomms4916.
- [174] H. Mogami, K. Nakano, A. V Tepikin, O.H. Petersen, Ca^{2+} flow via tunnels in polarized cells: recharging of apical Ca^{2+} stores by focal Ca^{2+} entry through basal membrane patch., *Cell.* 88 (1997) 49–55. <http://www.ncbi.nlm.nih.gov/pubmed/9019404>.
- [175] R. Courjaret, M. Dib, K. Machaca, Store-Operated Ca^{2+} Entry in Oocytes Modulate the Dynamics of IP_3 -Dependent Ca^{2+} Release From Oscillatory to Tonic., *J. Cell. Physiol.* 232 (2017) 1095–1103. doi:10.1002/jcp.25513.
- [176] S. Chakraborty, B.K. Deb, T. Chorna, V. Konieczny, C.W. Taylor, G. Hasan, Mutant IP_3 receptors attenuate store-operated Ca^{2+} entry by destabilizing STIM-Orai interactions in *Drosophila* neurons., *J. Cell Sci.* 129 (2016) 3903–3910. doi:10.1242/jcs.191585.
- [177] J.T. Smyth, J.G. Petranka, R.R. Boyles, W.I. DeHaven, M. Fukushima, K.L. Johnson, J.G. Williams, J.W. Putney Jr, Phosphorylation of STIM1 underlies suppression of store-operated calcium entry during mitosis, *Nat. Cell Biol.* 11 (2009) 1465–1472. doi:10.1038/ncb1995.
- [178] M. Bittremieux, R.M. la Rovere, H. Akl, C. Martines, K. Welkenhuyzen, K. Dubron, M. Baes, A. Janssens, P. Vandenberghe, L. Laurenti, K. Rietdorf, G. Morciano, P. Pinton, K. Mikoshiba, M.D. Bootman, D.G. Efremov, H. de Smedt, J.B. Parys, G. Bultynck, Constitutive IP_3 signaling underlies the sensitivity of B-cell cancers to the Bcl-2/ IP_3 receptor disruptor BIRD-2, *Cell Death Differ.* (2018) 1–17. doi:10.1038/s41418-018-0142-3.
- [179] L. Fan, A. Li, W. Li, P. Cai, B. Yang, M. Zhang, Y. Gu, Y. Shu, Y. Sun, Y. Shen, X. Wu, G. Hu, X. Wu, Q. Xu, Novel role of Sarco/endoplasmic reticulum calcium ATPase 2 in development of colorectal cancer and its regulation by F36, a curcumin analog., *Biomed. Pharmacother.* 68 (2014) 1141–8. doi:10.1016/j.biopha.2014.10.014.
- [180] P. Gélébart, T. Kovács, J.P. Brouland, R. Van Gorp, J. Grossmann, N. Rivard, Y. Panis, V. Martin, R. Bredoux, J. Enouf, B. Papp, Expression of endomembrane calcium pumps in colon and gastric cancer cells. Induction of SERCA3 expression during differentiation, *J. Biol. Chem.* 277 (2002) 26310–26320. doi:10.1074/jbc.M201747200.
- [181] J.P. Brouland, P. Gélébart, T. Kovács, J. Enouf, J. Grossmann, B. Papp, The loss of sarco/endoplasmic reticulum calcium transport ATPase 3 expression is an early event during the multistep process of colon carcinogenesis, *Am. J. Pathol.* 167 (2005) 233–242. doi:10.1016/S0002-9440(10)62968-9.
- [182] O. Dellis, A. Arbabian, J.P. Brouland, T. Kovács, M. Rowe, C. Chomienne, I. Joab, B. Papp, Modulation of b-cell endoplasmic reticulum calcium homeostasis by epstein-barr virus latent membrane protein-1, *Mol. Cancer.* 8 (2009) 59. doi:10.1186/1476-4598-8-59.
- [183] L. Aït Ghezali, A. Arbabian, H. Roudot, J.P. Brouland, F. Baran-Marszak, E. Salvaris, A. Boyd, H.G. Drexler, A. Enyedi, R. Letestu, N. Varin-Blank, B. Papp, Induction of endoplasmic reticulum calcium pump expression during early leukemic B cell differentiation, *J. Exp. Clin. Cancer Res.* 36 (2017) 1–12. doi:10.1186/s13046-017-0556-7.
- [184] K. Saito, K. Uzawa, Y. Endo, Y. Kato, D. Nakashima, K. Ogawara, M. Shiba, H. Bukawa, H. Yokoe, H. Tanzawa, Plasma membrane Ca^{2+} ATPase isoform 1 down-regulated in human oral cancer., *Oncol. Rep.* 15 (2006) 49–55. <http://www.ncbi.nlm.nih.gov/pubmed/16328033>.

- [185] J. VanHouten, C. Sullivan, C. Bazinet, T. Ryoo, R. Camp, D.L. Rimm, G. Chung, J. Wysolmerski, PMCA2 regulates apoptosis during mammary gland involution and predicts outcome in breast cancer, *Proc. Natl. Acad. Sci.* 107 (2010) 11405–11410. doi:10.1073/pnas.0911186107.
- [186] W.J. Lee, S.J. Roberts-Thomson, G.R. Monteith, Plasma membrane calcium-ATPase 2 and 4 in human breast cancer cell lines., *Biochem. Biophys. Res. Commun.* 337 (2005) 779–83. doi:10.1016/j.bbrc.2005.09.119.
- [187] P. Ribiczey, A. Tordai, H. Andrikovics, A.G. Filoteo, J.T. Penniston, J. Enouf, A. Enyedi, B. Papp, T. Kovács, Isoform-specific up-regulation of plasma membrane Ca²⁺-ATPase expression during colon and gastric cancer cell differentiation., *Cell Calcium.* 42 (2007) 590–605. doi:10.1016/j.ceca.2007.02.003.
- [188] C.S. Aung, W.A. Kruger, P. Poronnik, S.J. Roberts-Thomson, G.R. Monteith, Plasma membrane Ca²⁺-ATPase expression during colon cancer cell line differentiation, *Biochem. Biophys. Res. Commun.* 355 (2007) 932–936. doi:10.1016/j.bbrc.2007.02.050.
- [189] D.M. Grice, I. Vetter, H.M. Faddy, P.A. Kenny, S.J. Roberts-Thomson, G.R. Monteith, Golgi calcium pump secretory pathway calcium ATPase 1 (SPCA1) is a key regulator of Insulin-like Growth Factor Receptor (IGF1R) processing in the basal-like breast cancer cell line MDA-MB-231, *J. Biol. Chem.* 285 (2010) 37458–37466. doi:10.1074/jbc.M110.163329.
- [190] K.W. Kohn, B.M. Zeeberg, W.C. Reinhold, Y. Pommier, Gene expression correlations in human cancer cell lines define molecular interaction networks for epithelial phenotype, *PLoS One.* 9 (2014). doi:10.1371/journal.pone.0099269.
- [191] J. Tang, Y. Guo, X. Yu, W. Huang, M. Zheng, Z. Chen, CD147 reinforces [Ca²⁺] oscillations and promotes oncogenic progression in hepatocellular carcinoma, 6 (2015). doi:10.18632/oncotarget.5225.
- [192] H. Li, J. Xu, Y. Zhou, X. Liu, L.E. Shen, Y.U. Zhu, Z. Li, X. Wang, Q. Guo, H. Hui, PLSCR1/IP3R1/Ca²⁺ axis contributes to differentiation of primary AML cells induced by wogonoside, *Cell Death Dis.* 8 (2017) e2768-14. doi:10.1038/cddis.2017.175.
- [193] V. Prasad, G.P. Boivin, M.L. Miller, L.H. Liu, C.R. Erwin, B.W. Warner, G.E. Shull, Haploinsufficiency of Atp2a2, encoding the sarco(endo)plasmic reticulum Ca²⁺-ATPase isoform 2 Ca²⁺ pump, predisposes mice to squamous cell tumors via a novel mode of cancer susceptibility, *Cancer Res.* 65 (2005) 8655–8661. doi:10.1158/0008-5472.CAN-05-0026.
- [194] G. Roti, A. Carlton, K.N. Ross, M. Markstein, K. Pajcini, A.H. Su, N. Perrimon, W.S. Pear, A.L. Kung, S.C. Blacklow, J.C. Aster, K. Stegmaier, Complementary Genomic Screens Identify SERCA as a Therapeutic Target in NOTCH1 Mutated Cancer, *Cancer Cell.* 23 (2013) 390–405. doi:10.1016/j.ccr.2013.01.015.
- [195] S. Marchi, M. Corricelli, A. Branchini, V.A.M. Vitto, S. Missiroli, G. Morciano, M. Perrone, M. Ferrarese, C. Giorgi, M. Pinotti, L. Galluzzi, G. Kroemer, P. Pinton, Akt-mediated phosphorylation of MICU1 regulates mitochondrial Ca²⁺ levels and tumor growth, *EMBO J.* (2018) e99435. doi:10.15252/embj.201899435.
- [196] S. Hudecova, J. Markova, V. Simko, L. Csaderova, T. Stracina, M. Sirova, M. Fojtu, E. Svastova, P. Gronesova, M. Pastorek, M. Novakova, D. Cholujo, J. Kopacek, S. Pastorekova, J. Sedlak, O. Krizanov, Sulforaphane-induced apoptosis involves the type 1 IP3 receptor., *Oncotarget.* 7 (2016) 61403–61418. doi:10.18632/oncotarget.8968.
- [197] C. Giorgi, M. Bonora, G. Sorrentino, S. Missiroli, F. Poletti, J.M. Suski, F. Galindo Ramirez, R.

- Rizzuto, F. Di Virgilio, E. Zito, P.P. Pandolfi, M.R. Wieckowski, F. Mammano, G. Del Sal, P. Pinton, p53 at the endoplasmic reticulum regulates apoptosis in a Ca^{2+} -dependent manner, *Proc. Natl. Acad. Sci.* 112 (2015) 1779–1784. doi:10.1073/pnas.1410723112.
- [198] M.C. Curry, N.A. Luk, P.A. Kenny, S.J. Roberts-Thomson, G.R. Monteith, Distinct regulation of cytoplasmic calcium signals and cell death pathways by different plasma membrane calcium ATPase isoforms in MDA-MB-231 breast cancer cells, *J. Biol. Chem.* 287 (2012) 28598–28608. doi:10.1074/jbc.M112.364737.
- [199] X. Huang, M. Jin, Y.X. Chen, J. Wang, K. Zhai, Y. Chang, Q. Yuan, K.T. Yao, G. Ji, ERP44 inhibits human lung cancer cell migration mainly via IP3R2, *Aging (Albany, NY)*. 8 (2016) 1276–1286. doi:10.18632/aging.100984.
- [200] J. Prudent, N. Popgeorgiev, R. Gadet, M. Deygas, R. Rimokh, G. Gillet, Mitochondrial Ca^{2+} uptake controls actin cytoskeleton dynamics during cell migration, *Nat. Sci. Reports*. 6 (2016) 36570. doi:10.1038/srep36570.
- [201] C. Yu, Y. Wang, J. Peng, Q. Shen, M. Chen, W. Tang, X. Li, C. Cai, B. Wang, S.C. Cai, X. Meng, F. Zou, Mitochondrial calcium uniporter as a target of microRNA-340 and promoter of metastasis via enhancing the Warburg effect, *Oncotarget*. 8 (2017) 83831–83844. doi:10.1016/S1062-2896(49)80057-2.

Figure Legends

Fig. 1. An overview of Ca²⁺ signals. Ca²⁺ signals are created in the ON reactions (in green), where Ca²⁺ channels on the plasma membrane or on the membranes of intracellular stores open to increase the intracellular Ca²⁺ concentration. Ca²⁺ signals are terminated in the OFF reactions (in red), where Ca²⁺ pumps on the same membranes extrude Ca²⁺ outside the cytoplasm. Ca²⁺ effectors and buffers contribute to starting Ca²⁺-dependent processes and to the modulation of Ca²⁺ signals. Mitochondrial channels (in yellow) contribute to the regulation of Ca²⁺ signals as well as the cellular bioenergetics.

Ca ²⁺ channel	Up/down regulation	Cancer type	Refs	notes
INSP31	Up Up/high phosphorylation	colorectal cancer cell lines prostate cancer cell lines	[105] [114]	low ER content low ER content
INSP32	Up Up	breast cancer cell lines and patients diffuse large B-cell lymphoma cell lines (DLBCL)	[111] [178]	
INSP33	Down Up Up Up Up Up	colorectal cancer cell lines malignant gastric cancer cell lines small and non-small lung cancer cell lines colorectal cancer patients glioblastoma primary cells breast cancer cell lines and patients	[105] [107] [108,109] [110] [98] [111]	not in primary tumour-derived cells
STIM1	Up Up Up Down Up Same but SOCE diminished Up Up in invasive front Up Down	(advanced) hepatocellular carcinoma patients colorectal cancer patients colorectal cancer cell lines colorectal cancer cell lines (HCT116) colon cancer cell lines (DLD-1) invasive melanoma primary cells metastatic colorectal cancer primary cells cervical cancer cell lines Breast cancer patients (high STIM1/STIM2 ratio) Mammary epithelium during lactation	[137] [134] [135] [156] [156] [158] [136] [138] [139] [139]	PKC
STIM2	Up Up Down Up Down Up	Glioblastoma patients colorectal cancer patients colorectal cancer cells colorectal cancer cells (HCT116) colon cancer cells (DLD-1) Mammary epithelium during lactation	[160] [161] [135] [156] [156] [139]	
Orai1	Up Up Up Up	Malignant glioblastoma primary cells Malignant oesophageal cancer primary cells colon cancer cell lines breast cancer primary cells	[140] [141] [135] [139]	
Orai2	Up Up	colon cancer cell lines metastatic colorectal cancer primary cells	[135] [136]	
Orai3	Up Up	colon cancer cell lines prostate cancer cell lines	[135] [164]	
SERCA 2	Down	small/non-small lung cancer cell lines	[108]	lower ER content

	Down Up	colon cancer cell lines colon cancer cell lines and primary tumours	[105] [179]	
SERCA 3	Down Down Down Down	Colon and gastric cell lines Colon cancer primary cells Leukemia cells (immortalization) Leukemia primary cells	[180] [181] [182] [183]	
PMCA1	Down	oral cancer primary cells and cell lines	[184]	early epigenetic silencing
PMCA2	Up Up	breast cancer primary cells (normally decreased during weaning) breast cancer cell lines	[185] [186]	
PMCA4	Up during differentiation Up during differentiation	colon and gastric cancer cell lines colon cancer cell lines	[187] [188]	
SPCA1	Up	primary basal-like breast tumours	[189]	
SPCA2	Up	epithelial cancer cell lines vs mesenchymal cancer cell lines	[190]	EMT, metastasis

Table 1. Summary of alterations in expression of different Ca²⁺ toolkit proteins in different cancers.

Tumour stage	Ca ²⁺ channel	Event	Cancer type	Refs	Notes
Early: growth control independency	INSP31	Promotes Ca ²⁺ oscillations which sustain proliferation	hepatocellular carcinoma	[191]	
		Mediates wogonoside-induced differentiation	primary acute myeloid leukemia	[192]	
	INSP33	2-APB Inhibition blocks proliferation	gastric cancer	[107]	
		Silencing reduces proliferation	breast cancer	[113]	
		2-APB/XestC inhibition reduces growth	breast cancer	[99]	
	STIM1	STIM1/Orai1 siRNA blocks in G1/S	cervical cancer	[58]	passive G2/M
		Silencing blocks in S and G2/M	cervical cancer	[59]	
		siRNA impairs proliferation	clear cell renal carcinoma	[143]	
	Orai1	Inhibition suppresses growth	oesophageal cancer	[141]	
	Orai3	Inhibition blocks proliferation	Breast cancer	[163]	
	SERCA2	Heterozigosity confers tumorigenicity	squamous cell tumours	[193]	
		Silencing inhibits proliferation	CRC cells	[179]	
		SERCA inhibition provokes G ₀ /G ₁ arrest	leukemia (Notch +)	[194]	
	MCU/MICU1	MICU1 inhibitory phosphorylation by AKT sustains growth	Lung cancer	[195]	increased ROS

Table 2 (a). Implications of Ca²⁺ toolkit proteins at early stages of tumorigenesis (uncontrolled growth).

Tumour stage	Ca ²⁺ channel	Event	Cancer type	Ref.s	Notes
Early: apoptosis evasion	INSP3R1	Mediates sulforaphane-dependent apoptosis	ovarian cancer cell lines	[196]	nuclear translocation
		Mediates androgen removal-derived apoptosis	prostate cancer cell lines	[114]	
		Downregulation protects from cisplatin-induced apoptosis	bladder cancer cell lines	[115]	
		Silencing reduces apoptosis	colorectal cancer	[112]	
	INSP3R2	Mediates apoptosis at MAMs through interaction with Bcl-2	DLBCL/CLL cell lines	[119,178]	
	INSP3R3	2-APB inhibition induces apoptosis	gastric cancer cell lines	[107]	
		2-APB inhibition enhances apoptosis	colorectal cancer patients	[110]	
		Silencing inhibits tumour growth and induces apoptosis	colorectal cancer ovarian cancer renal cancer	[112]	
	STIM1	Inhibits extrinsic apoptosis	leukemic cell lines	[153,154]	
		Increased SOCE improves survival	Malignant melanoma cell lines	[152]	
	Orai1	Downregulation confers apoptosis resistance	prostate cancer cell lines	[155]	
	Orai3	Inhibition induces proliferation	Breast cancer cell lines	[163]	NFAT
		Mediates SOCE	ER+ breast cancer cells	[162]	
		Promotes proliferation via AA channels	prostate cancer cell lines	[164]	
	SERCA2	p53 stimulates activity inducing	Cervical, lung, breast,	[197]	

		mitochondrial pro-apoptotic uptake	colorectal cancer cells, fibroblasts		
	PMCA1	Silencing enhances ionomycin-induced necrosis	breast cancer cell lines	[198]	
	PMCA2	Overexpression protects from apoptosis	breast cancer cell lines	[185]	
	PMCA4	Silencing enhances ABT-263-dependent apoptosis	breast cancer cell lines	[198]	
	SPCA1	Inhibition decreases IGFR production	basal-like breast tumours	[189]	
	MCU/MICU1	Increased mitochondrial Ca^{2+} flux improves survival Inhibition of Ca^{2+} flux induces apoptosis	Malignant melanoma cell lines Breast and prostate cancer cells, transformed fibroblasts	[152] [74]	

Table 2 (b). Implications of Ca^{2+} toolkit proteins at early stages of tumorigenesis (apoptosis evasion).

Tumour stage	Ca ²⁺ channel	Event	Cancer type	Refs	Notes
Late: Angiogenesis Migration invasion metastasis	INSP3R2	2-APB inhibition impairs migration	lung adenocarcinoma cell lines	[199]	
	INSP3R3	Silencing favours rounded shape (non-migratory) and remodels cytoskeleton	breast cancer cell lines	[104]	EMT
		Caffeine inhibition reduces migration	primary glioblastoma cells	[98]	
		Silencing decreases migration	breast cancer cell lines	[103]	
	STIM1	siRNA blocks metastasis	breast cancer cell lines	[142]	
		siRNA reduces invasion	Glioblastoma primary cells	[140]	
		siRNA reduces focal adhesion turnover	hepatocellular carcinoma patients	[137]	
		siRNA impairs migration	primary clear cell renal carcinoma cells	[143]	
		shRNA inhibits migration	colorectal cancer patients	[134]	
		HIF-1 inhibition downregulates STIM1 and vice versa		[146]	
	STIM2	Inhibition reduces migration	Melanoma cell lines	[144]	
		Does not regulate migration	Cervical cancer cell lines	[138]	
	Orai1	Inhibition reduces migration	Melanoma cell lines	[144]	
	SERCA2	Silencing inhibits migration	Colorectal cancer cell lines	[179]	
	MCU	Silencing reduces migration	Cervical and breast cancer cell lines	[200]	miRNA-340
		Silencing reduces migration, overexpression enhances metastasis	breast cancer cell lines	[201]	

Table 2 (c). Implications of Ca²⁺ toolkit proteins at late stages of tumorigenesis (metastasis).