Failure to diverge in African Great Lakes: the case of

***Dolicirroplectanum lacustre* comb. nov. (Monogenea,**

Diplectanidae) infecting latid hosts

Nikol Kmentová^{1,2,3*}, Stephan Koblmüller⁴, Maarten Van Steenberge^{3,5,6}, Tom Artois²,

Fidel Muterezi Bukinga⁷, Théophile Mulimbwa N'sibula⁷, Donatien Muzumani Risasi⁷,

Pascal Masilya Mulungula⁷, Milan Gelnar¹, Maarten P.M. Vanhove^{1,2,3,8}

¹ Department of Botany and Zoology, Faculty of Science, Masaryk University, Kotlářská 2,
611 37 Brno, Czech Republic

² Hasselt University, Centre for Environmental Sciences, Research Group Zoology:
Biodiversity & Toxicology, Agoralaan Gebouw D, B-3590 Diepenbeek, Belgium

³ Laboratory of Biodiversity and Evolutionary Genomics, Department of Biology, University
of Leuven, Ch. Deberiotstraat 32, B-3000 Leuven, Belgium

⁴ Institute of Biology, University of Graz, Universitätsplatz 2, A-8010 Graz, Austria

⁵ Operational Directorate Taxonomy and Phylogeny, Royal Belgian Institute of Natural
Sciences, Vautierstraat 29, B-1000 Brussels, Belgium

⁶ Biology Department, Royal Museum for Central Africa, Leuvensesteenweg 13, 3080,
Tervuren, Belgium

⁷ Centre de Recherche en Hydrobiologie, Département de Biologie, B.P. 73 Uvira,
Democratic Republic of Congo

⁸ Finnish Museum of Natural History, Zoology Unit, University of Helsinki, Finland

* corresponding author: kmentovan@mail.muni.cz, +420 723 004 419

Abstract

Speciation of fish in the African Great Lakes has been widely studied. Surprisingly, extensive speciation in parasites was only recently discovered in these biodiversity hotspots, notably in monogeneans (Platyhelminthes) from Lake Tanganyika. *Diplectanum* is a monogenean genus of which only a single species is known from the Great Lakes: *Diplectanum lacustre* (Diplectanidae) living on latid perches of Lake Albert. Despite their primary marine origin, latids have diversified in African freshwaters including several Great Lakes. In better-studied marine diplectanid species, incongruence between morphological and genetic differentiation was documented. As freshwater systems provide more opportunities for speciation than the marine realm, we ask whether diplectanids of *Lates* spp. of the Great Lakes underwent similar diversification as their hosts.

Fresh and museum specimens of five African latid species (*Lates angustifrons*, *L. mariae*, *L. microlepis*, *L. niloticus*, *L. stappersii*) were examined for the presence of monogenean gill parasites. Monogeneans were characterised morphologically via morphometrics of sclerotised structures and genetically using nuclear ribosomal and mitochondrial markers.

Continuous morphological variation was documented in these parasites. In addition, the genetic distance, based on the COI region, between parasites of geographically isolated host species did not reach the level typically associated with distinct diplectanid species.

Therefore, a single species of a newly described genus, *Dolicirroplectanum lacustre* gen. nov. comb. nov. is suggested to infect latid species in the examined basins. We discuss this parasite's failure to diverge in the light of the congruence between the rate of molecular evolution in COI and host historical distribution.

Keywords: parasitic flatworm - *Lates* – DNA barcoding - evolutionary history - Nile perches

Introduction

African Great Lakes are known for their species rich flocks of cichlid fish, that are well established models in evolutionary biology (Salzburger, 2018). Remarkably, Lake Tanganyika is characterised by extraordinary diversity and high degrees of endemism not only of cichlids but also other fish families (Salzburger et al., 2014) as well as invertebrate taxa (Coulter, 1991) including parasitic flatworms (Pariselle et al., 2015). Monogeneans (Platyhelminthes) are mainly parasites of fish. They display a high level of host specificity believed to be connected with their direct life cycle (a single host needed) combined with the adaptive evolution of monogenean hardparts responsible for attachment (the haptor in the posterior part of the body) and reproduction (Poulin, 2002).

Parasite speciation mirroring host diversification was reported for monogeneans infecting tropheine cichlids in Lake Tanganyika (Vanhove et al., 2015). However, the history of monogenean interactions with their hosts does not always feature only co-speciation events. For example, in *Dactylogyrus* Diesing, 1850 infecting cyprinid fishes, diversification can be mainly explained by intrahost speciation (Šimková et al., 2004). In the case of *Cichlidogyrus casuarinus* Pariselle, Muterezi Bukinga & Vanhove, 2015, a monogenean infecting deepwater cichlids in Lake Tanganyika, no specificity or preference was detected towards its various well-diverged host species (Kmentová et al., 2016), a process called “failure to diverge” (Brooks, 1979).

Among the parasitic flatworms known to infect lates perches (Latidae) are diplectanids (Diplectanidae), a monogenean family with more than 250 species described worldwide, mainly from marine perciform fishes (Domingues and Boeger, 2008). The genus *Lates* L., 1758 consists of 11 species, seven of which inhabit African freshwaters, with the rest to be found in marine, brackish and freshwater habitats in the Indo-Pacific region (Otero, 2004). While seven diplectanid species from three different genera were documented from *Lates*

81 *calcarifer* (Bloch, 1790) from the Indo-Pacific region (Tingbao et al., 2006), only one species
82 was described from African *Lates* spp. so far: *Diplectanum lacustre* Thurston & Paperna,
83 1969 infecting *Lates niloticus* L. from Lake Volta and the Victoria Nile near Lake Albert
84 (Paperna and Thurston, 1969), and from near Cairo in Egypt (Ergens, 1981). The native range
85 of *L. niloticus* includes most major river basins and Great Lakes in the Nilo-Sudanic region
86 and large parts of the Congo basin (Paugy et al., 2003). Importantly, *L. niloticus* was
87 introduced to Lake Victoria for fisheries with a dramatic impact on the local environment
88 (Ogutu-Ohwayo, 1995). In Lake Tanganyika, four endemic latid fishes, *Lates angustifrons*
89 Boulenger, 1906, *Lates mariae* Steindachner, 1909, *Lates microlepis* Boulenger, 1898 and
90 *Lates stappersii* (Boulenger, 1914), with different habitat preferences, are present (Poll,
91 1953).

92 Small interspecific morphological differences and high levels of phenotypic plasticity render
93 the species status of some diplectanids questionable (Poisot et al., 2011; Schoelinck et al.,
94 2012; Wu et al., 2005) with unclear phylogenetic relationships within and between some of
95 the genera (Villar-Torres et al., 2019). In the 21st century, species delineation is often based
96 on a combination of morphological and molecular data (Schlick-Steiner et al., 2010).

97 Integrative techniques revealed problems in various taxonomic groups, especially in soft-
98 bodied organisms or lineages with heteromorphic life stages such as parasitic flatworms
99 (Georgieva et al., 2013; Rahmouni et al., 2017). Species identification using specific

100 molecular tags derived from the cytochrome *c* oxidase subunit 1 gene (COI) in the
101 mitochondrial DNA, known as DNA barcoding, was successfully implemented in many
102 taxonomic groups such as fishes (Hubert et al., 2008), mammals (Francis et al., 2010) and
103 lepidopteran insects (Hebert et al., 2003). However, this approach proved problematic in
104 many other taxa (DeSalle et al., 2005; Will and Rubinoff, 2004) including monogenean
105 flatworms (Vanhove et al., 2013). So far, little correlation between host specificity and

taxonomic diversification was found in diplectanid monogeneans (Desdevises et al., 2001; Villar-Torres et al., 2019). However, the latter studies were conducted in a marine system with no real geographic barrier between host species that even form mixed schools. As freshwater systems provide more opportunities for speciation than the marine realm, and given the age of Lakes Albert and Tanganyika, which are situated in different basins, these lakes are a perfect study system to investigate diplectanid evolution under allopatry. We hypothesize that diplectanid monogeneans infecting latids belong to different species in Lake Albert and Lake Tanganyika. If so, is there a congruence between the level of morphological and molecular diversification in diplectanid parasites infecting African of *Lates* spp.? Within Lake Tanganyika, we ask whether diplectanids of *Lates* spp. underwent similar diversification as their hosts, or whether they rather failed to diverge like the above-mentioned *C. casuarinus*, a monogenean infecting bathybatine cichlids. These are, like latids, non-littoral fishes, and this lack of parasite specificity is considered an adaptation of low host availability outside of the littoral zone (Kmentová et al., 2016).

Material & Methods

Sampling

Fish samples of five latid species (*Lates angustifrons*, *L. mariae*, *L. microlepis*, *L. niloticus*, *L. stappersii*) were examined in this study. Samples included specimens of all *Lates* species from the ichthyology collection of the Royal Museum for Central Africa (RMCA) (Tervuren, Belgium) and fresh specimens from recent field expeditions (2010, 2016, 2017 and 2018). At Lake Albert, fresh specimens of *L. niloticus* were obtained from local fishermen (Nzunzu, Uganda). For Lake Tanganyika, the four endemic latid species (*Lates angustifrons*, *L. mariae*, *L. microlepis* and *L. stappersii*) were either caught with gill nets from the experimental fishing unit of the Centre de Recherche en Hydrobiologia-Uvira (CRH) (Uvira, Democratic

Republic of the Congo) or obtained from local fish markets (see Table 1). To provide a broader geographical range for morphological comparison, fish specimens of *L. niloticus* from seven additional localities throughout the host's range were examined. In total, gills (one side in the case of museum specimens) of 158 fish specimens from 20 localities in African freshwaters (see Table 1) were examined following the standard protocol of Ergens & Lom (Ergens and Lom, 1970). In the field, fresh monogenean specimens were either mounted on slides using a solution of glycerine ammonium picrate (GAP) or using Hoyer's medium in the case of ethanol-fixed specimens from Lake Albert and specimens retrieved from the museum collection. Some of the individuals were cut in three parts with the anterior and posterior parts mounted on slides for morphological characterisation and the rest preserved in 99% ethanol for genetic analyses. To characterize internal anatomy, some specimens were stained using the Carmine method described by Justine (2005) without the initial step of putting a live parasite under a cover slip. Parasite identification and description were carried out using an Olympus BX51 microscope equipped with a drawing tube and OLYMPUS KL 1500 LED illumination. Specimens were compared with the holotype (MRAC MT.35572) and voucher material (MRAC MT.35573) of *D. lacustre*. Drawings were edited with a graphics tablet compatible with Adobe Illustrator CS6 16.0.0 and Adobe Photoshop CS6 13.0. Fish tissue samples were deposited in the ichthyology collection of the RMCA under collection number 2016.20.P for Lake Tanganyika and 2016.036.P for Lake Albert. Parasite voucher specimens are available from the invertebrate collection of the RMCA, the Iziko South African Museum (SAMC), Cape Town, Republic of South Africa; the Muséum national d'Histoire naturelle (MNHN), Paris, France; the Natural History Museum (NHMUK), London, United Kingdom; and the Finnish Museum of Natural History (MZH), Helsinki, Finland.

Morphometrics

Measurements of sclerotised structures were taken at a magnification of 1000 $\times$ (objective $\times$ 100 immersion, ocular $\times$ 10) using an Olympus BX51 microscope with incorporated phase contrast and the software Digital Image Analysis v4. In total, 29 parameters of the hardparts of the haptor and copulatory organs were measured for morphometric characterisation and a detailed redescription (see Fig. 1). Terminology was based on Justine & Henry (2010). To investigate the level of morphological differentiation (haptor morphology), raw measurements were analysed by multivariate statistical techniques in R (R Core Team, 2013). Principal component analyses (PCAs) were conducted with scaled variables on 17 morphological characters of the haptor using the package adegenet (Jombart, 2008). Results of the PCA were visualised with the packages ggplot2 (Wickham, 2009) and factoextra (Kassambara and Mundt, 2017). To visualise the variance in the total size of the ventral anchor, a density plot using uncorrected measurements was drawn using ggplot2 and factoextra. A Kruskal-Wallis test of multiple comparison with Bonferroni's post-hoc correction via Dunn's test, implemented in the package FSA (Ogle et al., 2019), respectively, was conducted to test the relation of the host species and the catch locality to copulatory organ measurements, respectively. The assumption of normality was tested by Shapiro-Wilk's W tests implemented in stats. The assumption of homogeneous variance within sample groups was tested by Levene's test in the R package lawstat (Gastwirth et al., 2017).

Molecular characterisation

Morphological characterisation was combined with genetic characterisation using tissue samples of the central part of some of the parasite individuals collected from fresh fish specimens from Lake Tanganyika and Lake Albert, as described above. No fresh material was available from other locations. To genetically verify parasite species delineation, we used three different nuclear sequence fragments, from the small and large ribosomal subunit gene (18 and 28 rDNA) and the first internal transcribed spacer region (ITS-1). To assess

180 intraspecific genetic diversity, part of the mitochondrial COI gene was used. Whole genomic
181 DNA was extracted using the Qiagen Blood and Tissue Isolation Kit following the
182 manufacturer's instructions with some modifications (samples in ATL buffer (180 µl) with
183 protein kinase (20 µl) were kept in 1.5 ml Eppendorf tubes overnight at room temperature).
184 The DNA extract was concentrated to a volume of 80 µl in 1.5 ml Eppendorf tubes using a
185 vacuum centrifuge and stored at a temperature of -20 °C. Partial 18S rDNA and ITS-1 were
186 amplified using the S1 (5'-ATTCCGATAACGAACGAGACT-3') (Sinnappah et al., 2001)
187 and Lig5.R (5'-GATACTCGAGCCGAGTGATCC-3') (Blasco-Costa et al., 2012) primers.
188 Each reaction mix contained 1.5 unit of *Taq* polymerase, 1X buffer containing 0.1 mg/ml
189 bovine serum albumin (BSA), 1.5 mM MgCl₂, 200 mM dNTPs, 0.8 mM of each primer and 3
190 µl of isolated DNA (concentration was not measured) in a total reaction volume of 30 µl
191 under the following conditions: 2 min at 95 °C, 39 cycles of 1 min at 95 °C, 1 min at 55 °C
192 and 1 min and 30 s at 72 °C, and finally 10 min at 72 °C. Primers C1 (5'-
193 ACCCGCTGAATTTAAGCAT-3') and D2 (5'-TGGTCCGTGTTTCAAGAC-3') (Hassouna
194 et al., 1984) were used for amplification of the partial 28S rDNA gene. Each PCR reaction
195 contained 1.5 unit of *Taq* polymerase, 1X buffer containing 0.1 mg/ml BSA, 1.5 mM MgCl₂,
196 200 mM dNTPs, 0.5 mM of each primer and 5 µl of isolated DNA (concentration was not
197 measured) in a total reaction volume of 30 µl under the following conditions: 2 min at 94 °C,
198 39 cycles of 20 seconds at 94 °C, 30 seconds at 58 °C and 1 min and 30 s at 72 °C, and finally
199 10 min at 72 °C. Part of the mitochondrial COI gene was amplified using ASmit1 (5'-
200 TTTTTTGGGCATCCTGAGGTTTAT-3') combined with Schisto3 (5'-
201 TAATGCATMGGAAAAAACA-3'), and with ASmit2 (5'-
202 TAAAGAAAGAACATAATGAAAATG-3') in case of nested PCR (Littlewood et al., 1997).
203 For both primer combinations, the amplification reaction contained 24 µl of PCR mix (one
204 unit of *Taq* polymerase, 1X buffer containing 2 mM MgCl₂, 0.1 mg/ml BSA, 0.2 mM dNTPs,

0.8 mM of each primer) with 1 µl of isolated DNA (concentration was not measured) in a total reaction volume of 25 µl and was performed under the following conditions: initial denaturation at 95°C for 5 min and then 40 cycles of 1 min at 94°C, 1 min at 50°C and 1 min at 72°C, and final elongation for 7 min at 72°C. Amplification success was checked by agarose gel electrophoresis and for positive samples, 2.5 µg of PCR product was enzymatically cleaned up using 1 µl of ExoSAP-IT reagent under the following conditions: 15 min at 37 °C and 15 min at 80 °C. After cycle sequencing of purified PCR products using BigDye v3.1, following the manufacturer's recommendations, fragments were cleaned up using the BigDye XTerminator® Purification Kit and visualized on an ABI3130 capillary sequencer. Electropherograms were visually inspected, corrected and sequences were aligned using MUSCLE (Edgar, 2004) under default settings as implemented in MEGA v7 (Kumar et al., 2016), together with selected previously published sequences of representatives of Diplectanidae (see Table S1). The newly obtained haplotype sequences were deposited in NCBI GenBank under the accession numbers MK937579-MK937581 (28S rDNA), MK937574-MK937576 (18S+ITS-1 rDNA) and MK908145- MK908196 (COI mtDNA).

Genetic distances and phylogeny

The consistency of all alignments was checked and corrected under the “automated 1” option in trimAL v1.2, which uses a heuristic search to find the best method for trimming the alignment (Capella-Gutiérrez et al., 2009). As there is a lack of available ITS sequences of diplectanid species, phylogenetic analyses were based on two regions: 18S and 28S rDNA. These two regions were analysed separately because of the lack of species for which both regions are available. Topali v2.5 (Milne et al., 2004) was used to identify the best fitting model of molecular evolution based on the Bayesian information criterion (28S rDNA: GTR + Γ , gamma shape parameter of 0.461; 18S rDNA: K2P + Γ , gamma shape parameter of

0.130). For each gene, pairwise distances were calculated using both the most appropriate evolutionary model and, to compare with previous studies, uncorrected pairwise distances. The number of haplotypes and polymorphic sites, haplotype diversity and nucleotide diversity were calculated using ARLEQUIN v3.5 (Excoffier and Lischer, 2010). Phylogenetic analyses were carried out using maximum likelihood (ML) and Bayesian inference (BI) in RAxML v8 (Stamatakis, 2014) and MrBayes v3.2.0 (Ronquist et al., 2012), respectively. A ML tree was inferred using RAxML's standard tree search algorithm and bootstrap support was calculated using the option with an automated number of replicates to obtain stable support values under the frequency stopping criterion (Stamatakis, 2014). Bayesian inference was based on two independent runs (100,000,000 generations, sampled every 1,000th generation following a burn-in of 10%). Parameter convergence and run stationarity were assessed in Tracer v1.6 (<http://beast.bio.ed.ac.uk>). As Dactylogyridae and Diplectanidae were shown to be sister taxa (Šimková et al., 2003), *Dactylogyrus extensus* (Mueller and Van Cleave, 1932) (sequence from: Šimková, Matějusková, & Cunningham, 2006) together with *Cichlidogyrus attenboroughi* Kmentová, Gelnar & Vanhove, 2016 (sequence from: Kmentová et al. (2018) in the case of 28S rDNA region were selected as outgroup for phylogenetic inference. Phylogenetic trees were edited in FigTree v1.4.2 (<http://tree.bio.ed.ac.uk/software/figtree/>) and Adobe Photoshop CS6. Phylogenetic relationships among COI haplotypes were inferred by means of a Median Joining network (Bandelt et al., 1999) in PopART 1.7104 (Leigh and Bryant, 2015).

Results

A single diplectanid species, morphologically identified as *Diplectanum lacustre* was recorded from three of the four species of *Lates* from Lake Tanganyika (*L. angustifrons*, *L. mariae*, *L. microlepis*) and from *L. niloticus* from Lakes Albert, Kossou, Nasser and Victoria,

from the Taja River in Sierra Leone and from the Bahr-Sara, mouth of the Mandoul River in Tchad. In total, 473 parasite specimens were collected (for more details see Table 1 and Fig. 2). Based on morphological characterisation and phylogenetic reconstruction (see Figs. 7&8), a new genus *Dolicirroplectanum* gen. nov. is described with *Dolicirroplectanum lacustre* comb. nov. as the type species. The internal anatomy is characterised, including the sclerotised vagina, prostatic reservoir and seminal vesicle, which were absent in the original description of *D. lacustre* comb. nov. Measurements of the parasite's internal organs and sclerotised haptoral and copulatory structures are presented in Table 2.

Taxonomy and species redescription

***Dolicirroplectanum* gen. nov. Kmentová, Gelnar & Vanhove (Fig. 3 - 5)**

Family: Diplectanidae Monticelli, 1903

Genus: *Dolicirroplectanum* gen. nov.

Type species: *Dolicirroplectanum lacustre* (Thurston & Paperna, 1969)

Type host: *Lates niloticus* L. (Latidae)

Type locality: Lake Volta, Ghana; Lake Albert, Uganda

Site: Gills

Additional hosts: *L. angustifrons*, *L. mariae*, *L. microlepis*

Other species: *Dolicirroplectanum penangi* comb. nov. for *Diplectanum penangi* Liang & Leong, 1991 (original designation)

Material examined: type material: MRAC MT. 35572, vouchers: MNHN HEL744-47 (4 specimens), USNPC 180-A 3-7; MRAC. MT. 38206-10, 38913-39058 (243 specimens), MZH 10067-71 (6 specimens), SAMC-A089971-72 (6 specimens), NHMUK 2018.4.13.4-13.7 (8 specimens)

278 **Zoobank registration:** To comply with the regulations set out in article 8.5 of the amended
279 2012 version of the International Code of Zoological Nomenclature (ICZN) (International
280 Commission on Zoological Nomenclature, 2012), details of the genus have been submitted to
281 ZooBank. The Life Science Identifier (LSID) of the article is
282 urn:lsid:zoobank.org:pub:209675D6-2EBE-4E37-84CB-DA59994F7B2. The LSID for the
283 new genus *Dolicirroplectanum* is urn:lsid:zoobank.org:act:89BFF3C5-271B-4482-98E0-
284 AC667AA6611D.

285 **Etymology:** The genus name derives from Latin and refers to the barrel shape of the male
286 copulatory organ, noticeably wider than in other diplectanid genera.

287 **Diagnosis:** Tegument smooth. Genital pore opening posterior to male copulatory organ (MCO).
288 Genital atrium sclerotised. MCO wide, robust, composed of two nested tubes. Prostatic
289 reservoir simple. Seminal vesicle sinistral. Accessory copulatory organ absent. Squamodiscs
290 ventral, dorsal; rows of bone-shaped rodlets with open rings. Superficial root of ventral anchor
291 reduced. Parasites of perciform fishes (*Lates* spp.). Vagina sclerotised or muscular.

292 **Description:**

293 Multiple pairs of head organs, two pairs of eye-spots. No tegument scales were observed.
294 *Dolicirroplectanum* gen. nov. is characterised by two pairs of dorsal and ventral anchors with
295 a regularly curved shaft point, a large and wide ventral bar and two dorsal bars. Dorsal anchors
296 smaller than ventral ones and without developed outer root. 14 marginal hooklets of similar size
297 and relatively small compared to other haptoral structures. Two squamodiscs, ventral and
298 dorsal, formed by concentric open rows of bone-shaped rodlets of similar width in all rows.
299 Intestinal bifurcation follows pharynx, oesophagus absent. Caeca simple, terminate blindly.
300 Testis spherical, intercaecal. Vas deferens emerges from anterior part of testis, enlarges into
301 seminal vesicle. Seminal vesicle single in the middle region of body, transforms into elongated

duct connected with sclerotized part of copulatory organ. Prostatic reservoir simple. Slightly sclerotized MCO composed of two straight tubes, one inside the other, almost as wide as long. Ovary intercaecal, pre-testicular, encircles right caecum. Oviduct passes medially to oötype, surrounded by Mehlis' gland, oötype short, enters into uterus. Uterus sinistral. Vaginal atrium sclerotised or muscular.

Discussion: Species of *Dolicirroplectanum* gen. nov. can be distinguished by the combination of: 1) presence of a robust barrel-shaped MCO formed by two narrow nested tubes, almost as wide as long, 2) absence of an accessory piece, 3) squamodiscs composed of bone-shaped rodlets forming open rings, 4) superficial roots of ventral anchor reduced, 5) a simple prostatic reservoir not separated into zones, 6) seminal vesicle as an expansion of vas deferens, 7) ovary intercaecal, pre-testicular, encircles right caecum and 8) a lack of tegumental scales. The status of *Dolicirroplectanum* gen. nov. is supported by its placement outside of the clade including *Diplectanum aequans* (Wagener, 1857), the type species of *Diplectanum* (Figs. 7&8). Particularly, *Dolicirroplectanum* gen. nov. differs from other diplectanids including *D. aequans* by the short but wide sclerotised part of the MCO. In contrast to *D. aequans*, a simple prostatic reservoir is present. Conversely, a prostatic reservoir separated into three zones is one of the specific characters for *Diplectanum* sensu stricto mentioned in Domingues & Boeger (2009). *Diplectanum penangi* has all the diagnostic features attributed to *Dolicirroplectanum* gen. nov. The position within the genus was supported by its position in a phylogenetic reconstruction, clustering with *Dolicirroplectanum lacustre* comb. nov. (Figs. 6&7). The holotype of *D. penangi* comb. nov. could not be verified as the specimen was not provided by Lee Kong Chian Natural History Museum in Singapore and as the digital pictures we received were taken at insufficient magnification/resolution. Therefore, voucher material deposited in the National Museum of Natural History in Washington and the National Museum of Natural History in Paris was checked instead, and the two nested copulatory tubes and simple prostatic reservoir

327 were found to be present in *D. penangi* comb. nov. together with other characteristics mentioned
328 in its original description (Fig. 5).

329 **Redescription**

330 **Family:** Diplectanidae Monticelli, 1903

331 **Genus:** *Dolicirroplectanum*

332 *Dolicirroplectanum lacustre* comb. nov. (Thurston & Paperna, 1969)

333 **Synonyms:** *Diplectanum lacustre*

334 **Zoobank registration:** To comply with the regulations set out in article 8.5 of the amended
335 2012 version of the International Code of Zoological Nomenclature (ICZN) (International
336 Commission on Zoological Nomenclature, 2012), details of the species have been submitted to
337 ZooBank. The Life Science Identifier (LSID) of the article is
338 urn:lsid:zoobank.org:pub:209675D6-2EBE-4E37-84CB-DA59994F7B2. The LSID for the
339 new name *Dolicirroplectanum lacustre* is urn:lsid:zoobank.org:act:423241C3-777D-4F86-
340 A70F-02B4D74F9E66.

341 **Figures:** 3, 44

342 **Material examined:** holotype: MRAC MT. 35572, paratype: MRAC MT. 35573

343 **Vouchers:** MRAC. MT. 38206-10, 38913-39058 (243 specimens), MZH 10067-71 (6
344 specimens), MNHN HEL744-47 (4 specimens), SAMC-A089971-72 (6 specimens), NHMUK
345 2018.4.13.4-13.7 (8 specimens)

346 **Type host:** *Lates niloticus* L. (Latidae)

347 **Type locality:** Lake Volta, Ghana; Lake Albert, Uganda

348 **Site:** Gills

349 **Additional hosts:** *L. angustifrons*, *L. mariae*, *L. microlepis*

Additional localities: Bahr-Sara, Tchad (08°56'N-17°58'E); Kisumu, Lake Victoria (00°06'S-34°45'E), Lake Kossou, Egypt (07°10'N-05°20'E), Lake Nasser, Egypt (24°05'N-33°00'E), Luxor market, Egypt (25°42'N 32°38'E), Njala, riv. Taja, Sierra Leone (08°06'N-12°04'E), Lake Albert – Nyawiega (01°28'N-30°56'E); Nzunzu (1°19'N, 30°72'E); Lake Tanganyika – Crock Island (8°42'S-31°07'E), Katukula (8°35'S-31°10'E), Mpulungu (8°46'S-31°07'E); Rumonge (3°97'S-29°43'E); Sumbu Bay (8°31'S-30°29'E); Bujumbura (3°23'S-29°22'E); Ilagala (5°12'S-29°50'E); Kilomoni (4°20'S, 29°09'E); Mulembwe (6°07'S, 29°16'E); Nyanza (4°20'S-29°35'E); Edith Bay (6°30'S-29°55'E); Uvira (3°22' S 29°08'E)

Infection parameters: 4 of 8 *Lates angustifrons* infected with 1 – 15 specimens, 15 of 23 *L. mariae* infected with 1 – 18 specimens, 21 of 31 *L. microlepis* infected with 1 – 53 specimens. 1 of 1 *L. niloticus* from Bahr-Sara infected with 2 specimens, 2 of 3 *L. niloticus* from Kisumu (Lake Victoria) infected with 1-7 specimens, 4 of 5 *L. niloticus* from Lake Kossou infected with 5-9 specimens, 1 of 5 *L. niloticus* from Lake Nasser infected with 1 specimen, 1 of 3 *L. niloticus* from Nyawiega (Lake Albert) infected with 2 specimens, 5 of 11 *L. niloticus* from Nzuzu (Lake Albert) infected with 2-10 specimens, 1 of 1 *L. niloticus* from Nzuzu (Lake Albert) infected with 2 specimens.

Diagnosis: *Dolicirroplectanum lacustre* comb. nov. is a monogenean infecting gills of freshwater African latid species distinguished from its congener by the width of the outer root of the ventral anchor. The copulatory tube is oriented anteriorly.

Description: Tegument thin, smooth. Three pairs of head organs, two pairs of eye-spots, the posterior ones larger and closer together. Two squamodiscs, ventral squamodisc larger than dorsal, both consist of 9-12 concentric rows of bone-shaped rodlets, the two distal rows of which are composed of only rudimentary rodlets. Two pairs of anchors, rudiment of inner root and wide base of outer root in dorsal anchor. Marginal hooklets (14) of similar size. Ventral bar tapering towards extremities with terminal auricles. Dorsal bar broadening towards the centre

of the haptor area. Testis post-ovarial, thin vas deferens along the dextral intestinal caecum. Single seminal vesicle in the middle of the body. Simple prostatic reservoir. MCO robust and formed by two nested tubes. Copulatory tube oriented anteriorly. Ovary looping around the left intestinal caecum towards the oviduct, surrounded by Mehlis' glands located near oötype. Uterus simple tube towards vagina. Vagina is formed by a complex of sclerotized structures consisting of an elongated primary canal followed by a secondary tube opening into an anterior duct; duct continues into distal sclerotized part ending in blade-shaped structure. Orientation of sclerotized vagina with blade-shaped end always anterior. Sclerotised vagina can be absent. Vitellaria dense, located around outer wall of intestinal caeca.

Discussion: *Dolicirroplectanum lacustre* comb. nov. resembles its congener *Dolicirroplectanum penangi* comb. nov. infecting *Lates calcarifer* in Asia. The type species of the genus can be easily distinguished from *D. penangi* comb. nov. by the comparative morphology of the anchors, especially the thinner outer root in *D. penangi* comb. nov. (see Fig. 5). Contrary to *D. lacustre* comb. nov., a sclerotised vagina was not observed in *D. penangi* comb. nov. (Liang and Leong, 1991). Our findings are based only on a combination of the original description of *D. penangi* comb. nov. and voucher material deposited in the National Museum of Natural History in Washington and the National Museum of Natural History in Paris as the Lee Kong Chian Natural History Museum in Singapore refused to provide the holotype material.

Morphometric variation

Morphological variation was visualized based on a PCA performed on 17 standardised haptoral morphometric parameters from 148 individuals. The first PC explained 48.4 % of the variation in the data, the second one 10.1 %. Results show a high level of variability in specimens of *D. lacustre* comb. nov. infecting *L. niloticus* and a continuous size gradient not

related to the locality of origin with an intermediate position of specimens from Lake Kossou and Lake Victoria along the first axis. Moreover, individuals collected in the Taja River seem to be separated from the others. Interestingly, two morphotypes were retrieved from different fish specimens in Lake Albert (Lake Albert1 and Lake Albert2). Therefore, the morphology of *D. lacustre* comb. nov. does not seem to be influenced by neither geographical nor host species origin (Fig. 8A). Moreover, two specimens from Lake Albert (belonging to Lake Albert2) were, based on the haptoral sclerotised structures and MCO, more similar to those collected outside the lake (see Table 2). The position of specimens in the scatterplot was mainly influenced by the size of dorsal anchors, maximum width of the dorsal bar and length of both squamodiscs. However, almost all parameters were correlated with the first axis. Other PCs did not show a clearer separation. The length of the dorsal anchor was shown to be related to the combination of host species and geographic origin, in a gradient, with two morphotypes recognised in Lake Albert (Lake Albert1 and Lake Albert2), as visualised in a density plot (Fig. 8B).

MCO parameters from 91 individuals of *D. lacustre* comb. nov. were compared. Significantly wider and longer copulatory organs were observed in specimens collected from *L. niloticus* (n=31), than in those collected from *L. microlepis* (n=38) (Bonferroni's post-hoc correction, MCO length $Z_{2,87}=-6.48$, $P<0.001$, MCO width $Z_{2,89}=-6.74$, $P<0.001$) and *L. mariae* (n=22) (Bonferroni's post-hoc correction, MCO length $Z_{2,87}=-4.98$, $P<0.001$, MCO width $Z_{2,89}=-4.25$, $P<0.001$). The influence of geographical origin was tested only for samples from these three host species from Lake Albert and Lake Tanganyika as there was an insufficient number of high-quality specimens from other localities and *L. angustifrons*, respectively. In both parameters of the MCO, a significantly larger size was observed in specimens from Lake Albert, morphotype Lake Albert1 (Bonferroni's post-hoc correction, MCO length – $Z_{1,89} = 6.61$, $P < 0.001$, MCO width – $Z_{1,87} = 6.41$, $P < 0.001$).

Apart from these size differences, the variable presence of a sclerotised vagina was documented (see Table 2), also including data from two previous records of the species (Ergens, 1981; Thurston and Paperna, 1969).

Genetic characterisation and phylogeography

Uncorrected p-distances between *D. lacustre* comb. nov. collected from Lake Tanganyika and Lake Albert, respectively, varied among the amplified regions from 0.5% in 18S rDNA (441 base pairs (bp)), 1.1% in 28S rDNA (810 bp) to 9% in ITS-1 rDNA (478 bp) and 9.0 – 10.2% in COI mtDNA (412 bp). In previous studies, the ability to align ITS-1 sequences was used as a criterion for diplectanid species delineation (Poisot et al., 2011; Wu et al., 2007). No intralacustrine variability in rDNA regions was detected. Sequences of the ITS-1 region of all populations of *D. lacustre* comb. nov. in our study were alignable and included 19 indels. For comparison with the threshold of 14.5% difference in the COI region to distinguish intra- and interspecific diversity proposed for diplectanids by Vanhove et al. (2013), genetic distances were also calculated using the K2P model (Kimura, 1980), under which they amounted to 9.6 – 10.7%. Intralacustrine variation in COI was higher in Lake Albert than in Lake Tanganyika (Table 3). The haplotype network showed two distinct haplogroups, corresponding to the two lakes (Fig. 9). Identical COI haplotypes were shared among individuals of *D. lacustre* comb. nov. collected from *L. mariae* originating from the central subbasin (Mulembwe) and *L. microlepis* collected from the northern and southern subbasins of Lake Tanganyika (Uvira and Mpulungu).

Phylogeny

Phylogenetic inference at the family level (Diplectanidae) was based on two separate alignments of the 28S and 18S nuclear rDNA with 33 and 15 taxa, respectively (Table S1). The alignment of 28S rDNA and 18S rDNA totalled 803 and 482 bp, respectively.

Phylogenetic analyses of 28S rDNA placed the haplotypes of *Dolicirroplectanum lacustre* comb. nov. in a monophyletic clade sister to *Dolicirroplectanum penangi* comb. nov. collected from *Lates calcarifer* in Asia (Fig. 6). The tree obtained from the 18S rDNA fragment placed *D. lacustre* comb. nov. in a poorly resolved clade with species of *Pseudorhabdosynochus* Yamaguti, 1958 and *Echinoplectanum* Justine and Euzet (2006) (Fig. 7). ML and BI produced the same topologies. In both phylogenetic trees, the previous notion of *Diplectanum* appeared polyphyletic with the type species, *Diplectanum aequans*, placed outside the clade including species of *Dolicirroplectanum* gen. nov., hence supporting the erection of a new genus.

Discussion

The main aim of this study was to examine the level of diversification in diplectanid parasites infecting latid hosts in two of the African Great Lakes, Lakes Albert and Tanganyika. Moreover, museum specimens from throughout the host's range were added to provide a broader geographical range for morphological comparison. Morphological and molecular characterisation identified a single species in both lakes, reassigned to *Dolicirroplectanum* gen. nov. Despite the persistent geographic separation between Lakes Albert and Tanganyika for 9 MYA (Cohen et al., 1993), and the speciation of the hosts, their respective populations of *D. lacustre* comb. nov. have not reached the level of morphological and genetic differentiation typically associated with distinct species. Hence, we conclude that this is an example of a lineage that failed to speciate.

Diplectanid species infecting latid fishes in Africa – molecular and morphological perspectives

The monophyly of *Diplectanum* was already rejected in previous studies (Chotnipat et al., 2015; Villar-Torres et al., 2019) with *Dolicirroplectanum lacustre* comb. nov. being classified

outside of *Diplectanum* sensu stricto (Chotnipat et al., 2015; Domingues and Boeger, 2008).

The phylogenetic reconstructions based on ribosomal regions place *D. lacustre* comb. nov. in a separate lineage together with *D. penangi* comb. nov. infecting an Asian latid species, *L. calcarifer*, but outside the clade that includes *D. aequans*, the type species of *Diplectanum*.

This, combined with a detailed morphological characterisation, leads us to propose the new genus *Dolicirroplectanum* gen. nov., now including *D. lacustre* comb. nov. and *D. penangi* comb. nov. Overall, the phylogenetic position of other diplectanid genera corresponds with the study by Villar-Torres et al. (2019). The genetic distance between *D. lacustre* comb. nov. from Lake Tanganyika and Lake Albert, and *D. penangi* comb. nov., is 7.9% based on the 28S rDNA fragment. This is comparable to the situation in *Laticola latesi* (Tripathi, 1957) and *L. paralatesi* (Nagibina, 1976), which infect *L. calcarifer* in Hainan province, China (Tingbao et al., 2006). However, these diplectanid species occur sympatrically, infecting a single host species, whereas there is no contact between *L. calcarifer* and the species of *Lates* from Lakes Albert and Tanganyika.

Interestingly, copulatory tube width and length differ between most of the parasite individuals collected from *L. niloticus* from Lake Albert and three of Lake Tanganyika's species, *L. angustifrons*, *L. microlepis* and *L. mariae*, respectively. Differences in the MCO may be the basis for species delineation in diplectanids (e.g. in *Echinoplectanum*: Sigura & Justine, 2008). However, the mean values of individuals of *L. niloticus* from other localities (Lake Kossou, Lake Victoria) do not differ from Lake Tanganyika's specimens. Moreover, two specimens from Lake Albert (belonging to Lake Albert2) are, based on the haptoral sclerotised structures and the MCO, more similar to those collected outside this lake (see Table 2). A complex pattern of morphological variation emerging from the other populations of *D. lacustre* comb. nov., with a lot of overlapping features between host species and localities (Fig. 8 and Table 2), does not suggest the existence of different species. The

intermediate position of some specimens, particularly from Lake Albert (referred to as Lake Albert2), prevents a clear correlation between parasite morphotype, host species identity and/or geographic origin. Future genetic characterisation of such morphotypes is needed to address diversification of *D. lacustre* comb. nov. in detail. Internal anatomy is documented only in fresh specimens from Lake Albert and Lake Tanganyika, with high levels of intralacustrine variation and without structural differences in organisation between these lakes.

Moreover, the impossibility to align ITS-1 rDNA sequences is generally considered as an indicator for diplectanid species delineation (Wu et al., 2005a; Poisot et al., 2011). Since haplotypes from the two populations of *D. lacustre* comb. nov. in our study are alignable, the hypothesis of a single species is supported. Also, as the model-corrected genetic distance of 10.7% over the COI fragment does not reach the “best-compromise threshold” (Meier et al., 2006) for barcoding of 14.5% proposed by Vanhove et al. (2013) for diplectanids infecting Indo-Pacific groupers, we consider all specimens in our study as conspecific and belonging to *D. lacustre* comb. nov. Its records therefore increased from four to ten areas (see Table 1 and taxonomic part of the result section).

Based on differences in the size of haptoral structures and the split ends of the internal tube of the MCO (see Table 2 and Fig. 4), specimens from Taja River might be considered as belonging to a different species. This could be explained by the long separation between the Upper Guinean and Nil ichthyofaunal provinces (Roberts, 1975). However, more samples and molecular data from Taja River are needed to confirm the identity of the species of *Dolicirroplectanum* infecting *L. niloticus* in the Upper Guinean province.

Host range of the diplectanid monogeneans infecting lates perches in Lake Tanganyika

520 Based on our results, the host species list of *D. lacustre* comb. nov. was extended with three
521 of the four latid species from Lake Tanganyika. No monogeneans were found on *L. stappersii*.
522 A potential reason could be its different life style compared to other latid species in the lake.
523 In contrast to its congeners, *L. stappersii* is truly pelagic throughout its life, usually not
524 moving into inshore waters (Mannini et al., 1999; Mulimbwa and Mannini, 1993). Short-lived
525 and slow swimming monogenean larvae (oncomiracidia) are assumed to infect fish hosts in
526 littoral habitats, typically synchronised with their hosts' period of reproduction (Whittington
527 et al., 1999). Therefore, there is less chance for parasite infection in *L. stappersii* compared to
528 other latid species (see Rohde, 1980; Rohde et al., 1995). Moreover, there is no sign of
529 species diversification in Lake Tanganyika as we found haplotypes shared between latid hosts
530 and between subbasins.

531 The African Great Lakes are highly biodiverse areas with a remarkable species richness and a
532 high level of endemism (Salzburger et al., 2014). Parasite diversification linked with host
533 speciation was recently discovered in monogeneans belonging to *Cichlidogyrus* Paperna,
534 1960 infecting littoral cichlids of Lake Tanganyika (Vanhove et al., 2015). Interestingly, also
535 in Lake Tanganyika's pelagic zone, the same pattern as in *D. lacustre* comb. nov., apparently
536 without host preference or host-related speciation processes, was observed in *Cichlidogyrus*
537 *casuarinus*, a parasite of bathybatine cichlids (Kmentová et al., 2016). However, the
538 haplotype and nucleotide diversity in the COI region are remarkably lower in *D. lacustre*
539 comb. nov. (0.517 and 0.001 compared to 0.987 and 0.0205, respectively). Host species
540 hybridisation might explain the more generalist life style of certain monogeneans due to an
541 influence of host genetics on the susceptibility to infection, host specificity, and parasite
542 speciation (Šimková et al., 2013; Vanhove et al., 2011). However, there are no reports of
543 hybridisation among latid species (Otero, 2004). Moreover, a lack of host-related speciation in
544 diplectanids was observed in *Pseudorhabdosynochus cyanopodus* Sigura & Justine, 2008

infecting two deep-sea grouper species in New Caledonia (Schoelinck et al., 2012) with a maximum intraspecific distance of 1.2% in the COI region compared to 0.7% in *D. lacustre* comb. nov. in Lake Tanganyika. Our results therefore correspond with previous studies in marine and freshwater habitats where decreased host specificity in pelagic ecosystems was proposed to increase the chance of finding a host if host species exhibit low population densities (Kmentová et al., 2016; Rohde, 1980; Schoelinck et al., 2012).

Despite the generally high degree of endemism of macrofauna in Lake Tanganyika (Coulter, 1991; Salzburger et al., 2014; Snoeks, 2000), this might not be reflected in all microinvertebrate taxa. *Dolicirroplectanum lacustre* comb. nov. and some other monogenean species are found to naturally occur both within and outside Lake Tanganyika. *Gyrodactylus sturmbaueri* Vanhove, Snoeks, Volckaert & Huyse, 2011 was described from *Simochromis diagramma* (Günther, 1894), but also parasitizes on *Pseudocrenilabrus philander* (Weber, 1897) in Lake Kariba and the River Nwanedi (Zahradníčková et al., 2016). *Cichlidogyrus mbirizei* Muterezi Bukinga, Vanhove, Van Steenberge & Pariselle 2012, *C. halli* (Price and Kirk 1967) and *Scutogyrus gravivaginus* (Paperna and Thurston 1969) are known from an endemic tilapia, *Oreochromis tanganyicae* (Günther, 1894), but were also reported from other species of *Oreochromis* Günther, 1889 and other cichlids in Africa (Douëllou, 1993; Pariselle and Euzet, 2009) as well as cage-cultured tilapia species in Asia (Lim et al., 2016; Mohd Agos et al., 2016). This highlights the ability of some monogenean species to survive in a wide range of environments and host species (Huyse et al., 2006).

Rate of molecular evolution of *Dolicirroplectanum lacustre* comb. nov. and its implications

For want of paleontological data, substitution rates in parasitic flatworms are typically estimated using host fossils or calibrated with paleogeographical events and assuming that

parasite speciation follows that of the hosts (Meinilä et al., 2004). With a mean distance of 10% between the Lake Tanganyika and Lake Albert populations, the substitution rate in our 412 bp COI mtDNA region, using the end of rifting in the eastern African Rift Valley 9 MYA (Cohen et al., 1993) for the age of the most recent common ancestor of their hosts, is estimated at 0.5%/MY.

Often, molecular evolution of parasites is considered and was proven to be faster than within the homologous loci of their hosts (Hafner et al., 1994; Huyse et al., 2005). Surprisingly, *D. lacustre* comb. nov. appears to have a slower rate of molecular evolution in its mitochondrial DNA than most fish taxa (1-4% in cytochrome *b*) (Bermingham et al., 1997; He and Chen, 2007; Muss et al., 2001). However, preliminary molecular data of African latids show even less difference over the COI region (4.5-6% uncorrected p-distance between Lake Tanganyika and Lake Albert, own unpublished data) than their monogenean parasite. Therefore, the widespread hypothesis of a faster evolutionary rate of parasites compared to hosts, based on their shorter generation time (Cable and Harris, 2002; Thomas et al., 2010) does hold in the case of *D. lacustre* comb. nov. The link between the rate of morphological and molecular evolution varies among different taxa (Omland, 1997) with studies showing either rate decoupling (Poisot et al., 2011) or rate correlation. However, it seems that in the case of *D. lacustre* comb. nov., there is a correlation between a slow rate of molecular evolution in the COI gene, which is a structural coding marker known to be under balancing selection (Wu et al., 1997), and failure to diverge seen in the lack of speciation.

There are three possible scenarios explaining the observed situation in *D. lacustre* comb. nov. First, the rate of evolution of latids and their parasites is slower in comparison to other fish (Bermingham et al., 1997; Muss et al., 2001) and other monogenean taxa. Indeed, the mutation rate of *D. lacustre* comb. nov. seems to be much lower than the 13.7 – 20.3% per million years estimated for *Gyrodactylus* von Nordmann, 1832 by Meinilä et al. (2004). This

can be explained by their different life history, as diplectanids lack the asexual reproduction that has also led to a high species richness promoted by host switches and peripatric speciation processes in *Gyrodactylus* (Boeger et al., 2003). The failure to diverge of *D. lacustre* comb. nov. in Lake Tanganyika then corresponds with the hypothesis suggesting that a lower rate of molecular evolution resulted in low diversification.

Secondly, the invasion of the studied latid lineage could be more recent than the lakes' formation, like in the case of the cichlid genera *Tylochromis* Steindachner, 1895 and *Oreochromis* (Klett and Meyer, 2002; Koch et al., 2007). This could explain the low level of genetic intralacustrine variation of *D. lacustre* comb. nov. reported in Lake Tanganyika (0.7% of uncorrected p-distance in COI) which contrasts with greater genetic variation seen in the host species (2-3% of uncorrected p-distance in COI, unpublished data). However, low intralacustrine genetic diversity in *D. lacustre* comb. nov. could also be caused by bottleneck events that have reduced the genetic variation present in the system.

A third possible scenario involves a latid origin in the proto-Tanganyikan region with more recent admixture of populations via lacustrine and riverine connections resulting in the polyphyly of latid species in Lake Tanganyika (suggested by Otero (2004) based on morphological data), as was documented for haplochromine cichlids in the lake (Meyer et al., 2015; Salzburger et al., 2005). Therefore, molecular and morphological similarity of *D. lacustre* comb. nov. in nowadays geographically isolated areas could be the result of recent and maybe multiple episodes of gene flow. In any case, phylogenetic reconstruction combined with the latids' fossil record is needed to discern between the above-mentioned hypotheses.

Conclusions

Diplectanid parasites occur primarily in marine environments. Discussion has arisen about the incongruence between their morphological species delineation and the level of molecular

differentiation. In our study, we focused on a unique allopatric situation of diplectanid parasites infecting latid species inhabiting freshwater lakes to study this incongruence. Based on morphological examination, a single diplectanid species was recorded from three of the four species of *Lates* from Lake Tanganyika (*L. angustifrons*, *L. mariae*, *L. microlepis*) and from *L. niloticus* from Lakes Albert, Kossou, Nasser and Victoria, from the Taja River in Sierra Leone and from Bahr-Sara in Tchad. Thus, similar to another monogenean species infecting pelagic host species of the cichlid tribe Bathybatini in Lake Tanganyika, this parasite on African *Lates* apparently failed to diverge. Results of phylogenetic reconstruction combined with detailed morphological characterisation led us to propose *Dolicirroplectanum* gen. nov. with *D. lacustre* comb. nov. as type species. Despite a persistent geographic barrier between Lake Albert and Lake Tanganyika and speciation in the hosts, their respective populations of *D. lacustre* comb. nov. infecting these lakes' latid fishes have not reached species-level distinction. We suggest a link between the lack of morphological differentiation between the parasite populations in both lakes, and the low rate of molecular evolution of the mitochondrial COI gene, estimated at 0.5%/MY (assuming *Lates* from Lakes Albert and Tanganyika diverged 9 MYA). As alternatives, scenarios proposing either more recent invasion of the latid lineage into Lake Tanganyika or recent gene flow among the latid lineages in Lakes Albert and Tanganyika could explain the apparently slow rate of the hosts' molecular evolution and lack of parasite differentiation. Therefore, detailed studies of host phylogeography, dated using the fossil record, are needed to discern between these scenarios. Although species-level differentiation can be expected in the future under persisting separation between the lakes, the question about the existence of genetically intermediate populations of *D. lacustre* comb nov. remains.

Acknowledgments

The authors would like to thank: Ludmila Raisingerová, Chahrazed Rahmouni, Mulongaibalu Mbalassa, Vercus Lumami Kapepula, Joseph Mbirize Ndaloziwba, Jos Snoeks and Christian Sturmbauer for help with obtaining or identifying samples. Fieldwork was carried out with the approval of the competent local authorities under mission statement 031/MINRST/CRH-U/2016 and an export permit from the CRH-Uvira and from the National Fish and Fisheries Research Institute (Nafirri, Jinja, Uganda). We also thank Tine Huyse, Nathan Vranken, Emmanuel Vreven, Miguël Parrent (RMCA), Filip Volckaert (KU Leuven) and people in the parasitological group, Masaryk University, Brno and Research Centre of Hydrobiology, Uvira for their hospitality and the RMCA for providing access to the collections.

References

- Bandelt, H.J., Forster, P., Rohl, A., 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.* 16, 37–48.
<https://doi.org/10.1093/oxfordjournals.molbev.a026036>
- Bermingham, E., McCafferty, S., Martin, A., 1997. Fish biogeography and molecular clocks: perspectives from the Panamanian Isthmus., in: Kocher, T., Stepien, C. (Eds.), *Molecular Systematics of Fishes*. Academic Press, pp. 113–128.
- Blasco-Costa, I., Míguez-Lozano, R., Sarabeev, V., Balbuena, J.A., 2012. Molecular phylogeny of species of *Ligophorus* (Monogenea: Dactylogyridae) and their affinities within the Dactylogyridae. *Parasitol. Int.* 61, 619–627.
<https://doi.org/10.1016/j.parint.2012.06.004>
- Boeger, W.A., Kritsky, D.C., Pie, M.R., 2003. Context of diversification of the viviparous Gyrodactylidae (Platyhelminthes, Monogeneoidea). *Zool. Scr.* 32, 437–448.
<https://doi.org/10.1046/j.1463-6409.2003.00130.x>
- Brooks, D.R., 1979. Testing the context and extent of host-parasite coevolution. *Syst. Biol.* 28, 299–307. <https://doi.org/10.1093/sysbio/28.3.299>
- Cable, J., Harris, P.D., 2002. Gyrodactylid developmental biology: historical review, current status and future trends. *Int. J. Parasitol.* 32, 255–80.
[https://doi.org/https://doi.org/10.1016/S0020-7519\(01\)00330-7](https://doi.org/https://doi.org/10.1016/S0020-7519(01)00330-7)
- Capella-Gutiérrez, S., Silla-Martínez, J.M., Gabaldón, T., 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinforma. Appl. Note* 25, 1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>
- Chotnipat, S., Miller, T.L., Knuckey, R.M., Hutson, K.S., 2015. Molecular and morphological evidence for the widespread distribution of *Laticola paralatesi* infecting wild and farmed *Lates calcarifer* in Australia. *Dis. Aquat. Organ.* 113, 195–205.
<https://doi.org/10.3354/dao02848>

- 678 Cohen, A.S., Soreghan, M.J., Scholz, C.A., 1993. Estimating the age of formation of lakes:
679 An example from Lake Tanganyika, East African Rift system. *Geology* 21, 511–514.
680 [https://doi.org/10.1130/0091-7613\(1993\)021<0511:ETAOFO>2.3.CO;2](https://doi.org/10.1130/0091-7613(1993)021<0511:ETAOFO>2.3.CO;2)
- 681 Coulter, G.W., 1991. Composition of the flora and fauna, in: Coulter, G.W. (Ed.), *Lake*
682 *Tanganyika and Its Life*. Natural History Museum Publications & Oxford University
683 Press, London Oxford & New York, pp. 200–274.
- 684 Danley, P.D., Husemann, M., Ding, B., Dipietro, L.M., Beverly, E.J., Peppe, D.J., 2012. The
685 impact of the geologic history and paleoclimate on the diversification of East African
686 cichlids. *Int. J. Evol. Biol.* 574851. <https://doi.org/10.1155/2012/574851>
- 687 DeSalle, R., Egan, M.G., Siddall, M., 2005. The unholy trinity: taxonomy, species
688 delimitation and DNA barcoding. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 360, 1905–
689 1916. <https://doi.org/10.1098/rstb.2005.1722>
- 690 Desdevises, Y., Morand, S., Oliver, G., 2001. Linking specialisation to diversification in the
691 Diplectanidae Bychowsky 1957 (Monogenea, Platyhelminthes). *Parasitol. Res.* 87, 223–
692 230. <https://doi.org/10.1007/s004360000325>
- 693 Domingues, M. V., Boeger, W.A., 2008. Phylogeny and revision of Diplectanidae Monticelli,
694 1903 (Platyhelminthes: Monogeneoidea). *Zootaxa* 1903, 1–40.
- 695 Douëllou, L., 1993. Monogeneans of the genus *Cichlidogyrus* Paperna, 1960 (Dactylogyridae:
696 Ancyrocephalinae) from cichlid fishes of Lake Kariba (Zimbabwe) with descriptions of
697 five new species. *Syst. Parasitol.* 25, 159–186. <https://doi.org/10.1007/BF00007007>
- 698 Edgar, R.C., 2004. MUSCLE: Multiple sequence alignment with high accuracy and high
699 throughput. *Nucleic Acids Res.* 32, 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- 700 Ergens, R., 1981. *Diplectanum lacustre* Thurston et Paperna, 1969 (Monogenea,
701 Diplectanidae) from the Egyptian *Lates niloticus*. *Folia Parasitol.* 28, 160.
- 702 Ergens, R., Lom, J., 1970. Causative agents of fish diseases. Academia, Prague.
- 703 Excoffier, L., Lischer, H.E.L., 2010. Arlequin suite ver 3.5: a new series of programs to
704 perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* 10,
705 564–567. <https://doi.org/10.1111/j.1755-0998.2010.02847.x>
- 706 Francis, C.M., Borisenko, A. V., Ivanova, N. V., Eger, J.L., Lim, B.K., Guillén-Servent, A.,
707 Kruskop, S. V., Mackie, I., Hebert, P.D.N., 2010. The role of DNA barcodes in
708 understanding and conservation of mammal diversity in Southeast Asia. *PLoS One* 5,
709 e12575. <https://doi.org/10.1371/journal.pone.0012575>
- 710 Gastwirth, J.L., Gel, Y.R., Hui, W.L.W., Lyubchich, V., Miao, W., Noguchi, K., 2017.
711 lawstat: Tools for Biostatistics, Public Policy, and Law.
- 712 Georgieva, S., Selbach, C., Faltýnková, A., Soldánová, M., Sures, B., Skirnisson, K.,
713 Kostadinova, A., 2013. New cryptic species of the ‘*revolutum*’ group of *Echinostoma*
714 (Digenea: Echinostomatidae) revealed by molecular and morphological data. *Parasit.*
715 *Vectors* 6, 64. <https://doi.org/10.1186/1756-3305-6-64>
- 716 Hafner, M.S., Sudman, P.D., Villablanca, F.X., Spradling, T.A., Demastes, J.W., Nadler,
717 S.A., 1994. Disparate rates of molecular evolution in cospeciating hosts and parasites.
718 *Science* 265, 1087–1090.
- 719 Hassouna, N., Michot, B., Bachellerie, J.-P., Narbonne, D., 1984. The complete nucleotide

720 sequence of mouse 28S rRNA gene. Implications for the process of size increase of the
721 large subunit rRNA in higher eukaryotes. *Nucleic Acids Res.* 12, 3563–3583.

722 He, D., Chen, Y., 2007. Molecular phylogeny and biogeography of the highly specialized
723 grade schizothoracine fishes (Teleostei: Cyprinidae) inferred from cytochrome b
724 sequences. *Chinese Sci. Bull.* 52, 777–788. <https://doi.org/10.1007/s11434-007-0123-2>

725 Hebert, P.D.N., Cywinska, A., Ball, S.L., DeWaard, J.R., 2003. Biological identifications
726 through DNA barcodes. *Proceedings. Biol. Sci.* 270, 313–321.
727 <https://doi.org/10.1098/rspb.2002.2218>

728 Hubert, N., Hanner, R., Holm, E., Mandrak, N.E., Taylor, E., BurrIDGE, M., Watkinson, D.,
729 Dumont, P., Curry, A., Bentzen, P., Zhang, J., April, J., Bernatchez, L., 2008. Identifying
730 Canadian freshwater fishes through DNA barcodes. *PLoS One* 3, e2490.
731 <https://doi.org/10.1371/journal.pone.0002490>

732 Huyse, T., Pampoulie, C., Audenaert, V., Volckaert, F.A.M., 2006. First report of
733 *Gyrodactylus* spp. (Platyhelminthes: Monogenea) in the Western Mediterranean Sea:
734 molecular and morphological descriptions. *J. Parasitol.* 92, 682–690.
735 <https://doi.org/10.1645/GE-690R.1>

736 Huyse, T., Poulin, R., Théron, A., 2005. Speciation in parasites: a population genetics
737 approach. *Trends Parasitol.* 21, 469–475. <https://doi.org/10.1016/j.pt.2005.08.009>

738 International Commission on Zoological Nomenclature, 2012. Amendment of articles 8, 9,
739 10, 21 and 78 of the International Code of Zoological Nomenclature to expand and refine
740 methods of publication. *Zootaxa* 3450, 1–7.

741 Jombart, T., 2008. Adegnet: A R package for the multivariate analysis of genetic markers.
742 *Bioinformatics* 24, 1403–1405. <https://doi.org/10.1093/bioinformatics/btn129>

743 Justine, J.-L., 2005. Species of *Pseudorhabdosynochus* Yamaguti, 1958 (Monogenea:
744 Diplectanidae) from *Epinephelus fasciatus* and *E. merra* (Perciformes: Serranidae) off
745 New Caledonia and other parts of the Indo-Pacific Ocean, with a comparison of
746 measurements. *Syst. Parasitol.* 62, 1–37. <https://doi.org/10.1007/s11230-005-5480-0>

747 Justine, J.-L., Euzet, L., 2006. Diplectanids (Monogenea) parasitic on the gills of the
748 coralgroupers *Plectropomus laevis* and *P. leopardus* (Perciformes, Serranidae) off New
749 Caledonia, with the description of five new species and the erection of *Echinoplectanum*
750 n. g. *Syst. Parasitol.* 64, 147–172. <https://doi.org/10.1007/s11230-006-9028-8>

751 Justine, J.-L., Henry, É., 2010. Monogeneans from *Epinephelus chlorostigma* (Val.)
752 (Perciformes: Serranidae) off New Caledonia, with the description of three new species
753 of diplectanids. *Syst. Parasitol.* 77, 81–105. <https://doi.org/10.1007/s11230-010-9263-x>

754 Kassambara, A., Mundt, F., 2017. factoextra: extract and visualize the results of multivariate
755 data analyses.

756 Kimura, M., 1980. A simple method for estimating evolutionary rate of base substitutions
757 through comparative studies of nucleotide sequences. *J. Mol. Evol.* 16, 111–120.

758 Klett, V., Meyer, A., 2002. What, if anything, is a Tilapia?—mitochondrial ND2 phylogeny
759 of tilapiines and the evolution of parental care systems in the African cichlid fishes. *Mol.*
760 *Biol. Evol.* 19, 865–883. <https://doi.org/10.1093/oxfordjournals.molbev.a004144>

761 Kmentová, N., Gelnar, M., Mendlová, M., Van Steenberge, M., Koblmüller, S., Vanhove,

762 M.P.M., 2016. Reduced host-specificity in a parasite infecting non-littoral Lake
763 Tanganyika cichlids evidenced by intraspecific morphological and genetic diversity. Sci.
764 Rep. 6, 39605. <https://doi.org/10.1038/srep39605>

765 Kmentová, N., Van Steenberge, M., Raeymaekers, J.A.R., Koblmüller, S., Hablützel, P.I.,
766 Muterezi Bukinga, F., Mulimbwa N'sibula, T., Masilya Mulungula, P., Nzigidahera, B.,
767 Ntakimazi, G., Gelnar, M., Vanhove, M.P.M., 2018. Monogenean parasites of sardines in
768 Lake Tanganyika: diversity, origin and intra-specific variability. Contrib. to Zool. 87,
769 105–132.

770 Koch, M., Koblmüller, S., Sefc, K.M., Duftner, N., Katongo, C., Sturmbauer, C., 2007.
771 Evolutionary history of the endemic Lake Tanganyika cichlid fish *Tylochromis polylepis*:
772 a recent intruder to a mature adaptive radiation. J. Zool. Syst. Evol. Res. 45, 64–71.
773 <https://doi.org/10.1111/j.1439-0469.2006.00381.x>

774 Kumar, S., Stecher, G., Tamura, K., Gerken, J., Priesse, E., Quast, C., Schweer, T., Peplies,
775 J., Ludwig, W., Glockner, F., 2016. MEGA7: Molecular evolutionary genetics analysis
776 Version 7.0 for bigger datasets. Mol. Biol. Evol. 33, 1870–1874.
777 <https://doi.org/10.1093/molbev/msw054>

778 Leigh, J.W., Bryant, D., 2015. PopART: full-feature software for haplotype network
779 construction. Methods Ecol. Evol. 6, 1110–1116.

780 Liang, K., Leong, T., 1991. A redescription of *Pseudorhabdosynochus latesi* (Tripathi 1955)
781 and description of *Diplectanum penangi* n. sp. from *Lates calcarifer* cultured in floating
782 cages in Malaysia and Thailand. J. Biosci. 2, 77–84.

783 Lim, S.-Y., Ooi, A.-L., Wong, W.-L., 2016. Gill monogeneans of Nile tilapia (*Oreochromis*
784 *niloticus*) and red hybrid tilapia (*Oreochromis* spp.) from the wild and fish farms in
785 Perak, Malaysia: infection dynamics and spatial distribution. Springerplus 5, 1609.
786 <https://doi.org/10.1186/s40064-016-3266-2>

787 Littlewood, D.T.J., Rohde, K., Clough, K.A., 1997. Parasite speciation within or between host
788 species? - Phylogenetic evidence from site-specific polystome monogeneans. Int. J.
789 Parasitol. 27, 1289–1297. [https://doi.org/10.1016/S0020-7519\(97\)00086-6](https://doi.org/10.1016/S0020-7519(97)00086-6)

790 Mannini, P., Katonda, I., Kissaka, B., Verburg, P., 1999. Feeding ecology of *Lates stappersii*
791 in Lake Tanganyika. Hydrobiologia 407, 131–139.
792 <https://doi.org/10.1023/A:1003714203943>

793 Meier, R., Shiyang, K., Vaidya, G., Ng, P.K.L., 2006. DNA barcoding and taxonomy in
794 Diptera: A tale of high intraspecific variability and low identification success. Syst. Biol.
795 55, 715–728. <https://doi.org/10.1080/10635150600969864>

796 Meinilä, M., Kuusela, J., Ziętara, M.S., Lumme, J., 2004. Initial steps of speciation by
797 geographic isolation and host switch in salmonid pathogen *Gyrodactylus salaris*
798 (Monogenea: Gyrodactylidae). Int. J. Parasitol. 34, 515–526.
799 <https://doi.org/10.1016/j.ijpara.2003.12.002>

800 Meyer, B.S., Indermaur, A., Ehrensperger, X., Egger, B., Banyankimbona, G., Snoeks, J.,
801 Salzburger, W., 2015. Back to Tanganyika: a case of recent trans-species-flock dispersal
802 in East African haplochromine cichlid fishes. R. Soc. open Sci. 2, 140498.
803 <https://doi.org/10.1098/rsos.140498>

804 Milne, I., Wright, F., Rowe, G., Marshall, D.F., Husmeier, D., McGuire, G., 2004. TOPALi:

805 software for automatic identification of recombinant sequences within DNA multiple
806 alignments. *Bioinformatics* 20, 1806–1807.
807 <https://doi.org/10.1093/bioinformatics/bth155>

808 Mohd Agos, S., Shaharom-H, F., Ihwan Zaka, M., Hassan, M., 2016. Morphological study of
809 *Cichlidogyrus mbirizei* (Ancyrocephalidae) monogenean gill parasite on red tilapia
810 (*Oreochromis* sp.) from Como River Kenyir Lake, Terengganu Malaysia. *J. Fish. Aquat.*
811 *Sci.* 11, 432–436. <https://doi.org/10.3923/jfas.2016.432.436>

812 Mulimbwa, N., Mannini, P., 1993. Demographic characteristics of *Stolothrissa tanganyicae*,
813 *Limnothrissa miodon* and *Lates stappersii* in the Northwestern (Zairean) waters of Lake
814 Tanganyika. CIFA Occas. Pap.

815 Muss, A., Robertson, D.R., Stepien, C.A., Wirtz, P., Bowen, B.W., 2001. Phylogeography of
816 *Ophioblennius*: the role of ocean currents and geography in reef fish evolution. *Evolution*
817 (N. Y). 55, 561–572.

818 Ogle, D.H., Wheeler, P., Dinno, A., 2019. FSA: Fisheries Stock Analysis.

819 Ogutu-Ohwayo, R., 1995. Diversity and stability of fish stocks in Lakes Victoria, Kyoga and
820 Nabugabo after establishment of introduced species, in: Hart, P., Pitcher, T.J. (Eds.), *The*
821 *Impact of Species Changes in African Lakes*. Springer Netherlands, Dordrecht, pp. 59–
822 81. https://doi.org/10.1007/978-94-011-0563-7_5

823 Omland, K.E., 1997. Correlated rates of molecular and morphological evolution. *Evolution*
824 (N. Y). 51, 1381–1393.

825 Otero, O., 2004. Anatomy, systematics and phylogeny of both recent and fossil latid fishes
826 (Teleostei, Perciformes, Latidae). *Zool. J. Linn. Soc.* 81–133.
827 <https://doi.org/10.1111/j.1096-3642.2004.00111.x>

828 Paperna, I., Thurston, J.P., 1969. Monogenetic trematodes collected from cichlid fish in
829 Uganda; including the description of five new species of *Cichlidogyrus*. *Rev. Zool. Bot.*
830 *Africaines* 79, 15–33.

831 Pariselle, A., Euzet, L., 2009. Systematic revision of dactylogyridean parasites (Monogenea)
832 from cichlid fishes in Africa, the Levant and Madagascar. *Zoosystema* 31, 849–898.
833 <https://doi.org/10.5252/z2009n4a6>

834 Pariselle, A., Muterezi Bukinga, F., Van Steenberge, M., Vanhove, M.P.M., 2015.
835 Ancyrocephalidae (Monogenea) of Lake Tanganyika: IV: *Cichlidogyrus* parasitizing
836 species of Bathybatini (Teleostei, Cichlidae): reduced host-specificity in the deepwater
837 realm? *Hydrobiologia* 748, 99–119. <https://doi.org/10.1007/s10750-014-1975-5>

838 Paugy, D., Leveque, C., Teugels, G.G., 2003. The fresh and brackish water fishes of West
839 Africa - Volume 2. Royal Museum for Central Africa, Tervuren.

840 Poisot, T., Verneau, O., Desdevises, Y., Kitchener, A., Daniels, M., 2011. Morphological and
841 molecular evolution are not linked in *Lamellodiscus* (Platyhelminthes, Monogenea).
842 *PLoS One* 6, e26252. <https://doi.org/10.1371/journal.pone.0026252>

843 Poll, M., 1953. Exploration hydrobiologique du lac Tanganika, in: *Poissons Non Cichlidae*.
844 Institut Royal des Sciences Naturelles de Belgique, Brussels, p. 251.

845 Poulin, R., 2002. The evolution of monogenean diversity. *Int. J. Parasitol.* 32, 245–254.
846 [https://doi.org/10.1016/S0020-7519\(01\)00329-0](https://doi.org/10.1016/S0020-7519(01)00329-0)

847 R Core Team, 2013. R: A language and environment for statistical computing. R Foundation
848 for Statistical Computing, Vienna, Austria.

849 Rahmouni, I., Řehulková, E., Pariselle, A., Rkhami, O.B., Šimková, A., 2017. Four new
850 species of *Dactylogyirus* Diesing, 1850 (Monogenea: Dactylogyridae) parasitising the
851 gills of northern Moroccan *Luciobarbus* Heckel (Cyprinidae): morphological and
852 molecular characterisation. Syst. Parasitol. 94, 575–591. [https://doi.org/10.1007/s11230-](https://doi.org/10.1007/s11230-017-9726-4)
853 [017-9726-4](https://doi.org/10.1007/s11230-017-9726-4)

854 Roberts, T.R., 1975. Geographical distribution of African freshwater fishes. Zool. J. Linn.
855 Soc. 57, 249–319. <https://doi.org/10.1111/j.1096-3642.1975.tb01893.x>

856 Rohde, K., 1980. Diversity gradients of marine Monogenea in the Atlantic and Pacific
857 Oceans. Experientia 36, 1368–1369. <https://doi.org/10.1007/BF01960102>

858 Rohde, K., Hayward, C., Heap, M., 1995. Aspects of the ecology of metazoan ectoparasites of
859 marine fishes. Int. J. Parasitol. 25, 945–970. [https://doi.org/10.1016/0020-](https://doi.org/10.1016/0020-7519(95)00015-T)
860 [7519\(95\)00015-T](https://doi.org/10.1016/0020-7519(95)00015-T)

861 Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B.,
862 Liu, L., Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: Efficient Bayesian
863 phylogenetic inference and model choice across a large model space. Syst. Biol. 61, 539–
864 542. <https://doi.org/10.1093/sysbio/sys029>

865 Salzburger, W., 2018. Understanding explosive diversification through cichlid fish genomics.
866 Nat. Rev. Genet. 1. <https://doi.org/10.1038/s41576-018-0043-9>

867 Salzburger, W., Mack, T., Verheyen, E., Meyer, A., 2005. Out of Tanganyika: genesis,
868 explosive speciation, key-innovations and phylogeography of the haplochromine cichlid
869 fishes. BMC Evol. Biol. 5, 17. <https://doi.org/10.1186/1471-2148-5-17>

870 Salzburger, W., Van Bocxlaer, B., Cohen, A.S., 2014. Ecology and evolution of the African
871 Great Lakes and their faunas. Annu. Rev. Ecol. Evol. Syst. 45, 519–545.
872 <https://doi.org/10.1146/annurev-ecolsys-120213-091804>

873 Schlick-Steiner, B.C., Steiner, F.M., Seifert, B., Stauffer, C., Christian, E., Crozier, R.H.,
874 2010. Integrative taxonomy: A multisource approach to exploring biodiversity. Annu.
875 Rev. Entomol. 55, 421–438. <https://doi.org/10.1146/annurev-ento-112408-085432>

876 Schoelink, C., Cruaud, C., Justine, J.-L., 2012. Are all species of *Pseudorhabdosynochus*
877 strictly host specific? - A molecular study. Parasitol. Int. 61, 356–359.
878 <https://doi.org/10.1016/j.parint.2012.01.009>

879 Sigura, A., Justine, J.-L., 2008. Monogeneans of the speckled blue grouper, *Epinephelus*
880 *cyanopodus* (Perciformes, Serranidae), from off New Caledonia, with a description of
881 four new species of *Pseudorhabdosynochus* and one new species of *Laticola*
882 (Monogenea: Diplectanid. Zootaxa 44, 1–44.

883 Šimková, A., Dávidová, M., Papoušek, I., Vetešník, L., 2013. Does interspecies hybridization
884 affect the host specificity of parasites in cyprinid fish? Parasit. Vectors 6, 95.
885 <https://doi.org/10.1186/1756-3305-6-95>

886 Šimková, A., Matějusková, I., Cunningham, C.O., 2006. A molecular phylogeny of the
887 Dactylogyridae sensu Kritsky & Boeger (1989) (Monogenea) based on the D1-D3
888 domains of large subunit rDNA. Parasitology 133, 43–53.
889 <https://doi.org/10.1017/S0031182006009942>

- 890 Šimková, A., Morand, S., Jobet, E., Gelnar, M., Verneau, O., 2004. Molecular phylogeny of
891 congeneric monogenean parasites (*Dactylogyrus*): a case of intrahost speciation.
892 Evolution 58, 1001–1018. <https://doi.org/10.1111/j.0014-3820.2004.tb00434.x>
- 893 Šimková, A., Plaisance, L., Matějusková, I., Morand, S., Verneau, O., 2003. Phylogenetic
894 relationships of the Dactylogyridae Bychowsky, 1933 (Monogenea: Dactylogyroidea):
895 The need for the systematic revision of the Ancyrocephalinae Bychowsky, 1937. Syst.
896 Parasitol. 54, 1–11. <https://doi.org/10.1023/A:1022133608662>
- 897 Sinnappah, N.D., Lim, L.-H.S., Rohde, K., Tinsley, R., Combes, C., Verneau, O., 2001. A
898 paedomorphic parasite associated with a neotenic amphibian host: phylogenetic evidence
899 suggests a revised systematic position for sphyranuridae within anuran and turtle
900 polystomatoineans. Mol. Phylogenet. Evol. 18, 189–201.
901 <https://doi.org/10.1006/mpev.2000.0877>
- 902 Snoeks, J., 2000. Ancient lakes: biodiversity, ecology and evolution. Adv. Ecol. Res. 31, 17–
903 38. [https://doi.org/10.1016/S0065-2504\(00\)31005-4](https://doi.org/10.1016/S0065-2504(00)31005-4)
- 904 Stamatakis, A., 2014. RAxML version 8: A tool for phylogenetic analysis and post-analysis
905 of large phylogenies. Bioinformatics 30, 1312–1313.
906 <https://doi.org/10.1093/bioinformatics/btu033>
- 907 Thomas, J.A., Welch, J.J., Lanfear, R., Bromham, L., 2010. A generation time effect on the
908 rate of molecular evolution in invertebrates. Mol. Biol. Evol. 27, 1173–1180.
909 <https://doi.org/10.1093/molbev/msq009>
- 910 Thurston, J.P., Paperna, I., 1969. *Diplectanum lacustris* sp. nov. (Dactylogyroidea:
911 Diplectanidae), a Monogenetic Trematode from the gills of the Nile Perch. Proc.
912 Helminthol. Soc. 214–218.
- 913 Tingbao, Y., Kritsky, D.C., Yuan, S., Jianying, Z., Suhua, S., Agrawal, N., 2006. Diplectanids
914 infesting the gills of the barramundi *Lates calcarifer* (Bloch) (Perciformes:
915 Centropomidae), with the proposal of *Laticola* n. g. (Monogenoidea: Diplectanidae).
916 Syst. Parasitol. 63, 127–41. <https://doi.org/10.1007/s11230-005-9006-6>
- 917 Vanhove, M.P.M., Pariselle, A., Van Steenberge, M., Raeymaekers, J.A.M., Hablützel, P.I.,
918 Gillardin, C., Hellemans, B., Breman, F.C., Koblmüller, S., Sturmbauer, C., Snoeks, J.,
919 Volckaert, F.A.M., Huyse, T., 2015. Hidden biodiversity in an ancient lake: phylogenetic
920 congruence between Lake Tanganyika trophic cichlids and their monogenean
921 flatworm parasites. Sci. Rep. 5, 13669, 10.1038/srep13669.
922 <https://doi.org/10.1038/srep13669>
- 923 Vanhove, M.P.M., Tessens, B., Schoelinck, C., Jondelius, U., Littlewood, D.T.J., Artois, T.,
924 Huyse, T., 2013. Problematic barcoding in flatworms: A case-study on monogeneans and
925 rhabdocoels (Platyhelminthes). Zookeys 365, 355–379.
926 <https://doi.org/10.3897/zookeys.365.5776>
- 927 Vanhove, M.P.M., Volckaert, F.A.M., Pariselle, A., 2011. Ancyrocephalidae (Monogenea) of
928 Lake Tanganyika: I: Four new species of *Cichlidogyrus* from *Ophthalmotilapia ventralis*
929 (Teleostei: Cichlidae), the first record of this parasite family in the basin. Zool. 28, 253–
930 263. <https://doi.org/10.1590/S1984-46702011000200016>
- 931 Villar-Torres, M., Repullés-Albelda, A., Montero, F.E., Raga, J.A., Blasco-Costa, I., 2019.
932 Neither *Diplectanum* nor specific: a dramatic twist to the taxonomic framework of
933 *Diplectanum* (Monogenea: Diplectanidae). Int. J. Parasitol. 49, 365–374.

<https://doi.org/10.1016/J.IJPARA.2018.11.003>

- Whittington, I.D., Chisholm, L.A., Rohde, K., 1999. The larvae of Monogenea (Platyhelminthes). *Adv. Parasitol.* 44, 139–232. [https://doi.org/10.1016/S0065-308X\(08\)60232-8](https://doi.org/10.1016/S0065-308X(08)60232-8)
- Wickham, H., 2009. *ggplot2*. Springer, New York. <https://doi.org/10.1007/978-0-387-98141-3>
- Will, K.W., Rubinoff, D., 2004. Myth of the molecule: DNA barcodes for species cannot replace morphology for identification and classification. *Cladistics* 20, 47–55. <https://doi.org/10.1111/j.1096-0031.2003.00008.x>
- Wu, W., Goodman, M., Lomax, M.I., Grossman, L.I., 1997. Molecular evolution of cytochrome c oxidase subunit IV: Evidence for positive selection in simian primates. *J. Mol. Evol.* 44, 477–491. <https://doi.org/10.1007/PL00006172>
- Wu, X.Y., Chilton, N.B., Zhu, X.Q., Xie, M.Q., Li, A.X., 2005. Molecular and morphological evidence indicates that *Pseudorhabdosynochus lantauensis* (Monogenea: Diplectanidae) represents two species. *Parasitology* 130, 669–677. <https://doi.org/10.1017/S0031182004007152>
- Wu, X.Y., Zhu, X.Q., Xie, M.Q., Li, A.X., 2007. The evaluation for generic-level monophyly of Ancyrocephalinae (Monogenea, Dactylogyridae) using ribosomal DNA sequence data. *Mol. Phylogenet. Evol.* 44, 530–544. <https://doi.org/10.1016/j.ympev.2007.03.025>
- Zahradníčková, P., Barson, M., Luus-Powell, W.J., Přikrylová, I., 2016. Species of *Gyrodactylus* von Nordmann, 1832 (Platyhelminthes: Monogenea) from cichlids from Zambezi and Limpopo river basins in Zimbabwe and South Africa: evidence for unexplored species richness. *Syst. Parasitol.* 93, 679–700. <https://doi.org/10.1007/s11230-016-9652-x>

Funding

Research was supported by the Czech Science Foundation (GBP505/12/G112 – ECIP) and GA19-13573S), Belgian Science Policy (BELSPO), EMBRC Belgium - FWO project GOH3817N as well as by a joint program between the Austrian agency for international mobility and cooperation in education, science and research and the Ministry of Education, Youth and Sports (project number 8J18AT007), the Austrian Agency for International Cooperation in Education and Research (OEAD; project number CZ 08/2018; to SK) and the Systematics Research Fund (SRF; to SK).

967 Table 1: An overview of host species examined for monogenean parasites with locality and
 968 country.

Host species	Locality (geographic coordinates, year)	Locality – subbasins (Danley et al., 2012) or country	Number of fish specimens examined (accession number in RMCA)	Number of infected fish specimens	Number of monogenean individuals*
Lake Tanganyika					
<i>L. angustifrons</i>	Mpulungu (08°46'S- 31°07'E, 27.7.1967)	The southern subbasin	1 (MRAC 190480)	1	2
	Mpulungu (12.4.2018)	The southern subbasin	4(-)	1	15
	Rumonge (03°58'S- 29°25'E, 30.6.1967)	The northern subbasin	2 (MRAC 94069.0052-53)	2	3
	Sumbu Bay (08°31'S-30°29'E, 31.3.1947)	The southern subbasin	1 (MRAC 90850)	1	1
<i>L. mariae</i>	Bujumbura (03°23'S- 29°22'E, 5.5.1947)	The northern subbasin	5 (MRAC 90908-912)	5	14
	Ilagala (05°12'S- 29°50'E, 20.8.1993)	The northern subbasin	3 (MRAC 93152.0318-20)	2	10
	Kilomoni (04°20'S, 29°09'E, 12.8.2016)	The northern subbasin	2 (-)	1	1
	Mpulungu (27.7.1967)	The southern subbasin	2 (MRAC 190493-94)	1	3
	Mpulungu (16.4.2018)	The southern subbasin	7 (-)	0	0
	Mulembwe (06°07'S 29°16'E, 9.4.2010)	The central subbasin	7 (-)	3	19
	Nyanza (04°20'S- 29°35'E, 1.1.1997)	The northern subbasin	1 (MRAC 53738)	1	23
	Rumonge (30.6.1994)	The southern subbasin	1 (MRAC 94069.0067)	0	0
	Sumbu Bay (31.3.1947)	The southern subbasin	2 (MRAC 90878-79)	2	22
	Bujumbura (4.5.1947)	The northern subbasin	5 (MRAC 90805-9)	5	24
	Crock Island (08°42'S-31°07'E, 16.4.2018)	The southern subbasin	8 (-)	2	13
	Edith Bay (06°30'S- 29°55'E, 30.5.1947)	The southern subbasin	3 (MRAC 90833-35)	5	10
<i>L. microlepis</i>	Katukula (08°35'S- 31°10'E, 14.4.2018)	The southern subbasin	5 (-)	4	22
	Moba Bay (30.12.1995)	The central subbasin	2 (MRAC 90725-6)	0	0
	Mpulungu (13.4. – 17.4. 2018)	The southern subbasin	13 (-)	6	29
	Nyanza Lac (1.1.1937)	The northern subbasin	11 (MRAC 53698- 703; 53725-29)	11	181
	Sumbu Bay (9.4.1995)	The southern subbasin	3 (MRAC 95096.1192,98,99)	1	1
	Uvira (03°22' S 29°09'E, 12.8.2016)	The northern subbasin	7 (-)	2	17
	Karala (05°33'S- 29°28'E, 10.4.1947)	The northern subbasin	1 (MRAC P90928)	0	0
	Kasasa (08°31'S- 30°42'E, 6.9.1967)	The southern subbasin	3 (MRAC 190126;35- 6)	0	0
	Mpulungu (12.4.2008)	The southern subbasin	3 (-)	0	0
	Mpulungu (6.4.2018)	The southern subbasin	3 (-)	0	0
<i>L. stappersii</i>					

	Uvira (12.8.2016)	The northern subbasin	28 (-)	0	0
	Other localities				
<i>L. niloticus</i>	Bahr-Sara (08°56'N-17°58'E, 1. -. 31.3.1965)	Tchad	1 (MRAC 154006)	1	2
	Kisumu, Lake Victoria (00°06'S-34°45'E, 17.12.1991)	Kenya	3 (MRAC 91104.37-39)	2	8
	Kossou (07°10'N-05°20'E, 17.12.1973)	Ivory Coast	5 (MRAC 74014.328-29; 2755-56)	4	22
	Lake Nasser (24°05'N-33°00'E, 26.2. - 11.3.1984)	Egypt	3 (MRAC 84006.0116-18)	0	0
	Lake Nasser (1.9.-30.9.1983)	Egypt	2 (MRAC 83030.0114-15)	1	1
	Luxor market (25°42'N 32°38'E, 24.11.2000)	Egypt	1 (MRAC 190480)	0	0
	Njala, riv. Taja (08°06'N-12°04'e, 12.4.1969)	Sierra Leone	5 (MRAC 73010.7057-61)	1	8
	Nyawiega, Lake Albert (01°28'N-30°56'E, 21.11.-6.12.1989)	Uganda	3 (MRAC 89059.0279)	1	2
	Nzunzu, Lake Albert (01°19'N-30°72'E, 5.4.-6.4.2017)	Uganda	1 (MRAC 2016.036.P)	1	2
	Nzunzu, Lake Albert (5.4.-6.4.2017)	Uganda	11 (MRAC 2016.036.P)	2	18

* Only one gill arch examined in the case of specimens retrieved from the RMCA

973 Table 2: Comparison of measurements performed on haptoral and genital hardparts of *Dolicirroplectanum lacustre* comb. nov. reported in Thurston
974 and Paperna (1969) from Lake Volta, in Ergens (1981) from Cairo and in this study with host species and locality (a – mean value±standard
975 deviation, b – range).

Parameters (µm)	<i>L. niloticus</i> , Lake Volta	<i>L. niloticus</i> , Cairo	<i>L. angustifrons</i> , Lake Tanganyika	<i>L. mariae</i> , Lake Tanganyika	<i>L. microlepis</i> , Lake Tanganyika
Total length	650-1000	-	544,2 (n=1)	522,0±35,4 ^a (487,4 - 587,3) ^b ; n=11	675,0±66,9 (588,1 - 813,1); n=13
Total width	150-250	-	188,7 (n=1)	178,8±39,5 (115,5 - 243,3); n=11	183,2±15,1 (4,7 - 204,1); n=13
Ventral anchor					
Length to notch	19-20	20-22 (n=3)	20,4 (19,5 - 21,3); n=7	19,3±1,4 (16,3 - 24,4); n=25	18,7±0,9 (16,4 - 21,2); n=61
Total length	70-80	53-57 (n=3)	43,2 (41,4 - 44,1); n=7	42,6±2,0 (41,1 - 47,2); n=28	43,2±2,4 (40,5 - 48,3); n=62
Length to inner root	-	-	22,8 (20,9 - 24,8); n=7	22,6±1,6 (18,3 - 26,9); n=27	22,2±1,1 (18,7 - 24,9); n=52
Inner root length	10-20	10-12 (n=3)	8,7 (7,2 - 9,8); n=7	9,8±0,8 (7,7 - 11,2); n=28	9,2±1,1 (6,1 - 11,4); n=54
Outer root length	40-60	32-35 (n=3)	22,6 (20,1 - 24,4); n=7	23,7± (19,1 - 27,9); n=24	24,8±2,6 (19,2 - 30,1); n=52
Point length	5-7	9-10 (n=3)	8,2 (7,3 - 9,5); n=6	7,2±1,1 (5,4 - 9,2); n=25	7,0±1,1 (4,9 - 9,0); n=45
Dorsal anchor					
Total length	40-50	44-47 (n=3)	35,3 (32,6 - 36,6); n=6	32,3±2,0 (28,4 - 35,6); n=25	33,4±2,1 (26,0 - 38,6); n=47
Point length	-	9-10 (n=3)	7,4 (6,9 - 8,0); n=6	6,3±0,9 (4,7 - 8,1); n=13	6,0±0,9 (4,1 - 8,2); n=31
Ventral bar					
Straight length	50-60	38-44 (n=3)	44,8 (40,7 - 51,5); n=4	50,8±6,4 (39,2 - 63,3); n=30	46,6±4,5 (36,4 - 55,5); n=58
Maximum width	-	8-11 (n=3)	8,2 (7,6 - 9,2); n=6	11,4±2,1 (7,2 - 15,3); n=27	9,3±1,7 (6,1 - 13,2); n=52
Dorsal bar					
Straight length	35-40	34-38 (n=3)	31,5 (29,9 - 33,5); n=6	37,6±3,9 (29,6 - 46,1); n=29	32,8±3,5 (24,3 - 39,5); n=60
Maximum width	-	9-12 (n=3)	8,9 (7,7 - 9,8); n=5	7,2±1,4 (5,1 - 9,6); n=28	6,5±0,9 (4 - 9,0); n=59
Ventral squamodisc					
Length	30-40	38-44 (n=3)	41,6 (35,2 - 48,6); n=4	38,4±7,6 (32,3 - 57,4); n=15	38,3±6,5 (26,7 - 57,2); n=22
Width	50-70	38-51 (n=3)	47,8 (46,2 - 50,0); n=4	41,9±7,3 (29,8 - 60,4); n=16	42,5±4,4 (33,3 - 53,1); n=22
Dorsal squamodisc					
Length	-	38-44 (n=3)	35,9 (33,6 - 39,6); n=4	34,8±4,8 (29,5 - 43,4); n=6	34,9±5,8 (21,3 - 42,9); n=16
Width	-	38-51 (n=3)	42,9 (41,5 - 45,0); n=4	38,0±7,9 (33,4 - 57,2); n=8	40,4±4,3 (30,0 - 46,2); n=16
Hook					
Copulatory tube straight length	21-23	-	10,5 (9,5 - 11,3); n=4	10,4±0,6 (9,4 - 11,8); n=27	10,0±0,7 (7,9 - 11,7); n=49
			27,6 (24,5 - 29,2); n=3	33,9±5,2 (24,1 - 43,6); n=20	32,3±3,9 (21,4 - 41,5); n=38

Copulatory tube width	-	-	20,6 (19,6 - 21,3); n=3	20,6±4,0 (13,5 - 29,1); n=22	33,4±2,1 (26,0 - 38,6); n=47
Vagina	Not reported	Not reported	Reported in 0 out of 5 available specimens	Reported in 2 out of 24 available specimens	Reported in 0 out of 38 available specimens
Total length	-	-	-	22,1±0,8 (21,5 - 22,6); n=2	-
Tube length	-	-	-	5±0,6 (4,6 - 5,4); n=2	-
Point length	-	-	-	6±0,3 (5,8 - 6,2); n=2	-
Eyes spots					
Smaller pair distance	-	-	44,2 (40,0 - 48,4); n=2	37,3±6,3 (30,0 - 50,0); n=14	-
Larger pair distance	-	-	42,6 (42,2 - 43,0); n=2	32,9±5,1 (26,8 - 46,7); n=14	-
Pharynx length	-	-	-	34,3±9,9 (22,9 - 53,0); n=8	-
Testes					
Length	-	-	-	94,3 (n=1)	-
Width	-	-	-	42,7±13,9 (29,7 - 78,1); n=13	-
Ovary width	-	-	-	31,0±10,9 (31,0 - 64,6); n=14	-

976

977

Parameters (µm)	<i>L. niloticus</i> , Lake Albert1	<i>L. niloticus</i> , Lake Albert2	<i>L. niloticus</i> , Lake Victoria	<i>L. niloticus</i> , river Taja	<i>L. niloticus</i> , Lake Kossou
Total length	736,5±115,3 ^a (644,1 - 931,3) ^b ; n=5	553,1±52,1 (497,2 - 600,3); n=3	537,9±34,8 (509,2 - 576,6); n=3	662,1±113,8 (510,9 - 787,0); n=5	574,7±22,2 (550,3 - 593,7); n=3
Total width	282,0±57,6 (191,4 - 333,9); n=5	182,8±46,9 (151,7 - 236,7); n=3	220,5±11,4 (210,6 - 233,0); n=3	185,0±56,4 (141,7 - 277,8); n=5	217,3±13,6 (208,3 - 233,0); n=3
Ventral anchor					
Length to notch	21,1±1,1 (19,3 - 23,1); n=18	19,3±1,2 (18,2 - 20,6); n=3	20,5±0,6 (19,7 - 21,1); n=4	17,1±0,7 (16,2 - 18,1); n=6	20,9±1,0 (19,5 - 23,1); n=10
Total length	53,4±2,9 (50,0 - 66,4); n=18	42,1±2,1 (40,2 - 44,0); n=3	43,1±1,4 (42,7 - 45,8); n=5	34,4±1,2 (32,8 - 36,5); n=6	43,9±1,8 (40,5 - 45,9); n=10
Length to inner root	24,7±1,9 (21,0 - 29,4)	23,8±0,8 (22,9 - 24,3); n=3	24,9±1,4 (23,3 - 25,8); n=3	21,0±1,1 (19,6 - 22,5); n=5	23,7±1,2 (21,4 - 25,2); n=8
Inner root length	11,3±0,7 (10 - 12,9); n=18	10,7±1,0 (9,8 - 11,8); n=3	10,5±0,9 (9,5 - 11,5)	8,9±0,7 (8,1 - 9,7); n=6	10,0±0,9 (8,0 - 11,1); n=10
Outer root length	32,5±3,3 (29,7 - 43,2); n=18	23,3±1,6 (22,0 - 25,0); n=3	24,1±1,5 (22,7 - 25,7); n=3	17,5±13,4 (15,6 - 19,4); n=6	23,5±1,3 ((21,5 - 25,7); n=10
Point length	7,9±1,1 (6 - 9,4); n=18	8,5±0,7 (7,9 - 9,3); n=3	9,5±0,9 (8,8 - 10,7)	7,4±1,1 (5,6 - 8,3); n=5	8,0±1,3 (5,4 - 9,6); n=9
Dorsal anchor					
Total length	47,3±4,0 (42,7 - 60,2); n=17	39,0±1,0 (37,6 - 39,9); n=3	40,7±0,4 (40,2 - 41,0); n=3	28,1±1,0 (27,0 - 29,6); n=6	39,9±1,8 (36,7 - 42,4); n=10
Point length	7,9±1,0 (5,9 - 9,2); n=12	7,7±1,2 (6,8 - 8,5); n=2	8,9±0,6 (8,6 - 9,6); n=3	6,6±1,4 (4,8 - 8,6); n=5	7,8±0,6 (7,3 - 8,5); n=3
Ventral bar					
Straight length	72,6±4,8 (61,5 - 78,8); n=16	59,1±11,5 (52,2 - 72,4); n=3	45,8±6,4 (36,5 - 50,0); n=4	40,0±0,9 (39,2 - 40,5); n=2	50,1±1,6 (47,8 - 52,3); n=7
Maximum width	18,5±1,9 (13,5 - 21,0); n=11	15,3±0,4 (15,0 - 15,6); n=2	15,3 (n=1)	10,2±2,7 (8,3 - 12,1); n=2	13,2±1,4 (11,2 - 14,7); n=7

Dorsal bar						
Straight length	55,4±3,2 (47,6 - 59,9); n=17	41,0±2,1 (39,5 - 42,4); n=2	39,0±2,8 (36,7 - 42,1); n=3	30,0±2,9 (26,1 - 33,5); n=6	37,4±2,5 (32,7 - 40,6); n=10	
Maximum width	18,4±2,3 (13,5 - 22,0); n=15	16,5±2,7 (13,8 - 19,6); n=4	7,4±0,5 (6,9 - 7,8); n=3	6,3±2,1 (4,3 - 9,8); n=5	9,3±2,7 (6,5 - 13,9); n=8	
Ventral squamodisc						
Length	64,1±7,2 (51,8 - 80,4); n=14	38,4 (n=1)	35,0±2,8 (32,4 - 39,6); n=5	32,1±5,7 (27,9 - 38,6); n=3	34,6±4,4 (29,8 - 38,6); n=3	
Width	78,3±13,0 (62,5 - 118,4); n=14	44,0 (n=1)	51,8±3,5 (48,1 - 55,9); n=5	36,4±6,4 (31,5 - 43,7); n=3	50,9±3,0 (47,5 - 52,9); n=3	
Dorsal squamodisc						
Length	61,4±11,1 (45,4 - 85,6); n=11	39,0±1,8 (36,9 - 40,1); n=3	32,0±2,4 (29,6 - 35,7); n=5	24,3±2,8 (21,3 - 26,8); n=3	35,3±4,3 (30,3 - 40,1); n=4	
Width	63,9±6,3 (49,9 - 71,2); n=11	48,8±10,5 (42,3 - 60,9); n=3	42,9±3,5 (38,8 - 46,9); n=5	24,9±0,8 (24,2 - 25,8); n=3	41,9±2,4 (38,4 - 43,5); n=4	
Hook	11,2±0,8 (10,3 - 14,0); n=18	10,4±1,0 (9,3 - 11,0); n=3	11,4± (11,0 - 12,2); n=4	9,7±0,7 (8,4 - 10,3); n=6	11,3±0,9 (10,1 - 13,2); n=11	
Copulatory tube straight length	58,3±8,8 (44,5 - 83,1); n=18	44,3±3,8 (40,8 - 49,5); n=4	43,7±11,8 (36,1 - 64,6); n=5	27,3±2,9 (24,2 - 31,2); n=4	35,5±4,8 (29,2 - 42,0); n=6	
Copulatory tube width	36,2±4,3 (27,2 - 44,5); n=18	27,6±5,3 (21,0 - 33,2); n=4	26,1± (24,5 - 28,0); n=5	26,1±3,6 (21,2 - 30); n=4	22,1±2,3 (18,2 - 24,5); n=6	
Vagina	Reported in 18 out of 18 available specimens	Reported in 4 out of 4 available specimens	Reported in 1 out of 6 available specimens	Reported in 0 out of 6 available specimens	Reported in 3 out of 11 available specimens	
Total length	41,1±4,5 (34,0 - 47,3); n=14	35,7±5,6 (30,1 - 41,3); n=3	-	-	-	
Tube length	7,8±1,1 (4,8 - 9,3); n=14	6,9±1,4 (5,3 - 7,9); n=3	6,7 (n=1)	-	-	
Point length	7,0±0,7 (6,0 - 8,5); n=16	8,0±1,1 (6,7 - 8,7); n=3	8,9 (n=1)	-	-	
Eyes spots						
Smaller pair distance	56,7±10,2 (42,8 - 80,4); n=16	-	-	-	-	
Larger pair distance	47,2±8,5 (34,9 - 64,6); n=17	-	-	-	-	
Pharynx length	56,3±11,0 (39,9 - 73,2); n=10	-	-	-	-	
Testes						
Length	102,1±46,7 (57,9 - 150,9); n=3	-	-	-	-	
Width	70,3±16,2 (50,5 - 103,2); n=9	-	-	-	-	
Ovary width	72,7±11,1 (50,7 - 88,9); n=9	-	-	-	-	

978

979

980

981

982 Table 3: Genetic intraspecific variability indices in a 412 bp portion of COI mtDNA region of *Dolicirroplectanum lacustre* comb. nov.

	Maximum uncorrected p-distance (number of individuals)	Nucleotide diversity	Haplotype diversity	Number of polymorphic sites
Lake Albert	1.2% (14)	0.0036+/-0.0026	0.8022+/-0.0936	6
Lake Tanganyika	0.7% (38)	0.0019+/-0.0016	0.5747+/-0.0713	4

983

984 **Figure captions**

986 Figure 1: Measurements for sclerotized structures of haptor and reproductive organs of
 987 *Dolicirroplectanum lacustre* comb. nov. A Ventral Anchor: 1—Total length, 2—Length to
 988 notch, 3—Length to inner root, 4—Inner root length, 5—Outer root length, 6—Point length;
 989 B Dorsal anchor: 7—Total length, 8—Point length; C Ventral bar: 9—Straight length 10—
 990 Maximum width; D Dorsal bar: 11—Straight length, 12—Maximum width; E Hook: 13—
 991 Hook length; F Squamodisc: 14—Squamodisc length, 15—Squamodisc width; G Male
 992 copulatory organ: 16—Copulatory tube length, 17—Copulatory tube width; H Vagina: 18—
 993 Total length, 19—Tube length, 20—Point length.

Figure 2: Localities with confirmed presence of *Dolicirroplectanum lacustre* comb. nov. Star – localities sampled for the original description of *D. lacustre* by Thurston & Paperna, 1969, triangle – locality documented by Ergens, 1981, circle – localities sampled in this study. M – Mpulungu, C – Crocodile Island, K – Katukula. Colours denote host species: black – *Lates niloticus*, blue (number 2) – *L. angustifrons*, green (number 1) – *L. mariae*, red (number 3) – *L. microlepis*. Map created using SimpleMappr software v7.0.0. (available at <http://www.simplemappr.net>. Accessed February 25, 2018).

1002

1003 Figure 3: *Dolicirroplectanum lacustre* comb. nov. collected from *Lates niloticus* in Lake Albert.

1004 Specimen drawn from the ventral view. e, eye spots; i, intestine; mg, Mehlis' glands; o, ovary;

1005 oö, oötype; ov, oviduct; p, pharynx; pr, prostatic reservoir; sv, seminal vesicle; t, testes; u,

1006 uterus; v, vittellaria; vd, vas deferens, A, ventral anchor; B, dorsal anchor; C, male copulatory

1007 organ; D, vagina, E, ventral bar; F, dorsal bar; G, hook; H, ventral squamodisc; I, dorsal
1008 squamodisc.

1010 Figure 4: Haptoral and male genital sclerotised structures of *Dolicirroplectanum lacustre* comb.
 1011 nov. from different host species and localities collected in this study (scale bars A-F: 25 µm;
 1012 G-N: 10 µm). A) Opisthaptor, *L. niloticus* in Lake Albert B) Opisthaptor, *L. niloticus* in Taja
 1013 River C) Opisthaptor, *L. microlepis* in Lake Tanganyika D) Opisthaptor, *L. niloticus* in Lake
 1014 Victoria E) Opisthaptor, *L. niloticus* in Lake Kossou F) Opisthaptor, *L. mariae* in Lake
 1015 Tanganyika G) Male copulatory organ, *L. niloticus* in Lake Albert H) Male copulatory organ,
 1016 *L. mariae* in Lake Tanganyika I) Male copulatory organ, *L. niloticus* in Lake Kossou J)
 1017 Sclerotised vagina, *L. mariae* in Lake Tanganyika K) Sclerotised vagina, *L. niloticus* in Lake
 1018 Albert L) Male copulatory organ, *L. niloticus* in Taja River M) Male copulatory organ, *L.*
 1019 *microlepis* N) Sclerotised vagina, *L. niloticus* in Lake Victoria. Pictures were stacked.

1020
 1021 Figure 5: Sclerotised structures of *Dolocirroplectanum penangi* comb. nov.. A) Ventral anchor
 1022 (MNHN HEL1086), Hainan, China B) Male copulatory organ (USNPC 180-A 6), Zhanjiang,
 1023 China. Scale bar: 20 µm; several layers in the picture were combined.

Figure 6: Bayesian inference phylogram based on 28S fragments from 33 haplotypes of different diplectanid species. Posterior probabilities for Bayesian inference (before slashes) and bootstrap percentages for maximum likelihood (behind slashes) are shown. The values lower than 90 of posterior probability and 80 for maximum likelihood are marked with an asterisk. The clade containing two lineages of *Dolicirroplectanum lacustre* comb. nov. is boxed. The scale-bar indicates the expected number of substitutions per site.

1031

1032 Figure 7: Bayesian inference phylogram based on 18S fragments from 15 haplotypes of
 1033 different diplectanid species. Posterior probabilities for Bayesian inference (before slashes)
 1034 and bootstrap percentages for maximum likelihood (behind slashes) are shown. The values
 1035 lower than 90 of posterior probability and 80 for maximum likelihood are marked with an
 1036 asterisk. The clade containing two lineages of *Dolicirroplectanum lacustre* comb. nov. is
 1037 boxed. The scale-bar indicates the expected number of substitutions per site.

1038

1039 Figure 8: Morphometric variability of haptoral structures of *Dolicirroplectanum lacustre* comb.

1040 nov. A) biplot of PCA (first two axes) based on measurements of haptoral sclerotized structures.

B) Density plots depicting the total size of the dorsal anchor. Colours and signs denote host species and locality of specimens.

Figure 9: Haplotype network of *Dolicirroplectanum lacustre* comb. nov. COI sequences (n = 52). The circles represent different haplotypes with the size proportional to the number of individuals sharing this haplotype. Haplotypes are connected with lines, indicating the number of substitutions between haplotypes. Colours correspond to A) the host species and B) geographic origin (subbasins in Lake Tanganyika).