Determinants of dietary and physical activity behaviours among women of reproductive age in urban sub-Saharan Africa: a systematic review

Peter Yiga^{1,2}, Jan Seghers³, Patrick Ogwok², and Christophe Matthys^{1,4*}

¹Nutrition & Obesity Unit, Clinical and Experimental Endocrinology, Department of Chronic Diseases, Metabolism and Aging, KU Leuven, Belgium

²Department of Food Technology, Kyambogo University, Kampala, Uganda

³Department of Movement Sciences, KU Leuven, Leuven, Belgium;

⁴Clinical Nutrition Unit, Department of Endocrinology, University Hospitals Leuven, Leuven, Belgium

Corresponding author: Christophe Matthys, DIEET-Herestraat 49, 3000 Leuven, TEL. +3216342655, christophe.matthys@uzleuven.be

Key-words: Urban sub-Saharan Africa, Physical activity, Dietary Behaviour, determinants, Women of reproductive age

Running Title: Determinants of diet and physical activity

This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114520001828

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Abstract

Urban sub-Saharan Africa is in a nutrition transition shifting towards consumption of energy dense nutrient poor diets and decreasing physical activity. Determinants of nutrition transition in sub-Saharan Africa are presently not well understood. The objective of this review was to synthesize available data on determinants of dietary and physical activity behaviours among women of reproductive age in urban sub-Saharan Africa according to the socio-ecological framework. We searched MEDLINE, EMBASE, Scopus, Web of Science and bibliographies of included articles for qualitative, observational and randomized controlled studies published in English from January 2000 to September 2018. Studies conducted within general population of women aged 18 to 49 years were included. Searches were according to a predefined protocol published on PROSPERO (ID=CRD42018108532). Two reviewers independently screened identified studies. From a total of 9853 unique references, 23 studies were retained, and were mainly from South and West Africa. No rigorous designed quantitative study was identified. Hence, data synthesis was narrative. Notable determinants of dietary behaviour included; convenience, finances, social network, food skills and knowledge gaps, food deserts and culture. Cultural beliefs include strong relationship between high social status and weight gain, energy dense confectionary, salt or fat rich foods. Physical activity is influenced by fast-changing transport environment and cultural beliefs which instigate unfavourable gender stereo types. Studies with rigorous qualitative and quantitative designs are required to validate and develop the proposed frameworks further especially within East Africa. Nevertheless, available insights suggest a need for comprehensive skill-based interventions focusing on socio-cultural misconceptions and financial limitations.

Key words; Urban Sub Saharan Africa, Physical activity, Dietary Behaviour, Determinants, Women of reproductive age

INTRODUCTION

Lifestyle related non-communicable diseases (NCD) account for almost half of all deaths and disability in low middle-income countries (LMIC) ⁽¹⁾. It is projected that NCD cases will overtake infectious diseases by 2030 in sub-Saharan Africa (SSA) ^(2; 3; 4). Even with the worrying trend, NCD are not given due attention as a public health issue in SSA ^(5; 6). The key determinants of NCD are overweight and obesity. Prevalence of overweight and obesity is increasing at a high rate in urban SSA particularly among women of reproductive age (WRA). The prevalence is higher among women at a rate that is almost twice as high as in the men ⁽⁷⁾. For example, while 18.9% of urban Ugandan men are overweight/obese, levels among urban women stand at 44% ⁽⁸⁾. In Tanzania, an estimated 42% of urban women are either overweight or obese ⁽⁹⁾ while in South Africa the prevalence is at 70% ⁽¹⁰⁾. Maternal overweight and obesity poses health risks to mothers as well as their offspring ^(11; 12). The key modifiable risk factors for overweight and obesity are unhealthy dietary pattern and physical inactivity ^(13; 14). Studies depict an increase in physical inactivity in urban Africa, especially among women ⁽¹⁵⁾. Projections show that 2018 to 2030 WHO global action plan on physical activity (PA) target (a 15% relative reduction in insufficient PA) cannot be met ⁽¹⁶⁾. Dietary patterns observed among populations of urban low/middle income countries show poor alignment with current recommendations ^(17; 18; 19; 20; 21).

At the base of unhealthy dietary and PA behaviours are socio-cultural and environmental determinants which are not only intertwined conceptually but also vary with context ⁽²²⁾. In SSA, these determinants are largely shaped by women ^(23; 24; 25; 26). Hence, understanding factors influencing and informing dietary and PA decisions in women is vital. A comprehensive and systematically synthesized overview of determinants of dietary and PA behaviours of WRA is lacking in urban SSA. In this study, a systematic review was conducted to synthesis and collate determinants of dietary and PA behaviour among WRA in urban SSA according to the socio- ecological framework.

Methodology

Systematic searches of literature were conducted according to a predefined protocol published on PROSPERO, ID: CRD42018108532 ⁽²⁷⁾. The PRISMA was followed as a guideline to conduct and report this systematic review ⁽²⁸⁾.

Search strategy

Electronic databases of MEDLINE, EMBASE, Scopus and Web of Science were searched. Additional eligible studies were identified from the bibliographies of included articles (snowballing). The initial search string (**table 2**) was developed in EMBASE by using the Population, Intervention, Comparison and Outcome model, combining population (WRA (18 to 49 years) living in urban SSA), intervention (factors influencing dietary and PA behaviours as risk factors for nutrition related NCD), comparison (not applicable), and outcome (determinants of dietary and PA behaviours) terms, **table 1**. The search string was refined for use in the different databases. Searches were conducted from August to September 2018. All identified studies were imported to endnote.

Eligibility criteria

The review was restricted to studies published in English from 2000 to September 2018. This timeframe was chosen because the world is changing at a fast rate and determinants before the turn of the century may no longer be relevant. Observational studies (cross-sectional, cohort, and case-control), qualitative and randomized controlled trial designs were included. Studies with a population aged <18 or >49 years and those in rural SSA were excluded. Studies conducted in population /samples with clinical diagnosis directly related to nutrition related NCD were as well excluded.

Study selection

Two reviewers working independently screened the identified studies according to the inclusion criteria to select relevant ones. Prior to this, duplicates were removed. First, one reviewer (PY) screened the titles. Abstracts of selected studies were then reviewed independently by PY and CM. A full text was read in cases of insufficient information in the title and abstract. The full text of the selected studies was then retrieved and read to determine whether they met the inclusion criteria by one reviewer (PY). Excluded full-text studies were listed with reasons for their exclusion. See PRISMA diagram, **figure 1**.

Data extraction

Data extraction; study design, study setting (Countries and City), sample size (number of participants), study information (title, author(s), journal, year of publication), and characteristics of participants (type of community, age, sex and social economic status) was done by a single reviewer. Data extraction for determinants of PA and dietary behaviour (components of specific dietary and PA behaviours studied, how they were measured and

outcomes) was initially accomplished by one reviewer (PY). Extracted data was then scrutinized by a second reviewer (CM). Data extraction sheets were pilot tested on 5 papers before applying them to all included studies. Extracted data is presented in **supplementary Tables 1 & 2**. Identified studies could not allow quantifying the magnitude of identified determinants⁽²⁹⁾. Thus, data synthesis was narrative. To structure the narrative synthesis, thematic analysis approach was used⁽³⁰⁾. Thematic analysis involved identifying key concepts from studies and organizing them into one overall data scheme. The identified key concepts were categorized into themes by constant comparative analysis. Comparative analysis involved taking concepts from one study and recognising the same concepts in other studies. Qualitative research depicts the scope and variation in views, and the uncommon views in a few studies might represent important features⁽³¹⁾. Therefore, findings mentioned by a few studies were considered relevant in this review. Themes were structured according to the socio ecological model. The results from all identified studies were summarized irrespective of methodological quality.

Quality assessment

Standardized quality assessment tools were used to assess the quality of the studies. The ‘standard for reporting of reporting qualitative research tool’ was used to assess qualitative studies⁽³²⁾ while the National Heart, Lung, and Blood Institute (NHLBI) quality assessment tool for observational cohort and cross sectional studies was used for cross sectional studies⁽³³⁾. To arrive at quality judgement, we critically appraised how methodological fatal flaws contributed to overall risk of bias.

Results

Description of included studies

The systematic search yielded 9853 studies after duplicate removal. From these studies, 93 articles (47 for dietary behaviour and 46 for PA) were screened for full text review. From the articles, 23 studies (13 investigated dietary behaviour only, 7 PA, while 3 investigated both behaviours) were included in quality assessment and analysis (**figure 1**). The characteristics of studies included in analysis are presented in **supplementary Tables 1 & 2**. Searches identified only qualitative studies and descriptive cross-sectional studies. A total of 11 studies used a descriptive cross-sectional design, nine used qualitative while three used a mixed design. In terms of study setting, the majority were from South Africa and West Africa. Of

the 16 studies on dietary behaviour, six were from South Africa^(23; 24; 34; 35; 36; 37), four from Ghana^(19; 38; 39; 40), and two from Burkina Faso^(17; 41). Four studies were from Kenya⁽²⁶⁾, Tanzania⁽²¹⁾, Senegal⁽²⁵⁾ and Cameroon⁽²⁰⁾. Studies on PA were from South Africa^(23; 37; 42; 43; 44; 45; 46), Nigeria^(47; 48) and Cameroon⁽²⁰⁾.

Quality appraisal

Overall, 4/11 qualitative studies were rated as good quality, four were of poor quality while three were of moderate quality, **supplemental table 3**. Strength was well elaborated data analysis and data collection methods. The main weaknesses were lack of specification of qualitative approach and research paradigm used as well as data management techniques. Majority of studies did not use a theoretical framework as well. 7/12 descriptive cross-sectional studies were of moderate/poor methodological quality **supplemental table 4**. The main limitations were insufficient description of the study population and sample size. These methodological gaps signify a potential bias and it is important to interpret study findings in while considering the potential bias.

Data synthesis

The determinants identified for both behaviours included factors that cut across all the five levels of the socio-ecological framework⁽⁴⁹⁾ (**figures 2 & 3**). Dietary behaviours studied included; food purchasing⁽⁴⁰⁾, food choice^(24; 26; 34; 37; 39), healthy eating^(20; 23; 25; 38), food environment characteristics⁽¹⁹⁾, nutrition knowledge^(35; 36) and dietary diversity and pattern^(17; 21; 41). Components of PA studied included; PA pattern^(44; 45; 48), PA beliefs^(23; 37), PA knowledge and socio-cultural barriers^(20; 46) and experiences of PA during pregnancy and postpartum^(42; 43; 47). Qualitative studies uncovered more of knowledge, attitude and socio-cultural mediators while cross sectional studies, reported on majorly socio-economic determinants.

Determinants of dietary behaviour

Table 2 below gives an overview of determinants of dietary behaviours among WRA in urban SSA

Factors at intra-personal level

Food and Nutrition Knowledge

Nine studies reported food and nutrition knowledge to have an influence on dietary behaviours^(17; 20; 23; 24; 25; 35; 36; 38; 39). Several misconceptions about healthy eating were

reported in qualitative terms. Healthy eating was interpreted in terms of moderate intake of fat with unclear understanding of the benefits of fruits and vegetables, and diet diversification. Dietary diversity was understood from perspective of consuming different foods rather than from concept of food group diversity⁽³⁸⁾. Knowledge of the role of healthy food to NCD prevention was scanty^(25; 36). There was a link between fat and NCD but salt and sugar linkage along with preventive role of fruits and vegetables was not understood. The role of food and nutrition knowledge on dietary behaviours was inconclusive^(20; 23; 39). A study among young adults in South Africa reported that despite having a good understanding of healthy eating it was not a priority when making dietary decisions⁽²³⁾. In contrast, a study among Ghanaian street food consumers found that among the reasons people choose street foods was to compose a healthy diet from the available diversity of foods⁽³⁹⁾.

Food attitude

Ten studies reported attitude to be a determinant of dietary behaviours^(20; 21; 23; 24; 26; 34; 36; 37; 38; 39). All studies reported a positive attitude towards energy dense diets that are rich in sugar, salt or fat (unhealthy). Unhealthy foods were perceived to have a better taste and to require minimal or no time for preparation^(20; 23; 26; 34; 36; 37; 38; 39). Healthy foods, including whole grains, fruits and vegetables, were not preferred by consumers. The high cost of food, long time involved in food preparation and undesirable sensory attributes (tasteless, dry and insipid), explained the negative attitude towards healthy foods^(20; 24; 26; 34; 36; 38). Perceived health concerns also influenced dietary behaviours^(23; 24; 34; 36; 38). Health concerns (adverse health risks to an individual or loved ones) were discussed as elicitors of a positive attitude towards healthy foods^(23; 24; 34; 36; 38). However, this was discussed to influence older adults. Young adults perceived that health concerns only manifest in old age.

Psychological and physiological factors

Five studies found emotional satisfaction to be a determinant for consumption of foods rich in fat, salt and sugar^(23; 24; 34; 37; 38). Fruits and vegetables were also perceived to be non-satiating^(24; 38).

Time and financial barriers

Financial difficulties and lack of time were noted as major barriers influencing dietary behaviours^(20; 21; 24; 26; 34; 36; 37; 38; 39). Financial difficulties were mentioned as determinants of low fruit and vegetable intake^(20; 21; 36; 38). Studies on consumption of street foods in Kenya, Ghana and South Africa found that ability to get food on credit and savings on fuel when food preparation is avoided partly explains the increased consumption of these foods^(26; 37; 39).

In South Africa and Cameroon participants perceived that a balanced diet was hard to sustain due to budget constraints.

Food skills and self-efficacy

Dietary behaviours were discussed to be influenced by food preparation skills and self-efficacy^(20; 24; 26; 36; 37; 38). Participants in a qualitative study in Ghana reported that the ability to improvise with available resources to prepare healthy tasty meals informs their dietary decisions⁽³⁸⁾. Everett-Murphy *et al.*⁽²⁴⁾ in South Africa found that many individuals lack the skills and confidence to prepare healthy tasty meals. Participants reported a difficulty in breaking unhealthy entrenched dietary habits⁽²⁴⁾. The difficulty was mediated by rejection of healthy foods by family especially when perceived to be un-tasty.

Food habits (habit strength)

Five studies reported food habits to have an influence on dietary behaviours^(17; 20; 24; 35; 39). High consumption of carbohydrate rich foods and low consumption of fruits and vegetables is habitually ingrained^(17; 20; 35; 39). In South Africa, sweet treats are social rituals on weekends while deep frying are entrenched habits nurtured from childhood⁽²⁴⁾.

Age

Four studies reported age to have an influence on dietary behaviours^(17; 20; 23; 41). In Cameroon, young adults (18 to 40 years) regarded fruits and vegetables to be for the older adults (≥ 40 years)⁽²⁰⁾. Younger Burkinabe adults compared to older adults have a snacking behaviour with mainly energy dense foods that are nutrient poor^(17; 41).

Socio economic status

Seven studies reported on influence of socio-economic status indices; income/economic, education and occupation on dietary behaviours^(17; 21; 26; 34; 39; 40; 41). Findings are mixed on the direction of influence. A low-income status was found to be a predictor for high consumption of unhealthy street foods in Ghana, Kenya and South Africa^(26; 34; 39). In contrast, a high income status in Tanzania was a predictor for consumption of unhealthy foods⁽²¹⁾. Three studies found higher education level to be associated with higher dietary diversity and increased likelihood to choose healthy food offers^(17; 34; 41). Women working in informal sector (street and market vendors) were reported to depend on majorly street foods⁽³⁹⁾. In urban Burkina Faso, working in the formal sector (professional/ corporate jobs) was associated with higher snacking habits⁽¹⁷⁾.

Factors at interpersonal level**Household composition**

Household size and composition, according to three studies, had an influence on dietary behaviours^(17; 26; 38). The presence of a housewife was associated with decreased dependence on out of home eating⁽²⁶⁾. A large household size was associated with increased prioritization of satiety giving carbohydrate rich foods over other food groups^(26; 38).

Family and peer influence

Six studies found dietary decisions to be mediated by the family and friends^(23; 24; 34; 36; 37; 38). Dietary patterns are adopted from largely family networks^(20; 23; 24; 38). At the same time, the preferences of family members are a vital consideration in dietary choices^(24; 36). Two qualitative studies in South Africa and Ghana found that due to peer pressure young adults (≤ 35 years) eat more energy dense street/fast foods^(34; 38). Community groups were also reported as determinants of dietary behaviours.

Factors at community level**Cultural beliefs**

Several cultural beliefs influenced dietary behaviours^(17; 20; 23; 24; 35; 37; 38; 39). Three studies reported perception of body image to have an influence on food intake^(20; 23; 37). Overweight is associated with beauty, dignity, health, wealth and good treatment by husband while weight loss is a source of stigma and a sign of HIV/AIDS^(20; 23; 37). A positive weight gain perception is a motivation for unhealthy dietary behaviours. Other reported cultural beliefs included low social status accorded to fruits and vegetables, legumes and minimally refined cereals, and high social-status accorded to unhealthy fast foods and eating out^(24; 37; 38; 39).

Physical environment

Eleven studies reported food choices to be influenced by ubiquitous cheap high energy dense nutrient poor foods and food deserts^(17; 19; 20; 21; 24; 26; 34; 37; 38; 39; 40). An abundance of ready to eat street food offers was reported^(19; 24; 26; 37; 39). These street offers are limited in fruits and vegetables but provide mostly refined carbohydrate rich foods. Food purchases are from mainly three retail sources; traditional markets including farmers' weekly market days, super markets and convenience retail stores that offer mainly processed food^(19; 40). Studies reported that majority of the population still prefer traditional markets because they offer low-priced fresh food. It was also found that proximity to food retail stores determines purchasing behaviour⁽⁴⁰⁾. Participants in these studies reported a high dependence on convenience stores due to their ubiquity in the communities^(19; 38; 40). Some participants

reported weekly food purchasing behaviour to counter accessibility difficulties to traditional markets⁽⁴⁰⁾.

Virtual/information environment

Three studies reported virtual environment (media) to influence food decisions^(23; 34; 36). A strong trust in credibility of nutrition information shared from main stream media (TV and radio) was reported^(34; 36).

Factors at institutional level

Religion and food deserts within workplaces influenced dietary choices^(17; 36; 41). Dietary advice shared through religious institutions was reported to highly influence dietary decisions.

Figure 2 below gives a schematic overview of determinants of dietary behaviour among WRA in Urban SSA arranged according to the socio ecological framework.

Determinants of PA behaviour

Table 3 below gives an overview of determinants of PA behaviour among WRA in Urban SSA

Determinants at intra-personal level

Awareness

Seven studies reported key determinants of PA as being knowledge/awareness of PA recommendations, its health benefits and how one might engage in PA.^(20; 23; 37; 42; 43; 46; 47). Researchers found a high proportion of participants un-aware of the health benefits of PA^(20; 37; 47). Health benefits of PA were related to mainly body cleansing^(20; 43). Knowledge on the role of PA in NCD prevention was lacking. Adeniyi *et al.*⁽⁴⁷⁾ found that an estimated 61.0% of Nigerian Women were not aware of PA benefits towards enhancing postpartum health. Studying PA in pregnancy, Muzigaba *et al.*⁽⁴²⁾ reported that women were not aware of PA recommendations.

Intra individual barriers

Barriers included lack of time, feeling tired (laziness), preference for sedentary activities (TV viewing) and financial constraints (money to buy bike, to pay at gyms)^(20; 23; 42; 43; 45; 47; 48). Long working hours and need to prioritize social obligations (family needs) were reported to be a major basis for insufficient time^(42; 43; 47). Physiological changes (increased stomach

size) during pregnancy and psychological fear that increased PA may hurt the baby or cause miscarriage were reported to be barriers to PA ^(42; 43).

PA attitude

Perceived severity, benefits and positive experience were reported to have an influence on PA attitude ^(20; 23; 42; 43; 46; 47). Perceived PA benefits included positive pregnancy outcomes, body freshening, keeping in shape and feel-good mood ^(20; 23; 42; 43). Likewise, health concerns (adverse health risks to an individual or loved one) were discussed to increase likelihood to engage in PA ^(20; 23; 42; 46). Positive experience, for example blood pressure normalization resulting from a past PA program was reported as a motivator to continue with PA ⁽²³⁾. The PA perceived to be simple and socially engaging was described to contribute to a positive attitude ^(42; 47).

PA self-efficacy

Two studies reported self-efficacy to be a determinant of PA behaviour ^(42; 47). Several factors were described to determine PA self-efficacy. Social support, having a PA buddy or belonging to a group increased PA self-efficacy ^(42; 47).

Socio economic status

Four descriptive cross-sectional studies found socioeconomic indices; asset ownership and occupation to be associated with PA behaviours ^(44; 45; 47; 48). A study among Nigerian women working in informal markets reported that of the total 16 hours spent on daily activities, 11.5 hours are spent on predominantly sedentary market vending activities ⁽⁴⁸⁾.

Determinants at Interpersonal Level

Social environment

Eight studies found family, friends and community groups to have an influence on PA ^(20; 23; 37; 42; 43; 44; 46; 47). Direction of social environment influence was inconclusive. Family members were reported to discourage their relatives from PA they don't approve ^(23; 37; 42; 46). Membership in a community health club/group positively influenced PA ^(37; 43) while lack of access to sports teams/community groups had a negative influence ^(23; 42; 43; 47). Community groups were reported to offer practical solutions to overcome PA barriers, emotional support and an opportunity to participate in group PA.

Determinants at community level

Cultural beliefs

Six studies reported cultural misconceptions to be determinants of PA^(20; 23; 37; 43; 46; 47). PA is viewed as unfeminine and in conflict with the perceived weight gain prestige^(20; 23; 37; 46). Walking is considered a sign of poverty and idleness^(20; 37; 43). Two studies reported a gender stereo type that women by tradition should not engage in outdoor related PA^(23; 46). Dress code also had an influence on PA, exercise attires (tight and short) are perceived negatively⁽⁴⁶⁾.

Physical environment

Insufficient infrastructure (playgrounds, parks, pavements for pedestrians and safe roads for cyclists) and safety concerns (violence against women) were the major impediments to outdoor PA^(20; 37; 42; 45). Three studies reported cheap motorized transportation (motor bikes) to influence PA^(20; 37; 45).

Figure 3 below gives a schematic overview of determinants of physical activity among WRA in Urban SSA arranged according to the socio ecological framework.

Discussion

The review identified 23 studies from settings of majorly South and West Africa. In the search, we did not find any randomized control trial, quasi-experiment or observational study (e.g. cohort and case-control). A similar finding was reported by Gissing SC and colleagues⁽⁵⁰⁾ in their mapping review of drivers of dietary behaviour in urban Africa. Across studies, the heterogeneity in methodologies does not allow quantification of determinant's effect on dietary and PA behaviour. Within the socio-ecological framework, included studies reported factors on mainly intra/interpersonal and community level. Few studies reported factors on the community, institutional and public policy level.

Dietary behaviour

Notable determinants of dietary behaviour at intra-individual level were food knowledge, attitude, skills, self-efficacy and psychological factors. In total, 9 of 16 identified studies reported healthy eating to be interpreted in terms of mainly fat intake moderation with unclear understanding of the role of sugar, salt, fruits and vegetables, and diet diversification^(17; 20; 23; 24; 25; 35; 36; 38; 39). Majority of countries in SSA lack contextualized food based dietary guidelines while those in place are poorly communicated^(51; 52). This partly explains the

misconceptions about healthy eating. Research from Northern settings show that food and nutrition knowledge may inform decisions about food intake and differentiating between healthy and unhealthy foods ^(53; 54; 55). Thus, the misconceptions about healthy eating in urban SSA could unknowingly be translating into unhealthy dietary behaviour.

Emotional and satiety satisfaction are important determinants of food choices. Sensory attributes instigate emotional needs ^(23; 24; 34; 37; 38). Due to financial difficulties, hunger satisfaction is a major consideration during food decisions in urban SSA. These findings reinforce recommendations by Perry EA ⁽⁵⁶⁾ and Vidgen HA ⁽⁵³⁾ that healthy food decisions need not only consider nutrition but also gastronomic needs and available resources (time and money). This signifies a need for knowledge, skills and self-efficacy to plan, manage and deliver healthy but feasible food decisions. In particular, skill to plan healthy food choices on time and budget is paramount to address the financial and time insufficiency barriers. In Northern settings ⁽⁵⁷⁾, in urban SSA, female labour market opportunities have increased and does not only limit available time for household domestic roles but also cooking skills ⁽²¹⁾.

Beyond intra personal levels; family, peer and community groups influence are notable determinants at interpersonal level ^(23; 24; 34; 36; 37; 38). At community level; cultural beliefs, physical and virtual environments are important determinants ^(17; 19; 20; 21; 23; 24; 35; 37; 38; 39; 40). In line with findings from high income countries, unhealthy food choices are linked to endorsement from peers while family network influences largely cooking methods and dietary patterns ^(58; 59; 60; 61). A high regard for community health groups was found. Community health groups offer emotional and valuable support necessary to overcome barriers to unhealthy dietary behaviour ⁽³⁷⁾. Several healthy eating recommendations now recognize the importance of social support in shaping dietary behaviour ^(53; 56).

Culture informs food beliefs while communities align these beliefs with healthy dietary recommendations ^(56; 60). The prevailing food beliefs within urban SSA promote dietary behaviours that deviates from healthy recommendations. Next to food beliefs, cultural norms perpetuate gender constructs which variably influence dietary behaviour. These norms include culturally constructed gender roles and overweight prestige. Overweight among women is perceived as a sign of good treatment by the husband. Because of prestige ensuing from weight gain, husbands are inclined to endorse behaviours that promote overweight ⁽⁵⁰⁾. In line with findings from other low- and middle-income settings, food shopping and cooking

are customarily a woman's role ^(38; 62). These culturally defined gender roles have implications in dietary behaviours for both men and women. Firstly, it restricts men's food preparation skills making them dependent on women. Meanwhile, female labour markets across urban SSA are on the rise which ultimately affects their involvement in household roles like food preparation. Our findings demonstrate increased out of home eating among members in households without housewives ^(21; 26). Studies from other settings report unhealthy dietary practices among men compared to women ^(62; 63). It is therefore important to understand gender roles for effective behaviour change within the households.

Eleven studies reported food choices in this setting to be influenced by food deserts and ubiquitous availability of cheap high energy dense nutrient poor foods ^(17; 19; 20; 21; 24; 26; 34; 37; 38; 39; 40). Over the last decade, supermarkets, street foods and fast food chains with less healthy offers have increased within urban Africa ^(19; 39; 40). Between 1992 and 2014, per capita calorie availability grew by over 10% in middle and low income countries ⁽⁶⁴⁾. Topical studies in urban SSA have found an association between supermarket purchases and unhealthy weight gain ^(65; 66; 67). Present review has found a high regard of information shared through religious institutions. Use of religious institutions as vehicles of health promotion has shown potential among black American communities ⁽⁶⁸⁾.

PA behaviour

Awareness, attitude and self-efficacy were identified as important determinants of PA at intra-personal level. The awareness of the benefits and different types of PA is still low in SSA. Accordingly, awareness is an important determinant of PA in SSA setting compared to developed countries. Majority of SSA countries lack action-based PA guidelines. Studies also report inadequate capacity of health personnel ^(43; 51). For example, in South Africa only 19% of health workers provide PA related information during antenatal visits ⁽⁶⁹⁾.

The review found lack of time to be a major barrier to PA ^(20; 23; 42; 43; 47; 48). Culturally in SSA setting, women oversee domestic affairs of the home ⁽⁴⁶⁾. This coupled with the improving female labour market opportunities and longer commuting hours (traffic jam and distinct separation of home and work location) greatly diminishes available time for structural related PA.

Several cultural misconceptions relating to PA were found. The current review found a preference for PA perceived to be simple; like walking. Presently, over 75% of PA in urban SSA is contributed by travel related walking ⁽¹⁵⁾. However, participation in walking is threatened by cultural misconceptions which depict walking as a sign of poverty and idleness. With increasing evolution of cheap motorized transportation within urban SSA ^(16; 20; 37; 45), negative perception towards walking may decrease PA. PA is viewed as unfeminine since it affects glossy attractiveness of women due to excess body fat and their traditional domestic roles ^(20; 23; 37; 46). Traditionally, women are socialized into gender stereotypical roles by their mothers. Automation and technological advancements are gradually replacing previously active domestic chores ^(16; 44; 46). Time spent on passive forms of recreation (e.g. television and social media) is increasing ⁽⁴⁴⁾. The growing trend in sedentary life require innovative approach to address gender stereotypes for women being restricted to indoor activities. Initiating a culture of PA opportunities in post high school period (e.g. adult amateur netball community clubs) could be valuable ⁽²³⁾. The adjustment of the physical education curriculum could as well prepare communities to accept women participation in exercise related PA ^(15; 63).

Social environment, particularly community health groups and PA role models positively influence PA ^(23; 37; 42; 43; 47). Membership in a community health group may help increase awareness of importance of PA ⁽⁴⁶⁾. These groups may as well offer practical solutions to PA barriers and misconceptions through experience sharing. According to sport and social sciences people socialize into PA ^(70; 71). Central agents of socialization are the community groups. Potential of group activities to increase PA among adults has been reported by Seghers *et al.* ⁽⁷²⁾ in Belgium. Socialization through community groups could address cultural gender stereotypes around PA in urban SSA.

Physical environment was an important PA determinant in this setting. Built environmental variables like public and private recreational facilities, paved sidewalks and neighbourhood aesthetics enhance recreational and travel related PA ^(73; 74; 75). Safety concerns related to increasing crime rates in developing countries have been shown to negatively influence PA in developing settings ^(76; 77). Feeling unsafe may diminish confidence in the ability to be physically active or willingness to go outdoors ⁽⁷⁸⁾. SSA is rapidly undergoing urbanization. This urbanization is largely un-organized and has brought along with it several undesirable effects; chief among them being youth unemployment, a catalyst for crime rates.

Limitations

The review identified mainly studies from South and West Africa. SSA is a very diverse community, thus, generalization of the current findings to East and Central African parts of SSA may be challenging. Restriction of the review to English publications could also have limited the scope leaving out publications in French yet countries of West and Central SSA are predominantly French speaking. Additionally, this review identified majorly cross-sectional studies making causality deduction difficult. Furthermore, a bigger proportion of the qualitative studies identified did not include theoretical framework in their design. Across studies, there was considerable heterogeneity in study designs, assessed indicators, measurement and behaviour components assessed, making comparability of indicators difficulty.

Conclusion

This review has proposed socio-ecological frameworks for potential determinants of dietary and PA behaviour among WRA in urban SSA. It highlights salient potential determinants; importance of traditional weekly farmers' markets, food deserts within work environments, cultural misconceptions (gender stereotypes), virtual environment (media), sedentary home/work environments and cheap motorized transport (motorcycles). However, evidence is from mainly cross-sectional studies and on intra and inter-personal levels. Research into determinants of dietary and PA behaviour in context of urban SSA is minimal, particularly on the community, institutional and public policy level. Studies with rigorous qualitative and quantitative designs are required to validate and develop the proposed frameworks further especially within East Africa. The cross-sectional nature of available evidence makes policy recommendation challenging. Nevertheless, available insights suggest a need for comprehensive skill-based interventions focusing on socio-cultural misconceptions and financial limitations.

Conflicts of interest

None declared

Acknowledgements

Financial support; the work was funded by the Belgian Directorate General for Development Cooperation and Humanitarian Aid (DGD), an initiative implemented by VLIR-UOS. We acknowledge Constance Herrebout who identified studies from Scopus database and Merel Leyman, Anke Vandycke, Sigrid Devloo and Diane Berardis for their input in quality assessment.

Authors contribution

CM and PY were responsible for the design of the study and development of search string. PY carried out the search and performed the initial screening. CM and PY independently reviewed abstracts of selected studies. PY extracted the data and CM helped with interpretation of the data. PY drafted the initial manuscript. All authors contributed to interpretation of results and critical review of the manuscript and have approved the final manuscript.

References

1. Gakidou E, Afshin A, Abajobir AA *et al.* (2017) Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet* **390**, 1345–1422.
2. Haddad L, Hawkes C, Udomkesmalee E *et al.* (2016) Global Nutrition Report 2016: From Promise to Impact: Ending Malnutrition by 2030. *Washington, DC: International Food Policy Research Institute.*
3. Yach D, Hawkes C, Gould CL *et al.* (2004) The global burden of chronic diseases: overcoming impediments to prevention and control. *Jama* **291**, 2616–2622.
4. Mathers CD, Loncar D (2006) Projections of global mortality and burden of disease from 2002 to 2030. *PLoS medicine* **3**, e442.
5. Ministry of Health Uganda (2015) Health sector development plan. Kampala.
6. Geneau R SD, Stachenko S, McKee M, Ebrahim S, Basu S, Chockalingham A, Mwatsama M, Jamal R, Alwan A, *et al.* (2010) Raising the priority of preventing chronic diseases: a political process. *Lancet* **376**, 1689–1698.
7. Hawkes C, Fanzo J (2017) Nourishing the SDGs: Global Nutrition Report 2017.
8. Ministry of Health Uganda, World Diabetes Foundation, World Health Organisation, United Nations Development Programme (2014) Non-Communicable Disease Risk Factor Baseline Survey Report.
9. Ministry of Health, Gender, Elderly and Children, Ministry of Health, National Bureau of Statistics, Office of the Chief Government Statistician (2016) *Tanzania Demographic and Health Survey and Malaria Indicator Survey (TDHS-MIS) 2015-16*
10. Cois A, Day C (2015) Obesity trends and risk factors in the South African adult population. *BMC obesity* **2**, 42.
11. Hanson M, Barker M, Dodd JM *et al.* (2017) Interventions to prevent maternal obesity before conception, during pregnancy, and post partum. *The Lancet Diabetes & Endocrinology* **5**, 65–76.
12. Godfrey KM, Reynolds RM, Prescott SL *et al.* (2017) Influence of maternal obesity on the long-term health of offspring. *The lancet Diabetes & endocrinology* **5**, 53–64.
13. Alwan A (2011) *Global status report on noncommunicable diseases 2010*: World Health Organization.

14. Cecchini M, Sassi F, Lauer JA *et al.* (2010) Tackling of unhealthy diets, physical inactivity, and obesity: health effects and cost-effectiveness. *The Lancet* **376**, 1775-1784.
15. Guthold R, Stevens GA, Riley LM *et al.* (2018) Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1· 9 million participants. *The Lancet Global Health* **6**, e1077-e1086.
16. World Health Organisation (2018) Global action plan on physical activity 2018–2030: more active people for a healthier world. . Geneva: World Health Organization.
17. Becquey E, Savy M, Danel P *et al.* (2010) Dietary patterns of adults living in Ouagadougou and their association with overweight. *Nutrition journal* **9**, 13.
18. Frank LK, Kröger J, Schulze MB *et al.* (2014) Dietary patterns in urban Ghana and risk of type 2 diabetes. *Br J Nutr* **112**, 89-98.
19. Dake FA, Thompson AL, Ng SW *et al.* (2016) The local food environment and body mass index among the urban poor in Accra, Ghana. *Journal of Urban Health* **93**, 438-455.
20. Kiawi E, Edwards R, Shu J *et al.* (2006) Knowledge, attitudes, and behavior relating to diabetes and its main risk factors among urban residents in Cameroon: A qualitative survey. *Ethnicity & Disease* **16**, 503-509.
21. Cockx L, Colen L, De Weerd J (2018) From corn to popcorn? Urbanization and dietary change: Evidence from rural-urban migrants in Tanzania. *World Development* **110**, 140-159.
22. Eldredge LKB, Markham CM, Ruiter RA *et al.* (2016) *Planning health promotion programs: an intervention mapping approach*: John Wiley & Sons.
23. Phillips EA, Comeau DL, Pisa PT *et al.* (2016) Perceptions of diet, physical activity, and obesity-related health among black daughter-mother pairs in Soweto, South Africa: A qualitative study. *BMC Public Health* **16**.
24. Everett-Murphy K, De Villiers A, Ketterer E *et al.* (2015) Using formative research to develop a nutrition education resource aimed at assisting low-income households in South Africa adopt a healthier diet. *Health Educ Res* **30**, 882-896.
25. Holdsworth M, Delpeuch F, Landais E *et al.* (2006) Knowledge of dietary and behaviour-related determinants of non-communicable disease in urban Senegalese women. *Public Health Nutrition* **9**, 975-981.
26. Van 'T Riet H, Den Hartog AP, Mwangi AM *et al.* (2001) The role of street foods in the dietary pattern, of two low-income groups in Nairobi. *Eur J Clin Nutr* **55**, 562-570.
27. Peter Yiga, Christophe Matthys, Patrick Ogwok. (2018) Mediators of dietary and physical activity behaviors among women of reproductive age in Urban Sub Saharan Africa - a systematic review. . PROSPERO CRD42018108532.

28. Moher D LA, Tetzlaff J, Altman DG. (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. . *PLoS Med* 2009 **6**, e1000097.
29. Bauman AE, Sallis JF, Dzewaltowski DA *et al.* (2002) Toward a better understanding of the influences on physical activity: the role of determinants, correlates, causal variables, mediators, moderators, and confounders. *American journal of preventive medicine* **23**, 5-14.
30. Thomas J, Harden A (2008) Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC medical research methodology* **8**, 45.
31. Fade SA (2003) Communicating and judging the quality of qualitative research: the need for a new language. *Journal of Human Nutrition and Dietetics* **16**, 139-149.
32. O'brien BC, Harris IB, Beckman TJ *et al.* (2014) Standards for reporting qualitative research: a synthesis of recommendations. *Academic Medicine* **89**, 1245-1251.
33. Bethesda M (2014) National Heart, Lung, and Blood Institute (NHLBI) Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies. Available at: <https://www.nhlbi.nih.gov/health-pro/guidelines/in-develop/cardiovascular-risk-reduction/tools/cohortNational>.
34. Van Zyl MK, Steyn NP, Marais ML (2010) Characteristics and factors influencing fast food intake of young adult consumers in Johannesburg, South Africa. *S Afr J Clin Nutr* **23**, 124-130.
35. Van Den Berg VL, Okeyo AP, Dannhauser A *et al.* (2012) Body weight, eating practices and nutritional knowledge amongst university nursing students, Eastern Cape, South Africa. *Afr J Prim Health Care Fam Med* **4**.
36. Charlton KE, Brewitt P, Bourne LT (2004) Sources and credibility of nutrition information among black urban South African women, with a focus on messages related to obesity. *Public health nutrition* **7**, 801-811.
37. Stern R, Puoane T, Tsolekile L (2010) An exploration into the determinants of noncommunicable diseases among rural-to-urban migrants in periurban South Africa. *Prev Chronic Dis* **7**.
38. Boatemaa S, Badasu DM, De-Graft Aikins A (2018) Food beliefs and practices in urban poor communities in Accra: Implications for health interventions. *BMC Public Health* **18**.
39. Hiamey SE, Amuquandoh FE, Boison GA (2013) Are we indeed what we eat? Street food consumption in the Market Circle area of Takoradi, Ghana. *Nutr Health* **22**, 215-235.
40. Aryeetey R, Oltmans S, Owusu F (2016) Food retail assessment and family food purchase behavior in Ashongman Estates, Ghana. *Afr J Food Agric Nutr Dev* **16**, 11386-11403.

41. Savy M, Martin-Prével Y, Danel P *et al.* (2008) Are dietary diversity scores related to the socio-economic and anthropometric status of women living in an urban area in Burkina Faso? *Public Health Nutrition* **11**, 132-141.
42. Muzigaba M, Kolbe-Alexander TL, Wong F (2014) The perceived role and influencers of physical activity among pregnant women from low socioeconomic status communities in South Africa. *Journal of physical activity & health* **11**, 1276-1283.
43. Watson ED, Norris SA, Draper CE *et al.* (2016) "Just because you're pregnant, doesn't mean you're sick!" A qualitative study of beliefs regarding physical activity in black South African women. *BMC Pregnancy Childbirth* **16**.
44. Gradidge PJJ, Crowther NJ, Chirwa ED *et al.* (2014) Patterns, levels and correlates of self-reported physical activity in urban black Soweto women. *BMC Public Health* **14**.
45. Hattingh Z, Le Roux M, Nel M *et al.* (2014) Assessment of the physical activity, body mass index and energy intake of HIV-uninfected and HIV-infected women in Mangaung, Free State province. *S Afr Fam Pract* **56**, 196-200.
46. Walter CM, du Rosa R (2011) Socio-cultural barriers to physical activity among black isixhosa speaking professional women in the nelson mandela metropolitan municipality. *S Afr J Res Sport Phys Educ Recreat* **33**, 143-155.
47. Adeniyi AF, Ogwumike OO, Bamikefa TR (2013) Postpartum exercise among nigerian women: Issues relating to exercise performance and self-efficacy. *ISRN Obstetrics and Gynecology* **2013**.
48. Afolabi WAO, Addo AA, Sonibare MA (2004) Activity pattern, energy intake and obesity among Nigerian urban market women. *International Journal of Food Sciences and Nutrition* **55**, 85-90.
49. Glanz K, Bishop DB (2010) The role of behavioral science theory in development and implementation of public health interventions. *Annual review of public health* **31**, 399-418.
50. Gissing SC, Pradeilles R, Osei-Kwasi HA *et al.* (2017) Drivers of dietary behaviours in women living in urban Africa: a systematic mapping review. *Public health nutrition* **20**, 2104-2113.
51. Lachat C, Otchere S, Roberfroid D *et al.* (2013) Diet and physical activity for the prevention of noncommunicable diseases in low-and middle-income countries: a systematic policy review. *PLoS medicine* **10**, e1001465.
52. Parker W-a, Steyn NP, Levitt NS *et al.* (2012) Health promotion services for patients having non-communicable diseases: Feedback from patients and health care providers in Cape Town, South Africa. *BMC Public Health* **12**, 503.

53. Vidgen HA GD (2014) Defining food literacy and its components. . *Appetite* **76**, 50-59.
54. Vaitkeviciute R BLHN (2015) The relationship between food literacy and dietary intake in adolescents: a systematic review. . *Public health nutrition* **18**, 649–658.
55. Scripa I (2012) Development and implementation of nutrition education classes and cooking sessions into an existing life skills program for young mothers at the YWCA of Greensboro, NC. The University of North Carolina at Greensboro.
56. Perry EA, Samra HR, Edmonstone S, Davidson L, Faulkner A, Petermann L, Manafò E, Kirkpatrick SI. (2017) Identifying attributes of food literacy: A scoping review. . *Public health nutrition* **20**, 24062415.
57. Pettinger C HM, Gerber M (2006) Meal patterns and cooking practices in Southern France and Central England. . *Public health nutrition* **9**, 1020– 1026.
58. Sleddens EF, Kroeze W, Kohl LF *et al.* (2015) Determinants of dietary behavior among youth: an umbrella review. *International Journal of Behavioral Nutrition and Physical Activity* **12**, 7.
59. Story M, Kaphingst KM, Robinson-O'Brien R *et al.* (2008) Creating healthy food and eating environments: policy and environmental approaches. *Annu Rev Public Health* **29**, 253-272.
60. Larson N, Story M (2009) A review of environmental influences on food choices. *Annals of Behavioral Medicine* **38**, s56-s73.
61. Deliens T, Clarys P, De Bourdeaudhuij I *et al.* (2014) Determinants of eating behaviour in university students: a qualitative study using focus group discussions. *BMC Public Health* **14**, 1-12.
62. Caperon L, Arjyal A, Puja K *et al.* (2019) Developing a socio-ecological model of dietary behaviour for people living with diabetes or high blood glucose levels in urban Nepal: A qualitative investigation. *PloS one* **14**.
63. Spencer RA, Rehman L, Kirk SF (2015) Understanding gender norms, nutrition, and physical activity in adolescent girls: a scoping review. *International Journal of Behavioral Nutrition and Physical Activity* **12**, 6.
64. Steyn NP MZ (2014) Obesity and the nutrition transition in sub-Saharan Africa. *Annals of the New York academy of sciences* **1311**, 88-101.
65. Demmler KM, Klasen S, Nzuma JM *et al.* (2017) Supermarket purchase contributes to nutrition-related non-communicable diseases in urban Kenya. *PloS one* **12**, e0185148.
66. Rischke R, Kimenju SC, Klasen S *et al.* (2015) Supermarkets and food consumption patterns: The case of small towns in Kenya. *Food Policy* **52**, 9-21.

67. Kimenju SC, Rischke R, Klasen S *et al.* (2015) Do supermarkets contribute to the obesity pandemic in developing countries? *Public health nutrition* **18**, 3224-3233.
68. Newton Jr RL, Carter LA, Johnson W *et al.* (2018) A Church-Based Weight Loss Intervention in African American Adults using Text Messages (LEAN Study): Cluster Randomized Controlled Trial. *Journal of medical Internet research* **20**.
69. Watson ED OB, Constantinou D. (2015) Exercise during pregnancy: knowledge and beliefs of medical practitioners in South Africa: a survey study. . *BMC Pregnancy Childbirth* **15**.
70. Moore LLL, White M, Campbell JL, Oliveria SA & Curtis Ellison R. (1991) Influence of parents' physical activity levels on activity levels of young children. . *The Journal of Pediatrics*, 118, 215-219
71. Coakley J (2007) *Sports in Society: Issues and Controversies*. 9th ed. ed. New York: McGraw Hill.
72. Seghers J, Van Hoecke A-S, Schotte A *et al.* (2014) The added value of a brief self-efficacy coaching on the effectiveness of a 12-week physical activity program. *Journal of Physical Activity and Health* **11**, 18-29.
73. Owen N HN, Leslie E, Bauman A, Sallis JF (2004) Understanding environmental influences on walking: review and research agenda. . *Am J Prev Med* **27**, 67–76.
74. Gebel K, Bauman AE, Petticrew M (2007) The physical environment and physical activity: a critical appraisal of review articles. *American journal of preventive medicine* **32**, 361-369. e363.
75. Saelens BE, Handy SL (2008) Built environment correlates of walking: a review. *Medicine and science in sports and exercise* **40**, S550.
76. Parra DC, Gomez LF, Sarmiento OL *et al.* (2010) Perceived and objective neighborhood environment attributes and health related quality of life among the elderly in Bogota, Colombia. *Social Science & Medicine* **70**, 1070-1076.
77. Gomes GA, Reis RS, Parra DC *et al.* (2011) Walking for leisure among adults from three Brazilian cities and its association with perceived environment attributes and personal factors. *International journal of behavioral nutrition and physical activity* **8**, 111.
78. Bennett GG, McNeill LH, Wolin KY *et al.* (2007) Safe to walk? Neighborhood safety and physical activity among public housing residents. *PLoS medicine* **4**, e306.

Table 1: PICO criteria used to define research question.

Participants	Women within reproductive age (18 to 49 years) living in urban Sub Saharan Africa
Intervention	Factors influencing dietary and physical activity behaviours as risk factors for diet related non-communicable diseases. Studies investigating supplementation or fortification will be excluded.
Comparisons	Not applicable
Outcomes	Primary outcome(s) Mediators of dietary and physical activity behaviours (self-efficacy, knowledge, skills, perceived barriers, attitudes and socio-economic factors) Secondary outcome(s) Not applicable
Study design	Observational studies (cross-sectional, cohort, and case-control), qualitative and randomized controlled trial designs were included. Exclusion Criteria <ul style="list-style-type: none"> • Studies with a population aged <18 or >49 years • Studies in rural sub Saharan Africa • Population/samples with clinical diagnosis directly related to non-communicable diseases as aim was to ascertain influencing factors in the general population not those with specific medical conditions

Table 2: Search String developed in Embase

Search strategy – developed in Embase
'africa south of the sahara'/exp AND women AND ('dietary behaviour*' OR ('diet'/exp OR 'food intake'/exp OR 'feeding behaviour'/exp OR 'food choice'/exp OR 'dietary intake'/exp OR 'eating habit'/exp OR 'obesity'/exp))
Embase search strategy for physical activity studies; africa south of the sahara'/exp AND women AND ((('television watching' OR 'television viewing'/exp OR 'television viewing' OR 'video viewing' OR 'tv' OR 'video games'/exp OR 'video games' OR 'electronic game playing' OR 'computer gaming' OR 'computer time' OR 'computer use'/exp OR 'computer use' OR 'media time' OR 'media use' OR 'web browsing' OR 'screen time'/exp OR 'screen time' OR 'bedroom media' OR 'electronic media' OR 'pc' OR 'pc use' OR 'occupational sitting' OR 'deskbound' OR 'reading'/exp OR 'reading' OR 'motor* transport') OR (('physical performance'/exp OR 'physical performance') OR ('physical exercise*' OR (('sport'/exp OR 'physical activity'/exp OR 'leisure time physical activity'/exp OR 'leisure'/exp OR 'recreation'/exp OR 'exercise'/exp OR 'physical inactivity'/exp OR 'lifestyle'/exp OR 'sedentary behaviour'/exp OR 'sedentary time'/exp OR 'daily life activity'/exp) OR 'physical movement*')))))

Table 2: Determinants of dietary behaviour arranged according socio-ecological framework

Level	Mediator (Number of studies)	Study references
Intra-Individual Level	Food & Nutrition Knowledge (n = 9)	(17; 20; 23; 24; 25; 35; 36; 38; 39)
	Food attitude (n=10)	(20; 21; 23; 24; 26; 34; 36; 37; 38; 39)
	Organoleptic properties (n = 8)	(20; 23; 26; 34; 36; 37; 38; 39)
	Perceived convenience (n = 8)	(20; 23; 26; 34; 36; 37; 38; 39)
	Perceived expensiveness (n = 6)	(20; 24; 26; 34; 36; 38)
	Physiological (satiating) (n = 2)	(24; 38)
	Health concerns (n = 5)	(23; 24; 34; 36; 38)
	Barriers (n = 9)	(20; 21; 24; 26; 34; 36; 37; 38; 39)
	Financial difficulties (n = 9)	(20; 21; 24; 26; 34; 36; 37; 38; 39)
	Lack of time (n = 9)	(20; 21; 24; 26; 34; 36; 37; 38; 39)
	Psychological factors (n = 5)	(23; 24; 34; 37; 38)
	Food skills & self-efficacy (n = 6)	(20; 24; 26; 36; 37; 38)
	Food habits (n = 4)	(17; 20; 35; 39)
	Traditionally oriented to carbohydrate rich foods (n = 4)	(17; 20; 35; 38; 39)
	Age (n = 4)	(17; 20; 23; 41)
	Socio economic status (n = 7)	(17; 21; 26; 34; 39; 41)
	Economic status (n = 4)	(21; 26; 34; 39)
	Education (n=3)	(17; 34; 41)
	Occupation (n = 3)	(17; 39; 41)
Interperson al level	Household composition (n = 3)	(17; 26; 38)
	Presence of a woman with domestic role (n = 1)	(26)
	Household size (n = 2)	(26; 38)
	Family and peer influence (n = 6)	(23; 24; 34; 36; 37; 38)
	Preference of family members (n = 4)	(20; 24; 36; 37)
	Dietary patterns/cooking methods adopted from family network (n=4)	(20; 23; 24; 38)
	Peer pressure (n =2)	(34; 38)
	Community groups (n = 2)	(20; 37)
Community	Cultural beliefs (n = 8)	(17; 20; 23; 24; 35; 37; 38; 39)

level	Weight gain prestige (n = 3)	(20; 23; 37)
	High social status accordance fast foods (n = 2)	(37; 39)
	Low social status accordance to fruits & vegetables, legumes & minimally processed grains (n = 2)	(24; 38)
	Physical environment (n = 11)	(17; 19; 20; 21; 24; 26; 34; 37; 38; 39; 40)
	Purchase source (food retail source) (n = 3)	(19; 38; 40)
	Distance to food retail stores (n=3)	(19; 38; 40)
	Food deserts (n = 11)	(17; 19; 20; 21; 24; 26; 34; 37; 38; 39; 40)
	Abundance of ready to eat energy dense street foods (n = 5)	(19; 24; 26; 37; 39)
	Virtual environment (n = 3)	(23; 34; 36)
	Seasonality (n = 1)	(20)
Institutional /organizational level	Religion (n = 3)	(17; 36; 41)
	Workplace (n = 1)	(17)
	Rural to urban migration (n = 2)	(19; 21)

Table 3: Determinants of physical activity behaviour arranged according socio-ecological framework

Level	Determinants (Number of studies)	Study references
Intra-Individual Level	Knowledge/awareness (n = 7)	(20; 23; 37; 42; 43; 46; 47)
	Health importance (n = 5)	(20; 23; 37; 43; 47)
	PA recommendations (n = 1)	(42)
	Understanding what PA entails (n=5)	(20; 23; 42; 43; 46)
	Barriers (n = 7)	(20; 23; 42; 43; 45; 47; 48)
	Lack of time/family obligations (n = 6)	(20; 23; 42; 43; 47; 48)
	Pregnancy related misconceptions (n = 2)	(42; 43)
	Financial constraints (n=4)	(20; 23; 42; 43)
	Preference for sedentary activities (n=2)	(23; 45)
	Attitude (n = 6)	(20; 23; 42; 43; 46; 47)
	Perceived benefits (n = 4)	(20; 23; 42; 43)
	Positive severity (n = 4)	(20; 23; 42; 46)
	Positive experience (n = 1)	(23)
	Simple and social engaging (n = 2)	(42; 47)
	Self-efficacy (n = 2)	(42; 47)
	Socio economic status (n = 4)	(44; 45; 47; 48)
	Economic (asset ownership) (n = 3)	(44; 45; 47)
	Occupation (n = 2)	(47; 48)
Inter - personal level	Social environment (n= 8)	(20; 23; 37; 42; 43; 44; 46; 47)
	Family (n = 4)	(23; 37; 42; 46)
	Friends (1)	(37)
	Significant others (3)	(20; 43; 44)
	Community groups (n = 5)	(23; 37; 42; 43; 47)
Community level	Cultural beliefs (n = 6)	(20; 23; 37; 43; 46; 47)
	Weight gain prestige (n = 4)	(20; 23; 37; 46)
	Negative perception towards walking (n = 3)	(20; 37; 43)
	PA dress code (n = 1)	(46)
	Outdoor PA unfeminine (n = 2)	(23; 46)
	Physical environment (n = 4)	(20; 37; 42; 45)
	Insufficient infrastructure (n = 2)	(20; 42)
	Safety concerns (n = 2)	(37; 42)
Institutional level	Cheap motorized transportation (n = 3)	(20; 37; 45)
	No PA education in health education system (n = 3)	(42; 43; 47)
	Sedentary worksite environment (n = 1)	(37)
	Health information from school (n = 1)	(23)

Figure 1: PRISMA flowchart showing study selection process ^{PA (physical activity)}

Figure 2: Determinants of dietary behaviour arranged according to socio ecological framework

Figure 3: Determinants of physical activity behaviour arranged according to socio ecological framework