3.1 Early pregnancy HbA1c and GDM prediction

Table 2 shows the test characteristics of an early pregnancy HbA_{1c} for detecting GDM using the IADPSG criteria. The baseline HbA_{1c} showed a poor area under the ROC curve (AUC) for identifying women with GDM. An HbA_{1c} threshold of 5.7% (39 mmol/mol) showed low sensitivity (15.9%) but high specificity (89.4%) for GDM at any time during pregnancy. Overall, 51.4% of the women in the HbA_{1c} ≥ 5.7% (39 mmol/mol) group developed GDM, and 72% of these cases were detected before 20 weeks (in the < 5.7% (39 mmol/mol) group, this was 61% (184/302) (*p* = 0.14)). Women with a higher HbA_{1c} in early pregnancy at ≥ 5.7% (39 mmol/mol) had a 1.7 times higher risk for GDM sometime in pregnancy compared to women with an HbA_{1c} of < 5.7% (39 mmol/mol) (adjusted odds ratio (aOR) of 1.72 (1.02–2.89)). The aORs for GDM before 20 weeks, at 24–28 weeks (after excluding early GDM), at 35–37 weeks (after excluding early and 24–28 week GDM) among women in the higher HbA_{1c} group were 2.77 (1.39–5.51), 0.68 (0.26–1.76), and 1.96 (0.84–4.58), respectively. Using fasting glucose of ≥ 6.1 mmol/l (110 mg/dl) in the first trimester gave better ROC characteristics (0.70 (0.53–0.86)) for GDM, but still with poor sensitivity (0% of 3 cases) (**Table 2**).

Table 2							
The table layout displayed in this section is not how it will appear in the final version. The representation below is solely purposed for providing corrections to the table. To preview the actual presentation of the table, please view the Proof.							
Test characteristics of HbA _{1c} at a threshold of ≥ 5.7% (39 mmol/mol) for detecting GDM.							
GDM criteria	Gestation	GDM Prevalence (%)	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)	AUC (CI), <i>p</i> value *
IADPSG	Throughout gestation	41.3 (359/869)	15.9	89.4	51.4	60.2	0.55 (0.51–0.59), 0.01
	Before 20 weeksFirst trimester13–20 weeks	25.9 (225/869)32.1 (50/156)24.5 (175/713)	18.214.019.4	89.186.889.6	36.933.337.8	75.768.177.4	0.55 (0.50–0.59), 0.04
	24–28 weeks	14.3 (75/524) [†] (8.6 of total n)	8.0	88.6	10.5	85.2	0.54 (0.47–0.61), 0.27
	35–37 weeks	14.5 (59/407) [‡] (6.8 of total n)	16.9	89.7	21.7	86.4	0.57 (0.49–0.65), 0.09
Fasting 6.1–6.9 mmol/l (110–125 mg/dl)	First trimester	1.9 (3/156)	0.0	86.3	0.0	97.8	0.70 (0.53–0.86), 0.24
	13–20 weeks	1.0 (7/712)	28.6	87.5	2.2	99.2	0.69 (0.54–0.83), 0.09
The table shows the test characteristics of early pregnancy HbA _{1c} for detecting GDM at a threshold of ≥ 5.7% (39 mmol/mol). *The area under the receiver operating characteristic curve shows the performance of early pregnancy HbA _{1c} for detecting GDM using all possible HbA _{1c} thresholds; [†] after excluding early GDM; [‡] after excluding early and 24–28 week GDM; GDM = Gestational diabetes mellitus; PPV = Positive predictive value; NPV = Negative predictive value; AUC = Area under the curve; CI = Confidence interval, IADPSG = International Association of Diabetes in Pregnancy Study Groups.							

3.2 Early pregnancy HbA_{1c} as a predictor of adverse pregnancy outcomes

There was no significant association between a higher HbA_{1c} ≥ 5.7% (39 mmol/mol) and the risk of adverse pregnancy outcomes (**Table 3**). In a subgroup analysis stratified by GDM status, women with higher HbA_{1c} in the non-GDM group were less likely to have cesarean section and GWG > 5 kg (Supplemental **Table 1**). The ROC analysis showed poor performance of HbA_{1c} for predicting adverse pregnancy outcomes regardless of GDM status (Supplemental **Table 2**).

Table 3

The table layout displayed in this section is not how it will appear in the final version. The representation below is solely purposed for providing corrections to the table. To preview the actual presentation of the table, please view the Proof.

Pregnancy outcomes between HbA_{1c} categories.

Pregnancy outcomes	≥ 5.7% (39 mmol/mol), n = 111 n/N (%)	< 5.7% (39 mmol/mol), n = 758 n/N (%)	<i>p</i> value*	OR (95% CI)	Adjusted OR (95% CI)
Spontaneous abortion	0/88 (0)	2/623 (0.3)	1.00	–	–
Birth weight over 4 kg	15/88 (17.0)	103/628 (16.4)	0.88	1.05 (0.58–1.90)	0.94 (0.46–1.92)
Large for gestational age	12/83 (14.5)	79/566 (14.0)	0.87	1.04 (0.54–2.01)	1.39 (0.64–3.03)
Small for gestational age	3/83 (3.6)	45/568 (7.9)	0.26	0.44 (0.13–1.44)	0.82 (0.22–3.09)
Preeclampsia	3/82 (3.7)	18/554 (3.2)	0.74	1.13 (0.32–3.93)	0.77 (0.15–3.82)
Pregnancy induced hypertension	7/82 (8.5)	69/552 (12.5)	0.37	0.65 (0.29–1.48)	0.44 (0.15–1.27)
Preterm birth	6/83 (7.2)	26/586 (4.4)	0.27	1.68 (0.67–4.21)	1.57 (0.48–5.21)
Cesarean section	29/88 (33.0)	233/622 (37.5)	0.48	0.82 (0.51–1.32)	0.55 (0.30–1.00)
Hyperbilirubinemia	7/81 (8.6)	21/515 (4.1)	0.09	2.23 (0.91–5.42)	1.56 (0.55–4.41)
Neonatal Intensive Care Unit stay	6/76 (7.9)	47/509 (9.2)	0.83	0.84 (0.35–2.04)	0.64 (0.23–1.78)
Weight gain under 5 kg	22/65 (33.8)	121/498 (24.3)	0.10	1.59 (0.92–2.77)	1.35 (0.72–2.52)
Induction of labour	37/82 (45.1)	281/553 (50.8)	0.35	0.80 (0.50–1.27)	0.81 (0.45–1.46)

The table shows ORs (adjusted and unadjusted) for the development of adverse pregnancy outcomes among women with an early pregnancy HbA_{1c} of ≥ 5.7% (39 mmol/mol) compared to those with an HbA_{1c} of < 5.7% (39 mmol/mol). Results were adjusted for maternal age, pre-pregnancy body mass index, booking body mass index, family history of diabetes, country, randomisation group, GDM status and GDM treatment. **p* value of the Fisher's Exact test; –not applicable due to small numbers; GDM = Gestational diabetes mellitus; OR = Odds ratio; CI = Confidence interval.

In the sensitivity analysis, an HbA_{1c} threshold of 5.9% (41 mmol/mol) showed a small increase in specificity with a decrease in sensitivity for detecting GDM (Supplemental **Table 3**). An HbA_{1c} ≥ 5.9% (41 mmol/mol) did not predict adverse pregnancy outcomes (Supplemental Table 4).

4 Discussion

Our study of overweight and obese women from a largely European background shows that while GDM was present in 36.9% of women with an HbA_{1c} ≥ 5.7% (39 mmol/mol) before 20 weeks of gestation, this was also the case in 24.3% of the remaining women, a numerically much larger group. Our results clearly show the poor sensitivity of an early pregnancy HbA_{1c} for detecting GDM. While the 5.7% (39 mmol/mol) cut-off was highly specific for GDM, this threshold could not correctly identify most of the cases of GDM with a false negative rate of 81.8% before 20 weeks and 84.1% for GDM at any time. Low ROC AUC values show the poor ability of the HbA_{1c} to detect GDM at any time regardless of the week of pregnancy. This low sensitivity was also shown by Hughes et al. [12] (18.8% for the threshold of ≥ 5.9% for early GDM) using New Zealand criteria, and Odsæter et al. [20] (0% sensitivity for the threshold of 5.8% (40 mmol/mol) after 18 weeks' gestation) using modified IADPSG criteria. Several other studies also reported poor sensitivity of early HbA_{1c} for diagnosing GDM in late pregnancy, with the value ranging between 0.9% and 25.7% for a higher threshold above ≥ 5.7–5.9% (39–41 mmol/mol) [11,13,20–22]. The AUC values in our results were similar to those reported in other studies with poor values at different pregnancy weeks and throughout gestation [20–23].

In our study, women with an HbA_{1c} of ≥ 5.7% (39 mmol/mol) were not at increased risk of adverse pregnancy outcomes. Among those with negative OGTT results using IADPSG criteria, there was no relationship between adverse pregnancy outcomes and the higher HbA_{1c}, except that women with higher values had less GWG and cesarean rate (possibly from lifestyle interventions). Several previous studies have shown mixed results. Similar to our findings, Fong et al. [10] and Osmundson et al. [11] found no adverse outcomes associated with higher HbA_{1c} ≥ 5.7% (39 mmol/mol). In contrast, a study by Hughes et al. [12] reported a strong relationship between higher HbA_{1c} values (≥ 5.9% (41 mmol/mol)) and adverse pregnancy outcomes such as preeclampsia, perinatal death, major congenital anomaly, and shoulder dystocia, but not LGA (customized) or macrosomia. The difference in ability to predict pregnancy outcomes in our study compared with the Hughes et al. study may be due to several factors including sample size and ethnic differences (the Hughes et al. study had a large number of Polynesian women with a higher HbA_{1c}), and perhaps using the New Zealand criteria, a large number of women with IADPSG-defined GDM remained undiagnosed and untreated but in the “non-GDM” analyses. Similarly, Mane et al. [13] found a higher risk of preeclampsia and macrosomia among participants with a higher HbA_{1c} in a multiethnic cohort using the National Diabetes Data Group (NDDG) OGTT criteria following a 50 g glucose challenge test. In contrast, two other studies [20,23] found no association between early HbA_{1c} values and outcome variables such as preeclampsia, birthweight, macrosomia, polyhydramnios, SGA, and hypertensive disorders. A direct comparison between previous studies was difficult as their population characteristics and sample sizes differed substantially. Also, there was variability in the criteria adopted for GDM diagnosis (those adopted included IADPSG criteria [10,11,21,23], World Health Organization (WHO) 1999 criteria [20], Modified IADPSG criteria [20], National Diabetes Data Group (NDDG) criteria [22], and Carpenter-Coustan (CC) criteria [10]).

Our data question the use of an early HbA_{1c} for detecting/predicting hyperglycemia below the cut point for diabetes in pregnancy. Based on our findings, we suggest that at these low levels, accuracy is severely influenced by the variability between individuals including red blood cell (RBC) longevity and ethnicity. The longer a RBC survives, the longer time there is for glycation, even with a normal blood glucose level. This non-glycaemic influence reduces the discriminative power of HbA_{1c} for detecting mild hyperglycemia as evidenced by low AUC values for GDM defined by the IADPSG criteria. The discordance in association between higher HbA_{1c} and GDM-associated adverse outcomes, is most likely due to the impact of GDM treatment/lifestyle intervention and overall glycemias, being less than the influence of inter-individual variation in the mean RBC age across the population. The differential effect of turnover on HbA_{1c}, reproducible individuals, has been shown outside of pregnancy [24,25]. On the other hand, the contribution of variability in mean RBC age to the HbA_{1c} level appears to be less at higher levels of glycemias. This is evident from the higher AUC for detecting GDM defined by the higher fasting glucose of ≥ 6.1 mmol/l (110 mg/dl).

We, therefore, suggest that an early pregnancy HbA_{1c} value should not be used as a biomarker for detecting mild hyperglycemia in pregnancy and that a threshold of ≥ 5.7% (39 mmol/mol) has limited utility in the prediction of poor pregnancy outcomes in obese pregnant women. Further studies are urgently needed to define the best strategies to diagnose and identify early GDM_i.

A major limitation of our study was the high percentage of missing data (17.6–32.7% for the majority of the outcome variables) which may have influenced our study findings. Secondly, In the higher HbA_{1c} group, 8.8% of the women diagnosed with GDM received medication therapy and 57% of the non-GDM women (excluding women in the usual care, placebo, and vitamin D limbs) received some kind of lifestyle intervention (Supplemental Table 5). This may have affected the outcome of the pregnancy and attenuated the effects of higher HbA_{1c} on adverse outcomes. Nevertheless, the analyses have been adjusted for lifestyle intervention and GDM treatment, and this made no difference to the results. Furthermore, in spite of the clinical treatment, pregnancy outcomes were still worse among women with GDM. We have previously shown that neither gestational weight gain limitation across the cohort, lifestyle change nor Vitamin D supplementation has a substantive impact on pregnancy outcomes or GDM incidence [17,18]. Further, the IADPSG criteria were developed for diagnosis of GDM at 24–28 weeks' gestation, not early pregnancy, and criteria for this time point are under active investigation [26,27]. Moreover, our participants were mainly overweight/obese European women (87%), which limits the generalizability of the study results. Also, all the women had accepted to participate in a life-style intervention study, and therefore these women might not be representative of an unselected population. Finally, full blood count data were not available for the identification of possible shorter or longer red blood cell longevity, so results were not adjusted for the presence of anaemia.

In conclusion, an early HbA_{1c} measurement was not useful in the prediction of GDM in obese European women. An HbA_{1c} with a threshold of 5.7% (39 mmol/mol) is a specific but insensitive biomarker for GDM and is not associated with adverse pregnancy outcomes. Because of the “dilution” effect of hemoglobin glycation by RBC turnover variation at these relatively low levels of HbA_{1c}, the proposed threshold has limited utility in screening for early GDM, in spite of its value for detecting diabetes in pregnancy. Future studies are needed to confirm the results in different populations.

Declaration of Competing Interest

The authors declared that there is no conflict of interest.

Acknowledgments

Professor Andre van Assche, PhD from the KU Leuven Department of Development and Regeneration: Pregnancy, Fetus and Neonate, Belgium and Gynaecology and Obstetrics, University Hospitals Leuven, Belgium contributed equally to the study but sadly passed away before submission.

Author Contributions

Jl undertook the statistical analyses, interpreted the data and drafted the manuscript. DS conceived the project, supervised Jl, interpreted the data, reviewed and edited the draft and provided critical input to the manuscript. DS, RC, RD, AVA, AL, EW, AK, FD, PD, ERM, DH and FS designed and executed the DALI study and provided input for the interpretation of the results. DS was the study coordinator. JMA, SG, MGD, AB, JH, AZ, DM, LTA, HS, JI contributed to the execution of the study and provided input for the interpretation of the results. GD designed and executed the study and provided input for the interpretation of the results. He was the principal investigator of the study. MVP designed and executed the study and provided input for the interpretation of the results. She was the sponsor of the study. All authors read and approved the final manuscript. JI, is the guarantor of this work and takes full responsibility for the contents of this article.

Funding

This project received funding from the European Community's 7th Framework Program (FP7/2007-2013; grant agreement no. 242187). In the Netherlands, additional funding was provided by the **Netherlands Organization for Health Research and Development** (ZonMw) (Grant nr **200310013**). In the UK, the DALI team acknowledges the support received from the NIHR Clinical Research Network: Eastern, especially the local diabetes clinical and research teams based in Cambridge. In Spain, additional funding was provided by CAIBER 1527-B-226. JI is supported by a postgraduate research scholarship from Western Sydney University.

Clinical trial registration number: ISRCTN70595832.

Paper presentation information: Parts of this study were presented at the Diabetes UK Professional Conference, Liverpool, UK, 6–8 March, 2019 and the Australian Diabetes in Pregnancy Society Annual Scientific Meeting, Sydney, Australia, 23–29 August, 2019.

D Simmons, J Immanuel, F Dunne, et al. An early pregnancy HbA_{1c} ≥ 41 mmol/mol (5.9%) is not a good biomarker for early or late Gestational Diabetes (GDM) and does not predict GDM associated complications in obese women. Diabetic medicine. Special issue: Abstracts of the Diabetes UK Professional Conference 2019, ACC Liverpool, 6-8 March 2019. A10 (P92). https://doi.org/10.1111/dme.1_13882

Appendix A Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.diabres.2020.108378>.

References

The corrections made in this section will be reviewed and approved by a journal production editor. The newly added/removed references and its citations will be reordered and rearranged by the production team.

[1] Leng J., Shao P., Zhang C., et al. Prevalence of gestational diabetes mellitus and its risk factors in Chinese pregnant women: a prospective population-based study in Tianjin. China. PLoS One 2015;10. doi:10.1371/journal.pone.0121029. e0121029.

[2] Lavery J.A., Friedman A.M., Keyes K.M., Wright J.D., Ananth C.V. Gestational diabetes in the United States: temporal changes in prevalence rates between 1979 and 2010. BJOG 2017;124:804–813. doi:10.1111/1471-0528.14236.

[3] Djelmis J., Pavic M., Mulliqi Kotori V., Pavlic Renar I., Ivanisevic M., Oreskovic S. Prevalence of gestational diabetes mellitus according to IADPSG and NICE criteria. Int J Gynaecol Obstet 2016;135:250–254. doi:10.1016/j.ijgo.2016.07.005.

[4] American Diabetes Association. 2. Classification and Diagnosis of Diabetes: Standards of Medical Care in Diabetes-2019. Diabetes Care 2019;42:S13–28. <https://doi.org/10.2337/dc19-S002>.

[5] Gallagher E.J., Le Roith D., Bloomgarden Z. Review of hemoglobin A(1c) in the management of diabetes. J Diabetes 2009;1:9–17. doi:10.1111/j.1753-0407.2009.00009.x.

[6] O'Connor C., O'Shea P.M., Owens L.A., Carmody L., Avalos G., Nestor L., et al. Trimester-specific reference intervals for haemoglobin A1c (HbA1c) in pregnancy. Clin Chem Lab Med 2011;50:905–909. doi:10.1515/ccm.2011.397.

[7] Mills J.L., Jovanovic L., Knopp R., et al. Physiological reduction in fasting plasma glucose concentration in the first trimester of normal pregnancy: the diabetes in early pregnancy study. Metabolism 1998;47(1140–44). 10.1016/S0026-0495(98)90290-6.

[8] Lurie S., Marnet Y. Red blood cell survival and kinetics during pregnancy. European J Obstet Gynecol Reprod Bio 2000;93:185–192. doi:10.1016/S0301-2115(00)00290-6.

[9] Hartland A.J., Smith J.M., Clark P.M.S., Webber J., Chowdhury T., Dunne F. Establishing trimester- and ethnic group-related reference ranges for fructosamine and HbA1c in non-diabetic pregnant women. Ann Clin Biochem 1999;36:235–237. doi:10.1177/000456329903600218.

[10] Fong A., Serra A.E., Gabby L., Wing D.A., Berkowitz K.M. Use of hemoglobin A1c as an early predictor of gestational diabetes mellitus. Am J Obstet Gynecol 2014;211(641):e1–e7. doi:10.1016/j.ajog.2014.06.016.

[11] Osmundson S.S., Zhao B.S., Kunz L., et al. First trimester hemoglobin A1c prediction of gestational diabetes. Am J Perinatol 2016;33:977–982. doi:10.1055/s-0036-1581055.

[12] Hughes R.C., Moore M.P., Gullam J.E., Mohamed K., Rowan J. An early pregnancy HbA1c >=41 mmol/mol (5.9%) is optimal for detecting diabetes and identifies women at increased risk of adverse pregnancy outcomes. Diabetes Care 2014;37:2953–2959. doi:10.2337/dc14-1312.

[13] Mañé L., Flores-Le Roux J.A., Benaiges D., et al. Role of First-Trimester HbA1c as a Predictor of Adverse Obstetric Outcomes in a Multiethnic Cohort. J Clin Endocrinol Metab 2017;102:390–397. doi:10.1210/je.2016-2581.

[14] Ministry of Health. 2014. Screening, Diagnosis and Management of Gestational Diabetes in New Zealand: A clinical practice guideline. Wellington: Ministry of Health. Available from <https://www.health.govt.nz/publication/screening-diagnosis-and-management-gestational-diabetes-new-zealand-clinical-practice-guideline>. Accessed January 29, 2019.

[15] Jelsma J.G., van Poppel M.N., Galjaard S., et al. DALI: Vitamin D and lifestyle intervention for gestational diabetes mellitus (GDM) prevention: an European multicentre, randomised trial-study protocol. BMC pregnancy and childbirth 2013;13:142. doi:10.1186/1471-2393-13-142.

[16] Metzger B.E., Gabbe S.G., Persson B., et al. International association of diabetes and pregnancy study groups recommendations on the diagnosis and classification of hyperglycemia in pregnancy. Diabetes Care 2010;33:676–682. doi:10.2337/dc09-1848.

[17] Simmons D., Devlieger R., van Assche A., et al. Effect of physical activity and/or healthy eating on GDM risk: the DALI lifestyle study. J Clin Endocrinol Metab 2017;102:903–913. doi:10.1210/je.2016-3455.

[18] Corcoy R., Mendoza L.C., Simmons D., et al. The DALI vitamin D randomized controlled trial for gestational diabetes mellitus prevention: No major benefit shown besides vitamin D sufficiency. Clin Nutr (in press) 2019. doi:10.1016/j.clnu.2019.04.006.

[19] Immanuel J., Simmons D. Screening and treatment for early-onset gestational diabetes mellitus: a systematic review and meta-analysis. Curr Diab Rep 2017;17:115. doi:10.1007/s11892-017-0943-7.

[20] Odsæter L.H., Åsberg A., Vanky E., et al. Hemoglobin A1c as screening for gestational diabetes mellitus in Nordic Caucasian women. Diabetol Metab Syndr 2016;8:43. doi:10.1186/s13098-016-0168-y.

[21] Wu K., Cheng Y., Li T., et al. The utility of HbA1c combined with haematocrit for early screening of gestational diabetes mellitus. Diabetol Metab Syndr 2018;10:14. doi:10.1186/s13098-018-0314-9.

[22] Benaiges D., Flores-Le Roux J.A., Marcelo L., et al. Is first-trimester HbA1c useful in the diagnosis of gestational diabetes? Diabetes Res Clin Pract 2017;133:85–91. doi:10.1016/j.diabres.2017.08.019.

[23] Amylidi S., Mosimann B., Stettler C., Fiedler G.M., Surbek D., Raio L. First-trimester glycosylated hemoglobin in women at high risk for gestational diabetes. Acta Obstet Gynecol Scand 2016;95:93–97. doi:10.1111/aogs.12784.

[24] Cohen R.M., Franco R.S., Smith E.P., Higgins J.M. When HbA1c and blood glucose do not match: How much is determined by race, by genetics, by differences in mean red blood cell age? J Clin Endocrinol Metab 2019;104:707–710. doi:10.1210/je.2018-02409.

[25] Cohen R.M., Franco R.S., Khera P.K., et al. Red cell life span heterogeneity in hematologically normal people is sufficient to alter HbA1c. Blood 2008;112:4284–4291. doi:10.1182/blood-2008-04-154112.

[26] Wexler D.J., Powe C.E., Barbour L.A., et al. Research gaps in gestational diabetes mellitus: executive summary of a national institute of diabetes and digestive kidney disease workshop. Obstet Gynecol 2018;132:496–505. doi:10.1097/AOG.0000000000000276.

[27] Simmons D., Hague W.M., Teede H.J., et al. Hyperglycemia in early pregnancy: the treatment of booking gestational diabetes mellitus (TOBOGM) study. A randomized controlled trial. Med J Aust 2018;209(405–6). doi:10.5694/mja17.01129.

Highlights

- Early pregnancy HbA1c is a poor predictor of gestational diabetes.
- An HbA_{1c} ≥ 5.7% (39 mmol/mol) has poor sensitivity for GDM regardless of gestation.
- An HbA_{1c} ≥ 5.7% (39 mmol/mol) was not associated with greater risk of adverse outcomes.

Appendix A Supplementary material

The following are the Supplementary data to this article:

[Multimedia Component 1](#)

Supplementary data 1

Queries and Answers

Query: Your article is registered as a regular item and is being processed for inclusion in a regular issue of the journal. If this is NOT correct and your article belongs to a Special Issue/Collection please contact s.hurren@elsevier.com immediately prior to returning your corrections.

Answer: Yes

Query: The author names have been tagged as given names and surnames (surnames are highlighted in teal color). Please confirm if they have been identified correctly.

Answer: Yes

Query: Please check the 'Declaration of Competing Interest' statement has been set correctly.

Answer: Checked. No conflict of interest

Query: Have we correctly interpreted the following funding source(s) and country names you cited in your article: Research and Development, United States; Health Research, United States; Western, Canada? /

Answer: