Journal Pre-proof

Congenital Disorders of Glycosylation in Portugal – Two Decades of Experience

Dulce Quelhas PharmaD, MSc, Esmeralda Martins, MD, PhD, Luísa Azevedo, PhD, Anabela Bandeira, MD, Luísa Diogo, MD, PhD, Paula Garcia, MD, Sílvia Sequeira, MD, Ana Cristina Ferreira, MD, Elisa Leão Teles, MD, Esmeralda Rodrigues, MD, Ana Maria Fortuna, MD, Carla Mendonça, MD, Helena Cabral Fernandes, MD, Ana Medeira, MD, Ana Gaspar, MD, Patrícia Janeiro, MD, Anabela Oliveira, MD, Francisco Laranjeira, MSc, Isaura Ribeiro, MSc, Erica Souche, MSc, Valérie Race, MSc, Liesbeth Keldermans, MSc, Gert Matthijs, PhD, Jaak Jaeken, MD, PhD

PII: S0022-3476(20)31508-0

DOI: <https://doi.org/10.1016/j.jpeds.2020.12.026>

Reference: YMPD 11977

To appear in: *The Journal of Pediatrics*

Received Date: 4 August 2020

Revised Date: 9 November 2020

Accepted Date: 10 December 2020

Please cite this article as: PharmaD DQ, Martins E, Azevedo L, Bandeira A, Diogo L, Garcia P, Sequeira S, Ferreira AC, Teles EL, Rodrigues E, Fortuna AM, Mendonça C, Fernandes HC, Medeira A, Gaspar A, Janeiro P, Oliveira A, Laranjeira F, Ribeiro I, Souche E, Race V, Keldermans L, Matthijs G, Jaeken J, Congenital Disorders of Glycosylation in Portugal – Two Decades of Experience, *The Journal of Pediatrics* (2021), doi: <https://doi.org/10.1016/j.jpeds.2020.12.026>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier Inc. All rights reserved.

Congenital Disorders of Glycosylation in Portugal – Two Decades of Experience

Authors

Dulce Quelhas, PharmaD, MSc^{1,2,3}, Esmeralda Martins, MD, PhD^{2,3}, Luísa Azevedo, PhD^{4,5,6}, Anabela Bandeira, MD³, Luísa Diogo, MD, PhD⁷, Paula Garcia, MD⁷, Sílvia Sequeira, MD⁸, Ana Cristina Ferreira, MD⁸, Elisa Leão Teles, MD⁹, Esmeralda Rodrigues, MD⁹, Ana Maria Fortuna, MD¹⁰, Carla Mendonça, MD¹¹, Helena Cabral Fernandes, MD¹², Ana Medeira, MD¹³, Ana Gaspar, MD¹⁴, Patrícia Janeiro, MD¹⁴, Anabela Oliveira, MD¹⁴, Francisco Laranjeira, MSc^{1,2}, Isaura Ribeiro, MSc^{1,2}, Erica Souche, MSc¹⁵, Valérie Race, MSc¹⁵, Liesbeth Keldermans, MSc¹⁵, Gert Matthijs, PhD¹⁵, Jaak Jaeken, MD, PhD¹⁶

¹ Unidade de Bioquímica Genética, Centro de Genética Médica, Centro Hospitalar Universitário do Porto, Porto, Portugal

² Unit for Multidisciplinary Research in Biomedicine, ICBAS, UP, Porto Portugal

³ Centro Referência Doenças Hereditárias do Metabolismo, Centro Hospitalar Universitário do Porto, Porto, Portugal

⁴ i3S- Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Population Genetics and Evolution Group, Porto, Portugal

⁵ IPATIMUP-Institute of Molecular Pathology and Immunology, University of Porto, Porto, Portugal

⁶ FCUP-Department of Biology, Faculty of Sciences, University of Porto, Porto, Portugal

⁷ Centro Referência Doenças Hereditárias do Metabolismo, Centro Hospitalar Universitário de Coimbra, Coimbra, Portugal

⁸ Centro Referência Doenças Hereditárias do Metabolismo, Centro Hospitalar Universitário de Lisboa Central, Lisboa, Portugal

⁹ Centro Referência Doenças Hereditárias do Metabolismo, Centro Hospitalar Universitário de S. João, Porto, Portugal

¹⁰ Unidade de Genética Médica, Centro Genética Médica, Centro Hospitalar do Porto, Porto, Portugal

¹¹ Centro de Neuropediatria e Desenvolvimento, Centro Hospitalar Universitário do Algarve, Faro, Portugal

¹² Centro de Desenvolvimento da Criança, Centro Hospitalar do Funchal, Funchal, Portugal

¹³ Centro Clínico dos SAMS, Lisboa, Portugal

¹⁴ Centro Referência Doenças Hereditárias do Metabolismo, Centro Hospitalar Universitário de Stª Maria, CHLN

¹⁵ Department of Human Genetics, KU Leuven, Leuven, Belgium

¹⁶ Center for Metabolic Diseases, KU Leuven, Leuven, Belgium

Supported by National Funds through the Fundação para a Ciência e a Tecnologia (Portuguese national funding agency for science, research and technology) in the frameworks of the UID/Multi/00215/2013 project–Unit for Multidisciplinary Research in Biomedicine–UMIB/ICB AS/UP. The authors declare no conflicts of interest.

Abbreviations

LLO Lipid-linked oligosaccharide

CDG Congenital disorder(s) of glycosylation

ER Endoplasmic Reticulum

MPI Mannose-6-phosphate isomerase

PMM2 Phosphomannomutase 2

Key words

CDG; congenital disorder(s) of glycosylation; transferrin, phenotype, *PMM2* genotype

Corresponding author

Dulce Quelhas

Unidade de Bioquímica Genética, Centro de Genética Médica, Centro Hospitalar

Universitário do Porto

Praça Pedro Nunes 88,

4099-028, Porto

PORTUGAL

E-mail: dulce.quelhas@chporto.min-saude.pt

Tel No: +351 22 607 0309

Fax No: +351 22 607 0399

Objective: To describe the clinical, biochemical, and genetic features of both new and previously reported patients with congenital disorders of glycosylation (CDG) diagnosed in Portugal over the last 20 years.

Study design: The cohort includes patients with an unexplained multisystem or single organ involvement, with or without psychomotor disability. Serum sialotransferrin isoforms and, whenever necessary, apolipoprotein CIII (ApoC3) isoforms and glycan structures were analyzed. Additional studies included measurement of phosphomannomutase (PMM) activity

and analysis of lipid-linked oligosaccharides in fibroblasts. Sanger sequencing and massive parallel sequencing were used to identify causal variants or the affected gene, respectively.

Results: 63 individuals were diagnosed covering 14 distinct CDG; 43 patients diagnosed postnatally revealed a type 1, 14 a type 2, and two a normal pattern on serum transferrin isoelectrofocusing. The latter patients were identified by whole exome sequencing. Nine of them presented also a hypoglycosylation pattern on ApoC3 isoelectrofocusing, pointing to an associated O-glycosylation defect. Most of the patients (62 %) are PMM2-CDG and the remaining carry pathogenic variants in *ALG1*, *ATP6AP1*, *ATP6AP2*, *ATP6V0A2*, *CCDC115*, *COG1*, *COG4*, *DPAGT1*, *MAN1B1*, *SLC35A2*, *SRD5A3*, *RFT1* or *PGM1*.

Conclusion: Portuguese CDG patients are here presented, some of them showing unique clinical phenotypes. Among the 14 genes mutated in Portuguese individuals, eight are shared with a previously reported Spanish cohort. However, regarding the mutational spectrum of PMM2-CDG, the most frequent CDG, a striking similarity between the two populations was found, as only one mutated allele found in the Portuguese group has not been reported in Spain.

Because glycosylation occurs in all cells of the organism and approximately one-half of all proteins are glycoproteins [1], it is not surprising that the congenital disorders of glycosylation (CDG) show highly diverse clinical presentations. Currently, they comprise over 130 genetic diseases, caused by any impairment in the assembly and attachment of glycoprotein and glycolipid glycans. There are two main categories of glycosylation pathways the N-glycosylation and the O-glycosylation. N-glycans are linked to the amide group of asparagine, and O-glycans are linked to the hydroxyl group of serine or threonine of the protein moiety. The synthesis of N-glycans proceeds in three stages: formation of

nucleotide-linked sugars in the cytosol, assembly in the cytosol and the endoplasmic reticulum (ER), and processing in the Golgi. The synthesis of O-glycans does not involve processing, and occurs mainly in the Golgi. Besides, there are also defects in lipid glycosylation and glycosylphosphatidylinositol anchor biosynthesis [2].

The standard test for the diagnosis of N-glycosylation disorders with sialic acid deficiency is still isoelectrofocusing of serum transferrin, a liver synthesized N-glycosylated protein. A type 1 pattern (decreased tetrasialotransferrin, increased disialo- and asialotransferrin) points to a glycan assembly defect or a defect in the transfer of the glycan to the peptide chain (CDG-I), and a type 2 pattern (increase also of threesialo- and monosialotransferrin) suggests a remodelling defect (CDG-II) [2].

Around 60 genetic defects have been described in the N-glycosylation pathway, either isolated or in combination with O-glycosylation defects. Phosphomannomutase 2 (PMM2)-CDG is the most common CDG and still underdiagnosed [3]. Signs and symptoms of a CDG can appear at any age, but usually come to medical attention during infancy. The wide-range of clinical presentations and severity of the symptoms, which often involve neurologic manifestations, justifies the need to screen for CDG-related genes in any unexplained disorder with a multisystem phenotype or even when only a single organ is involved. Clinical manifestations include involvement of the nervous system (global developmental disability, seizures, ataxia, hyporeflexia, polyneuropathy, olivopontocerebellar hypoplasia) musculature (generalized hypotonia, muscle weakness), cardiovascular system (cardiomyopathy, pericardial effusion), digestive system (feeding difficulties in infancy, hepatomegaly, elevated serum transaminases, hepatic fibrosis, steatosis), skeletal system (kyphosis, osteopenia), abnormalities of the eyes (nystagmus, strabismus, esotropia, retinopathy), head and neck (microcephaly, facial dysmorphism), skin (abnormal subcutaneous fat tissue distribution, cutis laxa, ichthyosis), endocrine system (hypergonadotropic hypogonadism,

hyperinsulinism), immune system (immunoglobulin deficiencies, B and T cell disorders), blood and blood-forming tissues (dyserythropoietic anemia, thrombocytopenia, low coagulation factors) and growth (failure to thrive). Nonimmune *hydrops fetalis* can be a predominant manifestation in the perinatal period.

We present the features of 63 Portuguese patients with a confirmed molecular diagnosis during the last 20 years.

Methods

Most of the patients presented in this paper were followed in hospitals from the Portuguese National Health Service, and had mainly a multisystem disease of unknown etiology. These patients were selected on a clinical suspicion of CDG (particularly developmental/intellectual disability and liver disease) and/or the finding of laboratory abnormalities such as persistently elevated serum transaminases, coagulation factor decreases (decrease of antithrombin activity, factor XI, protein C, and protein S concentrations), hypoglycemia, hypoproteinemia, hypo- or hypercholesterolemia, decreased thyroxin-binding globulin (TBG)/total thyroxine (T4) and hyperinsulinism. Ferreira et al reported the recognizable phenotypes in CDG (n=36), and Francisco et al enumerated the clinical features suggestive of (distinct CDG (n=73, thus about half of the known CDG number of 137). The conclusion is that CDG should be considered in any unexplained syndrome, with or without intellectual disability [4,5].

Data from major clinical signs and symptoms and diagnostic tests were collected by the referring clinicians

The Figure is a graphic representation of the main diagnostic steps in CDG. The percentage of carbohydrate-deficient serum transferrin (Tf) (sum of a-, mono-, and di-sialo Tf isoforms; % CDT) was calculated according to %CDT Bio-Rad™ HPLC protocol. When the percentage of the hypoglycosylated isoforms was abnormal, serum Tf isoelectrofocusing [6]

and/or capillary zone electrophoresis [7] were performed. In patients with a Tf isoelectrofocusing type 2 pattern, an O-glycosylation defect was investigated through the study of serum ApoC3 isoforms by isoelectrofocusing [8].

In some patients with a type 1 pattern, phosphomannomutase (PMM2; EC 5.4.2.8) and mannose-6-phosphate isomerase (MPI; EC 5.3.1.8) activities were determined in fibroblasts and/or leukocytes, according to Van Schaftingen and Jaeken [9]. After excluding phosphomannomutase 2 and phosphomannose isomerase deficiencies, analysis of dolichol-linked oligosaccharides (LLO) was carried out in fibroblasts of five patients with a type 1 pattern [10] and glycan structure analysis for six patients with a type 2 pattern [11]. Genetic testing was performed after written informed consent from the patients or their legal guardians. Pathogenic variants in *PMM2* were confirmed by Sanger sequencing, as previously reported [12].

Most patients with non-*PMM2*-CDG were examined by a capture targeted panel of 79 CDG associated genes. The mean coverage in the target region was about 600 x and a genotype (homozygous reference, heterozygous or homozygous variant) was called for more than 95 % of the targeted bases. Variant annotation and prioritization were done using the Cartagenia Bench tool. Analysis was done for the exonic sequences and 20 bp of the flanking intronic regions. In addition, deep intronic and promotor regions, in which variants were annotated in the HGMD database, were also scrutinized. An initial filtering of the variants was done on the basis of population frequency (<5 % in at least 2 population databases). Non-informative cases were submitted to whole exome sequencing. Genomic DNA was sheared by sonication, platform-specific adaptors were ligated, and the resulting fragments were size selected. The library was captured using the SeqCap® EZ Human Exome Library v3.0 (Roche NimbleGen), and paired end (2 × 101 bp) sequenced on a HiSeq2000 (Illumina). Reads were aligned to the human reference genome (hg19) using Burrows-Wheeler Aligner (BWA

v0.6.2), and duplicate reads were removed using Picard MarkDuplicates (v1.78). Local realignment around insertions and deletions, base quality score recalibration, and variant calling were performed using Genome Analysis Toolkit (GATK v2.4.9) RealignerTargetCreator, IndelRealigner, BaseRecalibrator, and UnifiedGenotyper. Variants were annotated using Annovar (v11-02-2013). Variant prioritization was applied, based on recessive inheritance, conservation, and pathogenicity prediction scores.

Results

Table I summarizes the main clinical and diagnostic data on the 63 patients, with 14 different disorders of protein N- and of protein N-combined with O-glycosylation (presented according to their subcellular location). 11 of the 24 non-PMM2-CDG patients and 15 of the 39 PMM2-CDG patients have been reported previously [13-32].

Apart from two SLC35A2-CDG patients (one male and one female) diagnosed through a NGS panel, the CDG patients included in this work presented an increased % CDT and an abnormal Tf isoform profile, even in adulthood

Forty-three patients had a type 1 Tf isoform profile, fourteen patients had a type 2 profile and from the latter, nine also showed an abnormal ApoC3 isoform profile suggestive of an associated O-glycosylation defect. PMM activity in fibroblasts was measured in 18 patients with CDG-I, with activities ranging from undetectable to normal [33]. Serum dolichols were measured in one ALG1-CDG patient revealing a very small increase of isoprenols. Sanger sequencing of *PMM2* confirmed the diagnosis of PMM2-CDG in 39 patients from 35 unrelated families. Seventeen pathogenic variants were detected: 15 resulting in an amino acid change (missense variants), one resulting in a stop codon (nonsense variant) and one causing a splice variant. The variants found in these PMM2-CDG patients have been reported (Table 2). All patients were compound heterozygotes except one that was homozygous for c.193G>T (p.D65Y) which is known to be a mild variant. These patients developed

symptoms early in life, mostly hypotonia and failure to thrive (Table 3). The phenotypic spectrum further comprised the known features with psychomotor disability, ataxia and intestinal, hepatic, cardiac, renal and coagulation involvement [34] [35][36] [37]. Overall mortality rate was 18% (7/39) between the age of one month and one year. As for the non-PMM2-CDG patients, a final diagnosis was obtained mostly by a combination of biochemical, molecular or glycobiological investigations [26] [27] [30] supplemented in recent years by massive parallel sequencing either in selected panels (CDG, epilepsy or clinical exome), or whole exome sequencing. The patients identified with this integrated approach included one ALG1-CDG, one ATP6AP1-CDG, one ATP2AP2-CDG, two ATP6V0A2-CDG, one CCDC115-CDG, one COG1-CDG, one COG4-CDG, two DPAGT1-CDG, four MAN1B1-CDG, one PGM1-CDG, three RFT1-CDG, four SLC35A2-CDG and two SRD5A3-CDG.

Discussion

We compared the mutational spectrum of CDG found in Portuguese families with the one previously reported in a Spanish cohort [38]. Only 8 out of 18 CDG found in Spanish patients are shared by the Portuguese cohort (ALG1-CDG, ATP6V0A2-CDG, DPAGT1-CDG, PGM1-CDG, PMM2-CDG, RFT1-CDG, SLC35A2-CDG and SRD5A3-CDG), which reveals a high level of genotypic differences between the two countries, despite their geographic proximity.

A more detailed analysis of the most frequent CDG, PMM2-CDG, revealed that there are 39 patients (from 35 families) with 17 different variants in the Portuguese cohort, compared with 71 patients with 37 variants in the Spanish cohort. The number of deleterious variants in distinct populations seems to reflect specific aspects of the population structure rather than

any biological aspect related to *PMM2* itself; therefore the genetic diversity associated with *PMM2* in each population is high [3][34][39].

Portuguese and Spanish groups have 15 *PMM2* variants in common. The two most prevalent mutations are the c.422G>A (p.R141H) (20 %), the most common variant worldwide and the c.193G>T (p.D65Y) (18 %), an Iberian founder variant [15]. In a Scandinavian CDG cohort there are also two nearly equally prevalent mutations: the c.422G>A (p.R141H) and the c.357C>A (p.F199L), the latter being a South Scandinavian founder variant [40]. Note that, contrary to the c.422G>A (p.R141H) [41] variant, neither c.193G>T (p.D65Y) or c.357C>A (p.F199L) are lethal in the homozygous form.

As to the clinical presentations, there are no marked differences with the CDG patients from other countries except for five patients. In our cohort of *PMM2*-CDG patients, P30 and 32 (Table 3) had an unusually mild/borderline phenotype. In addition, two RFT1-CDG adult patients with mild intellectual disability (from the same small region and thus possibly related), do not present deafness or epilepsy, contrary to all other reported patients, [22]. The fifth is a patient with cutis laxa due to ATP6V0A2-CDG but also corneal dystrophy necessitating repeated corneal transplantation [24]. This eye disorder has not been reported in other patients with this CDG.

Although the vast majority of the patients are still alive, 7 *PMM2*-CDG and 1 *CCDC115*-CDG patients died within their first year. One ATP6V02-CDG patient died in his twenties. Our oldest CDG patients are actually 37 and 39 years.

Due to their clinical and genetic heterogeneity, many CDG are difficult to recognize and to diagnose, whereas, vice versa, few types are sufficiently pathognomonic to drive specific genetic testing. Some phenotypes overlap with those of non-CDG disorders such as mitochondrial diseases [42]. There has been a significant increase in the number of detected CDG, particularly since the more generalized use of massive parallel sequencing.

Acknowledgments

We thank the patients and their families for their participation in this work, and the Portuguese Association for CDG and other Rare Metabolic Diseases (APCDG-DMR). We also thank Dirk J. Lefeber and Luisa Sturiale for glycan structure analysis, Christian Thiel, and the late Christian Körner for LLO studies and Sofia Duarte and Diana Antunes for referring the patients. We thank Eugénia Pinto, Sónia Rocha and Domingos Sousa for the technical support.

References

- [1] Varki A. Biological roles of oligosaccharides: all of the theories are correct. *Glycobiology* 1993;3(2):97-130.
- [2] Péanne R, De Lonlay P, Foulquier F, Kornak U, Lefeber DJ, Morava E, et al. Congenital disorders of glycosylation (CDG): Quo vadis ? *Eur J Med Genet* 2018;61:643–63.
- [3] Schiff M, Roda C, Monin M, Arion A, Barth M, Bednarek N, et al. Clinical , laboratory and molecular findings and long-term follow-up data in 96 French patients with PMM2-CDG (phosphomannomutase2-congenital disorder of glycosylation) and review of the literature. *J Med Genet* 2017;54:843-51.
- [4] Ferreira CR, Altassan R, Marques-Da-Silva D, Francisco R, Jaeken J, Morava E. Recognizable phenotypes in CDG. *J Inherit Metab Dis* 2018;41:541–53.
- [5] Francisco R, Marques-da-Silva D, Brasil S, Pascoal C, Dos Reis Ferreira V, Morava E, et al. The challenge of CDG diagnosis. *Mol Genet Metab* 2019;126:1–5.
- [6] van Eijk HG, van Noort WL, Dubelaar ML, van der Heul C. The microheterogeneity of human transferrins in biological fluids. *Clin Chim Acta* 1983;132:167-71.
- [7] Carchon HA, Chevigne R, Falmagne J, Jaeken J. Diagnosis of congenital disorders of glycosylation by capillary zone electrophoresis of serum transferrin. *Clin Chem* 2004;50:101-11.
- [8] Wopereis S, Grünewald S, Huijben KM, Morava E, Mollicone R, van Engelen BG, et al. Transferrin and apolipoprotein C-III isofocusing are complementary in the diagnosis of N- and O-glycan biosynthesis defects. *Clin Chem* 2007;53:180-7.
- [9] Van Schaftingen E, Jaeken J. Phosphomannomutase deficiency is a cause of

- carbohydrate-deficient glycoprotein syndrome type I. *FEBS Lett* 1995;377:318–20.
- [10] Chantret I, Dancourt J, Dupré T, Delenda C, Bucher S, Vuillaumier-Barrot S, et al. A deficiency in dolichyl-P-glucose:Glc₁Man₉GlcNAc₂-PP-dolichyl α 3-glucosyltransferase defines a new subtype of congenital disorders of glycosylation. *J Biol Chem* 2003;278:9962–71.
- [11] Guillard M, Gloerich J, Wessels HJ, Morava E, Wevers RA, Lefeber DJ. Automated measurement of permethylated serum N-glycans by MALDI-linear ion trap mass spectrometry. *Carbohydr Res* 2009;344:1550–7.
- [12] Matthijs G, Schollen E, Pardon E, Veiga-Da-Cunha M, Jaeken J, Cassiman JJ, van Schaftingen E. Mutations in PMM2, a phosphomannomutase gene on chromosome 16p13, in carbohydrate-deficient glycoprotein type I syndrome (Jaeken syndrome). *Nat Genet* 1997;16:88–92.
- [13] Gomes AL. CDG Síndrome: Uma nova doença multissistémica com atraso mental. *Acta Pediatr Port* 1997;28:355–60.
- [14] Pérez B, Briones P, Quelhas D, Artuch R, Vega AI, Quintana E, et al. The molecular landscape of phosphomannose mutase deficiency in iberian peninsula: Identification of 15 population-specific mutations. *JIMD Rep* 2011;1:117-23
- [15] Quelhas D, Quental R, Vilarinho L, Amorim A, Azevedo L. Congenital disorder of glycosylation type Ia: Searching for the origin of common mutations in PMM2. *Ann Hum Genet* 2007;71:348–53.
- [16] Pérez-Cerdá C, Quelhas D, Vega AI, Ecay J, Vilarinho L, Ugarte M. Screening using serum percentage of carbohydrate-deficient transferrin for congenital disorders of glycosylation in children with suspected metabolic disease. *Clin Chem* 2008;54:93–

- 100.
- [17] Nogueira C, Quelhas D, Vilarinho L. Prenatal diagnosis for CDG Ia based on post-mortem molecular study of Guthrie card. *Mol Genet Metab* 2006;87:379.
- [18] de la Morena-Barrio ME, Sevivas TS, Martínez-Martínez I, Miñano A, Vicente V, Jaeken J, et al. Congenital disorder of glycosylation (PMM2-CDG) in a patient with antithrombin deficiency and severe thrombophilia. *J Thromb Haemost* 2012;10:2625–7.
- [19] Quelhas D, Azevedo L, Freitas M C, Vasconcelos R, Sousa D, Matthijs G, et al. SRD5A3-CDG: a novel mutation. *J Inherit Metab Dis* 2015;38 (Suppl 1):S311.
- [20] Quelhas D, Santos M, Jaeken J, Matthijs G, Lacerda L, Martins E. Exuberant myopathic phenotype in a DPAGT1-CDG patient. *J Inherit Metab Dis* 2016;39 (S1), P-569.
- [21] Ng BG, Shiryaev SA, Rymen D, Eklund EA, Raymond K, Kircher M, et al. ALG1-CDG: Clinical and molecular characterization of 39 unreported patients. *Hum Mutat* 2016;37:653–60.
- [22] Quelhas D, Jaeken J, Fortuna A, Azevedo L, Bandeira A, Matthijs G, et al. RFT1-CDG: Absence of epilepsy and deafness in two Patients with novel pathogenic variants. *JIMD Rep* 2019;43:111-116.
- [23] Rymen D, Peanne R, Millón MB, Race V, Sturiale L, Garozzo D, et al. MAN1B1 deficiency: An unexpected CDG-II. *PLoS Genet* 2013;9(12):e1003989.
- [24] Kornak U, Reynders E, Dimopoulou A, van Reeuwijk J, Fischer B, Rajab A, et al. Impaired glycosylation and cutis laxa caused by mutations in the vesicular H⁺-ATPase subunit ATP6V0A2. *Nat Genet* 2008;40:32–4.

- [25] Leao-Teles E, Quelhas D, Vilarinho L, Jaeken J. De Barsy syndrome and ATP6V0A2-CDG [letter]. *Eur J Hum Genet* 2010;18:526.
- [26] Butler M, Quelhas D, Critchley AJ, Carchon H, Hebestreit HF, Hibbert RG, et al. Detailed glycan analysis of serum glycoproteins of patients with congenital disorders of glycosylation indicates the specific defective glycan processing step and provides an insight into pathogenesis. *Glycobiology* 2003;13:601-22.
- [27] Rujano MA, Cannata Serio M, Panasyuk G, Péanne R, Reunert J, Rymen D, et al. Mutations in the X-linked *ATP6AP2* cause a glycosylation disorder with autophagic defects. *J Exp Med* 2017; 214:3707-29.
- [28] Foulquier F, Vasile E, Schollen E, Callewaert N, Raemaekers T, Quelhas D, et al. Conserved oligomeric Golgi complex subunit 1 deficiency reveals a previously uncharacterized congenital disorder of glycosylation type II. *Proc Natl Acad Sci U S A* 2006;103:3764–9.
- [29] Reynders E, Foulquier F, Teles EL, Quelhas D, Morelle W, Rabouille C, et al. Golgi function and dysfunction in the first COG4-deficient CDG type II patient. *Hum Mol Genet* 2009;18:3244–56.
- [30] Jansen JC, Cirak S, van Scherpenzeel M, Timal S, Reunert J, Rust S, et al. *CCDC115* Deficiency causes a disorder of Golgi homeostasis with abnormal protein glycosylation. *Am J Hum Genet* 2016;98:310–21.
- [31] Lopes F, Barbosa M, Ameer A, Soares G, de Sá J, Dias AI, et al. Identification of novel genetic causes of Rett syndrome-like phenotypes. *J Med Genet* 2016;53(3):190-9.
- [32] Vals MA, Ashikov A, Ilves P, Looorits D, Zeng Q, Barone R, et al. Clinical,

- neuroradiological, and biochemical features of SLC35A2-CDG patients. *J Inherit Metab Dis* 2019;42(3):553-564.
- [33] Grünewald S, Schollen E, Van Schaftingen E, Jaeken J, Matthijs G. High residual activity of PMM2 in patients' fibroblasts: possible pitfall in the diagnosis of CDG-Ia (phosphomannomutase deficiency). *Am J Hum Genet* 2001;68(2):347-54.
- [34] Monin M-L, Mignot C, De Lonlay P, Héron B, Masurel A, Mathieu-Dramard M, et al. 29 French adult patients with PMM2-congenital disorder of glycosylation: outcome of the classical pediatric phenotype and depiction of a late-onset phenotype. *Orphanet J Rare Dis* 2014;9:207.
- [35] Serrano NL, Diego V De, Cuadras D, Monseny AFM, Velázquez-Fragua R, López L, et al. A quantitative assessment of the evolution of cerebellar syndrome in children with phosphomannomutase 2 deficiency (PMM2-CDG). *Orphanet J Rare Dis* 2017:10–5.
- [36] Wolthuis DFGJ, Janssen MC, Cassiman D, Lefeber DJ, Morava-Kozicz E. Defining the phenotype and diagnostic considerations in adults with congenital disorders of N-linked glycosylation. *Expert Rev Mol Diagn* 2014;14:217–24.
- [37] Pancho C, Garcia-cazorla A, Varea V, Artuch R, Ferrer I, Vilaseca MA, et al. Congenital disorder of glycosylation type Ia revealed by hypertransaminasemia and failure to thrive in a young boy with normal neurodevelopment. *J Pediatr Gastroenterol Nutr* 2005;40:230–2.
- [38] Pérez-Cerdá C, Girós ML, Serrano M, Ecay MJ, Gort L, Pérez Duenas B, et al. A population-based study on congenital disorders of protein N- and combined with O-glycosylation experience in clinical and genetic diagnosis. *J Pediatr* 2017;183:170-7.

- [39] Erlandson A, Bjursell C, Stibler H, Kristiansson B, Wahlström J, Martinsson T
Scandinavian CDG-Ia patients: genotype / phenotype correlation and geographic
origin of founder mutations. *Hum Genet* 2001;108:359–67.
- [40] Kjaergaard S, Schwartz M, Skovby F. Congenital disorder of glycosylation type Ia
(CDG-Ia): phenotypic spectrum of the R141H / F119L genotype. *Arch Dis Child*
2001;85:236–9.
- [41] Matthijs G, Schollen E, van Schaftingen E, Cassiman JJ, Jaeken J. Lack of
homozygotes for the most frequent disease allele in carbohydrate-deficient
glycoprotein syndrome type 1A. *Am J Hum Genet* 1998;62:542–50.
- [42] Gardeitchik T, Wyckmans J, Morava E. Complex phenotypes in inborn errors of
metabolism: overlapping presentations in congenital disorders of glycosylation and
mitochondrial disorders. *Pediatr Clin N Am* 2018;65:375–88.

Web Resources

1000 Genomes Project, <https://www.genome.gov/27528684/1000-genomes-project>

dbSNP, <http://www.ncbi.nlm.nih.gov/projects/SNP/>

Genome of the Netherlands (GoNL), <http://www.nlgenome.nl/>

Genome Analysis Toolkit, <https://software.broadinstitute.org/gatk/>

NHLBI Grand Opportunity Exome Sequencing Project (ESP),

<https://esp.gs.washington.edu/drupal/>

OMIM, <http://www.omim.org/>

The Human Gene Mutation Database, <http://www.hgmd.cf.ac.uk/ac/index.php>

The Human Phenotype Ontology, <https://hpo.jax.org/app/>

Table 3 – Updated version of the table presented in 2007 15 with Portuguese PMM2-CDG patients (including HPO severity classification)*

Patient	Gender	Age at diagnosis	Nucleotide change	Protein	Clinical severity
P1	-	PND	c.422G>A; c.484C>T	p.(Arg141His); p.(Arg162Trp)	-
P2	F	1.5 years	c.422G>A; c.677C>G	p.(Arg141His); p.(Thr226Ser)	moderate
P3	M	1 month	c.470T>C; c.691G>A	p.(Phe157Ser); p.(Val231Met)	profound
P4	-	PND	c.470T>C; c.691G>A	p.(Phe157Ser); p.(Val231Met)	-
P5	M	3 years	c.367C>T; c.484C>T	p.(Arg123*); p.(Arg162Trp)	moderate
P6	M	3 months	c.193G>T; c.368G>A	p.(Asp65Tyr); p.(Arg123Gln)	profound
P7	-	PND	c.193G>T; c.368G>A	p.(Asp65Tyr); p.(Arg123Gln)	-
P8	F	2.5 years	c.422G>A; c.484C>T	p.(Arg141His); p.(Arg162Trp)	moderate
P9	F	3 months	c.422G>A; c.691G>A	p.(Arg141His); p.(Val231Met)	profound
P10	M	1.5 years	c.422G>A; c.722G>C	p.(Arg141His); p.(Cys241Ser)	moderate
P11	F	3 months	c.193G>T; c.422G>A	p.(Asp65Tyr); p.(Arg141His)	profound
P12	F	4 months	c.193G>T; c.368G>A	p.(Asp65Tyr); p.(Arg123Gln)	profound
P13	F	4 years	c.193G>T; c.193G>T	p.(Asp65Tyr); p.(Asp65Tyr)	moderate
P14	F	2 years	c.470T>C; c.722G>C	p.(Phe157Ser); p.(Cys241Ser)	moderate
P15	F	1 month	c.193G>T; c.422G>A	p.(Asp65Tyr); p.(Arg141His)	profound
P16	M	4 years	c.422G>A; c.484C>T	p.(Arg141His); p.(Arg162Trp)	moderate
P17	F	16 years	c.338C>T; c.422G>A	p.(Pro113Leu); p.(Arg141His)	moderate
P18	M	2 years	c.368G>A; c.458T>C	p.(Arg123Gln); p.(Ile153Thr)	moderate
P19	M	3 months	c.470T>C; c.691G>A	p.(Phe157Ser); p.(Val231Met)	severe
P20	F	5 years	c.422G>A; c.484C>T	p.(Arg141His); p.(Arg162Trp)	moderate
P21	M	18 years	c.470T>C; c.484C>T	p.(Phe157Ser); p.(Arg162Trp)	mild
P22	F	1 year	c.422G>A; c.722G>C	p.(Arg141His); p.(Cys241Ser)	moderate
P23	M	2 years	c.422G>A; c.677C>G	p.(Arg141His); p.(Thr226Ser)	moderate
P24	M	1 month	c.193G>T; c.422G>A	p.(Asp65Tyr); p.(Arg141His)	profound
P25	M	1 month	c.193G>T; c.367C>T	p.(Asp65Tyr); p.(Arg123*)	profound
P26	-	PND	c.193G>T; c.367C>T	p.(Asp65Tyr); p.(Arg123*)	-
P27	M	11 years	c.193G>T; c.722G>C	p.(Asp65Tyr); p.(Cys241Ser)	mild
P28	M	1 month	c.193G>T; c.470T>C	p.(Asp65Tyr); p.(Phe157Ser)	profound
P29	M	1 month	c.193G>T; c.470T>C	p.(Asp65Tyr); p.(Phe157Ser)	profound
P30	M	4 years	c.353C>G+c.550C>A; c.484C>T	p.(Thr118Ser)+p.(Pro184Thr);	mild/borderline

				p.Arg162Trp	
P31	F	3 years	c.367C>T; c.722G>C	p.(Arg123*); p.(Cys241Ser)	mild
P32	F	3 years	c.193G>T; c.484C>T	p.(Asp65Tyr); p.(Arg162Trp)	mild/borderline
P33	M	1 year	c.395T>C; c.523+3A>G	p.(Ile132Thr)	moderate
P34	F	1 year	c.484C>T; c.710C>G	p.(Arg162Trp); p.(Thr237Arg)	moderate
P35	F	8 months	c.470T>C; c.710C>T	p.(Phe157Ser); p.(Thr237Met)	severe
P36	M	19 years	c.422G>A; c.722G>C	p.(Arg141His); p.(Cys241Ser)	moderate
P37	M	9 years	c.422G>A; c.484C>T	p.(Arg141His); p.(Arg162Trp)	mild
P38	F	3 years	c.422G>A; c.484C>T	p.(Arg141His); p.(Arg162Trp)	moderate
P39	M	1 month	c.193G>T; c.368G>A	p.(Asp65Tyr); p.(Arg123Gln)	profound

PND: prenatal diagnosis; *Clinical severity classification is based on the degree of CNS involvement (ID) and involvement of other organs (heart, kidneys, liver)

Table 2- Portuguese PMM2-CDG patients' allelic frequency

Nucleotide changes	Protein changes	Allelic frequency (%)
c.422G>A	p.Arg141His	20.5
c.193G>T	p.Asp65Tyr	19.3
c.484C>T	p.Arg162Trp	14.1
c.470T>C	p.Phe157Ser	10.3
c.722G>C	p.Cys241Ser	7.7
c.368G>A	p.Arg123Gln	6.4
c.691G>A	p.Val231Met	5.1
c.367C>T	p.Arg123*	5.1
c.677C>G	p.Thr226Ser	2.6
c.338C>T	p.Pro113Leu	1.3
c.353C>G	p.Thr118Ser	1.3
c.395T>C	p.Ile132Thr	1.3
c.458T>C	p.Ile153Thr	1.3
c.550C>A	p.Pro184Thr	1.3
c.710C>T	p.Thr237Met	1.3
c.710C>G	p.Thr237Arg	1.3
c.523+3A>G		1.3

Table 1- Core data of the patients (Human Phenotype Ontology (HPO) designations used)

CDG phenotype MIM nr	Number of patients; families (references)	Gender F;M	Age at diagnosis	Affected glycosylation pathway (serum Tf IEF pattern)	Detection method	Major clinical signs and symptoms	Results of laboratory tests, histology and brain imaging
Defects in the cytosol							
PMM2-CDG #212065	39 (35 families) [13][14][15][16][17][18]	16 F; 19 M, 4 PND	birth to 19 y	N (1)	38 Sanger; 1 WES	Nonimmune hydrops fetalis Microcephaly, abnormal facial shape, mild to severe global developmental delay, ataxia, muscular hypotonia, dystonia Abnormal eye movements, strabismus Abnormal subcutaneous fat tissue distribution, generalized oedema, cryptorchidism, inverted nipples Feeding difficulties, vomiting and diarrhoea in infancy Hypertrophic cardiomyopathy, ventricular hypertrophy, pericardial effusion Abnormality of the skeletal system, arachnodactyly	Hypoalbuminemia, proteinuria, elevated serum transaminases Reduced protein C and protein S activity MRI: olivopontocerebellar hypoplasia
PGM1 # 614921	1	M	10 Y	N (2)	Rhabdomyolysis gene panel	Hepatomegaly Submucous cleft hard palate Recurrent rhabdomyolysis associated with acute infections	Elevated serum transaminases, (asymptomatic) hypoglycaemia Elevated CK levels,
Defects in the ER							
SRD5A3-CDG # 612379	2 (1 family) [19]	1 F;1 M	8 y;14 y	N (1)	Sanger	Severe global developmental delay and intellectual disability, seizures, hyperreflexia Rod-cone dystrophy, nystagmus, alternating exotropia, strabismus and blue sclerae Short nose and gingival overgrowth, wide intermamillary distance and gynecomastia Abnormal subcutaneous fat tissue distribution and hirsutism	LLO: relatively low incorporation of radioactivity without specific changes Brain CT scan: normal
DPAGT1-CDG # 608093	2 (2 families) [20]	1 F;1 M	16 y; 5 y	N and O (1)	CDG and metabolic myopathies capture gene panel	Global developmental delay and intellectual disability, developmental regression, generalized myoclonic and tonic-clonic seizures, congenital muscular torticollis Myopathy, limb-girdle muscle weakness, fatigable weakness, proximal and distal amyotrophy, abnormal pyramidal sign and inability to walk Limited extraocular movements, ophthalmoparesis Inverted nipples, abnormal subcutaneous fat tissue distribution	Muscle biopsy: autophagic vacuoles, muscle fibre cytoplasmic inclusion bodies, muscle fibre tubular inclusions, type 1 muscle fibre predominance and decreased activity of mitochondrial respiratory chain MRI: global brain atrophy
ALG1-CDG # 608540	1 [21]	1 M	7 y	N (1)	Sanger	Severe global developmental delay, microcephaly, seizures, generalized hypotonia	MRI: cerebellar atrophy
RFT1-CDG # 612015	3 (3 families) [22]	1 F;2 M	4 y;20 y; 2 y	N (1)	CDG capture gene panel	Intrauterine growth retardation Global developmental delay and intellectual disability (borderline to severe), seizures (1/3), sensorineural hearing impairment (1/3), ataxia,	Increased thyroid-stimulating hormone level, normal serum free T3 and T4 levels Prolonged partial thromboplastin time

						stereotypy (1/3), muscular hypotonia Abnormal facial shape with prominent forehead, deeply set eyes, epicanthus Scoliosis	and prothrombin time Brain MRI: normal
Defects in the Golgi/ vacuolar membranes/ vesicular membranes							
MAN1B1- CDG * 604346 # 614202	4 (4 families) [23]	1 F; 3M	7 y; 5 y; 7 y; 4 y	N (2)	Sanger; WES	Global developmental delay, intellectual disability, moderate to severe, macrocephaly, generalized hypotonia Long face, macrotia, low-set ears, epicanthus, downslanted palpebral fissures, hypertelorism, prominent nose, thin upper lip vermilion, short philtrum Retinal coloboma Long slender fingers Inverted nipples, wide intermamillary distance Truncal obesity Aggressive behaviour and stereotypy	Echography: aortic aneurysm MRI: normal (1 /4) or cerebellar hypoplasia, cerebellar vermis hypoplasia
ATP6VOA2-CDG # 219200	2 (2 families) [24][25]	2 M	14 y; 3 m	N and O (2)	Sanger; CDG capture gene panel	Mild to severe global developmental delay, Normal to moderate intellectual disability, seizures Failure to thrive Cutis laxa, redundant skin, abnormality of connective tissue Delayed cranial suture closure, large fontanelles, abnormality of the face, dermatochalasis, downslanted palpebral fissures, microretrognathia, high palate, abnormal location of ears and abnormality of the pinna Strabismus and corneal dystrophy (requiring transplantation) Lumbar hyperlordosis, thoracic kyphosis, genu recurvatum, joint hypermobility, bilateral congenital hip dislocation Multiple carious teeth Splenomegaly	Elevated serum aspartate aminotransferase Leukopenia, thrombocytopenia Reduced factor XI activity, prolonged prothrombin time, reduced protein C and protein S activity MRI: pachygyria, cerebellar vermis hypoplasia, Dandy-Walker malformation
ATP6AP1-CDG # 300972	1	1 M	1.5 y	N and O (2)	Sanger	No intellectual disability Relative macrocephaly, facial dysmorphism Pectus excavatum, scoliosis, joint hypermobility Hepatosplenomegaly Cryptorchidism Recurrent respiratory infections	Decreased circulating IgG and total IgM, hyperphosphatasia, hypercholesterolemia, decreased serum ceruloplasmin Liver biopsy: microvesicular hepatic steatosis, cirrhosis
ATP6AP2-CDG * 300556 No MIM for the CDG phenotype	1 [26][27]	1 M	5 m	N and O (2)	WES	Mild cognitive impairment Abnormal facial shape Prolonged neonatal jaundice, persistent hepatosplenomegaly and ascites Recurrent severe infections (e.g. sepsis, peritonitis)	Leukocytosis, neutrophilia Elevated serum transaminases, hypoalbuminemia Prolonged partial thromboplastin time and prothrombin time Decreased antibody levels in blood, (IgG, IgM, IgA), specific anti- polysaccharide antibody deficiency, decreased proportion of CD4-positive T cells and CD8-positive T cells

							IgG hypoglycosylation Liver biopsy: macrovesicular hepatic steatosis, micronodular cirrhosis
COG1-CDG # 611209	1 [28]	1 F	7 m	N and O (2)	Sanger	Mild psychomotor disability, progressive microcephaly Generalized hypotonia, mainly in upper limbs, but with normal muscle strength Narrow forehead and downslanted palpebral fissures, small hands and feet Feeding difficulties since birth and failure to thrive, no hepatosplenomegaly in infancy Left ventricular hypertrophy with left ventricular diastolic dysfunction After infancy: growth delay with rhizomelia and mild hepatosplenomegaly	MRI: slight cerebral and cerebellar hypoplasia
COG4-CDG # 613489	1 [29]	1 M	1 y	N and O (2)	Sanger	Moderate psychomotor disability, postnatal microcephaly, complex seizures, mild ataxia, axial hypotonia, slight peripheral hypertonia, hyperreflexia, brisk reflexes, limb dysmetria and poor speech. Abnormality of the face (down sloping frontal area and thick hair)	Elevated serum transaminases and alkaline phosphatase, increased lactate dehydrogenase activity, hypercholesterolemia Leukocytosis, thrombocytopenia Brain MRI: normal
CCDC115-CDG # 616828	1 [30]	1 F	5 m	N and O (2)	WES	Generalized moderate hypotonia, with reduced muscle size Redundant skin Failure to thrive, jaundice, hepatosplenomegaly and progressive cholestatic liver disease Postmortem liver: severe cholestasis with intrahepatic portal vein sclerosis and cirrhosis	Hypoglycemia, hyperammonemia, elevated serum transaminases, alkaline phosphatase and total and LDL cholesterol Reduced factor VII activity, increased factor VIII activity Anemia with acanthocytes and reticulocytosis Bone marrow biopsy: hypoplastic anemia, some foam cells and hemophagocytosis Control biopsy after 2 months: bone-marrow foam cells and erythroid hypoplasia with perinuclear deposition of iron
SLC35A2-CDG # 300896	4 (4 families) [31] [32]	2 F; 2 M	5 m; 2 y; 3 y; 4 y	N and O (2)	CDG capture gene panel	Two distinct phenotypes: <u>1 without epilepsy</u> Minor neurological involvement Growth deficiency associated with decreased serum IGF1 Minor facial dysmorphism, camptodactyly of fingers and toes <u>3 with epileptic encephalopathy</u> Severe global developmental delay and intellectual disability Muscular hypotonia of the trunk, limb hypertonia Facial dysmorphism Yellow/white lesions of the retina, chorioretinal hyperpigmentation Hepatomegaly	Elevated serum- ransaminases (aspartate aminotransferase > alanine aminotransferase) EEG: burst suppression and hypsarrhythmia Echo: left ventricular hypertrophy, abnormal atrial septum morphology Brain MRI: normal

N: N-glycosylation pathway; O: O-glycosylation pathway; PND: prenatal diagnosis

Journal Pre-proof

Fig. 1- CDG diagnosis algorithm. 1: Transferrin, 2: Next generation sequencing (CDG panel analysis, WES, WGS), 3: Intellectual disability, 4: Cerebellar hypoplasia, 5: Glycosylphosphatidylinositol anchor synthesis defect, 6: Alkaline phosphatase. *Adapted from Francisco et al. [5], ** [4] [5]