1 **Phase 2 study of anastrozole in recurrent estrogen (ER) / progesterone**

2 **(PR) positive endometrial cancer: The PARAGON trial – ANZGOG 0903**

3 Linda Mileshekin^{1*}, Richard Edmondson^{2*}, Rachel L O'Connell³, Katrin M Sjoquist³,
4 John Andrews³, Rema Jyothirmayi⁴, Philip Beale⁵, Tony Bonaventura⁶, Jeffrey
5 Goh⁷, Marcia Hall⁸, Andrew Clamp⁹, John Green¹⁰, Rosemary Lord¹⁰, Frédéric
6 Amant¹¹, Laura Alexander¹², Karen Carty¹², James Paul¹², James Scurry⁶, David
7 Millan¹³, Steven Nottley¹³, Michael Friedlander¹⁴ on behalf of the PARAGON study
8 group

9 *Joint first author

10 ¹ Peter MacCallum Cancer Centre and The Sir Peter MacCallum Department of
11 Oncology, The University of Melbourne, Melbourne, VIC, Australia

12 ² Division of Cancer Sciences, Faculty of Biology, Medicine and Health, University
13 of Manchester, St Mary's Hospital, Manchester, UK; Department of Obstetrics and
14 Gynaecology, Manchester Academic Health Science Centre, St Mary's Hospital,
15 Central Manchester NHS Foundation Trust, Manchester Academic Health Science
16 Centre; Level 5, Research, Oxford Road, Manchester, UK;

17 ³ NHMRC Clinical Trials Centre, University of Sydney, Sydney, NSW, Australia

18 ⁴ Maidstone Hospital, Kent, UK

19 ⁵ Chris O'Brien Lifehouse, Sydney, NSW, Australia

20 ⁶ Pathology New South Wales, Hunter New England and Faculty of Health and
21 Medicine, University of Newcastle, Newcastle, NSW Australia

22 ⁷ Royal Brisbane and Women's Hospital, Brisbane, Australia

23 ⁸ Mount Vernon Cancer Centre, Middlesex, UK

24 ⁹ The Christie NHS Foundation Trust, Manchester, UK

25 ¹⁰ The Clatterbridge Cancer Centre, Liverpool and Wirral, UK

26 ¹¹ Division of Gynecologic Oncology, University Hospitals Gasthuisberg, Leuven,
27 Belgium

28 ¹² Cancer Research UK Clinical Trials Unit, Institute of Cancer Sciences, University
29 of Glasgow, United Kingdom

30 ¹³ Queen Elizabeth University Hospital, Glasgow, Scotland, UK

31 ¹⁴ Royal Hospital for Women/Prince of Wales Hospital and Prince of Wales Clinical
32 School, University of New South Wales, Sydney, Australia

33

34 **Corresponding Author**

35 Linda Mileshekin

36 Peter MacCallum Cancer Centre

37 305 Grattan St, Melbourne, VIC, 3000, Australia

38 Ph: +613 85595000

39 Fax: +613 85597739

40 Email: Linda.Mileshekin@petermac.org

41

42 **Acknowledgements**

43

44 This research was supported by Cancer Australia Grant 632740 (2009-2011),
45 Cancer Australia PdCCRS 1063014 (2014-2016), Cancer Australia Priority-

driven Collaborative Cancer Research Scheme Project Grant APP1142697,
Cancer Research UK (grants C22375/A12846 and C22375/A13784),
and Cancer Institute NSW Research Infrastructure Grant 15/RIG/1-16. UK Co-
Sponsors: NHS Greater Glasgow & Clyde and University of Glasgow.
FA is a senior researcher for the Flemish Research Fund Flanders (F.W.O.). LM
was supported by a Clinical Research Fellowship from the Victorian Cancer
Agency.

Anastrozole was provided by AstraZeneca.

Word count abstract = 248

Word count main text = 3537

Abstract

Background: The clinical benefit rate with aromatase inhibitors and the impact of treatment on quality of life (QOL) in endometrial cancer is unclear. We report the results of a phase 2 trial of anastrozole in endometrial cancer.

Methods: Investigator initiated single-arm, open label trial of anastrozole, 1 mg/d in patients with ER and /or PR positive hormonal therapy naive metastatic endometrial cancer. Patients were treated until progressive disease (PD) or unacceptable toxicity. The primary end-point was clinical benefit (response + stable disease) at 3 months. Secondary endpoints include progression-free survival (PFS), quality of life (QOL) and toxicity.

Results: Clinical benefit rate in 82 evaluable patients at 3 months was 44% (95% CI: 34-55%) with a best response by RECIST of partial response in 6 pts (7%; 95% CI: 3-15%). The median PFS was 3.2 months (95% CI: 2.8-5.4). Median duration of clinical benefit was 5.6 months (95% CI: 3.0-13.7). Treatment was well tolerated. Patients who had clinical benefit at 3 months reported clinically significant improvements in several QOL domains compared to those with PD; this was evident by 2 months including improvements in: emotional functioning (39 vs 6%: $p = 0.002$), cognitive functioning (45 vs 19%: $p = 0.021$), fatigue (47 vs 19%: $p = 0.015$) and global health status (42 vs 9%: $p = 0.003$).

Conclusion: Although the objective response rate to anastrozole was relatively low, clinical benefit was observed in 44% of patients with ER/PR positive metastatic endometrial cancer and associated with an improvement in QOL.

Introduction

The incidence of endometrial cancer is rising and in developed countries it is now the commonest gynaecological malignancy[1]. Although many patients are cured with surgery alone there remains a proportion who will relapse despite surgery and adjuvant therapy. Treatment of patients with relapsed or metastatic disease is challenging as standard cytotoxic chemotherapy rarely results in prolonged disease control and can be associated with significant toxicity in a population who are often elderly and have other significant co morbidities.

Estrogen, plays a key role in the pathogenesis of endometrial hyperplasia and endometrioid adenocarcinoma type I. Most of these Type 1 endometrioid carcinomas are associated with endometrial hyperplasia and are also ER/PR positive, p53 negative and have a low Ki-67. There is evidence that hormonal therapy can be associated with clinical benefit in patients with recurrent/metastatic EC and is widely used to treat a subset of patients. Most work to date has focussed on the role of progestogens and historically response rates of up to 70% were reported in women with PR-positive endometrial cancers compared with 12% in women with PR-negative tumors [2-4]. However, using more rigorous response criteria in clinical trials and institutional studies, the objective response rates are lower and range from 15% to 35% [2, 5]. Medroxyprogesterone is approved by the FDA for the treatment of women with

109 EC but can be associated with significant adverse effects, including weight gain,
110 hypertension, fluid retention, increased blood sugar, insomnia, tremor,
111 thrombosis, and pulmonary emboli. These can potentially worsen quality of life
112 and may be life threatening [2]. There has been interest in the potential role of
113 aromatase inhibitors in EC given their activity in ER positive breast cancer and
114 superiority to tamoxifen in breast cancer. In addition, aromatase is highly
115 expressed in the endometrial stroma, and is responsible for local synthesis of
116 estrogens which may promotes estrogen-induced proliferation of tumour
117 cells[3].

118 The reported response rates to aromatase inhibitors in recurrent and metastatic
119 endometrial cancer have generally been low. To date 6 studies have reported the
120 use of aromatase inhibitor therapy in advanced or recurrent endometrial cancer
121 [6-11] including a total of 104 patients. Response rates vary but are in the order
122 of 10% [12], and this almost certainly reflects the population who were treated
123 [7]. In the studies that have been reported, most patients have had high-grade,
124 hormone receptor-negative cancers, where a low likelihood of response would
125 be expected. There is still a need to evaluate aromatase inhibitors in women with
126 well-differentiated and/or hormone receptor-positive tumors, where the
127 expected response to an aromatase inhibitor is likely to be higher, as well as to
128 evaluate the impact of treatment on quality of life.

129 The PARAGON (ANZGOG-0903) trial is an investigator-initiated basket trial
130 investigating the activity of anastrozole in post-menopausal patients with a wide
131 range of ER or PR positive recurrent or metastatic gynaecological tumours. It
132 includes 7 separate phase 2 open label prospective trials in different
133 gynaecological cancer types embedded within the one protocol. Here we report

the results of the prospective trial in the endometrial cancer cohort which aimed to investigate the clinical benefit rate of the use of anastrozole, in women with hormone receptor positive recurrent or metastatic endometrial cancer with an additional focus on the impact on quality of life.

Materials and Methods

The study design was a single-arm, open label trial of anastrozole at a dose of 1mg daily in post-menopausal women with ER and/or PR positive hormonal therapy naïve recurrent endometrial cancer. Patients were treated until disease progression by RECIST version 1.1 criteria or unacceptable toxicity.

PARAGON is a Gynaecologic Cancer InterGroup trial led by the Australian New Zealand Gynecological Oncology Group (ANZGOG) and coordinated by the NHMRC Clinical Trials Centre, University of Sydney. The collaborating groups are the Scottish Gynaecological Cancer Trials Group (SGCTG) via Cancer Research UK Clinical Trials Unit, Glasgow and the Leuven Cancer Institute. The Study was performed in accordance to the NHMRC Statement on Ethical Conduct in Research Involving Humans and the Declaration of Helsinki. Ethical approval was obtained at all participating sites and all participants provided signed, written, informed consent.

PARAGON is registered on the Australian New Zealand Clinical Trials Registry (#ACTRN12610000796088)

Eligibility

Eligible patients had recurrent or metastatic endometrial cancer that was ER and/or PR positive based on local testing at sites. ER and/or PR positivity was defined as at least 10% of cells staining positive for ER and/or PR in their primary tumour or a biopsy of recurrent disease. In addition, eligible patients were postmenopausal, had no prior anti-cancer endocrine treatment, aged ≥ 18 , ECOG performance status 0 – 2, had a life expectancy of >3 months, and measurable disease by RECIST v 1.1 criteria. Women receiving hormone replacement therapy or those with significant hepatic (bilirubin >2 x upper limit of normal) or renal dysfunction (creatinine >3 x upper limit of normal) were excluded. The baseline evaluation included history, physical examination, ECOG performance status, abdominal and pelvic computed tomography (CT) scan, full blood count, blood chemistry and liver function tests. A CT chest was also required in patients with known or suspected pulmonary metastases.

Study objectives

The primary objective was to assess the clinical benefit rate, defined as the proportion of patients who had response or stable disease at 3 months by RECIST 1.1 criteria. Secondary objectives included progression free survival (PFS), response duration, quality of life, and toxicity. Quality of life (QOL) and tolerability was assessed using the EORTC QLQ-C30 and the FACT-ES subscale score. The proportion of patients experiencing grade 3 or 4 toxicities as well as number of patients who came off therapy because of adverse events were also documented. The QLQ-C30 is the core module of the EORTC's quality of life questionnaire (QLQ) suite [13] and contains 5 functional scales (physical, role,

emotional, cognitive and social functioning), overall health/global HRQL (global health status (GHS)), and 9 symptom/difficulties scales. Definitions for minimal important difference (MID) were adopted from Cocks et al 2011 for the EORTC QLQ-C30 [14]. Subjective endocrine-related symptoms were assessed by Functional Assessment of Cancer Therapy - Endocrine Symptoms (FACT-ES) subscale, a validated 5-point response scale developed for breast cancer research [15].

Response and Toxicity Assessment

Response status was assessed using RECIST v1.1 criteria. Clinical deterioration in the absence of proven progression was determined by the treating physician. CT scans were done at baseline and repeated every 3 months while patients remained on the trial. Adverse events were collected monthly for the first 3 months and then 3 monthly and toxicity was graded according to NCI CTCAE V4.0. Quality of life was measured at registration prior to commencing anastrozole, monthly for the first 3 months, and 3 monthly thereafter until progression.

Immunohistochemical staining for ER and PR

Central testing for ER and PR status was performed after trial completion in the UK for cases with sufficient tissue available (unstained slides or blocks were requested). Representative tumour sections (4 µm) were mounted on Superfrost Plus microscope slides and dried at 37°C for 24 hours before a final incubation of 30 minutes at 60 C before staining. A composite control slide for each staining run and control breast cancer sections were included on each test slide.

Immunohistochemical staining using primary antibodies to ER (Ventana SP1, Ref. 790-4324) and PR (Ventana 1E2, Ref. 790-2223) was performed using a Ventana BenchMark ULTRA IHC staining module according to the manufacturer's instructions (Ventana Medical Systems Inc., Tucson, Arizona, USA), visualized with diaminobenzidine and counterstained with haematoxylin. Staining was evaluated by 2 pathologists blinded to treatment response and scored according to the Allred protocol[16].

Statistical considerations

The expected clinical benefit rate at 3 months in the endometrial cancer cohort was 20% based on a review of the literature suggesting a 10% response rate in previously reported studies of aromatase inhibitors in endometrial cancer[8, 9]. To provide sufficient precision in the estimates of clinical benefit, a sample size of 75 patients was planned. The study had a stopping rule to allow for termination if there was lack of efficacy. There was a pre-planned interim analysis after 25 evaluable subjects had been on study for at least 3 months (and received at least 2 weeks of treatment). The data were reviewed by an Independent Data Monitoring and Safety Committee on 24th April 2013 who recommended continued recruitment to 75 patients since the minimum number of clinical benefit responses required to be consistent with the expected clinical benefit rate was observed (i.e., at least 2 in the first 25 patients). Analysis of the efficacy (overall RECIST response/clinical benefit) was performed using the proportion of patients who responded/experienced clinical benefit together with 95% confidence interval for the estimates. These rates were based on all patients receiving anastrozole for the first 2 weeks (intention

to treat population) as well as patients who were on study for at least 4 weeks (evaluable for response). Toxicity analysis was evaluated by treatment received. Comparisons were 2-tailed and a nominal significance level of 0.05 was applied. Progression-free survival and duration of clinical benefit were analysed using time-to-event methods, with Kaplan-Meier survival curves constructed for graphical display and unadjusted log-rank tests performed where appropriate. Death from any cause was considered an event. 95% CIs for proportions were constructed using the modified Wilson method [17]. The conditional binomial exact test was used to test for association between binary variables. Cut-points were chosen for the ER and PR scores with consideration to patient numbers in respective categories and using optimal cut-point selection based on both sensitivity and specificity. The paired t-test was used to compare baseline and on study QoL scores at individual time-points. In addition, change scores between baseline and on-study averaged scores were also computed and assessed using one-sample t tests. The proportion of patients whose score improved by ≥ 10 -points (considered clinically relevant) was calculated for each QLQ-C30 subscale. Linear regression was used to compare change in QoL scores between patients achieving a 3-month clinical benefit and those who progressed, with adjustment for the baseline score.

Results

84 eligible patients with ER and/or PR positive recurrent or metastatic endometrial cancer were enrolled in 31 centres in Australia (n = 40), the UK (n = 38), Belgium (n = 4) and New Zealand (n = 2) between Feb 2012 and March 2014

(Consort diagram, figure 1). Mean age was 68 years (range 36-89). 50% patients had received prior chemotherapy whilst 67% had received prior radiotherapy, table 1. There was a wide distribution of initial stage and grade of tumours with 24 (29%) tumours reported as grade 3 and 50 (60%) $\geq$ FIGO stage 2 at diagnosis. One patient withdrew before taking any study drug whilst one further patient took drug for less than 1 week leaving 82 patients eligible for analysis. All eligible patients received anastrozole 1mg daily until clinical progression, figure 1.

Clinical Response

Clinical benefit at 3 months was recorded in 36/82 patients (44%; 95% CI: 34-55%). A best RECIST response of partial response was observed in 6 patients (7%; 95% CI: 3-15%). The median PFS for the whole cohort was 3.2 months (95% CI: 2.8 – 5.4, Figure 2a). PFS was superior in patients with a treatment free interval prior to registration of greater than 12 months (Figure 2b). The median duration of clinical benefit was 5.6 months (95% CI: 3.0 – 13.7, Figure 2c). Fifteen patients (18%) remained on treatment for over a year. Partial responses were seen only in patients with grade 1 or 2 EC, but prolonged disease control was observed in 2 patients with grade 3 EC (figure 3). The clinical benefit rate at 3 months was not significantly different between normal weight, overweight and obese individuals (46%, 21% and 51% respectively: P-value (trend) = 0.603). Interestingly the clinical benefit rate was higher in those women who had received prior chemotherapy compared to those

who had not (56% vs 32%; $P = 0.026$). None of the partial responses were seen in women who had received prior chemotherapy.

The likelihood of achieving clinical benefit did not vary significantly by local testing of ER or PR status. However, partial responses were only seen in patients who had both ER and PR positive disease with no responses seen in patients who had ER+/PR- or ER-/PR+ disease. All 4 patients with ER-/PR+ disease progressed at their first re-staging.

Sufficient tumor for central testing of ER and PR was available for 53 patients. In 3 of these patients no ER or PR expression was seen. No strong association was seen between the clinical benefit rate and the Allred scores (Table 2). However, there was a trend towards a lower clinical benefit rates among patient with a PR proportion <4 compared to those with 4 or more (30% vs 58%; $p = 0.056$).

Safety

Toxicity data were available for 82 of the 83 patients who received treatment. In general, treatment was well tolerated with the toxicity profile being as expected for an aromatase inhibitor. The commonest side effects were hot flushes (44%), arthralgia (48%) and fatigue (69%). The 7 (9%) grade 3 toxicities were confined to fatigue, which could be related to treatment or disease. There were no grade 4 toxicities (table 3) and no patient stopped treatment due to adverse events.

Other adverse events reported during treatment were considered disease-related and included abdominal ascites, abdominal distension, small intestinal obstruction, cellulitis, abdominal pain, small intestine fistula, pulmonary embolus and pneumonia.

303

304 QoL

305 QoL data were available for 79 of the 84 registered patients; Belgium did not
306 participate (n=4) in the QoL component of PARAGON. Compliance with
307 completing QoL questionnaires was high: 95% completed QoL questionnaires at
308 baseline (76/80 patients), 93% at 1 month (70/75), 96% at 2 months (67/70)
309 and 95% at 3 months (53/56); 71 completed QoL at both baseline and follow-up.

310

311 For the total cohort there were no statistically significant changes from baseline
312 in the QLQ-C30 subscales averaged over the total time on-study. By time-point,
313 emotional functioning scores improved significantly albeit by a small amount at
314 months 1 and 2 i.e., (n = 67; mean change: 3.2 points; 95% CI: 0.3-6.1; p = 0.03)
315 and (n = 65; 4.7; 95% CI: 1.0-8.4; p = 0.02) respectively. For those still on study
316 at 6-months (n = 31), scores improved significantly for cognitive functioning
317 (8.1; 95% CI: 1.4-14.7; p = 0.02) and diarrhoea (-8.6; 95% CI: -15.6- -1.6; p =
318 0.02); these changes are defined as small (subtle but clinically relevant) and
319 medium respectively by Cocks et al[14]. At 3-months (n=51) nausea and
320 vomiting worsened significantly by a small but significant amount (4.9; 95% CI:
321 0.6-9.2; p=0.03).

322

323 No significant changes from baseline in FACT-ES total scores (range 0-180) were
324 seen at any time point or averaged over the total on-study period. For the
325 emotional well-being subscale (range 0-24) scores improved significantly by 1.1
326 points averaged over the total time on study (95% CI: 0.32-1.82; p = 0.006).

FACT-G (range 0-108) scores improved significantly at 6-months by 3.8 points on average (n = 29; 95% CI: 0.1-7.5; p = 0.05).

In analyses stratified by 3-month clinical benefit status, patients recorded as having a clinical benefit were more likely to report clinically significant improvements of at least 10 points in several QoL domains after the first 2 months of treatment including improvements in: emotional functioning (39 vs 6%: p = 0.002), cognitive functioning (45 vs 19%: p = 0.021), fatigue (47 vs 19%, p = 0.015) and global health status (42 vs 9%: p = 0.003). In addition, the difference in averaged on-study changes for months 1-3 significantly favoured patients with a clinical benefit for those domains (except fatigue), as well as for role functioning, social functioning, pain, constipation and financial problems (Table 4, Figures 4-5). For example, after 3 months of treatment the change in mean score on the pain sub-scale was an improvement of -4.4 points on average for those with clinical benefit compared to a deterioration of 6.1 in those who had progressed by 3 months (p = 0.003).

Discussion

Anastrozole was associated with clinical benefit at 3 months in 44% of patients with metastatic or recurrent ER and/PR positive endometrial cancer. Responses were durable in a small subset of patients and treatment was well tolerated in most patients. Importantly we also demonstrated that those patients who were categorized as having a clinical benefit based on RECIST response or stable disease at 3 months also had significant improvements in global QoL as well as several specific QoL domains compared to baseline scores, supporting the notion

that these patients did indeed derive “clinical benefit”. Reductions in the symptoms of pain and fatigue (at 2 months) that can cause women significant distress were also seen, supporting the clinical benefit end-point based on the standard criteria of RECIST response /stable disease. The response rate of 7% (95% CI: 3-15%) is relatively low compared to other studies, but to the best of our knowledge this is the largest phase 2 trial of an aromatase inhibitor reported in a well-defined subset of patients with metastatic EC that were ER and/or PR positive and had no prior hormonal therapy. The fact that there was evidence of clinical benefit in almost half the patients, with an associated improvement in HRQOL, underscores the importance of also including HRQOL endpoints in phase 2 trials to support the primary endpoint. “Clinical benefit” is now widely used as an endpoint in phase 2 trials and is based on the combination of disease stabilisation and RECIST response and arguably HRQOL data should also be collected to back this up.

Importantly, anastrozole was well tolerated and one of the observations in this study was the apparently lower incidence of aromatase inhibitor (AI)-induced musculoskeletal events in women with EC cancers compared to these adverse effects reported by women with breast cancers. We speculated if this might be due to a lesser effect of the treatment in the women in our study who were overweight or obese. However, no clear difference in the clinical benefit rate was seen between normal weight and obese women. Studies in women on adjuvant aromatase inhibitors for early breast cancer reported 40-70% of treatment related arthralgia with 20-30% of patients stopping treatment early because of adverse effects [18-20]. In this study no patient stopped anastrozole for adverse

effects. Given that many patients with endometrial cancer are older and frequently have co-morbidities related to obesity, the option of using a relatively non-toxic systemic treatment such as hormonal therapy is highly desirable.

It is not clear why the responses to aromatase inhibitors in endometrial cancer, are so much lower than in ER positive breast cancer. There are a number of possible explanations including the high rate of alterations leading to overactivation of the PI3 kinase/AKT/MTOR pathway in Type 1 endometrioid endometrial cancers compared with breast cancers. These can lead to cross talk with other signalling pathways which can cause resistance to aromatase inhibitors[21, 22]. For example, activation of the AKT pathway due to alterations in PI3K, PTEN or AKT have been reported in over 90% of endometrial cancers. This can have variable affects on ER α transcriptional activity as well as blunting PR action in endometrial cancer. In addition, it is reported that ER and PR expression can be silenced by DNA methylation in endometrial cancer[23].

Combining agents that disrupt ER signalling such as aromatase inhibitors with PI3K/AKT/MTOR pathway inhibitors could potentially be synergistic and would be a rational combination to investigate in endometrial cancer. Of interest, more recent trials have reported promising results with the combination of an aromatase inhibitor and a MTOR inhibitor in endometrial cancer to try and overcome the effect of this pathway on endocrine resistance. Slomovitz et al reported a response rate of 35% in 35 patients treated with the combination of Letrozole and Everolimus, including 9 complete responses[11]. An non-comparative randomised phase 2 trial (GOG 3007) is aimed to determine the outcomes of this combination as well as a control arm of alternating tamoxifen

and medroxyprogesterone acetate with women with recurrent endometrial cancer. They recently reported a response rate of 24% and a PFS of 6.4 months with Letrozole/Everolimus, and a response rate of 22% with a PFS of 3.8 months with alternating tamoxifen/medroxyprogesterone. The response rate was even higher at 53% with Letrozole/Everolimus in chemo-naïve patients, similar to the trend seen in our study[24]. In addition, on-going trials are examining the efficacy of combined aromatase inhibitors and CDK 4/6 inhibitors in endometrial cancer.

The strengths of our trial include that women were selected based on having the target for the endocrine therapy present in their tumour, namely ER and/or PR. However, a limitation is that this was based predominantly on assessment of archival specimens taken at the time of initial diagnosis rather than metastatic sites. We do not know if ER and/or PR expression was still present in metastatic tumours. This may be important as a recent study reported that ER and PR median H-scores were significantly lower in metastases than in matched primary endometrioid endometrial cancers[25]. In addition, they found a higher frequency of hormone receptor gene promoter hypermethylation in metastases compared to matched primary tumours. These changes in hormone receptor expression in metastases may have important implications for treatment and argues for attempting to biopsy tumours prior to any hormonal therapy.

It was of interest that the degree of benefit from anastrozole did not seem to correlate strongly with the degree of ER and/or PR expression based on the Allred scores performed by the central laboratory. However, a trend was seen for those with lower PR expression to benefit less. It was also noteworthy that 3

cases were found to have no significant ER or PR expression at central testing. However, in some cases a significant degree of heterogeneity of staining was noted, likely related to fixation artefact in the slides sent centrally. It has also been described that there can be significant technical and inter-observer variability between laboratories performing immunohistochemistry[26]. This highlights the value of central testing and confirmation of biomarker testing being used to select patients for treatment with novel targeted therapies.

Although objective response rates were low, almost half of the patients derived clinical benefit from treatment with anastrozole as a single agent. However, the single-arm design of our study is a limitation to being sure about the PFS time seen in our study, since PFS may be more reliably assessed in randomised studies. It would be rational to continue to explore combining aromatase inhibitors with agents that inhibit other signal transduction pathways such as PI3K/AKT/MTOR as well as CDK4/6 in patients with ER/PR positive endometrial cancers. These are being considered in the PARAGON2 trial, with a greater emphasis and effort to collection of tumour specimens where possible at study entry and progression.

Figure legends

Figure 1: Study consort diagram

Figure 2: Clinical outcomes. (a) Progression-free survival for ITT population (n=84). (b) Progression-free survival for ITT population stratified by treatment free interval (n=78). (c) Progression-free survival beyond three month assessment point for those patients who gained clinical benefit (n=36), defined as complete, partial or stable disease at three month assessment.

Figure 3: Individual swimmers plot including subjects who received treatment (n=83) colour coded by tumour grade: depicting time on treatment as well as time of partial response and/or progression

Figure 4: QLQ-C30 functioning domains and Global Health averaged on-study changes for months 1-3 by Clinical benefit status at 3 months.

Figure 5: QLQ-C30 symptom domains and financial problems averaged on-study changes for months 1-3 by Clinical benefit status at 3 months.

468 Table 1: Clinical Characteristics of 84 participating patients
469

		N	%
Age	Mean = 68	Range (36-89)	
ECOG performance status	0	40	48
	1	35	42
	2	9	11
Body Mass Index	Normal (<25)	26	31
	Overweight (25-30)	11	13
	Obese (>30)	41	49
	Missing	6	7
Tumour grade	1	21	25
	2	33	39
	3	24	29
	Not available	6	7
FIGO stage	1	28	33
	2	13	16
	3	22	26
	4	15	18
	unknown	6	7
Prior chemotherapy	Yes	42	50
	No	42	50
Prior radiotherapy	Yes	56	67
	No	28	33
Treatment free interval†	<6 months	26	33
	6-12 months	13	17
	>12 months	39	50
IHC status	ER+/PR-	28	33
	ER+/PR+	52	62
	ER-/PR+	4	5

†Time since previous surgery/chemotherapy. 4 patients had no prior therapy, 2 had only Radiotherapy for which the date was not collected on the CRF.

470

Table 2: Association between ER and PR analysis and 3-month clinical benefit rate

Biomarker	Result	Number	Clinical Benefit at 3 months = Yes: n (%)	P-value
Local ER/PR testing		82		Fisher's exact test
	ER+/PR-	27	12 (44.4%)	0.236
	ER+/PR+	51	24 (47.1%)	
	ER-/PR+	4	0 (0%)	
Central ER/PR testing		53		Conditional Binomial Exact test
ER proportion	<5	17	6 (35.3%)	0.240
	5+	36	19 (52.8%)	
ER intensity	<3	35	17 (48.6%)	0.798
	3+	18	8 (44.4%)	
ER Total	<7	17	7 (41.2%)	0.558
	7+	36	18 (50%)	
PR proportion	<4	20	6 (30%)	0.056
	4+	33	19 (57.6%)	
PR intensity	<3	33	15 (45.5%)	0.743
	3+	20	10 (50%)	
PR total	<6	19	6 (31.6%)	0.097
	6+	34	19 (55.9%)	

477 Table 3: Toxicity, data were available from 82 patients
478

Toxicity	Grade	N	%
Anorexia	1	18	22
	2	10	12
Headaches	1	12	15
Nausea	1	22	27
	2	10	12
Fatigue	1	33	40
	2	17	21
	3	7	9
Vomiting	1	10	12
	2	6	7
Hypercholesteremia	1	1	1
	2	1	1
Alopecia	1	5	6
Hot flushes	1	35	43
	2	1	1
Rash	1	5	6
	2	1	1
Arthralgia	1	32	39
	2	7	9
Vaginal dryness	1	9	11

479
480

481 Table 4: QLQ-C30 averaged on-study changes for months 1-3 by Clinical Benefit
482 status at 3-months. Positive numbers represent improvements for the
483 functioning domains and Global Health, deterioration for the symptom domains
484 and financial problems.
485

Domain	Clinical benefit at 3 months (N=34)		Progressed by 3 months (N=35)		Clinical benefit - Progressed	
	Mean (95% CI)	P*	Mean (95% CI)	P*	Difference (95% CI)	P†
Physical Funct.	0.8 (-2.9 - 4.5)	0.656	-2.7 (-6.7 - 1.2)	0.172	4.6 (-0.7 - 9.8)	0.088
Role Funct.	3.8 (-5.5 - 13.0)	0.413	-4.8 (-10.1 - 0.4)	0.068	9.4 (0.6 - 18.3)	0.037
Emotional Funct.	7.7 (3.3 - 12.1)	0.001	0.0 (-3.8 - 3.8)	0.983	7.5 (2.8 - 12.3)	0.002
Cognitive Funct.	7.1 (1.2 - 13.0)	0.019	-5.6 (-12.3 - 1.0)	0.094	12.4 (4.4 - 20.3)	0.003
Social Funct.	10.5 (0.8 - 20.2)	0.036	-1.9 (-8.1 - 4.3)	0.537	12.8 (3.1 - 22.6)	0.011
Global Health	8.4 (4.1 - 12.7)	<0.001	-4.0 (-9.8 - 1.7)	0.158	12.8 (6.6 - 19.0)	<0.001
Fatigue	-4.0 (-11.4 - 3.4)	0.275	3.9 (-2.5 - 10.2)	0.226	-7.4 (-16.0 - 1.1)	0.087
Nausea Vomiting	0.9 (-2.7 - 4.5)	0.614	0.2 (-8.6 - 9.1)	0.957	-3.9 (-10.9 - 3.0)	0.260
Pain	-4.4 (-10.0 - 1.2)	0.120	6.1 (0.7 - 11.5)	0.028	-10.6 (-17.3 - -3.9)	0.003
Dyspnoea	1.5 (-5.7 - 8.6)	0.678	-0.6 (-7.7 - 6.4)	0.855	-0.6 (-9.8 - 8.7)	0.901
Insomnia	-2.5 (-10.6 - 5.7)	0.544	1.3 (-5.2 - 7.7)	0.691	-4.5 (-12.8 - 3.9)	0.290
Appetite Loss	-3.6 (-10.6 - 3.4)	0.303	3.0 (-5.9 - 11.9)	0.497	-5.2 (-15.0 - 4.6)	0.296
Constipation	-5.7 (-12.7 - 1.3)	0.107	2.5 (-7.9 - 12.9)	0.623	-9.2 (-18.2 - -0.2)	0.045
Diarrhoea	-2.6 (-10.2 - 4.9)	0.487	0.2 (-6.4 - 6.8)	0.961	-1.4 (-9.0 - 6.1)	0.703
Financial Problems	-8.8 (-15.7 - -2.0)	0.013	2.1 (-3.1 - 7.3)	0.425	-11.6 (-17.9 - -5.3)	<0.001

486

487 *Paired t-test

488 †Based on regression model including adjustment for baseline

489

CONFLICT OF INTEREST DISCLOSURES

Dr. Philip Beale reports receiving personal fees from Astrazeneca, outside of the submitted work.

Dr. Rosemary Lord reports receiving personal fees from advisory work for Astra Zeneca and Tesaro, outside of the submitted work.

Dr. Katrin Sjoquist reports receiving other funding from IPSEN, other funding from AMGEN, other funding from Pfizer, and personal fees from Merck, outside of the submitted work.

Dr. Andrew Clamp reports receiving a separate grants and non-financial support from AstraZeneca, outside of the submitted work; .

Dr. Michael Friedlander reports non-financial support from Astra Zeneca in the form of provision of Anastrozole for patients on the study. He also reports receiving personal fees from ASTRA ZENECA, personal fees from MSD, and personal fees from LILLY, outside of the submitted work.

The other authors have nothing to disclose.

AUTHOR CONTRIBUTIONS

Conception and design: Linda Mileshtkin, Rachel O'Connell, Katrin Sjoquist, Michael Friedlander

Collection and assembly of data: Linda Mileshtkin, Richard Edmondson, Katrin Sjoquist, John Andrews, Rema Jyothirmayi, Philip Beale, Tony Bonaventura, Jeffrey Goh, Marcia Hall, Andrew Clamp, John Green, Rosemary Lord, Frederic Amant, Laura Alexander, Karen Carty, James Paul, James Scurry, David Millan, Steven Nottley, Michael Friedlander

Data analysis and interpretation: Linda Mileshtkin, Richard Edmondson, Rachel O/Connell, David Millan, Michael Friedlander

Manuscript writing: Linda Mileshtkin, Richard Edmondson, Rachel O/Connell, David Millan, Michael Friedlander

Contribution to and final approval of manuscript: all authors

REFERENCES

- [1] Cote ML, Ruterbusch JJ, Olson SH, Lu K, Ali-Fehmi R. The Growing Burden of Endometrial Cancer: A Major Racial Disparity Affecting Black Women. *Cancer Epidemiol Biomarkers Prev.* 2015;24:1407-15.
- [2] Lentz SS, Brady MF, Major FJ, Reid GC, Soper JT. High-dose megestrol acetate in advanced or recurrent endometrial carcinoma: a Gynecologic Oncology Group Study. *J Clin Oncol.* 1996;14:357-61.
- [3] Garrett A, Quinn MA. Hormonal therapies and gynaecological cancers. *Best Pract Res Clin Obstet Gynaecol.* 2008;22:407-21.
- [4] Decruze SB, Green JA. Hormone therapy in advanced and recurrent endometrial cancer: a systematic review. *Int J Gynecol Cancer.* 2007;17:964-78.
- [5] Pautier P, Vergote I, Melichar B, Kutarska E, Hall G, Lisysanskaya A. A Phase 2, Randomized, Open-Label Study of Urosustat Versus Megestrol Acetate in Advanced Endometrial Cancer. *Int J Gynecol Cancer.* 2017;27:258-66.
- [6] Barker LC, Brand IR, Crawford SM. Sustained effect of the aromatase inhibitors anastrozole and letrozole on endometrial thickness in patients with endometrial hyperplasia and endometrial carcinoma. *Curr Med Res Opin.* 2009;25:1105-9.
- [7] Bellone S, Shah HR, McKenney JK, Stone PJ, Santin AD. Recurrent endometrial carcinoma regression with the use of the aromatase inhibitor anastrozole. *Am J Obstet Gynecol.* 2008;199:e7-e10.
- [8] Rose PG, Brunetto VL, VanLe L, Bell J, Walker JL, Lee RB. A phase II trial of anastrozole in advanced recurrent or persistent endometrial carcinoma: a Gynecologic Oncology Group study. *Gynecol Oncol.* 2000;78:212-6.
- [9] Ma BB, Oza A, Eisenhauer E, Stanimir G, Carey M, Chapman W, et al. The activity of letrozole in patients with advanced or recurrent endometrial cancer and correlation with biological markers--a study of the National Cancer Institute of Canada Clinical Trials Group. *Int J Gynecol Cancer.* 2004;14:650-8.
- [10] Wheler JJ, Moulder SL, Naing A, Janku F, Piha-Paul SA, Falchook GS, et al. Anastrozole and everolimus in advanced gynecologic and breast malignancies: activity and molecular alterations in the PI3K/AKT/mTOR pathway. *Oncotarget.* 2014;5:3029-38.
- [11] Slomovitz BM, Jiang Y, Yates MS, Soliman PT, Johnston T, Nowakowski M, et al. Phase II study of everolimus and letrozole in patients with recurrent endometrial carcinoma. *J Clin Oncol.* 2015;33:930-6.
- [12] Bogliolo S, Gardella B, Dominoni M, Musacchi V, Cassani C, Zanellini F, et al. Effectiveness of aromatase inhibitors in the treatment of advanced endometrial adenocarcinoma. *Arch Gynecol Obstet.* 2016;293:701-8.
- [13] Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst.* 1993;85:365-76.
- [14] Cocks K, King MT, Velikova G, Martyn St-James M, Fayers PM, Brown JM. Evidence-based guidelines for determination of sample size and interpretation of the European Organisation for the Research and Treatment of Cancer Quality of Life Questionnaire Core 30. *J Clin Oncol.* 2011;29:89-96.
- [15] Fallowfield LJ, Leaity SK, Howell A, Benson S, Cella D. Assessment of quality of life in women undergoing hormonal therapy for breast cancer: validation of an

endocrine symptom subscale for the FACT-B. *Breast Cancer Res Treat.* 1999;55:189-99.

[16] Allred DC, Bustamante MC, Daniel CO, Gaskill HV, Cruz AB. Immunohistochemical analysis of estrogen in human breast carcinomas. Evaluation of 130 cases and review of the literature regarding concordance with biochemical assay and clinical relevance. *Arch Surg.* 1990;125:107-13.

[17] Korn EL, Graubard BI. *Analysis of health surveys*: John Wiley & Sons; 2011.

[18] Crew KD, Greenlee H, Capodice J, Raptis G, Brafman L, Fuentes D, et al. Prevalence of joint symptoms in postmenopausal women taking aromatase inhibitors for early-stage breast cancer. *J Clin Oncol.* 2007;25:3877-83.

[19] Goss PE, Ingle JN, Pritchard KI, Robert NJ, Muss H, Gralow J, et al. Extending Aromatase-Inhibitor Adjuvant Therapy to 10 Years. *New England Journal of Medicine.* 2016;375:209-19.

[20] Presant CA, Bosserman L, Young T, Vakil M, Horns R, Upadhyaya G, et al. Aromatase inhibitor-associated arthralgia and/ or bone pain: frequency and characterization in non-clinical trial patients. *Clin Breast Cancer.* 2007;7:775-8.

[21] Pavlidou A, Vlahos NF. Molecular alterations of PI3K/Akt/mTOR pathway: a therapeutic target in endometrial cancer. *ScientificWorldJournal.* 2014;2014:709736.

[22] Slomovitz BM, Coleman RL. The PI3K/AKT/mTOR pathway as a therapeutic target in endometrial cancer. *Clin Cancer Res.* 2012;18:5856-64.

[23] Nieves-Neira W, Kim JJ, Matei D. Hormonal strategies in gynecologic cancer: Bridging biology and therapy. *Gynecol Oncol.* 2018;150:207-10.

[24] Slomovitz BM, Filiaci VL, Coleman RL, Walker JL, Fleury AC, Holman LL, et al. GOG 3007, a randomized phase II (RP2) trial of everolimus and letrozole (EL) or hormonal therapy (medroxyprogesterone acetate/tamoxifen, PT) in women with advanced, persistent or recurrent endometrial carcinoma (EC): A GOG Foundation study. *Annual Meeting of the Society of Gynecologic Oncology.* New Orleans, LA2018.

[25] Bartosch C, Monteiro-Reis S, Vieira R, Pereira A, Rodrigues M, Jeronimo C, et al. Endometrial Endometrioid Carcinoma Metastases Show Decreased ER-Alpha and PR-A Expression Compared to Matched Primary Tumours. *PLoS One.* 2015;10:30134969.

[26] Sinn HP, Schneeweiss A, Keller M, Schlombs K, Laible M, Seltz J, et al. Comparison of immunohistochemistry with PCR for assessment of ER, PR, and Ki-67 and prediction of pathological complete response in breast cancer. *BMC Cancer.* 2017;17:124.