Single-cell profiling of myeloid cells in glioblastoma across species and disease stage reveals macrophage competition and specialization

Ana Rita Pombo Antunes^{1,2,*}, Isabelle Scheyltjens^{1,2,*}, Francesca Lodi^{3,4}, Julie Messiaen⁵, Asier Antoranz⁵, Johnny Duerinck⁶, Daliya Kancheva⁷, Liesbet Martens^{8,9,10}, Karen De Vlaminck^{1,2}, Hannah Van Hove^{1,2}, Signe Schmidt Kjølner Hansen^{1,2}, Francesca Maria Bosisio¹¹, Koen Van der Borgh⁷, Steven De Vleeschouwer^{12,13}, Raf Sciot¹¹, Luc Bouwens¹⁴, Michiel Verfaillie¹⁵, Niels Vandamme^{16,17}, Roosmarijn E. Vandenbroucke^{18,19}, Olivier De Wever^{20,21}, Yvan Saeys^{16,17}, Martin Guillems^{8,9}, Conny Gysemans²², Bart Neyns²³, Frederik De Smet⁵, Diether Lambrechts^{3,4}, Jo A. Van Ginderachter^{1,2,#}, Kiavash Movahedi^{1,2,#,\$}

¹ Myeloid Cell Immunology Lab, VIB Center for Inflammation Research, Brussels, Belgium

² Lab of Cellular and Molecular Immunology, Vrije Universiteit Brussel, Brussels, Belgium

³ Laboratory of Translational Genetics, Department of Human Genetics, KU Leuven, Leuven, Belgium

⁴ VIB-KU Leuven Center for Cancer Biology, VIB, Leuven, Belgium

⁵ Laboratory for Precision Cancer Medicine, Translational Cell & Tissue Research Unit, Department of Imaging & Pathology, KU Leuven, Leuven, Belgium

⁶ Department of Neurosurgery, UZ Brussels, Brussels, Belgium

⁷ Janssen Pharmaceuticals NV, Beerse, Belgium

⁸ Laboratory of Myeloid Cell Ontogeny and Functional Specialization, VIB Center for Inflammation Research, Ghent, Belgium

⁹ Department of Biomedical Molecular Biology, Ghent University, Ghent, Belgium

¹⁰ Laboratory of Myeloid Cell Heterogeneity and Function, VIB Center for Inflammation Research, Ghent, Belgium

¹¹ Translational Cell & Tissue Research Unit, Department of Imaging & Pathology, KU Leuven, Leuven, Belgium

¹² Department of Neurosurgery, UZ Leuven, Leuven, Belgium

¹³ Laboratory for Experimental Neurosurgery and Neuroanatomy, Department of Neurosciences and Leuven Brain Institute (LBI), KU Leuven, Leuven, Belgium

¹⁴ Cell Differentiation Lab, Vrije Universiteit Brussel, Brussels, Belgium

¹⁵ Department of Neurosurgery, Europe Hospitals Saint Elisabeth, Ukkel, Belgium

¹⁶ Data Mining and Modeling for Biomedicine, VIB Center for Inflammation Research, Ghent, Belgium

¹⁷ Department of Applied Mathematics, Computer Science and Statistics, Ghent University, Ghent, Belgium

¹⁸ VIB Center for Inflammation Research, Ghent, Belgium

¹⁹ Department of Biomedical Molecular Biology, Ghent University, Ghent, Belgium

²⁰ Cancer Research Institute Ghent (CRIG), Ghent University, Ghent, Belgium

²¹ Laboratory for Experimental Cancer Research, Ghent University, Ghent, Belgium

²² Clinical and Experimental Endocrinology (CEE), KU Leuven, Leuven, Belgium

²³ Department of Medical Oncology, UZ Brussels, Brussels, Belgium

* contributed equally

Co-senior author

\$ Correspondence: kiavash.movahedi@vub.be

Abstract

Glioblastomas are aggressive primary brain cancers that recur as therapy-resistant tumors. Myeloid cells control glioblastoma malignancy, but their dynamics during disease progression remain poorly understood. Here, we employed single-cell RNA sequencing and CITE-Seq to map the glioblastoma immune landscape in newly diagnosed and recurrent patients and in mouse tumors. This revealed a large and diverse myeloid compartment, with dendritic cell and macrophage populations that were conserved across species and were dynamic across disease stages. Tumor-associated macrophages (TAMs) consisted of microglia- or monocyte-derived populations, with both exhibiting additional heterogeneity, including subsets with conserved lipid and hypoxic signatures. Microglia- and monocyte-derived TAMs (Mo-TAMs) were self-renewing populations that competed for space and could be depleted via CSF1R blockade. Microglia-derived TAMs were predominant in newly diagnosed tumors but were outnumbered by Mo-TAMs upon recurrence, especially in hypoxic tumor environments. Our results unravel the glioblastoma myeloid landscape and provide a framework for future therapeutic interventions.

Introduction

The success of immune checkpoint inhibition has revealed the immense potential of mobilizing anti-tumor immunity in cancer treatment. This also holds promise for primary brain tumors, especially since the field is gradually uncovering the intricacies of brain immunology. Recent work has shown that the brain and its border regions contain a diverse immune compartment¹⁻³, combined with a lymphatic system that drains brain-derived antigens and assists in mounting efficient immune responses⁴. However, glioblastoma (GBM), the most aggressive primary brain tumor⁵, remains refractory to immunotherapy. The standard of care for newly diagnosed GBM patients is neurosurgical resection followed by adjuvant radiotherapy and temozolomide chemotherapy, but cancer recurrence and death are almost inevitable. Therapy resistance is thought to in part result from the molecular-genetic heterogeneity of GBM cancer cells, which exist in multiple cellular states or subtypes^{6, 7}. Additionally, anti-tumor immunity is impaired, so a better understanding of the GBM immune compartment is critical for delineating and reprogramming these suppressive immune circuits⁸. Furthermore, the GBM immune landscape may evolve upon treatment and subsequent recurrence, an important feature that remains incompletely explored.

Macrophages and dendritic cells (DCs) are important components of the GBM immune landscape that affect both anti-tumor immunity and intrinsic tumor growth characteristics, including angiogenesis and invasion⁹. Brain DCs are known to initiate T-cell activation during autoimmune disease of the central nervous system (CNS)¹⁰, but their presence and heterogeneity within GBM remains unclear. The homeostatic brain contains a heterogenous population of macrophages, consisting of microglia in the parenchyma¹¹, and border-associated macrophages or BAMs in the brain's border regions¹². While microglia are yolk-sac-derived, BAMs have a distinct transcriptomic profile and some populations have a bone marrow (BM) origin^{1, 13}. Similarly, GBM tumors have been shown to contain tumor-associated macrophages or TAMs that are either microglia- or BM-derived¹⁴⁻¹⁷. However, it is unknown whether GBM TAM ontogeny is altered depending on the disease stage and whether ontogenetically distinct TAM subsets are found in different tumor microenvironments and/or exhibit distinctive functionalities. Furthermore, the degree of GBM TAM heterogeneity within the microglia- or BM-derived TAM populations remains incompletely understood. To provide insights into these matters, we performed single-cell RNA sequencing (scRNA-Seq) and cellular indexing of transcriptomes and epitopes by sequencing (CITE-Seq)¹⁸ on the GBM immune landscape from human newly diagnosed and recurrent tumors, and in a widely used mouse model of GBM. Our results now delineate the heterogeneity of the GBM myeloid compartment across species and disease stage.

Results

scRNA-Seq analysis reveals a dynamic immune landscape in human and mouse glioblastoma tumors.

We performed scRNA-Seq on fresh isocitrate dehydrogenase (IDH) WT tumors obtained from seven newly diagnosed (ND) (ND1-7) and four recurrent (R1-4) GBM patients (Fig. 1a). All ND patients were treatment-naïve, while recurrent patients had previously been treated with surgical resection and adjuvant radio- and chemotherapy (Supplementary Table 1). Patients R1-3 also received one injection of nivolumab 24 hours prior to surgery, but it is unlikely that this would significantly have altered the tumor immune compartment (TIC) within this short time frame. The ND and recurrent samples were combined in two separate datasets (ND1-7 and R1-4). For R4 we included a highly necrotic core as a separate sample (R4 n.c.). After combining the individual samples and excluding CD45⁻ cells, we aimed to remove batch effects, without obscuring relevant cellular heterogeneity. The harmony algorithm¹⁹ efficiently integrated the datasets and accurately grouped the distinct cell types that were common to all tumors (Extended data Fig. 1a,b). Unsupervised clustering and UMAP projections were performed on 21,303 immune cells from ND tumors (Fig. 1b) or 42,870 immune cells from recurrent tumors (Fig. 1c). To perform cross-species comparisons and assess how mouse GBM models compare to the human disease, we performed scRNA-Seq on CD45⁺ immune cells from orthotopic GL261 tumors. GL261 tumors were also grown in CCR2-KO mice, to examine monocyte contribution to the TIC, as CCR2 deficiency significantly reduces blood monocyte numbers²⁰. Mouse tumors were micro-dissected to minimize contamination with healthy brain tissue. Three repeats were performed for each genotype, resulting in a total of 27,276 WT and 22,588 CCR2-KO immune cells that were combined in a single harmony-corrected mouse dataset (Fig. 1d). Human and mouse immune cell clusters were identified based on previously described gene expression patterns^{1, 21}. We correlated the immune subsets in the two species and compiled a list of signature genes for their identification (Fig. 1e,f, Supplementary Table 2-4). Most immune cells in ND human tumors were macrophages (82-97%), followed by T cells (2-20%) (Fig. 1b). TAMs consisted of two main populations grouped as TAM-1 (orange) and TAM-2 (red). The main difference between these two populations lies at the level of their ontogeny, as will be shown later. Interestingly, recurrent human tumors showed a more diverse immune compartment (Fig. 1c). We observed an increase in the percentage of lymphocytes, including T, NK and B cells and a greater contribution of monocytes (Extended Data Fig. 1c). R1, who had experienced a long progression-free survival after initial treatment, was an outlier in terms of its TIC, with 75% T cells and 5% TAMs. In the other recurrent tumors, TAMs were still a large population, but contrary to ND tumors, the TAM-1 subset outnumbered TAM-2 (Extended Data Fig. 1d).

The TIC of mouse GL261 tumors was diverse, with a large and heterogeneous lymphocyte population and a high percentage of monocytes and DCs (Fig. 1d). TAMs were the largest immune cell fraction and we again identified two main populations: TAM-1 and TAM-2, relating to their human counterparts. Tumors from WT and CCR2-KO mice showed comparable immune cell subsets, but the percentage of monocytes was strongly reduced in CCR2-KO tumors (Extended Data Fig. 1e).

Remarkably, the percentage of TAMs was not different between WT and CCR2-KO tumors. However, while in WT tumors TAM-1 outnumbered TAM-2, CCR2-KOs showed a relative decrease in TAM-1 (Extended Data Fig. 1f), suggesting that this subset relies on monocyte infiltration.

Single-cell analysis reveals the heterogeneity of intratumoral dendritic cells and identifies cDC2 subsets that are conserved in humans and mice.

To examine DC heterogeneity, we re-clustered human DCs from ND and recurrent tumors in a single dataset (Fig. 2a, Supplementary Table 5). This identified *XCR1*⁺*CLEC9A*⁺*CADM1*⁺ cDC1 and *FCER1A*⁺*CLEC10A*⁺*CD1C*⁺ cDC2 conventional DC subsets (Fig. 2b, Extended Data Fig. 2a). The *CCR7*⁺*LAMP3*⁺*SAMSN1*⁺ DCs in cluster 2 (Fig. 2c) exhibited a gene signature reminiscent of migratory DCs (MigDCs)²². These cells did not express cDC1 or cDC2 marker genes, suggesting that this form of activation overrules the cDC1/cDC2 transcriptional programs. Cluster 3 was identified as *IL3RA*⁺*LILRA4*⁺ plasmacytoid DCs (pDCs) (Fig. 2d). Cluster 4 expressed *CD5*, *AXL* and *SIGLEC6* (Fig. 2d) and corresponds to pre-DCs^{21, 23-25}. cDC2s exhibited additional heterogeneity. cDC2s in cluster 6 were *CD1A*⁺*CD207*⁺*CEACAM3*⁺ (Fig. 2d), while clusters 7,8 exhibited upregulation of *IL1B*, *ZEB2* and *TSPO*. Furthermore, cluster 8 cDC2s expressed monocyte-related genes (*FCN1*, *VCAN*, *CD14*). This is in agreement with recent observations of a human cDC2 subset expressing monocyte genes^{21, 24, 25,26}. Cells in cluster 5 expressed macrophage genes (*MAFB*, *C1QA*) (Extended Data Fig. 2a) and were considered TAMs. However, this shows that cDC2s and macrophages share part of their transcriptional identity. We termed the two main cDC2 subsets cDC2a (cluster 6) and cDC2b (cluster 8), in line with the recent nomenclature²⁶. A fraction of cDC2b also showed an IFN-induced signature (cluster 9, Fig. 2d). In all tumors, cDC2s were the most abundant subset (Extended Data Fig. 2b).

Reclustering of intratumoral mouse DCs (Fig. 2e, Supplementary Table 6) revealed subsets reminiscent of cDC1s, cDC2s, MigDCs and pDCs (Fig. 2f,g, Extended Data Fig. 2c). Many of the top differentially expressed (DE) genes were shared across species (Fig. 2h, Extended Data Fig. 2d,e, Supplementary Table 7). Mouse cDC2s also exhibited heterogeneity, with cluster 4 being *Mgl2*⁺*Cd9*⁺*Epcam*⁺ DCs, while cells in cluster 6 were *Cd209a*⁺ and expressed monocyte-related genes (*Ccr2*, *Lyz2*, *Ly6c2*) (Fig. 2i, Extended Data Fig. 2c). This subset was not significantly altered in CCR2-KO tumors (Extended Data Fig. 2f), arguing against a monocyte origin. Based on their shared gene signatures, mouse clusters 4 and 6 emerge as the putative counterparts for human cDC2a and cDC2b, respectively.

Both in human and mouse tumors we observed cDC1 and cDC2 clusters expressing cell cycle genes (e.g. *STMN1*, *MKI67*) (Extended Data Fig. 2a,c). In human recurrent tumors 15±5% of cDC1s and 6±2% of cDC2s associated with a proliferation cluster (Fig. 2j). In mice, this was further increased both for cDC1s and cDC2s (Fig. 2k). To confirm that cDCs can proliferate in GBM tumors, mice were injected with bromodeoxyuridine (BrdU) and two hours later BrdU incorporation was assessed. Tumor cDC1 and cDC2s were identified via flow cytometry (Extended Data Fig. 2g) and exhibited a clear incorporation of BrdU (Fig. 2l), suggesting that cDCs actively proliferate within GBM tumors.

Glioblastoma TAMs are derived from microglia or classical monocytes, which differentially contribute to the macrophage pool in newly diagnosed versus recurrent tumors.

Monocytes and macrophages were re-clustered for human ND (Fig. 3a), recurrent (Fig. 3b) and mouse (Fig. 3c) tumors. The majority of human tumor monocytes were of the classical subset ($FCN1^+VCAN^+CD14^{int}FCGR3A^{lo}$) (Fig 3d,e). Recurrent tumors also contained a small cluster of non-classical monocytes ($FCN1^+CDKN1C^+VCAN^+CD14^{lo}FCGR3A^{hi}$) (Supplementary Fig. 1a,b). Similarly, mouse tumors mostly contained classical ($Ly6c2^+Chil3^+Ccr2^+$) and a small percentage of non-classical ($Ear2^+Ace^+$) monocytes (Fig. 3f, Supplementary Fig. 1c,d). TAMs downregulated $FCN1/VCAN$ (human) or $Ly6c2/Chil3$ (mouse) and upregulated macrophage markers such as $C1QA$. A fraction of TAMs expressed cell cycle genes ($STMN1$, $MKI67$) indicating proliferation (Fig. 3d-f). Importantly, TAMs consisted of two main populations, encircled by blue and red dotted lines (corresponding to TAM-1 and TAM-2 in Fig. 1, respectively). Comparing these two groups suggested a distinction at the level of ontogeny. The blue encircled clusters expressed genes that were previously associated with monocyte-derived brain macrophages^{1, 14, 15, 27-29} (e.g. $TGFBI$, $CLEC12A$, $FXYD5$) (Fig 3g,h, Supplementary Fig. 1e). These cells were termed Mo-TAMs. In contrast, Mg-TAMs expressed microglial signature genes (e.g. $SALL1$, $TMEM119$, $P2RY12$), suggesting a microglial ontogeny. While multiple ontogeny-defining genes were shared across species (Fig. 3i and Supplementary Fig. 1f), there were important exceptions. For example, $Cst7$, $Hexb$ and $Sparc$ were highly differential between Mg-TAMs and Mo-TAMs in mice, but not humans. Conversely, $TMIGD3$, $APOC2$, and $SCIN$, were Mg-TAM markers restricted to humans. Interestingly, while Mg-TAMs formed the major TAM fraction in ND patients, the opposite was observed in recurrent tumors (Fig. 3j and Supplementary Fig. 1g), revealing a switch in TAM ontogeny. In mouse tumors the majority of macrophages were Mo-TAMs. This fraction was significantly reduced in CCR2-KO mice, in line with the proposed monocytic origin for this subset (Fig. 3j and Supplementary Fig. 1h).

To further confirm the differential ontogeny of GBM TAMs, we relied on $Cx3cr1^{CreER}:R26-YFP$ mice for fate mapping of microglia³⁰. Microscopy confirmed that a fraction of F4/80⁺ TAMs expressed YFP, indicating a microglial ontogeny (Fig. 4a). As both classical and non-classical monocytes were detected in tumors, we investigated whether each could act as precursors for Mo-TAMs. Blood monocytes were purified from $Cx3cr1^{Gfp/+}$ mice and classical (1×10^6) or non-classical (7×10^5) monocytes were adoptively transferred in CCR2-KO mice harboring GL261 tumors (Fig. 4b). For both groups, GFP⁺ cells were detected in tumors five days post transfer (Fig. 4c), while being absent in non-injected mice (Fig. 4d). However, only classical monocytes were differentiating into macrophages, as indicated by their expanded scatter profile, which was also accompanied by a downregulation of Ly6B and upregulation of MHCII (Fig. 4c). This indicates that classical monocytes are the precursors of Mo-TAMs. To corroborate this conclusion, we performed adoptive transfers of classical (7×10^5) or non-classical (7×10^5) monocytes in WT mice, and in NR4A1-KO mice, which have a compromised non-classical monocyte compartment³¹. This confirmed that only classical monocytes were differentiating into tumor

macrophages, as indicated by an enlarged scatter profile and upregulation of F4/80 and MHCII, both in WT (Extended Data Fig. 3a-d) and NR4A1-KO (Extended Data Fig. 3e-h) mice.

To investigate the spatial distribution of Mg-TAMs and Mo-TAMs in human GBM, we performed multiplex immunohistochemistry using the MILAN method, mapping cancer cells (SOX2⁺), blood vessels/endothelial cells (CD34⁺) and macrophages (CD68⁺) (Fig. 4e). TMEM119^{hi} cells (defined as 1 SD higher than the overall average) were considered as Mg-TAMs. Tumor sections were stained from 2 ND and 5 recurrent patients (Fig. 4e-h, Supplementary Fig. 2). This showed that CD68⁺TMEM119^{hi} cells were often observed in patches covering areas ranging from 16-50% throughout the whole tumor section, both in regions with high and low density of SOX2⁺ cancer cells. This suggests that Mg-TAMs and Mo-TAMs are found throughout the whole tumor.

Monocyte- and microglia-derived TAMs exhibit a spectrum of activation states.

Mo-TAMs showed a high degree of heterogeneity. In human recurrent tumors we identified six main subsets (Fig. 5a, Supplementary Table 8). The "Transitory Mo-TAMs" in cluster 1 still expressed several monocyte-related genes (e.g. *EREG*, *S100A6*, *LYZ*) and were lower for mature macrophage markers (e.g. *C1QA*, *IGF1*) (Fig. 5b,c), suggesting an ongoing monocyte-to-macrophage differentiation. Cluster 2 Mo-TAMs exhibited a gene signature that associated with phagocytosis and lipid metabolism³² (e.g. *GPNUMB*, *LGALS3*, *FABP5*) (Fig. 5b,d), which was also confirmed by Gene Ontology (GO) analysis (Supplementary Fig. 3a, Supplementary Table 9). We suggest that this signature may associate with TAMs that are actively phagocytosing cells, debris and/or lipid-rich molecules. Interestingly, cluster 3 Mo-TAMs exhibited a clear hypoxic and glycolytic gene expression profile as indicated by GO analysis, with signature expression of *BNIP3*, *ADAM8*, *MIF* and *SLC2A1* (Fig. 5b,e; Supplementary Fig. 3b). Remarkably, Mo-TAMs in cluster 4 (*SEPP1*^{lo} Mo-TAMs) expressed many microglial signature genes (e.g. *CX3CR1*, *BIN1*, *SCIN*) (Fig. 5b,c). Furthermore, GO analysis identified "synapse pruning" as the top enriched GO term (Supplementary Fig. 3c). This suggests that these Mo-TAMs have adopted a microglia- or "resident-like" phenotype. In contrast, macrophages in cluster 5 were low for microglial genes and showed high expression of *SEPP1* (*SEPP1*^{hi} Mo-TAMs), *SLC40A1*, *FOLR2*, *MRC1* and *RNASE1* (Fig. 5b, Extended Data Fig. 4a), genes that are often associated with an anti-inflammatory activation. To corroborate this, we relied on the study from Xue et al., which measured the transcriptional changes of human macrophages exposed to various *in vitro* stimuli³³. We took the top enriched genes from cluster 5 (Extended Data Fig. 4a) and assessed whether these are induced or repressed in the Xue et al. dataset. This revealed a clear glucocorticoid (GC)-induced signature (Extended Data Fig. 4b). Of note, none of these recurrent patients had received synthetic GCs since their initial diagnosis and treatment. Finally, Mo-TAMs in cluster 6 showed an IFN-induced signature (Extended Data Fig. 4c,d, Supplementary Fig. 3d).

Each of these Mo-TAM subsets were also observed in ND tumors (Extended Data Fig. 5a-f, Supplementary Table 10). However, in these samples *SEPP1*⁺ Mo-TAMs did not clearly segregate into a *SEPP1*^{lo} and *SEPP1*^{hi} subset. Upon comparing *SEPP1*⁺ Mo-TAMs from ND (cluster 5) and recurrent samples (cluster 4 and cluster 5), *SEPP1*^{hi} genes such as *SEPP1* and *FOLR2* were found to be

increased in recurrent tumors, while *BIN1* and *SCIN* (*SEPP1*^{lo} genes) were downregulated (Extended Data Fig. 5g). This suggests that the *SEPP1*^{hi} signature is enhanced in recurrent tumors.

Interestingly, most of the Mo-TAM heterogeneity seen in human tumors was recapitulated in mouse GL261 (Fig. 5f-h, Supplementary Table 11). This included Transitory Mo-TAMs in cluster 1 (Fig. 5g), which were significantly reduced in CCR2-KO tumors (p value = 0,0325) (Extended Data Fig. 5h, Extended Data Fig. 6a). Mouse tumors also contained Mo-TAM clusters with Phagocytic/lipid (Extended Data Fig. 6b), IFN and hypoxic (Extended Data Fig. 6c) gene signature, which were conserved across species (Extended Data Fig. 6d-g, Supplementary Table 12). While the hypoxic gene signature was highly conserved in humans and mice, one exception was *Arg1*, a top DE gene in mouse hypoxic Mo-TAMs (Extended Data Fig. 6c,h), but absent in humans. To confirm that Mo-TAMs with a hypoxic signature were associated with hypoxia, mice were injected with pimonidazole (PIMO) to visualize hypoxic microenvironments in combination with Arginase1 (ARG1) staining. This showed that ARG1⁺ TAMs were enriched in PIMO⁺ hypoxic tumor regions (Extended Data Fig. 6i). Finally, Mo-TAMs in cluster 5, which exhibited an enriched expression of *Sepp1*, *Ms4a7* and *Apoe* (Extended Data Fig. 6a), likely represent the mouse counterpart for human *SEPP1*⁺ Mo-TAMs, as suggested by cross-species comparison (Extended Data Fig. 6g).

To explore Mg-TAM heterogeneity, we first assessed the transcriptional changes that brain microglia undergo as they infiltrate the tumor. Therefore, scRNA-Seq was performed on CD45⁺ cells isolated from the surrounding brain tissue of GL261 tumor-bearing mice and trajectory analysis was performed on microglia and Mg-TAMs (Extended Data Fig. 7a). This revealed the downregulation of homeostatic signature genes and induction of Disease-Associated Microglia markers³⁴ (e.g. *Cst7*, *Apoe*, *Axl*). A downregulation of homeostatic genes in Mg-TAMs as compared to parenchymal microglia has recently also been reported in human GBM¹⁷. We identified additional cellular states that were superimposed upon Mg-TAMs. Both in ND, recurrent and mouse tumors, clusters were observed that exhibited an IFN or phagocytic/lipid signature (Fig. 5i-k, Extended Data Fig. 7b-d). We also identified a Mg-TAM cluster with a hypoxic signature in human recurrent and mouse tumors (Extended Data Fig. 7e,f). However, in both cases the fraction of hypoxic Mg-TAMs was considerably smaller than that of hypoxic Mo-TAMs, (Fig. 5l,m). Conversely, the fraction of IFN-signature Mo- and Mg-TAMs in these tumors was more comparable (Fig. 5l,m). Additionally, no hypoxic Mg-TAMs were detected in the ND dataset. This suggests that Mg-TAMs are outcompeted by Mo-TAMs in the hypoxic tumor niche.

Altered transcriptional states in TAM subsets from newly diagnosed versus recurrent tumors.

We performed DE analysis between TAM subsets from ND and recurrent patients, to assess for transcriptional changes that may correlate with disease-stage and/or drug treatment. The analysis was first performed at the single-cell level (Supplementary Table 13) and subsequently the mean normalized expression levels of selected genes were plotted for individual patients (Extended data Fig. 8a,b). We focused the analysis on Mg-TAMs, *SEPP1*⁺ Mo-TAMs and hypoxic Mo-TAMs. Notable genes that were enriched in recurrent vs. ND TAMs related to monocyte chemotaxis, interferon signaling and phagocytosis (Extended Data Fig. 8a), as also indicated by GO analysis (Extended Data Fig. 8c).

Remarkably, many of the genes that were enriched in ND tumors are known to be induced by GCs (e.g. *TSC22D3*, *RHOB*, *NR4A1*), which was also highlighted by GO analysis (Extended Data Fig. 8d). Therefore, at least part of this signature in the ND patients may result from GC drug treatment that they had received prior to surgery, and which was absent in the recurrent cohort (Supplementary Table 1). We also identified TAM-expressed genes that exhibited a high variation in expression across both ND and recurrent patients (Extended Data Fig. 8e). This revealed patient-specific gene expression variation in TAM subsets. The association of these genes with GBM patient survival was assessed using PRECOG, which offers a prognostic score by integrating 6 GBM expression datasets³⁵. Notable genes that are linked to reduced patient survival, included the cytokines *IL1B*, *CXCL8*, *IL6* and *SPP1* (Extended Data Fig. 8e).

CITE-Seq with extensive antibody panels provides broad protein validation and identifies key surface markers of TAM and DC subsets.

To provide protein validation for the gene expression signatures observed in GBM myeloid cells, we performed simultaneous quantification of RNA and cell surface proteins using the CITE-Seq approach¹⁸. A panel of 268 and 174 barcoded antibodies (Supplementary Table 14) were used for profiling human and mouse tumors, respectively. CITE-Seq was performed on the CD45⁺ immune compartment of 1 ND (ND8) and 2 recurrent (R2, R5) human GBM tumors, after which monocytes, TAMs and DCs were selected and combined in a single dataset (14.793 cells), followed by unbiased clustering of the transcriptomic data (Fig. 6a). Similarly, following CITE-Seq of CD45⁺ cells from WT (WT4) and CCR2-KO (KO4) GL261 tumors, monocytes, TAMs and DCs were selected and pooled (13.635 cells, Fig. 6b). Both for human and mouse tumors, all previously identified TAM and DC subsets were reobserved (Fig 6a,b Extended Data Fig. 9a,b), indicating that CITE-Seq implementation did not introduce a transcriptional distortion. Harmony-corrected UMAP projections based on protein data showed a segregation of most TAM and DC subsets (Fig. 6a,b, Extended Data Fig. 9c,d). Many gene-protein combinations showed an excellent correlation, both in human (Fig. 6c, Extended Data Fig. 9e,f) and mouse (Extended Data Fig. 9g,h). However, some genes that were not differential at the RNA level showed differential antibody staining (Extended Data Fig. 9e-g). This highlights that CITE-Seq in conjunction with large unbiased antibody panels can reveal important new surface markers that are not always obvious candidates based on transcript expression. Our analysis now provides an extensive list of protein markers that are differentially expressed in the distinct human TAM subsets (Fig. 6c,d, Supplementary Table 15). These include markers that capture monocyte-to-macrophage differentiation (e.g. CD36, CD64), markers that are differential between Mo-TAMs and Mg-TAMs (CD44/CD49d/CD9 vs. CD69/CD151/TMEM119) and markers that are enriched in specific Mo-TAM subsets, including *SEPP1*^{hi} Mo-TAM (FolateRB, CD206, CD209), hypoxic Mo-TAMs (CD184, CD354) and Phagocytic/Lipid Mo-TAMs (CD71, CD72). Subset-specific surface markers were also identified for mouse GBM TAMs (Extended Data Fig. 9h, Supplementary Table 16)

To further assess protein expression in DCs, these cell populations were selected and re-clustered separately (Fig 6.e,f), revealing key signature proteins for all of the previously identified DC

subsets, both in human (Fig. 6g, Supplementary Table 17) and mouse (Fig. 6h, Supplementary Table 18). This includes new surface markers for MigDCs (CD273 in human, CD200 in mouse). Interestingly, a fraction of MigDCs still exhibited cDC1 or cDC2-related proteins, while this was hardly observed at the transcript level (Extended Data Fig. 10a,b). This suggests that following migratory cDC activation and suppression of the cDC1/cDC2 transcriptional programs, some cDC1 or cDC2-related proteins may persist longer due to a slower protein turnover. Remarkably, mouse MigDCs also exhibited high MHCII protein staining, while having low *H2-Aa* transcript levels (Extended Data Fig. 10c). The distinction between the cDC2a and cDC2b subsets was also recapitulated at the protein level, both in human (Fig. 6i) (e.g. CD1c/CD1a/CD117 vs. CD14/CD64) and mouse (Extended Data Fig. 10d) (e.g. CD326/CD301b/CD117 vs. CD62L/Ly6C).

***Ex vivo* functional profiling of monocyte- and microglia-derived TAMs reveals a convergent angiogenic and T-cell suppressive capacity.**

To investigate the functionality of Mo- and Mg-TAMs, we assessed their ability to secrete cytokines, induce angiogenesis and modulate T-cell responses. A FACS-gating strategy was developed to sort TAM subsets from mouse tumors (Fig. 7a). Monocytes and TAMs were selected by gating CD45⁺CX3CR1⁺F4/80⁺ cells. Next, classical monocytes and differentiating macrophages (Transitory Mo-TAMs) were separated from mature TAMs based on their expression of Ly6B and MHCII. Within the Ly6B⁻ mature TAM gate, Mg-TAMs were distinguished from Mo-TAMs based on CD11b vs. CD45 expression. Microglia from the surrounding brain tissue could be selected using the same gating as for Mg-TAMs (Fig. 7a). Fate mapping in *Cx3cr1*^{CreER};R26-YFP mice confirmed that YFP⁺ brain microglia and YFP⁺ Mg-TAMs were contained in the CD45^{low/int} gate (Fig. 7b).

Our scRNA-Seq analysis showed that Mo-TAMs and Mg-TAMs exhibited a distinct cytokine profile, notable examples being *Tnf*, *Ccl4* and *Ccl2* that were enriched in Mg-TAMs, while *Il10* was not significantly different (Supplementary Fig. 4). To investigate the differential secretion of these cytokines, sorted TAM subsets were cultured *in vitro*. This confirmed that secretion of TNF, CCL4 and CCL2 was significantly higher in Mg-TAMs (Fig. 7c). To broadly assess how the secretome of Mo-TAMs and Mg-TAMs affects angiogenesis, TAM subsets were cultured on a chick chorioallantoic membrane and their effect on blood vessel development was quantified (Fig. 7d). Mo- and Mg-TAMs exhibited a comparable pro-angiogenic activity, which was significantly higher than that of brain microglia or Transitory Mo-TAMs. To better understand how these TAM subsets may modulate adaptive immunity, we first assessed their T-cell stimulatory potential via the mixed leukocyte reaction assay. This showed that TAMs were unable to induce allogeneic CD4⁺ or CD8⁺ T-cell proliferation, while this was readily observed with naive splenic cDC subsets (Fig. 7e). This suggests that all GBM TAMs have a poor antigen-presenting capacity. In contrast, Mg-TAMs strongly suppressed anti-CD3 induced T-cell proliferation, while brain microglia did not (Fig. 7f). Mo-TAMs also suppressed T-cell proliferation, although less efficient than Mg-TAMs. Together, these data suggest that while Mo-TAMs and Mg-TAMs may exhibit some level of functional specialization, both subsets may be detrimental with regard to adaptive anti-tumor immunity.

Monocyte and microglia-derived TAMs self-renew, compete for space and both can be depleted via CSF1R inhibition.

To blunt the pro-tumoral activity of TAMs, most current strategies either rely on direct TAM depletion or on impairing monocyte influx. However, our scRNA-Seq data suggests that the latter may not reduce TAM numbers, but merely alters its ontogenetic composition (Extended Data Fig. 1e,f). To confirm this, we investigated TAM numbers and ontogeny in tumors from WT and CCR2-KO mice via flow cytometry (Fig. 8a-e). Monocytes and Transitory Mo-TAMs were strongly reduced in CCR2-KO tumors (Fig. 8a,b), resulting in a significant decrease of CX3CR1⁺F4/80⁺ cells (Fig. 8c). However, the percentage of CX3CR1⁺F4/80⁺Ly6B⁻ mature TAMs was not significantly different between WT and CCR2-KO tumors (Fig. 8d). This suggests that impairing monocyte influx results in compensatory mechanisms that maintain TAM numbers. CCR2-KO tumors showed a significantly higher percentage of Mg-TAMs, indicating that a reduction in Mo-TAMs is partly compensated by an increase in Mg-TAMs (Fig. 8e). This is likely due to an increased infiltration of microglia into the tumor, as pulse-chase BrdU analysis did not reveal a significant increase in Mg-TAM proliferation (Fig. 8f). BrdU analysis did show that Mo-TAMs proliferated significantly more in CCR2-KO tumors (Fig. 8f), in agreement with scRNA-Seq data (Fig. 8g). This indicates that a decrease in monocyte influx results in a compensatory increase of Mo-TAM proliferation.

An attractive approach to target TAMs in brain tumors is CSF1R inhibition³⁶. To investigate whether blocking CSF1R may deplete TAMs and whether this differentially affects Mo- vs. Mg-TAMs, WT and CCR2-KO tumor-bearing mice received PLX3397, an orally administered inhibitor of CSF1R³⁷. PLX3397 significantly reduced the percentage of mature TAMs in WT mice (Fig. 8d). However, the relative contribution of monocytes and Transitory Mo-TAMs within the CX3CR1⁺F4/80⁺ population increased (Fig. 8b), and a significantly higher percentage of monocytes was observed within tumors (Fig. 8e). This indicates that CSF1R inhibition does not impair monocyte influx into GBM tumors, but rather blocks monocyte-to-macrophage differentiation. Importantly, PLX3397 treatment significantly reduced tumor growth, confirming that TAMs play a pro-tumoral role (Fig. 8h). The PLX3397-mediated reduction in tumor growth was comparable in WT and CCR2-KO mice (Fig. 8h), which both contained a similar number of mature TAMs. This suggests that monocytes and Transitory Mo-TAMs, which accumulate in PLX-treated WT mice, but are low in PLX-treated CCR2-KOs, have a limited influence on primary tumor growth.

Discussion

DCs are promising targets for cancer immunotherapy³⁸⁻⁴⁰. The healthy mouse brain contains various DC subsets, which are mostly found in its border regions^{1, 2}. cDC2s are enriched in the dura mater, while cDC1s are dominant in the choroid plexus^{1, 10}. These border regions also contain DCs with a migratory gene signature¹, which was first reported for peripheral lymph node DCs²². Our results now show that DC diversity is also evident in glioblastoma tumors. In human tumors we identified 5 main subsets: cDC1, cDC2, MigDC, pre-DC and pDCs, with additional heterogeneity observed in cDC2s. A migratory signature does not necessarily imply DC migration, which remains to be shown. The recent emergence of cDC2 heterogeneity reveals a new layer of complexity within the cDC compartment. A subset of cDC2s with monocyte-related genes has been described in human blood^{21, 24, 25} and spleen²⁶, which we now also report for human GBM tumors. Putative counterparts for the human cDC2a and cDC2b subsets were also observed in mouse GBM tumors. However, it is likely that cDC2s exhibit additional tissue or disease-specific signatures, as was shown for tumor vs. adjacent-tissue DCs⁴¹. Tumor cDCs were actively proliferating in both human and mouse GBM tumors. This proliferative potential was recently also reported for peripheral tissue cDCs^{26, 42}, highlighting that cDC expansion does not solely rely on BM precursors.

Microglia are yolk-sac-derived cells that self-maintain throughout life. BM engraftment is only observed when embryonic microglia are compromised, for example following irradiation^{28, 29}. Previous work has shown that GBM TAMs have a dual ontogeny, either being microglia- or BM-derived^{14, 15}. Our results confirm this and suggest that in GBM tumors microglia- and monocyte-derived TAMs compete for space. Impairing tumor monocyte infiltration resulted in a compensatory increase of Mg-TAMs. Furthermore, recurrent tumors showed a switch in TAM ontogeny. One explanation is that adjuvant treatment, and especially radiotherapy, may compromise the ability of resident microglia to infiltrate and expand in brain tumors. The loss of microglia may also result from a progressive change in the microenvironment and loss of resident brain cells. The nature of the tumor microenvironment was recently shown to be an important determinant for TAM ontogeny, as the fraction of Mo-TAMs is higher in brain metastases as compared to primary gliomas^{43, 44}. Therapy-induced inflammation may also boost monocyte attraction to the tumor. Radiotherapy may elevate CXCL12, thereby facilitating monocyte infiltration via CXCR4 signaling⁴⁵. Recent work has also shown that radiotherapy increases the fraction of Mo-TAMs in recurrent mouse GBM tumors⁴⁶. Intriguingly, we observed that TAMs in hypoxic regions were primarily monocyte-derived. This may result from a higher plasticity of monocyte-derived macrophages, which may better adapt to such environments.

At least three GBM subtypes exist: proneural (TCGA-PN), classical (TCGA-CL) and mesenchymal (TCGA-MES)⁷. ScRNA-Seq studies indicate additional complexity, with GBM cancer cells that exist in four interchangeable cellular states⁶. There are indications of a two-way interaction between GBM cells and TAMs⁷. Future work will need to assess whether individual TAM subsets preferentially adhere to specific GBM subtypes and delineate their interactions.

While TAMs were historically often described in the context of M1/M2 activation, scRNA-Seq is revealing the complexity of macrophage responses and is moving us beyond the linear M1/M2 activation paradigm. Targeting specific TAM subsets may hold significant therapeutic potential and our results now provide a resource that can aid in this effort. This may lead to new therapeutic targets⁴⁷ or biomarkers for predicting therapy response. Since we observed conservation between the human and mouse GBM immune compartment, it highlights the value of preclinical models. A similar conclusion was drawn in a human and mouse comparison study for lung cancer⁴⁸. However, the GL261 immune landscape seemed to more closely resemble that of human recurrent tumors. This may result from the distinct growth patterns of human GBM as compared to transplantable mouse models. In contrast to the human situation, transplanted GL261 shows a clear demarcation between the healthy brain tissue and tumor. The latter may form a boundary which microglia may need to cross, while in humans microglia may be overgrown by tumor tissue. In this regard, spontaneous mouse glioma models may more accurately mimic the human situation. Understanding the strengths and pitfalls of these models will be important for the development of new therapies for glioblastoma.

Accession codes

scRNA-Seq/CITE-Seq sequencing data are deposited at the Gene Expression Omnibus (GEO) with the accession code GSE163120. and in the controlled access public repository European Genome-phenome Archive (EGA), under study accession code EGAS00001004871.

Acknowledgements

This work was supported by Innoviris (Attract grant, BB2B 2015-2) to KM; VLAIO (ImmCyt HBC.2016.0889) to K.M. and J.A.V.G.; KULeuven C-funding (grant C14/17/084) to F.D.S.; the Research Foundation Flanders (FWO, grant G011118N to F.D.S. and the fellowships of J.M. and I.S.); the Leuven Kankerinstituut and VIB. F.L. is supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 766069 ("GLIOTRAIN"). We thank Yvon Elkrim, Aleksandar Murgaski, Gert van Isterdael, the VIB flow core, the VIB Nucleomics core and the VIB BioImaging Core for technical assistance, Janik Mathys for advice regarding statistical analysis, VIB Tech Watch for support regarding single-cell RNA sequencing technologies, Lien Solie and Marleen Derweduwe for the collection and processing of fresh GBM tumor samples, and the KULeuven histology core for its support. We thank the VIB Single-cell accelerator program, Viacheslav Mylka, Niels Vandamme, Bavo Vanneste, Yvan Saeys, Charlotte Scott and Martin Guillems for setting up CITE-Seq at the VIB.

Author contributions

A.R.P.A., I.S., F.L., D.L., J.A.V.G. and K.M. designed experiments; A.R.P.A., I.S., F.L., J.M., K.D.V., H.V.H., S.S.K.H., N.V., R.E.V., O.D.W., C.G., F.D.S. and K.M. performed experiments; A.R.P.A., I.S., F.D.S. and K.M. carried out data analysis; I.S., F.L., A.A., D.K., L.M., K.V.d.B., N.V., Y.S., D.L. and K.M.

performed or helped with bioinformatics analysis; J.D., S.D.V., R.S., M.V. and B.N. provided clinical samples; F.M.B., L.B. and F.D.S. helped with imaging and its analysis; J.D., S.D.V., L.B., O.D.W., Y.S., M.G., B.N., F.D.S., D.L. and J.A.V.G. provided advice on experimental design, data analysis and interpretation; A.R.P.A., I.S., J.A.V.G. and K.M. wrote the manuscript; A.R.P.A., I.S., J.A.V.G. and K.M. revised the manuscript; K.M. conceived the study; J.A.V.G. and K.M. supervised the study.

Competing interests

The authors declare no competing interests.

References

1. Van Hove, H., *et al.* A single-cell atlas of mouse brain macrophages reveals unique transcriptional identities shaped by ontogeny and tissue environment. *Nat Neurosci* **22**, 1021-1035 (2019).
2. Mrdjen, D., *et al.* High-Dimensional Single-Cell Mapping of Central Nervous System Immune Cells Reveals Distinct Myeloid Subsets in Health, Aging, and Disease. *Immunity* **48**, 599 (2018).
3. Jordao, M.J.C., *et al.* Single-cell profiling identifies myeloid cell subsets with distinct fates during neuroinflammation. *Science* **363** (2019).
4. Louveau, A., *et al.* CNS lymphatic drainage and neuroinflammation are regulated by meningeal lymphatic vasculature. *Nat Neurosci* **21**, 1380-1391 (2018).
5. Omuro, A. & DeAngelis, L.M. Glioblastoma and other malignant gliomas: a clinical review. *JAMA* **310**, 1842-1850 (2013).
6. Neftel, C., *et al.* An Integrative Model of Cellular States, Plasticity, and Genetics for Glioblastoma. *Cell* **178**, 835-849 e821 (2019).
7. Wang, Q., *et al.* Tumor Evolution of Glioma-Intrinsic Gene Expression Subtypes Associates with Immunological Changes in the Microenvironment. *Cancer Cell* **32**, 42-56 e46 (2017).
8. Pombo Antunes, A.R., *et al.* Understanding the glioblastoma immune microenvironment as basis for the development of new immunotherapeutic strategies. *Elife* **9** (2020).
9. Quail, D.F. & Joyce, J.A. The Microenvironmental Landscape of Brain Tumors. *Cancer Cell* **31**, 326-341 (2017).
10. Mundt, S., *et al.* Conventional DCs sample and present myelin antigens in the healthy CNS and allow parenchymal T cell entry to initiate neuroinflammation. *Sci Immunol* **4** (2019).
11. Masuda, T., *et al.* Spatial and temporal heterogeneity of mouse and human microglia at single-cell resolution. *Nature* **566**, 388-392 (2019).
12. Kierdorf, K., Masuda, T., Jordao, M.J.C. & Prinz, M. Macrophages at CNS interfaces: ontogeny and function in health and disease. *Nat Rev Neurosci* **20**, 547-562 (2019).
13. Goldmann, T., *et al.* Origin, fate and dynamics of macrophages at central nervous system interfaces. *Nat Immunol* **17**, 797-805 (2016).
14. Bowman, R.L., *et al.* Macrophage Ontogeny Underlies Differences in Tumor-Specific Education in Brain Malignancies. *Cell Rep* **17**, 2445-2459 (2016).
15. Muller, S., *et al.* Single-cell profiling of human gliomas reveals macrophage ontogeny as a basis for regional differences in macrophage activation in the tumor microenvironment. *Genome Biol* **18**, 234 (2017).
16. Darmanis, S., *et al.* Single-Cell RNA-Seq Analysis of Infiltrating Neoplastic Cells at the Migrating Front of Human Glioblastoma. *Cell Rep* **21**, 1399-1410 (2017).
17. Sankowski, R., *et al.* Mapping microglia states in the human brain through the integration of high-dimensional techniques. *Nat Neurosci* **22**, 2098-2110 (2019).
18. Stoeckius, M., *et al.* Simultaneous epitope and transcriptome measurement in single cells. *Nat Methods* **14**, 865-868 (2017).
19. Korsunsky, I., *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat Methods* **16**, 1289-1296 (2019).
20. Serbina, N.V. & Pamer, E.G. Monocyte emigration from bone marrow during bacterial infection requires signals mediated by chemokine receptor CCR2. *Nat Immunol* **7**, 311-317 (2006).
21. Günther, P.C., B; Baßler, K; Händler, K; Becker, M; Dutertre, CA; Bigley, V; Newell, E; Collin, M; Ginhoux, F; Schlitzer, A; Schultze, J.L. A rule-based data-informed cellular consensus map of the human mononuclear phagocyte cell space. *bioRxiv* (2019).
22. Miller, J.C., *et al.* Deciphering the transcriptional network of the dendritic cell lineage. *Nat Immunol* **13**, 888-899 (2012).
23. See, P., *et al.* Mapping the human DC lineage through the integration of high-dimensional techniques. *Science* **356** (2017).

24. Dutertre, C.A., *et al.* Single-Cell Analysis of Human Mononuclear Phagocytes Reveals Subset-Defining Markers and Identifies Circulating Inflammatory Dendritic Cells. *Immunity* **51**, 573-589 e578 (2019).
25. Villani, A.C., *et al.* Single-cell RNA-seq reveals new types of human blood dendritic cells, monocytes, and progenitors. *Science* **356** (2017).
26. Brown, C.C., *et al.* Transcriptional Basis of Mouse and Human Dendritic Cell Heterogeneity. *Cell* **179**, 846-863 e824 (2019).
27. Bennett, F.C., *et al.* A Combination of Ontogeny and CNS Environment Establishes Microglial Identity. *Neuron* **98**, 1170-1183 e1178 (2018).
28. Cronk, J.C., *et al.* Peripherally derived macrophages can engraft the brain independent of irradiation and maintain an identity distinct from microglia. *J Exp Med* **215**, 1627-1647 (2018).
29. Shemer, A., *et al.* Engrafted parenchymal brain macrophages differ from microglia in transcriptome, chromatin landscape and response to challenge. *Nat Commun* **9**, 5206 (2018).
30. Goldmann, T., *et al.* A new type of microglia gene targeting shows TAK1 to be pivotal in CNS autoimmune inflammation. *Nat Neurosci* **16**, 1618-1626 (2013).
31. Hanna, R.N., *et al.* The transcription factor NR4A1 (Nur77) controls bone marrow differentiation and the survival of Ly6C- monocytes. *Nat Immunol* **12**, 778-785 (2011).
32. Jaitin, D.A., *et al.* Lipid-Associated Macrophages Control Metabolic Homeostasis in a Trem2-Dependent Manner. *Cell* **178**, 686-698 e614 (2019).
33. Xue, J., *et al.* Transcriptome-based network analysis reveals a spectrum model of human macrophage activation. *Immunity* **40**, 274-288 (2014).
34. Keren-Shaul, H., *et al.* A Unique Microglia Type Associated with Restricting Development of Alzheimer's Disease. *Cell* **169**, 1276-1290 e1217 (2017).
35. Gentles, A.J., *et al.* The prognostic landscape of genes and infiltrating immune cells across human cancers. *Nat Med* **21**, 938-945 (2015).
36. Pyonteck, S.M., *et al.* CSF-1R inhibition alters macrophage polarization and blocks glioma progression. *Nat Med* **19**, 1264-1272 (2013).
37. Elmore, M.R., *et al.* Colony-stimulating factor 1 receptor signaling is necessary for microglia viability, unmasking a microglia progenitor cell in the adult brain. *Neuron* **82**, 380-397 (2014).
38. Binnewies, M., *et al.* Unleashing Type-2 Dendritic Cells to Drive Protective Antitumor CD4(+) T Cell Immunity. *Cell* **177**, 556-571 e516 (2019).
39. Laoui, D., *et al.* The tumour microenvironment harbours ontogenically distinct dendritic cell populations with opposing effects on tumour immunity. *Nat Commun* **7**, 13720 (2016).
40. Salmon, H., *et al.* Expansion and Activation of CD103(+) Dendritic Cell Progenitors at the Tumor Site Enhances Tumor Responses to Therapeutic PD-L1 and BRAF Inhibition. *Immunity* **44**, 924-938 (2016).
41. Michea, P., *et al.* Adjustment of dendritic cells to the breast-cancer microenvironment is subset specific. *Nat Immunol* **19**, 885-897 (2018).
42. Cabeza-Cabrerizo, M., *et al.* Tissue clonality of dendritic cell subsets and emergency DCpoiesis revealed by multicolor fate mapping of DC progenitors. *Sci Immunol* **4** (2019).
43. Friebel, E., *et al.* Single-Cell Mapping of Human Brain Cancer Reveals Tumor-Specific Instruction of Tissue-Invasive Leukocytes. *Cell* **181**, 1626-1642 e1620 (2020).
44. Klemm, F., *et al.* Interrogation of the Microenvironmental Landscape in Brain Tumors Reveals Disease-Specific Alterations of Immune Cells. *Cell* **181**, 1643-1660 e1617 (2020).
45. Kioi, M., *et al.* Inhibition of vasculogenesis, but not angiogenesis, prevents the recurrence of glioblastoma after irradiation in mice. *J Clin Invest* **120**, 694-705 (2010).
46. Akkari, L., *et al.* Dynamic changes in glioma macrophage populations after radiotherapy reveal CSF-1R inhibition as a strategy to overcome resistance. *Sci Transl Med* **12** (2020).
47. Goswami, S., *et al.* Immune profiling of human tumors identifies CD73 as a combinatorial target in glioblastoma. *Nat Med* **26**, 39-46 (2020).

579 48. Zilionis, R., *et al.* Single-Cell Transcriptomics of Human and Mouse Lung Cancers Reveals
580 Conserved Myeloid Populations across Individuals and Species. *Immunity* **50**, 1317-1334 e1310 (2019).
581
582

Figure legends

Fig. 1. scRNA-Seq of the human and mouse glioblastoma immune compartment.

a, Schematic overview of the 10x chromium scRNA-Seq setup used on tumor tissue isolated from human or mouse brain. Tumors were resected from 7 ND and 4 recurrent patients. Mouse data contained GL261 tumors micro-dissected from WT C57BL/6J (n = 3) or CCR2-KO mice (n=3). **b-d**, Harmony-corrected UMAP-projection of 21.303 CD45⁺ immune cells from human ND GBM tumors (**b**, top panel), 42.870 CD45⁺ immune cells from human recurrent GBM tumors (**c**, top panel), and 49.864 CD45⁺ immune cells from WT and CCR2-KO GL261 tumors (**d**, top panel). **b-d**, bottom panels: pie charts for each individual human ND tumor (**b**); recurrent tumor (**c**) and mouse tumor (**d**), visualizing the relative contribution of each CD45⁺ immune cell population. Percentages are indicated in the pie charts. The colors match the cluster colors used in the UMAP-projections. **e,f**, Heatmaps representing relevant subset-specific genes that are expressed in immune cell subsets found in (**e**) human recurrent GBM tumors and (**f**) in mouse WT and CCR2-KO GL261 tumors. Colors and numbers match that of the clusters shown in **b-d**. B = B cells, DC = Dendritic cells, Mast = Mast cells, Mo = Monocytes, n.c. = necrotic core, NK = Natural Killer cells, Pl. B = Plasma B cells, prol. TAM = proliferating tumor-associated macrophages, T = T cells, TAM = tumor-associated macrophages, Treg = regulatory T cells.

Fig. 2. Single-cell analysis of tumor DCs and identification of distinct cDC2 subsets that are conserved in humans and mice.

a, 2.417 DCs were selected and pooled from human ND and recurrent tumors, followed by re-clustering and Harmony-corrected UMAP-projection. The distinct DC subsets that were identified are annotated. **b**, Volcano plot showing genes that are differentially expressed (DE) (in red: $-\log_{10}(\text{adjusted } P) > 20$, $\log_2(\text{FC}) > 0.5$) between cDC2 (clusters 6-9 in **a**) and cDC1 (cluster 1 in **a**). P value adjustment was performed using Bonferroni correction. **c**, Volcano plot showing genes that are DE (in red: $-\log_{10}(\text{adjusted } P) > 20$, $\log_2(\text{FC}) > 0.5$) between MigDC (cluster 2 in **a**) and cDC1 (cluster 1 in **a**). P value adjustment was performed using Bonferroni correction. **d**, Heatmap of normalized expression of selected marker genes in specific human DC subsets that were identified in **a**. **e**, 5.500 DCs were selected from the mouse GL261 datasets (Fig. 1d), followed by re-clustering and Harmony-corrected UMAP-projection. The distinct DC subsets that were identified are annotated. **f**, Volcano plot showing genes that are DE (in red: $-\log_{10}(\text{adjusted } P) > 20$, $\log_2(\text{FC}) > 0.5$) between cDC2 (clusters 4-6 in **e**) and cDC1 (cluster 1 in **e**). P value adjustment was performed using Bonferroni correction. **g**, Volcano plot showing genes that are DE between MigDC (cluster 2 in **e**) and cDC1 (cluster 1 in **e**). P value adjustment was performed using Bonferroni correction. **h**, Scatter plot comparing the $\log_{10}(\text{adjusted } P)$ value of the 150 most significantly up- or downregulated genes in mouse cDC2 versus cDC1 (x-axis) and human cDC2 versus cDC1 (y-axis). The sign of the $\log_{10}(\text{adjusted } P)$ corresponds to up- or downregulation of genes. Common up- and downregulated genes are represented in red and blue, respectively. Genes that are only up- or downregulated within 1 species (human or mouse), but are not differential in the

other species, are represented in black. Genes that are upregulated in mouse and downregulated in human, or vice versa, are represented in pink. P value adjustment was performed using Bonferroni correction. **i**, Heatmap of normalized expression of selected marker genes in specific mouse DC subsets. **j**, Proliferating human DCs (cluster 10 in panel **a**) were mapped onto the non-proliferating DCs to determine the subpopulation they belong to (see methods). Bar graphs represent the percentage of cDC1 (green bar) and cDC2 (red bar) that associated with the proliferation cluster, in individual patients (ND7, R1-4). Mean $\pm$ s.e.m. Significance was evaluated using a paired two-tailed *t*-test. **k**, The percentages of mouse cDC1 (green bars) and cDC2 (red bars) that associated with a proliferation cluster in individual WT and CCR2-KO scRNA-Seq samples. Bar graphs represent mean $\pm$ s.e.m. Significance was evaluated using a two-way ANOVA with Sidak's multiple comparisons test. **l**, The percentages of mouse cDC1 (green bars) and cDC2 (red bars) that are BrdU positive in WT and CCR2-KO GL261 tumors, two hours post-BrdU injection, as measured via flow cytometry. Bar graphs represent mean $\pm$ s.e.m. of *n* = 7 (WT control), *n* = 7 (CCR2-KO), which originate from 2 independent experiments. Significance was evaluated using a RM two-way ANOVA model with Sidak's multiple comparisons test. ns *P* > 0.05; ** *P* < 0.01; ****, *P* < 0.0001. Exact *P* values can be found in Supplementary table 19.

Fig. 3. GBM TAMs exhibit a mixed ontogeny which is significantly altered upon human disease stage.

a-c, Monocytes and macrophages were selected from the human and mouse datasets and re-clustered. Shown are Harmony-corrected UMAP-projections from human ND GBM tumors (18.157 cells, **a**), human recurrent GBM tumors (19.541 cells, **b**) and mouse GL261 tumors (29.309 cells, **c**). Mo-TAM clusters are encircled in blue, Mg-TAM clusters in red. **d-f**, UMAP plots showing the expression of selected genes in ND tumors (**d**), recurrent tumors (**e**) and mouse tumors (**f**). **g,h**, Volcano plot showing genes that are DE ($-\log_{10}(\text{adjusted } P) > 20$, $\log_2(\text{FC}) > 0.5$) between Mg-TAMs and Mo-TAMs in human recurrent tumors (**g**) and mouse tumors (**h**). P value adjustment was performed using Bonferroni correction. **i**, Scatter plot comparing the $\log_{10}(\text{adjusted } P)$ value of the 150 most significantly up- or downregulated genes in mouse Mg-TAMs versus Mo-TAMs (x-axis) and in human Mg-TAMs versus Mo-TAMs (y-axis). The sign of the $\log_{10}(\text{adjusted } P)$ corresponds to up- or downregulation of genes. Common up- and downregulated genes are represented in red and blue, respectively. Genes that are only up- or downregulated within 1 species (human or mouse) are represented in black. P value adjustment was performed using Bonferroni correction. **j**, Bar graphs representing the ratios of Mo-TAMs (blue bars) and Mg-TAMs (red bars) in human ND GBM tumors (patients ND1-7), human recurrent GBM tumors (patients R1-4), and in mouse GL261 tumors in WT (WT 1-3) and CCR2-KO (KO 1-3) mice. c.monocytes = classical monocytes, FC = fold change, Mg-TAM = microglia-derived TAM, Mo-TAM = monocyte-derived TAM, nc. monocytes = non-classical monocytes, prol. TAM = proliferating TAM.

Fig. 4. Fate mapping and spatial analysis reveals the distribution of Mo-TAMs and Mg-TAMs in mouse and human tumors.

a, *Cx3cr1^{CreER}:R26-YFP* mice received tamoxifen injections followed by a 4-week washout period prior to GL261 injection. Coronal brain sections of fate-mapped tumor-bearing mice, were stained for F4/80 (red), YFP (green) and DAPI (white). DAPI was used to delineate the border (dashed line) between tumor and brain. In the tumor, Mo-TAMs were identified as F4/80⁺YFP⁻ cells, while Mg-TAMs were F4/80⁺YFP⁺ (yellow). Insets (i, ii, iii) of regions of interest are shown and the percentage of YFP⁺ cells within the IBA1⁺/F4/80⁺ population was quantified and is represented in the bar graphs on the right. Graphs represent mean $\pm$ s.e.m. of $n = 3$ mice. Representative images of $n = 3$ mice. Scale bars: 50 μ m. **b**, Schematic overview of the adoptive transfer experiment. **c**, Adoptively transferred cells were identified as GFP⁺CX3CR1⁺F4/80⁺ cells (green) (pre-gated on CD45⁺ live single cells) and were overlaid on GFP⁺CX3CR1⁺F4/80⁺ cells (blue). This showed that adoptively transferred cells were contained in the CD11b⁺CD45^{hi} gate, where Mo-TAMs are present. However, adoptively transferred non-classical monocytes retain a monocytic scatter profile, while classical monocytes show an expanded profile characteristic of TAMs. Transferred classical monocytes also downregulate Ly6B expression. For reference the arrow indicates Ly6B expression levels of classical monocytes as they are found in blood. Plots are representative of $n = 2$ mice. **d**, GFP⁺ cells are not detected in tumors of CCR2-KO mice that did not receive an adoptive transfer of monocytes. Plots are representative of $n = 2$ mice. **e**, Fluorescent images of a recurrent GBM FFPE tumor section (R5) stained for various markers (DAPI = blue, CD68 = green, TMEM119 = red, SOX2 = magenta, CD34 = cyan; all images are at the same scale; scale bar in DAPI image = 50 μ m). The “overlay” image combines all single channel images. Finally, a digitized image in which the unequivocally identified cell types are highlighted with a colored dot, is shown in the last panel (color representations are indicated in the image). Results are representative for $n = 7$ GBM patients. **f**, Overlay image of the CD68 (green) and TMEM119 (red) staining showing the presence of both Mo- and Mg-TAMs in a GBM tumor section. Results are representative for $n = 7$ GBM patients. scale bar = 50 μ m **g**, Digital representation of the entire tumor section (square indicates the highlight of panel **e**) in which cancer cells, endothelial and TAMs are highlighted across the entire tissue (gray background are all detected DAPI⁺ cells; red dots = indicated cell type). **h**, (left panel) Digital representation of the entire tumor section, in which the TAMs are colored by their relative TMEM119 expression (0-5). (right panel) In each square of 100x100pixels, the number of TMEM119^{hi} and TMEM119^{lo} TAMs was counted: if TMEM119^{hi} cells were >50% of all detected TAMs in that square, it is colored red; otherwise the square is colored cyan. c. Mono = classical monocytes; nc. Mono = non-classical monocytes; MFs = macrophages.

Fig. 5. scRNA-Seq reveals the heterogeneity of Mo-TAMs and identifies subsets that are conserved across species.

a, Overview of Mo-TAM subsets in human recurrent GBM tumors, with 6 subclusters that were discerned: (1) Transitory Mo-TAM (green), (2) Phagocytic/lipid signature Mo-TAM (purple), (3) Hypoxic

Mo-TAM (blue), (4) *SEPP1*^{lo} Mo-TAM (dark orange), (5) *SEPP1*^{hi} Mo-TAM (pale orange), and (6) IFN signature Mo-TAM (pink). **b**, UMAP plots showing the expression of key signature genes that are differential between the various subsets. Dashed lines encircle the subsets as annotated in panel a. **c-e**, DE analysis in human recurrent Mo-TAMs. **c**, Volcano plot showing genes that are DE between *SEPP1*^{lo} Mo-TAMs (cluster 4) and transitory Mo-TAMs (cluster 1). **d**, Volcano plot showing genes that are DE between Phagocytic/lipid Mo-TAMs (cluster 2) and transitory Mo-TAMs (cluster 1). **e**, Volcano plot showing genes that are DE between hypoxic Mo-TAMs (cluster 3) and all other Mo-TAMs (clusters 1, 2, 4, 5, 6). DE threshold: $-\log_{10}(\text{adjusted } P) > 20$, $\log_2(\text{FC}) > 0.5$; P value adjustment was performed using Bonferroni correction (**c-e**). **f**, Overview of Mo-TAM subsets in mouse GL261 tumors, with 6 subclusters that were discerned. The cluster numbers and names indicate similarity to their human counterparts. **g**, UMAP plots showing the expression of key signature genes that are differential between the various subsets. **h**, Overview of the distinct Mo-TAM subsets present in human (top) and mouse (bottom) tumors. **i**, Overview of Mg-TAM subsets in human ND tumors. **j**, Mg-TAMs from human recurrent tumors were selected and re-clustered (3.914 cells). **k**, Mg-TAMs from mouse tumors were selected and re-clustered (4.504 cells). **l**, Bar graphs showing the percentage of hypoxic TAMs (left panel) and IFN signature TAMs (right panel) in human recurrent Mo-TAM (blue) and Mg-TAM (red) populations per patient. **m**, Bar graphs showing the percentage of hypoxic TAMs (left panel) and IFN-signature TAMs (right panel) in mouse Mo-TAM (blue) and Mg-TAM (red) populations per mouse sample in WT tumors (WT 1-3) and CCR2-KO tumors (KO 1-3).

Fig. 6. CITE-Seq analysis validates TAM and DC subsets at the protein level and reveals subset-specific cell surface protein markers.

a, Human CITE-Seq dataset consisting of GBM-associated monocytes, macrophages and DCs. Shown are Harmony-corrected UMAP-projections based on RNA expression (left panel) or cell surface protein expression (right panel) (14.793 cells). Colors are derived from RNA-based clustering. **b**, Mouse CITE-Seq dataset consisting of monocytes, macrophages and DCs derived from WT and CCR2-KO GL261 tumors. Shown are Harmony-corrected UMAP-projections based on RNA expression (left panel) or cell surface protein expression (right panel) (13.635 cells). Colors are derived from RNA-based clustering. **c**, Human RNA-based UMAP plots, showing the expression of selected genes (top plots) and their corresponding proteins based on CITE-Seq antibody staining (Antibody-Derived Tag or ADT signals) (bottom plots). **d**, Heatmap of normalized expression of selected cell surface proteins in human monocyte and TAM subsets. **e-f**, DCs were selected from the human CITE-Seq dataset (1333 cells) (**e**) or the mouse CITE-Seq dataset (2761 cells) (**f**), and re-clustered. Shown are Harmony-corrected UMAP-projections based on RNA expression (left panels) or cell surface protein expression (right panels). Colors are derived from RNA-based clustering. **g-h**, Heatmaps representing relevant proteins expressed in human DCs (**g**) or mouse DCs (**h**). **i**, Human Protein-based UMAP plots, showing the expression of selected genes (top) and their corresponding proteins (bottom).

Fig. 7. Functional profiling of Mo-TAMs and Mg-TAMs reveals their T-cell suppressive potential.

a, Representative contour plots depicting the gating strategy for selecting monocytes and TAM subsets in GL261 tumors (a1) and microglia in the surrounding brain (a2). Cells were pre-gated as live and single cells. (a3) Contour plots where brain microglia, Mg-TAMs and Mo-TAMs were overlaid to reveal their differential expression profile of CX3CR1 vs. F4/80 and Ly6B vs. MHCII. Representative plots of 4 independent experiments with $n = 14$ to 15 mice. **b**, Tumors and brains of *Cx3cr1^{CreER}:R26-YFP* mice, which were injected with GL261 cells 4 weeks post tamoxifen treatment, were processed individually for flow cytometry. The flow cytometry plots confirm the presence of YFP⁺ microglia-derived Mg-TAMs (red) and YFP⁻ monocyte-derived Mo-TAMs (blue) in tumors. YFP⁺ microglia from adjacent brain are indicated in green. YFP⁺ brain microglia, YFP⁺ Mg-TAMs and YFP⁻ Mo-TAMs were overlaid to reveal their differential expression profile of CD11b vs. CD45. Results are representative of 3 independent experiments with $n = 2$ mice. **c**, FACS-sorted Mg-TAMs, Mo-TAMs and Transitory Mo-TAMs were cultured for 20h, after which the presence of IL10, TNF, CCL4 and CCL2 in the cell culture medium was quantified. Bars represent mean $\pm$ s.e.m. $n = 6$ experiments (IL10, CCL4 and CCL2), $n = 5$ experiments (TNF), where in each experiment TAMs were sorted from a pool of $n = 10$ mice. Significance was evaluated using a one-way ANOVA and Sidak's multiple comparisons test. **d**, FACS-sorted TAM subsets, brain microglia or PBS/BSA were grafted on Chick Chorioallantoic membranes. The numbers of implant-directed vessels were quantified. Bars represent mean $\pm$ s.e.m of $n = 6$ (Brain Mg), $n = 18$ (Mg-TAMs), $n = 13$ (Mo-TAMs), $n = 12$ (Transitory Mo-TAMs) mice. Each data point represents an individual egg and data were pooled from 3 experiments. TAMs and microglia were always sorted from a pool of 6-10 mice. Significance was evaluated using a mixed effects ANOVA model and Sidak's multiple comparisons tests. **e**, Allogeneic mixed leukocyte reaction (MLR) assay, where sorted TAM subsets or splenic cDCs were cultured with purified CD4⁺ or CD8⁺ T cells from naive BALB/c mice. T-cell proliferation was measured using [³H]thymidine incorporation. Graphs represent the average level of [³H]thymidine incorporation $\pm$ s.e.m, expressed as counts per minute (CPM). Representative results of $n = 3$. **f**, Sorted TAM subsets or brain microglia were added to naive C57BL/6J splenocytes, followed by anti-CD3 treatment to activate T cells. T cell proliferation was measured using [³H]thymidine incorporation. Bars represent the % suppression of T cell proliferation (see methods). Each data point represents an independent experiment ($n = 6$ for TAM subsets, $n = 3$ for brain microglia). TAMs or microglia were always sorted from a pool of 10 mice. Significance was evaluated with a mixed effects ANOVA model and Sidak's multiple comparisons tests. ns $P > 0.05$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$. Exact P values can be found in Supplementary Table 19.

Fig. 8. CCR2-deficiency results in a switch in TAM ontogeny and boosts Mo-TAM proliferation, while CSF1R-blockade depletes TAMs.

a, Representative contour plots of tumors from control and PLX3397-treated C57BL/6J or CCR2-KO mice. Cells were pre-gated as CD45⁺ live single cells. **b**, Pie charts for the mouse groups, according to the order in panel **a**, showing the distribution of TAM subsets and classical monocytes within the

CX3CR1⁺F4/80⁺ compartment. **c,d**, Percentage of CX3CR1⁺F4/80⁺ (**c**) or CX3CR1⁺F4/80⁺Ly6B⁻ (**d**) cells within all live tumor cells of the indicated mouse groups. Significance was evaluated with an ordinary two-way ANOVA and Sidak's multiple comparisons tests. **e**, Percentage of TAM subsets and classical monocytes within all live tumor cells, for each group. Significance was evaluated with an ordinary two-way ANOVA and Sidak's multiple comparisons tests within each population. (**a-e**) Graphs represent mean $\pm$ s.e.m. of n = 15 (WT control), n = 17 (WT + PLX3397), n = 16 (CCR2-KO control), n = 24 (CCR2-KO + PLX3397) mice, which originate from 4 independent experiments. **f**, Percentages of BrdU-positive Mg-TAMs (red) and Mo-TAMs (blue) in WT and CCR2-KO (KO) GL261 tumors, two hours post-BrdU injection, as measured via flow cytometry. Bars represent mean $\pm$ s.e.m of n = 7 (WT control), n = 6 (CCR2-KO), which originate from 2 independent experiments. Significance was evaluated using a mixed effects ANOVA model and Sidak's multiple comparisons tests. **g**, Proliferating mouse TAMs were mapped onto the non-proliferating TAMs to determine the subpopulation they belong to (see methods). Bar graphs represent the percentage of Mg-TAMs (red) and Mo-TAMs (blue) that associated with a proliferation cluster, in individual WT and CCR2-KO scRNA-Seq samples. Significance was evaluated using a two-way ANOVA and Sidak's multiple comparisons tests. **h**, Tumor weights of control and PLX3397-treated C57BL/6J and CCR2-KO mice. Bar graphs represent mean $\pm$ s.e.m. of n = 25 (WT control), n = 27 (WT + PLX3397), n = 25 (CCR2-KO control), n = 25 (CCR2-KO + PLX3397) mice, which originate from 4 independent experiments. Significance was evaluated with an ordinary two-way ANOVA and Sidak's multiple comparisons tests. ns $P > 0.05$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$. Exact P values can be found in Supplementary Table 19.

Legends Extended Data Figures

Extended Data Fig. 1. Harmony efficiently corrects for patient-, batch-, and platform-induced gene expression noise without obscuring relevant heterogeneity.

a, UMAP-projections without (left) and with (right) Harmony-correction of the entire CD45⁺ immune compartment of human recurrent GBM tumors (top) and of a subsetting and re-clustered dataset of monocytes and macrophages only (bottom). Clusters are colored based on patient identity (R1-4), illustrating a clear patient-specific segregation without Harmony-correction, but the absence of this segregation after Harmony-correction. **b**, Harmony-corrected UMAP-projections of patients R1 to R4 represented separately on each plot, indicating that each cluster carries cells from every patient. The UMAP clusters are colored based on cluster annotations in Fig. 1c. Only cells belonging to one individual patient are colored per panel, cells belonging to the other patients, are colored in grey. n.c. = necrotic core. **c**, Percentage of the indicated populations within the CD45⁺ compartment of human ND and recurrent tumors, as measured via scRNA-Seq. Each datapoint stems from an individual patient: ND1-7 and R1-4. Bar graphs represent mean $\pm$ s.e.m. Significance was evaluated using the Mann-Whitney test (two-tailed). **d**, The percentage of cluster TAM 1 within (TAM 1 + TAM 2) in human ND and recurrent tumors. Each datapoint stems from an individual patient: ND1-7 and R1-4. Bar graphs represent mean $\pm$ s.e.m. Significance was evaluated using the Mann-Whitney test (two-tailed). **e**, Percentage of the indicated populations within the CD45⁺ compartment of WT and CCR2-KO GL261 tumors, as measured via scRNA-Seq. Each datapoint stems from an individual scRNA-Seq sample: WT 1-3 and CCR2-KO 1-3. Bar graphs represent mean $\pm$ s.e.m. Significance was evaluated using an unpaired two-tailed *t*-test with Welch's correction. **f**, The percentage of cluster TAM 1 within (TAM 1 + TAM 2) in WT and CCR2-KO GL261 tumors. Each datapoint stems from an individual scRNA-Seq sample: WT 1-3 and CCR2-KO 1-3. Bar graphs represent mean $\pm$ s.e.m. Significance was evaluated using an unpaired two-tailed *t*-test with Welch's correction. ns $P > 0.05$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. Exact P values can be found in Supplementary Table 19.

Extended Data Fig. 2. DC heterogeneity in human and mouse GBM tumors.

a, UMAP feature plots based on the Harmony-corrected UMAP-projection of DCs from human ND and recurrent GBM tumors of Fig. 2a, showing the expression levels of the indicated genes. **b**, Pie charts visualizing the percentage of the indicated DC subsets in ND and recurrent GBM tumors, represented per patient. The relative abundance of the various DC subsets was calculated for the individual patients for which sufficient numbers of DCs were captured (ND7 and R1-4). Colors match the cluster colors of Fig. 2a. **c**, UMAP feature plots based on the Harmony-corrected UMAP-projection of DCs from mouse GL261 tumors in Fig. 2e, showing the expression levels of the indicated genes. **d**, Scatter plot comparing the $\log_{10}(\text{adjusted } P)$ value of the most significantly up- or downregulated genes in mouse MigDC versus other DC populations (x-axis) and in human MigDC versus other DC populations (y-axis). The sign of the $\log_{10}(\text{adjusted } P)$ corresponds to up- or downregulation of genes. Common up- and downregulated genes are represented in red and blue, respectively. Genes that are only up- or

downregulated within 1 species (human or mouse), but are not differential in the other species, are represented in black. Genes that are upregulated in mouse and downregulated in human, or vice versa, are represented in pink. P value adjustment was performed using Bonferroni correction. (e), Similar to (d) but for pDCs. f, Bar graphs illustrating the percentages of DCs belonging to the distinct DC subpopulations for WT (WT 1-3) and CCR2-KO (KO 1-3) samples. Bar graphs represent mean $\pm$ s.e.m. Significance was evaluated using an unpaired two-tailed *t*-test with Welch's correction. ns $P > 0.05$; * $P < 0.05$. Exact P values can be found in Supplementary Table 19. g, Representative flow cytometry plots depicting the gating strategy for selecting cDC1 and cDC2 subsets in GL261 tumors. Cells were pre-gated as live and single cells (top). Plots show BrdU staining in cDC1s or cDC2s from control (non-injected) or BrdU injected mice (bottom). Representative plots of 2 independent experiments with $n = 7$ mice.

Extended Data Fig. 3. Adoptive transfer of classical and non-classical monocytes in C57BL/6 or NR4A1-KO GL261 tumor-bearing mice.

a, Representative flow cytometry plots showing GFP⁺ cells in GL261 tumors from C57BL/6 mice, 5 days after the adoptive transfer of classical (c. Mono) or non-classical (nc. Mono) monocytes that were purified from *Cx3cr1*^{GFP/+} mice. Plots are gated on live single CD45⁺CD11b⁺ cells. GFP⁺ cells were not detected in tumors of mice that did not receive an adoptive transfer of monocytes (Control). b, In mice that had received an adoptive transfer of classical monocytes, GFP⁺CD11b⁺CD45⁺ cells (red) were overlaid on GFP⁺CD11b⁺CD45⁺ cells (grey). The representative plots show how the transferred cells (red) relate to the host cells (grey) in terms of scatter profile and expression of the indicated markers. c, In mice that had received an adoptive transfer of non-classical monocytes, GFP⁺CD11b⁺CD45⁺ cells (blue) were overlaid on GFP⁺CD11b⁺CD45⁺ cells (grey). The representative plots show how the transferred cells (blue) relate to the host cells (grey) in terms of scatter profile and expression of the indicated markers. d, Overlay of GFP⁺ cells from (b) and (c). The results show that 5 days post adoptive transfer, classical monocytes (red) exhibited an enlarged scatter profile and upregulated CX3CR1, F4/80 and MHCII expression, while this was not seen for adoptively transferred non-classical monocytes (blue). e-h, Similar to a-d except that adoptive transfers were performed in tumor-bearing NR4A1-KO mice. Plots are representative of two independent experiments $n = 2$ to 4 mice. c. Mono = classical monocytes; nc. Mono = non-classical monocytes.

Extended Data Fig. 4. Delineating the gene signatures of human Mo-TAMs.

a, Volcano plot showing DE genes ($-\log_{10}(\text{adjusted } P) > 20$, $\log_2(\text{FC}) > 0.5$) between *SEPP1*^{hi} Mo-TAMs (cluster 5 in Fig. 5a) and *SEPP1*^{lo} Mo-TAMs (cluster 4 in Fig. 5a). P value adjustment was performed using Bonferroni correction. *SEPP1*^{hi} TAM enriched genes are marked in green, if their expression was reported to be affected by a panel of cytokines/stimuli in the study of Xue et al., 2014³³. b, For the *SEPP1*^{hi} TAM enriched genes marked in green in (a), a heatmap was made using the data from Xue et al., 2014, visualizing the normalized gene expression levels of human macrophages treated with the

indicated stimuli. **c**, Volcano plot showing DE genes (in red: $-\log_{10}(\text{adjusted } P) > 20$, $\log_2(\text{FC}) > 0.5$) between IFN sign. Mo-TAMs (cluster 6 in Fig. 5a) and *SEPP1*^{lo} Mo-TAMs (cluster 4 in Fig. 5a). P value adjustment was performed using Bonferroni correction. We marked IFN sign. enriched genes in pink, if their expression was reported to be affected by a panel of cytokines/stimuli in the study of Xue et al., 2014. **d**, For the IFN sign. enriched genes marked in pink in (**c**), a heatmap was made using the data from Xue et al., 2014, visualizing the normalized gene expression levels of human macrophages treated with the indicated stimuli. GC = glucocorticoid, HDL = high-density lipoprotein, upLPS = ultrapure lipopolysaccharide, LA = lauric acid, LiA = linoleic acid, OA = oleic acid, P3C = Pam3CSK4, PA = palmitic acid, PGE2 = prostaglandine E2, SA = stearic acid, sLPS = standard lipopolysaccharide, TPP = TNF + PGE2 + P3C.

Extended Data Fig. 5. Mo-TAM heterogeneity in GBM tumors.

a, Overview of Mo-TAM subsets in human ND GBM tumors, with 5 subclusters that were discerned: (1) Transitory Mo-TAM (green), (2) Phagocytic/lipid signature Mo-TAM (purple), (3) Hypoxic Mo-TAM (blue), (5) *SEPP1*⁺ Mo-TAM (pale orange), and (6) IFN signature Mo-TAM (pink). **b**, UMAP plots showing the expression of key signature genes that are differential between the various subsets. Dashed lines encircle the subsets as annotated in panel a. **c-d**, DE analysis in human ND Mo-TAMs. **c**, Volcano plot showing genes that are DE between Hypoxic Mo-TAMs (cluster 3) and *SEPP1*⁺ Mo-TAMs (cluster 5). **d**, Volcano plot showing genes that are DE between Phagocytic/lipid Mo-TAMs (cluster 2) and *SEPP1*⁺ Mo-TAMs (cluster 5). In red are shown genes with $-\log_{10}(\text{adjusted } P) > 20$, $\log_2(\text{FC}) > 0.5$; P value adjustment was performed using Bonferroni correction. **e-f**, Pie charts showing the percentage of Mo-TAM subsets in human ND tumors (**e**), and human recurrent tumors (**f**), per individual patient sample. **g**, Bar graphs showing the mean normalized expression of selected genes in ND *SEPP1*⁺ Mo-TAMs (ND cluster 5), Recurrent *SEPP1*^{lo} Mo-TAMs (R cluster 4) and Recurrent *SEPP1*^{hi} Mo-TAMs (R cluster 5), for the individual patient samples. Bar graphs represent mean $\pm$ s.e.m. Significance was evaluated using an ordinary One-way ANOVA and Tukey's multiple comparisons tests. ns $P > 0.05$; * $P < 0.05$; ** $P < 0.01$. Exact P values can be found in Supplementary Table 19. **h**, Pie charts showing the percentage of Mo-TAM subsets in mouse GL261 tumors per individual mouse sample.

Extended Data Fig. 6. Human GBM Mo-TAM heterogeneity is recapitulated in mouse GL261 tumors with conserved gene signatures between species.

a-c, DE analysis in mouse GL261 Mo-TAMs. **a**, Volcano plot showing genes that are DE between mouse GL261 *SEPP1*⁺ Mo-TAMs and Transitory Mo-TAMs. **b**, Volcano plot showing genes that are DE between mouse GL261 Phagocytic/lipid Mo-TAMs and Transitory Mo-TAMs. **c**, Volcano plot showing genes that are DE between Hypoxic Mo-TAMs and other Mo-TAMs. In red are shown genes with $-\log_{10}(\text{adjusted } P) > 20$, $\log_2(\text{FC}) > 0.5$; P value adjustment was performed using Bonferroni correction. **d**, Scatter plot comparing the $\log_{10}(\text{adjusted } P)$ value of the most significantly up- or downregulated

genes in mouse GL261 Lipid Mo-TAM versus other Mo-TAM populations (x-axis) and in human recurrent GBM Lipid Mo-TAM versus other Mo-TAM populations (y-axis). The sign of the $\log_{10}(\text{adjusted } P)$ corresponds to up- or downregulation of genes. Common up- and downregulated genes are represented in red and blue, respectively. Genes that are only up- or downregulated within 1 species (human or mouse), but are not differential in the other species, are represented in black. Genes that are upregulated in mouse and downregulated in human, or vice versa, are represented in pink. P value adjustment was performed using Bonferroni correction. **e**, Similar to (d) but for mouse and human IFN Mo-TAM **f**, Similar to (d) but for mouse and human Hypoxic Mo-TAM. **g**, Similar to (d) but for mouse *Sepp1*⁺ Mo-TAM and human *SEPP1*^{hi} Mo-TAM. **h**, UMAP plot showing the expression of *Arg1*, which was enriched in mouse hypoxic Mo-TAMs. **i**, GL261 tumor-bearing mice were injected with pimonidazole 2 hours prior to sacrifice. Coronal cryosections of GL261 brain tumor tissue, were stained for ARG1 (red), F4/80 (green) and pimonidazole (blue) to visualize hypoxia. ARG1 specifically colocalizes with F4/80⁺ macrophages (yellow) and is enriched in PIMO⁺ hypoxic tumor regions. Image is representative for n = 3 mice. Scalebar = 50 μm .

Extended Data Fig. 7. Mg-TAM heterogeneity in human and mouse GBM.

a, SCORPIUS trajectory inference was run on a dataset containing Mg-TAMs and microglia that were isolated from the surrounding brain tissue of GL261 tumor-bearing WT mice (WT5 w.b.). Cells were automatically ordered along a linear trajectory, where green cells represent brain microglia and blue cells Mg-TAMs. The top genes were clustered into three gene modules (normalized expression) that are down- or upregulated as brain microglia transition towards Mg-TAMs. **b**, Volcano plot showing genes that are DE (in red: $-\log_{10}(\text{adjusted } P) > 20$, $\log_2(\text{FC}) > 0.5$) between human ND GBM Mg-TAM cluster 3 (Phagocytic/lipid Mg-TAM) and Mg-TAM cluster 1. P value adjustment was performed using Bonferroni correction. **c-d**, UMAP plots of signature genes for phagocytic/lipid, IFN and hypoxic subsets in human recurrent (**c**) and mouse GL261 tumors (**d**) are shown. **e-f**, Volcano plots showing genes that are DE (in red: $-\log_{10}(\text{adjusted } P) > 20$, $\log_2(\text{FC}) > 0.5$) in hypoxic Mg-TAMs in human recurrent (**e**) and mouse GL261 (**f**) tumors. P value adjustment was performed using Bonferroni correction.

Extended Data Fig. 8. Altered transcriptional states in TAM subsets from human ND versus recurrent tumors.

a, b, Bar plots showing the mean normalized expression levels of a selection of genes that are enriched in human recurrent versus ND tumors (**a**), or that are enriched in human ND versus recurrent tumors (**b**), in Mg-TAM (left), *SEPP1*⁺ Mo-TAM (middle) and Hypoxic Mo-TAM (right), for each patient sample. Each data point corresponds to the individual human scRNA-seq sample, n= 7 for ND and n=5 for recurrent. Bar graphs represent mean $\pm$ s.e.m. Significance was evaluated using unpaired two-tailed t test with Welch's correction for Mg-TAM and hypoxic Mo-TAM and Ordinary One-way ANOVA for *SEPP1*⁺ Mo-TAM. ns $P > 0.05$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$. Exact P values can be found in Supplementary Table 19. **c, d**, Gene ontology analysis on DE genes between human

recurrent GBM *SEPP1*⁺ Mo-TAMs and ND GBM *SEPP1*⁺ Mo-TAMs showing the top 20 enriched GO terms in recurrent versus ND GBM TAMs (c) and enriched GO terms in ND versus recurrent GBM TAMs (d) (adjusted P-value cutoff of 10⁻¹⁰, log₂FC cutoff of 0.379). e, Bar graphs showing the mean normalized expression values for a selection of genes that exhibited a high variation in expression across individual ND and recurrent patients. Meta-z-score values for these genes were obtained via PRECOG³⁵ which represents the association of the genes with GBM patient survival based on public datasets. The values are colored according to the color-scale at the bottom.

Extended Data Fig. 9. CITE-Seq protein validation for human and mouse GBM TAMs.

a-b, RNA-based UMAP plots showing expression of signature genes for Mg-TAM, Monocytes, Transitory Mo-TAMs, Phagocytic/lipid Mo-TAMs, Hypoxic Mo-TAMs, *SEPP1*⁺ Mo-TAMs and IFN Mo-TAMs in human (a) and mouse (b) tumors. c-d, Protein-based UMAP plots in which the TAM clusters identified based on RNA-expression are highlighted in human (c) and mouse (d) tumors. e-g, Scatter plots comparing the log₁₀(adjusted P) value of the most significantly up- or downregulated proteins (ADT values) (x-axis) and genes (y-axis) in human GBM Mg-TAMs versus other TAM subsets (e), in human GBM *SEPP1*^{hi} TAMs versus other TAM subsets (f) and in mouse GL261 mg-TAMs versus other TAM subsets (g). The sign of the log₁₀(adjusted P) corresponds to up- or downregulation of protein/gene combinations. Common up- and downregulated protein/gene combinations are represented in red and blue, respectively. Protein/gene combinations that are only up- or downregulated on the protein level but not on the gene expression level, are represented in green. Genes that are upregulated on the protein level and downregulated on the gene expression level, or vice versa, are represented in pink. P value adjustment was performed using Bonferroni correction. h, Mouse RNA-based UMAP plots showing the expression of the indicated genes and their corresponding cell surface protein markers.

Extended Data Fig. 10. CITE-Seq analysis reveals the persistence of cDC1 or cDC2-related proteins in a fraction of MigDCs in human and mouse GBM.

a, Protein-based UMAP plot of human DCs, in which MigDCs are highlighted in purple (left). The rectangle delineates the insets shown in the right panels, where the expression of cDC1- or cDC2-related marker genes and their corresponding cell surface proteins are plotted. b, RNA-based UMAP plot of mouse DCs, in which MigDCs are highlighted in purple (left). The rectangle delineates the insets shown in the right panels, where the expression of cDC1- or cDC2-related marker genes and their corresponding cell surface proteins are plotted. c, RNA-based UMAP plots of mouse DCs showing the expression level of *H2-Aa* in the top panel and the corresponding protein IA-IE in the bottom panel. d, RNA-based UMAP plots of mouse DCs showing the expression of the indicated genes and their corresponding cell surface protein markers.

Methods

Patients, human tumor tissue acquisition and single-cell isolation

Fresh tumor tissue was obtained from 13 patients undergoing surgical resection at three Belgian hospitals: UZ Brussel (ND7, R1-5), KU Leuven (ND1-5) and St. Elisabeth (ND6, ND8). The study was approved by the local ethics committees of the UZ Brussel/Vrije Universiteit Brussel (Brussels, Belgium), UZ Leuven/KU Leuven (Leuven, Belgium) and Medical Ethical Committee of the Europe Hospitals (Brussels, Belgium). All patients provided informed consent. Clinical patient information can be found in Supplementary table 1. After resection, samples were immediately transported on ice to research facilities at the Vrije Universiteit Brussel or KU Leuven. For ND7, ND8 and R1-5, tumor samples were cut into small pieces (2-3 mm) and dissociated using Enzyme Mix A (30 U.ml⁻¹ DNase I (Roche), 10 U.ml⁻¹ collagenase type I (Worthington) and 400 U.ml⁻¹ collagenase type IV (Worthington) diluted in 1X HBSS (Gibco)) for 20 min at 37°C. Tumor tissue was crushed with a syringe plunger and homogenized via trituration using standard Serological Pipettes and filtered twice using a nylon filter (100 µm). Next, the samples were centrifuged (515 g, 5 min, 4°C) and the cell pellet was resuspended with red blood cell (RBC) lysis buffer (ammonium chloride (NH₄Cl) 155 mM, sodium hydrogencarbonate (NaHCO₃) 12 mM, EDTA 0,1 mM). After a 3 min incubation, 9 ml of RPMI (Gibco) was added and the samples were centrifuged and resuspended in FACS buffer. A fraction of the single cell suspension from the samples ND7 and R1-4 were stained with anti-CD45 Alexa Fluor 647 (clone HI30) for 20 min on ice. The remaining fraction of unstained cells was frozen, in 90% heat-inactivated fetal calf serum (Gibco) and 10% DMSO (SeccoSolv, Merck Millipore, Cat. No.: 102931) and kept in liquid nitrogen for future studies. CITE-Seq experiments were performed on fresh samples ND8 and R5 and on frozen sample R2. Hereto, one million cells were counted, isolated and centrifuged. The cell pellet was resuspended and incubated for 30 min on ice with 25 µL of staining mix in PBS + 0.04% BSA containing Human TruStain FcX (BioLegend), anti-CD45 Alexa Fluor 647 (clone HI30) and the human cell surface protein antibody panel containing 268 oligo-conjugated antibodies (see Supplementary Table 14). Cells were washed and CD45⁺ immune cells were sorted using a BD FACS Aria II using an 85 µm nozzle. 7-AAD (Biolegend) was used to exclude dead cells. Typically, between 50 000 - 100 000 cells were sorted per sample. The complete dissociation and cell sorting procedure took approximately 2.5 hours and cells were maintained on ice whenever possible. For ND1-6, tumor samples were dissociated using the MACS Brain Tumor Dissociation Kit (P) (Miltenyi Biotec) according to the manufacturer's instructions. Briefly, tumor tissues were transferred into preheated C tubes containing the provided buffers and enzymes to enzymatically and mechanically dissociate the tissue in single cells. Cells were subsequently passed through a 70 µm cell strainer and red blood cells were removed using the RBC lysis buffer (Roche). Finally, following centrifugation, samples were washed in culture medium and eventually resuspended in 1 ml PBS containing 8 µl UltraPure BSA (50 mg/ml; ThermoFisher Scientific) and filtered over Flowmi 40 µm cell strainers (VWR) using wide-bore 1 ml low-retention filter tips (Mettler-Toledo). Next, live cells were counted using an automated cell counter (Luna). The entire procedure was completed in less than 2 hours and cells were maintained on ice whenever possible.

Mice

C57BL/6J and BALB/c mice were purchased from Janvier or bred in house. The following gene targeted mouse strains were used: CCR2-KO⁴⁹, *Cx3cr1*^{CreER}³⁰, *Cx3cr1*^{GFP}⁵⁰, R26-YFP⁵¹, NR4A1-KO⁵² which were bred in house. For 10x chromium scRNA-Seq and CITE-Seq, 7-12-week-old female CCR2-KO and C57BL/6J mice were used. Additional details regarding the strain, sex and age of mice used within this study can be found within the Reporting Summary. Mice were kept in standard housing conditions with 12h light and 12h dark. Food and water were provided ad libitum. The temperature was kept between 20-24°C and the humidity between 45%-65%. Animals were checked daily. A logbook was kept, each abnormal observation was recorded on an individual basis and the motility of the mice, posture and grooming behavior were carefully monitored. All mouse experiments were performed according to the regulations of the Ethische Commissie Dierproeven at the Vrije Universiteit Brussel.

Stereotactic intracerebral tumor cell inoculation

The GL261 cell line⁵³ was cultured in DMEM/F12 (1:1) (1x) - Dulbecco's Modified Eagle Medium: Nutrient Mixture F-12 (Gibco; catalog nr 11320-0033) containing 10% (v/v) heat-inactivated fetal calf serum (Gibco), 300 mg.ml⁻¹ L-glutamine (Gibco), 100 U.ml⁻¹ penicillin and 100 mg.ml⁻¹ streptomycin (Gibco). For intracranial injection cells were harvested via trypsinization (Gibco), brought to a concentration of 1x10⁵ cells/μl and injected as previously described⁵⁴. Briefly, mice were anesthetized (ketamine (140 mg/kg, Ketamidol, Ecuphar)/xylazine (10 mg/kg, Rompun, Bayer)), and immobilized in a stereotactic frame. A midline incision was made on the skin to expose the scalp and with a microdrill a craniotomy was made 2 mm anterior and 1.5 mm lateral to the bregma. 5x10⁵ cells in a total volume of 5 μl were slowly injected over 1 min at a depth of 3 mm with a Hamilton syringe (ga26s/51mm/pst2). Tumors were allowed to grow for 21 days. Mice were randomly distributed into the relevant experimental groups and injections with GL261 were alternated across these groups.

Mouse tumor and brain single-cell isolation, flow cytometry and cell sorting

For tumor single-cell isolation, mice were deeply anaesthetized and transcardially perfused with 20 ml of ice-cold PBS and brains were collected. The tumors were carefully micro-dissected using a stereo microscope to avoid contamination with healthy brain tissue. Tumor samples were cut into small pieces (2-3 mm) and Enzyme Mix A (30 U.ml⁻¹ DNase I (Roche), 10 U.ml⁻¹ collagenase type I (Worthington) and 400 U.ml⁻¹ collagenase type IV (Worthington) diluted in 1X HBSS (Gibco)) was added. Following 20 min at 37°C, tumor tissue was crushed with a syringe plunger and homogenized via trituration using standard Serological Pipettes and filtered twice using a nylon filter (100 μm). Next, the samples were centrifuged (515 g, 5 min, 4°C) and the pellet was resuspended with red blood cell (RBC) lysis buffer (ammonium chloride (NH₄Cl) 155 mM, sodium hydrogencarbonate (NaHCO₃) 12 mM, EDTA 0,1 mM). After a 3 min incubation, 9 ml of RPMI (Gibco) was added and the samples were centrifuged and

resuspended in FACS buffer (2 mM EDTA (Duchefa), 2% heat-inactivated fetal calf serum (Gibco) dissolved in 1X HBSS).

For the processing of GL261 surrounding brain tissue for single-cell isolation, brain tissue from which the tumor was removed, was minced and dissociated using Enzyme Mix A for 20 min at 37°C. Cells were passed through a cell strainer (100 μ m), centrifuged and resuspended in 5 ml 70% Standard Isotonic Percoll (SIP) (GE Healthcare) and gently overlaid with 5 ml of 37% SIP and 5 ml of 30% SIP (centrifuged at 800 g, 4°C, 30 min). The 70/37% interphase containing the immune cells was collected and after centrifugation resuspended in FACS buffer.

For the scRNA-Seq experiments (KO1-3, WT1-3) cells were stained with anti-CD45-APC/Cy7 (clone 30-F11) for 20 minutes on ice. For the CITE-Seq experiments (KO4, WT4), one million cells per sample were stained in 25 μ L PBS + 0.04% BSA staining buffer containing mouse TruStain FcX (Biolegend), anti-CD45-APC/Cy7 (clone 30-F11) and the mouse cell surface protein antibody panel containing 174 oligo-conjugated antibodies (see Supplementary Table 14), for 30 min on ice. Subsequently, the samples were washed and CD45⁺ immune cells were sorted using a BD FACS Aria II using an 85 μ m nozzle and collected in ME medium (RPMI containing 10% (v/v) heat-inactivated fetal calf serum (Gibco), 300 mg.ml⁻¹ L-glutamine (Gibco), 100 U.ml⁻¹ penicillin, 100 mg.ml⁻¹ streptomycin (Gibco), 1% Minimum essential medium non-essential aminoacids (100x) (Gibco), 1% pyruvate 100 mM (100x) (Gibco) and 1% β -mercaptoethanol (Sigma)). 7-AAD (Biolegend) was used to exclude dead cells. Typically, around 100,000 cells were sorted per sample. The complete dissociation and cell sorting procedure took approximately 2.5 hours. The single cell suspension was centrifuged for 6 min at 4°C at 400 g and resuspended in PBS + 0.04% bovine serum albumin in a final concentration of 1000 cells μ L⁻¹.

For multicolor flow cytometry cells were incubated for 15 min with Zombie Aqua Fixable Live-dead (BioLegend) at a concentration of 1/1000. Next, cells were washed with FACS buffer, centrifuged and blocked with rat anti-mouse CD16/CD32 (clone: 2.4G2) for 15 min on ice and subsequently stained for 20 min. All antibodies used are listed in Supplementary Table 20 and in the Reporting Summary. For measuring BrdU incorporation C57BL/6J and CCR2-KO tumor-bearing mice (d21 post GL261 inoculation), were injected intraperitoneally with 1 mg of BrdU (FITC BrdU Flow Kit, BD Biosciences, cat nr 559619) 2 hours prior to sacrifice. BrdU incorporation was assessed via flow cytometry on tumor single-cell suspensions following the manufacturers' instructions. Flow cytometry data was acquired using a BD FACS Canto II (BD Biosciences) using the BD FACSDiva v4.0 software and analyzed using FlowJo v10 (Tree Star, Inc.).

Single cell RNA-Seq and CITE-Seq using the 10x Genomics platform

Single-cell suspensions were loaded on a GemCode Single Cell Instrument (10x Genomics) to generate single-cell gel beads-in-emulsion (GEM). For the mouse GL261 samples (KO1-3, WT1-3) and for patients ND6-7 and R1-4, GEMs and scRNA-Seq libraries were prepared using the GemCode Single Cell 3' Gel Bead and Library kit, version 2 (10x Genomics, No. 120237) and the Chromium i7 Multiplex Kit (10x Genomics, No 120262) according to the manufacturer's instructions. For ND1-5 samples,

libraries were prepared using the GemCode Single Cell 5' Gel Bead and Library kit (10x Genomics, No. 1000002) and the Chromium i7 Multiplex Kit (10x Genomics, No. 120262) according to the manufacturer's instructions. For the mouse GL261 samples (KO4, WT4) and for patients ND8 and R5, GEMs and CITE-Seq libraries were prepared using the Chromium Next GEM Single Cell 3' Gel Bead and Library kit, version 3.1 (10x Genomics, No. 1000121). Briefly, GEM reverse-transcription incubation was performed in a 96-deep-well reaction module at 53°C for 45 min, 85°C for 5 min and ending at 4°C. Next, GEMs were broken and complementary DNA (cDNA) was cleaned up with DynaBeads MyOne Silane Beads (Thermo Fisher Scientific, No. 37002D) and SPRIselect Reagent Kit (Beckman Coulter, No. B23318). Full-length, barcoded cDNA originating from the mRNA, and from the oligonucleotide-labeled cell surface protein antibodies, was PCR amplified with a 96-deep-well reaction module at 98°C for 3 min, fourteen cycles at 98°C for 15 s, 67°C for 20 s and 72°C for 1 min, one cycle at 72°C for 1 min and ending at 4°C. For the 5' samples, the PCR amplification was performed according to the following protocol: 98°C for 45 s, fourteen cycles at 98°C for 20 s, 67°C for 30 s and 72°C for 1 min, one cycle at 72°C for 1 min and ending at 4°C. Size selection with the SPRIselect Reagent Kit was used to separate the amplified cDNA molecules for 3' gene expression and cell surface protein library construction. Gene expression library construction to generate Illumina-ready sequencing libraries was performed after cleanup with the SPRIselect Reagent Kit and enzymatic fragmentation, by adding R1 (read 1 primer), P5, P7, i7 sample index and R2 (read 2 primer sequence) via end-repair, A-tailing, adapter ligation, post-ligation SPRIselect cleanup/size selection and sample index PCR. For cell surface protein library construction, sample index PCR and SPRIselect size selection was performed. The cDNA content of pre-fragmentation and post-sample index PCR samples was analyzed using the 2100 BioAnalyzer (Agilent). Sequencing libraries were loaded on an Illumina HiSeq4000 or an Illumina NovaSeq flow cell with sequencing settings according to the recommendations of 10x Genomics (for 3' gene expression and cell surface protein libraries: read 1: 26 cycles; read 2: 98 cycles; index i7: eight cycles; index i5: no cycles, pooled in a 80:20 ratio for the combined 3' gene expression and cell surface protein samples, respectively; for 5' gene expression libraries: read 1: 26 cycles; read 2: 91 cycles; index i7: eight cycles; index i5: no cycles).

Single-cell and CITE-Seq gene expression data processing

The Cell Ranger pipeline (10x Genomics) was used to perform sample demultiplexing and to generate FASTQ files for read 1, read 2 and the i7 sample index for the gene expression and cell surface protein libraries, respectively. Read 2 of the gene expression libraries, containing the cDNA, was mapped to the reference genome (mouse mm10, or human GRCh38) using STAR. Subsequent barcode processing, unique molecular identifiers filtering and single-cell 3' or 5' gene counting was performed using the Cell Ranger suite and Seurat v.3.0.1. The average of the mean reads per cell across all gene expression libraries was 53,228, with an average sequencing saturation of 67.6%, as calculated by Cell Ranger. In total 19 individual scRNA-Seq libraries were created in this study, totaling 141,708 cells of which 119,377 were CD45⁺ immune cells. In addition, five CITE-Seq libraries were generated,

consisting of 45,987 CD45⁺ cells. Aggregation of sample conditions was done using the Cell Ranger Aggr software from 10x Genomics. Digital gene expression matrices were preprocessed and filtered using SCRAN and Scater R packages⁵⁵. Outlier cells were first identified based on three metrics (library size, number of expressed genes and mitochondrial proportion); cells were tagged as outliers when they were three median absolute deviations distant from the median value of each metric across all cells. Secondly, a principal component analysis plot was generated based on the following metrics: 'pct_counts_in_top_100_features', 'total_features_by_counts', 'pct_counts_feature_control', 'total_features_by_counts_feature_control', 'log10_total_counts_endogenous' and 'log10_total_counts_feature_control'. Outlier cells in this principal component analysis plot were identified using the R package mvoutlier. Low-abundance genes were removed using the 'calcAverage' function and the proposed workflow. The raw counts were normalized in Seurat by a global-scaling normalization and log-transform method 'LogNormalize' that normalizes the gene expression measurements for each cell by the total expression, and multiplies it by a scale factor (10,000), and log-transforms the result. Highly variable genes were detected in Seurat according to the method described in Stuart et al.⁵⁶ and the data was scaled by linear transformation. Subsequently, the highly variable genes were used for unsupervised dimensionality reduction. A previously described list of dissociation-induced genes¹, was removed from the highly variable gene list, to limit their effect on dimensionality reduction and clustering. Differential expression analysis between different cell clusters and between ND and recurrent GBM samples was performed using Wilcoxon Rank Sum test through the Seurat "FindMarkers" function. P value adjustment was performed using Bonferroni correction. For the differential expression analysis between ND and recurrent GBM samples, identified immunoglobulin genes were excluded from the results as potential artefacts, possibly due to ambient RNA contamination from dying plasma cells.

The samples for scRNASeq belonging to different patients and mice were processed, sorted and their libraries constructed on different days. Furthermore, the human samples were processed with either the 10x Genomics Chromium Single Cell 3' or 5' platforms. To remove batch effects and technical/biological noise the Harmony algorithm¹⁹ was used in Seurat v.3.0.1. Harmony projects cells into a shared embedding in which cells group by cell type rather than dataset-specific conditions. After soft k-means clustering which assigns cells to clusters while favoring a mixed dataset representation, cluster centroids for each dataset are used to compute cluster-specific linear correction factors which correct the assignment of the cluster-weighted average of each cell. This process is iterated until all cell-cluster assignments are stable. Default parameters for Harmony were used, the θ -value was adapted per dataset. This parameter decides the degree of penalty for dependence between batch membership and cluster assignment. A higher θ -value favors more independence between batch and cluster assignment. The θ parameter was set to one, with the exception of four datasets, where it was varied depending on their sensitivity to batch differences: the macrophage/monocyte/DC subset of the human GBM CITE-Seq aggregate dataset ($\theta = 0$), the macrophage/monocyte subset of the newly

diagnosed human GBM aggregate dataset ($\theta = 2$), the microglia subset of the recurrent human GBM aggregate dataset ($\theta = 2$), and the microglia subset of the newly diagnosed human GBM aggregate dataset ($\theta = 3$).

The corrected Harmony embeddings were next used, rather than principal components, to perform unsupervised clustering of the cells using graph-based clustering based on SNN-Cliq and PhenoGraph as implemented in the Seurat v.3.0.1 R package (default parameters). Clustering was visualized in two-dimensional scatter plots (via UMAP) using the Seurat v.3.0.1 package.

CITE-Seq cell surface protein (ADT) data processing

CITE-Seq ADT reads were quantified using the feature-barcoding functionality of Cell Ranger. All ADT reads associated with low-quality cells and doublets based on the CITE-Seq gene expression data processing, have been discarded from further analysis. On average across the 5 CITE-Seq libraries, the mean number of ADT reads per cell-associated barcode, which also contained a recognized antibody barcode and a valid UMI, was 2668. The average sequencing saturation of the ADT reads was 49.8%.

ADT count data has been normalized using ASINH_GEOM transformation: inverse hyperbolic sine (asinh) transformation with an additional cofactor, which is a modified version of the geometric mean of the counts per antibody¹⁸. The normalization takes as input the ADT count matrix:

$$X \in \mathbb{N}_0^{n \times p},$$

where n is the number of cells, p is the number of antibodies, and $\mathbb{N}_0$ is the set of natural numbers $\{0, 1, 2, \dots\}$. Each column is a n -dimensional vector:

$$x^{(j)} = [x_1^{(j)} \ x_2^{(j)} \ \dots \ x_n^{(j)}]^T,$$

where $j = 1, \dots, p$ is the ADT index, $x_i^{(j)}$ is the UMI count of antibody j in cell i . The ASINH_GEOM normalization is applied element-wise over the ADT count vector:

$$\text{ASINH_GEOM}(x^{(j)}) = \text{asinh}\left(\frac{x^{(j)}}{M^{(j)}}\right),$$

The cofactor is a p -dimensional vector $M = [M^{(1)} \ M^{(2)} \ \dots \ M^{(p)}]$, where $M^{(j)}$ for antibody j equals:

$$M^{(j)} = \sinh\left(\frac{1}{n} \sum_{i=1}^n \text{asinh}\left(x_i^{(j)}\right)\right).$$

Asinh is a widely used transformation in flow and mass cytometry data processing⁵⁷. An advantage of applying this transformation for proteomic data analysis is a better delineation of positive and negative populations. In the high-value range it has similar behavior to a log-transformation, while for values approaching zero it is close to linear, therefore it can handle zero and small counts⁵⁸. The cofactor M

1204 defines the width of the linear or less log-like region of the transformed values. For the cofactor M we
 1205 did not use the standard definition of geometric mean in $\mathbb{R}_0^+$: the n^{th} root of the product of n numbers,
 1206 which becomes zero in presence of a single zero value. Instead, we made use of the fact that the
 1207 geometric mean can also be expressed as the exponential of the arithmetic mean of logarithms (in
 1208 $\mathbb{R}_{>0}$):

$$1209 \exp \left[\frac{1}{n} \sum_{i=1}^n \ln(a_i) \right],$$

1210 for a set of numbers $\{a_1, a_2, \dots, a_n\}$. This allowed us to calculate the cofactor M analogously to the
 1211 geometric mean in asinh scale, rather than in logarithmic scale. Therefore, we named our
 1212 transformation ASINH_GEOM.

1213 To visualize the ADT data, UMAP projections were calculated using the normalized ADT count matrix
 1214 that was centered and scaled. Antibodies with low expression in the studied samples were filtered out
 1215 before calculating the UMAP projections, based on inspection of the density plots and feature plots for
 1216 each antibody. For the human ADT data, library-specific batch effect was clearly present on the
 1217 UMAP plots, therefore Harmony correction was applied on the normalized ADT count matrix, followed
 1218 by centering and scaling. The default parameters of Harmony were applied, except the θ value, which
 1219 was set to zero to apply moderate batch correction.

1220 *Human antibodies used for UMAP projection calculation:*

1221 C5L2, CD10, CD102, CD103, CD105, CD106, CD107a, CD109, CD112, CD115, CD117, CD119,
 1222 CD11a, CD11b-A0014, CD11b-A0161, CD11c, CD123, CD126, CD127, CD13, CD134, CD137, CD14-
 1223 A0051, CD14-A0081, CD141, CD142, CD144, CD146, CD15-A0076, CD15-A0392, CD150, CD151,
 1224 CD155, CD158, CD158b, CD158e1, CD16, CD161, CD162, CD163, CD164, CD169, CD172a, CD177,
 1225 CD18, CD184, CD19, CD192, CD194, CD195, CD196, CD198, CD1a, CD1c, CD1d, CD2, CD20,
 1226 CD201, CD203c, CD204, CD205, CD206, CD209, CD21, CD218a, CD22, CD226-A0368, CD23, CD24,
 1227 CD244, CD25, CD26, CD267, CD27-A0154, CD27-A0191, CD270, CD271, CD272, CD273, CD275-
 1228 A0172, CD276, CD278, CD279, CD28, CD29, CD3-A0034, CD3-A0049, CD303, CD304, CD305,
 1229 CD307e, CD309, CD31, CD314, CD32, CD325, CD33, CD335, CD35, CD354, CD36, CD366, CD371,
 1230 CD38-A0389, CD38-A0410, CD39, CD4-A0045, CD4-A0072, CD40, CD41, CD43, CD44-A0125,
 1231 CD45-A0048, CD45-A0391, CD45RA, CD45RO, CD46, CD48, CD49a, CD49b, CD49d, CD49f, CD5,
 1232 CD52, CD54, CD55, CD56-A0047, CD57, CD58, CD59, CD62L, CD62P, CD63, CD64, CD66a-c-e,
 1233 CD66b, CD69, CD7, CD71, CD72, CD73, CD8, CD81, CD83, CD85g, CD85j, CD86, CD8a, CD9,
 1234 CD90, CD93, CD94, CD95, CD98, EGFR, ERK1, FcεR1a, FolateReceptorBeta, HLA-A2, HLA-ABC,
 1235 HLA-DR, IgD, IntegrinB7, IRF4, KLRG1-A0153, LOX-1, MERTK, Podoplanin, TCR-Va24-Ja18, TCR-
 1236 Va7.2, TCRab, Tim-4, TMEM119, TSLPR

1237 *Mouse antibodies used for UMAP projection calculation:*

CD102, CD103, CD105, CD106, CD107a, CD115, CD117, CD11a, CD11b, CD11c, CD124, CD127, CD134, CD137, CD138, CD14, CD150, CD16-CD32, CD169, CD172a, CD19, CD195, CD1d, CD2, CD200, CD200R, CD200R3, CD201, CD204, CD206, CD207, CD21-CD35, CD22, CD223, CD226, CD24, CD25, CD26, CD27, CD270, CD272, CD274, CD278, CD279, CD28, CD29, CD3, CD300c-d, CD300LG, CD301a, CD301b, CD304, CD31, CD317, CD326, CD357, CD36, CD366, CD371, CD38, CD39, CD4, CD41, CD43, CD45R-B220, CD48, CD49b, CD49d, CD49f, CD5, CD54, CD55, CD62L, CD63, CD64, CD69, CD71, CD73, CD80, CD86, CD8a, CD8b, CD9, CD90-2, CD95, CX3CR1, ENPP1, F4-80, FcεR1a, FRB, IA-IE, IgD, IL33Ra, Integrin-b7, JAML, KLRG1, Ly49D, Ly49H, Ly6A-Ly6E, Ly6C, Ly6G, Mac2, MERTK, NK1-1, P2X7R, PIRA-PIRB, SiglecH, TCRb, TIGIT, Tim4, TLR4, XCR1

Human-mouse comparison of the differentially expressed genes in TAM and DC subsets

Differential expression analysis was performed in GBM recurrent and mouse TAM aggregates for the TAM subsets and in the human and mouse DC aggregates for the DC subsets, respectively. The Seurat “FindMarkers” function was used, by applying the Wilcoxon Rank Sum test and setting the logfc.threshold (minimum logFC of a gene) and the min.pct (minimum fraction of cells, in which the gene is expressed) thresholds to zero in order to get an output for all genes in the respective human and mouse datasets. P value adjustment was performed using Bonferroni correction. The gene symbols were converted between human and mouse based on one-to-one orthology, using biomaRt v.2.44.1. The top 150 most significantly differentially expressed genes were manually curated based on the ortholog information in the Mouse Genome Informatics website (http://www.informatics.jax.org/faq/ORTH_gene.shtml).

SCORPIUS trajectory inference

SCORPIUS trajectory inference was performed on Mg-TAM clusters of mouse tumors and microglia from surrounding healthy brain tissue. SCORPIUS was run on the log₂ normalized counts using the proposed workflow⁵⁹.

Proliferation mapping

For the proliferation mapping we used the FindTransferAnchors function of the Seurat R package (v3.1.0). We first created a Seurat object containing all proliferating cells and a second Seurat object containing all the remaining cells. Next, we applied the FindTransferAnchors function with the non-proliferating cells as the ‘reference’ and the proliferating cells as the ‘query’. Predicting to which cluster the proliferating cells belong was done using the TransferData function with the output of FindTransferAnchors for the ‘anchorset’ parameter and the cluster IDs of the non-proliferating cells for the ‘refData’ parameter.

Gene ontology enrichment analysis and heatmaps

To predict putative biological functions based on differential gene expression, we performed a gene ontology analysis. Genes that were differentially expressed in human recurrent tumors between lipid

sign. Mo-TAMs and transitory Mo-TAMs (adjusted P-value cutoff of 10^{-10} , $\log_2$ FC cutoff of 0.5), between hypoxic sign. Mo-TAMs and the other Mo-TAM subsets (adjusted P-value cutoff of 10^{-20} , $\log_2$ FC cutoff of 0.5), between Resident-like Mo-TAMs and Transitory Mo-TAMs (adjusted P-value cutoff of 10^{-20} , $\log_2$ FC cutoff of 0.5), and between IFN-sign. Mo-TAMs and resident-like Mo-TAMs (adjusted P-value cutoff of 10^{-20} , $\log_2$ FC cutoff of 0.8) were inserted into the Metascape (<http://metascape.org/>)⁶⁰ online tool with default parameters.

The heatmaps of Fig. 2, 6d and Extended Data Fig. 4 were made using Morpheus (<https://software.broadinstitute.org/morpheus>).

Tamoxifen treatment

3-4-week old *Cx3cr1^{CreER}*:R26-YFP mice (double heterozygous) were injected with 4 mg tamoxifen (Sigma-Aldrich), dissolved in 200 μ l cornoil (Sigma-Aldrich). Injections were performed subcutaneously near the fore and hind limbs (4 x 50 μ l) and repeated three times at 48-hour intervals. GL261 tumor inoculations were performed 4 weeks after the final tamoxifen injection.

Adoptive transfer experiments

Blood was collected from naive *Cx3cr1^{GFP/+}* mice via heart puncture using heparin-filled syringes. RBCs were lysed using RBC lysis buffer followed by neutralization with RPMI and resuspension in FACS buffer. Classical and non-classical monocytes were sorted as shown in Supplementary Fig. 5a and were intravenously injected in tumor bearing CCR2-KO, C57BL/6J and NR4A1-KO mice (at d16 post GL261 inoculation). CCR2-KO mice were injected with 7×10^5 non-classical monocytes or 1×10^6 classical monocytes. C57BL/6J and NR4A1-KO mice were injected with 7×10^5 non-classical or 7×10^5 classical monocytes. At day 21 mice were sacrificed and tumors were processed for flow cytometry.

TAM cytokine secretion

Sorted TAM subsets were cultured in ME medium, in 96-well flat bottom plates. 20 hours later the plates were spun down (805 g, 5 min, 4°C), the supernatant was collected and Luminex technology was used to detect the cytokines IL10, TNF, CCL4 and CCL2 (Bio-Rad) according to the manufacturers' instructions.

Chorioallantoic membrane assay

Chorioallantoic membrane (CAM) assays were performed as described earlier⁶¹. Gelatin sponges (1–2 mm³; Hospithera) with 2×10^4 sorted TAM subsets or brain microglia were placed on the CAM. PBS/0.1% bovine serum albumin (BSA; 50 μ g/CAM) served as control. At day 13, membranes were fixed and analyzed using a Zeiss Lumar V.12 stereomicroscope with NeoLumar S 1.5 $\times$ objective (15 $\times$ magnification).

T-cell suppression assay

Splenocytes were isolated from the spleen of naive female C57BL/6J mice. The spleen was crushed using a syringe plunger and homogenized via trituration using standard Serological Pipettes. The sample was then lysed with RBC lysis buffer (5 ml, 2 min), blocked with 30ml of RPMI, filtered and centrifuged. The pellet was resuspended in ME medium and was plated in flat-bottom 96-well plates (Falcon) at 2×10^5 splenocytes/well. Sorted TAM subsets were added at 1×10^5 TAMs/well (in triplicate) and the cultures were treated with 1 μ g/ml of anti-CD3 (clone 145-2C11). 24 hours later [3 H]thymidine was added and cells were allowed to proliferate for another 18 hours prior to measurement on a Wallac 1450 MicroBeta TriLux Liquid Scintillation Counter. % suppression was calculated as: $(1 - [\text{proliferation with TAMs}] / [\text{proliferation without TAMs}]) \times 100$.

Mixed leukocyte reaction (MLR) assay

CD4⁺ and CD8⁺ T cells were isolated from spleen single-cell suspensions (see T-cell suppression assay) of naive BALB/c mice via successive steps of MACS sorting. First, CD11c⁺ (CD11c beads - catalog nr. 130-108-338, lot nr. 5190903182, Miltenyi Biotec) and CD19⁺ cells (CD19 beads – catalog nr. 130-052-201, lot nr. 5190306670, Miltenyi Biotec) were depleted. Next, CD8⁺ T cells were purified (CD8 beads - catalog nr. 130-049-401, lot nr. 5180308206, Miltenyi Biotec), followed by CD4⁺ T cells (CD4 beads - catalog nr. 130-108-043, lot nr. 5190903039, Miltenyi Biotec). 2×10^5 CD4⁺ or CD8⁺ T cells were co-cultured, in ME medium, in 96-well round bottom plates, with different ratios of sorted TAM subsets or splenic C57BL/6J cDCs (1:2, 1:4, 1:8 and 1:16 TAM/cDC:T cell ratio). 3 days later cultures were pulsed with [3 H]thymidine and cells were allowed to proliferate for another 18 hours prior to measurement on a Wallac 1450 MicroBeta TriLux Liquid Scintillation Counter. The splenic cDCs (cDC1 and cDC2) were obtained from 10-15 spleens of GL261 tumor-bearing C57BL/6J mice. Spleens were flushed with 200 U.ml⁻¹ of collagenase III (Worthington) and incubated at 37°C for 30 min. Next, spleens were processed to single-cell suspensions and CD11c⁺ cells were enriched using MACS-sorting with CD11c beads. cDC1 and cDC2 subsets were subsequently FACS sorted based on the gating strategy shown in Supplementary Fig. 5b.

PLX3397 treatment of tumor-bearing mice

C57BL/6J and CCR2-KO mice were injected with GL261 after which half of the mice were fed control chow and the others PLX3397 chow. PLX3397 (Advanced Chemblock Inc.) was incorporated in AIN-76A chow by Research Diets inc. at a concentration of 600 mg/kg. At day 21, mice were sacrificed, and the tumors were micro-dissected, weighed and processed for flow cytometry.

Immunohistochemistry of mouse brain sections

Tumor-bearing mice were deeply anaesthetized and transcardially perfused with ice-cold PBS followed by 4% ice-cold paraformaldehyde (PFA)/PBS solution. Brains were removed, post-fixed in 4% PFA/PBS at 4°C for 4 hours and dehydrated with sucrose/PBS solution (15% o/n, 30% o/n). Brains were embedded in Tissue-Tek O.C.T™ compound (Sakura Finetek Europe B.V. Alphen aan den Rijn, NL) and frozen using dry ice. 12 μ M cryosections were made using a Leica CM1850 UV cryostat. Sections

were blocked and permeabilized with 0.1% PBS-Triton™X-100 (PBS-T) (Sigma-Aldrich) containing 10% Normal Donkey Serum (NDS) (v/v) (Sigma) for 1 hour. Sections were incubated with primary antibodies (diluted in a PBS-T solution containing 3% NDS) o/n at 4°C in a moist chamber. For visualizing hypoxia, tumor-bearing mice were injected ip with pimonidazole (Hypoxyprobe™ Omni Kit, catalog nr HP3-100) at a concentration of 60 mg/Kg of body weight, 2 hours prior to sacrifice. The following primary antibodies were used: chicken anti-GFP (id. Ab13970, polyclonal, lot nr. GR3190550-3, Abcam, 1/500), rabbit anti-IBA-1 (id. 019-19741, polyclonal, lot nr. WDE1198, Wacko, 1/250), rat anti-F4/80 (id. MCA497GA, monoclonal, lot nr. 1610, Biorad, 1/250), goat anti-arginase1 (id. NB100-59740, polyclonal, lot nr. S2C2E08719, NovusBio, 1/500), rabbit anti-pimonidazole (id. Pab2617(AP) rabbit anti-sera, lot nr. 022319, Hypoxyprobe™ Omni Kit, catalog nr HP3-100, 1/100). The following secondary antibodies were used: goat anti-chicken Alexa Fluor 488 (id. A11039, polyclonal, lot nr. 1691381, Life Technologies, 1/500), donkey anti-rabbit Alexa Fluor 647 (id. A31573, polyclonal, lot nr. 1693297, Life Technologies, 1/500) and goat anti-rat Alexa Fluor 555 (id. A21434, polyclonal, lot nr. 1722994, Life Technologies, 1/500), donkey anti-goat Alexa Fluor 568 (id. A11057, polyclonal, lot nr. 1975005, Invitrogen, 1/500), donkey anti-rat Alexa Fluor 488 (id. A21208, polyclonal, lot nr. 1789917, Life Technologies, 1/500). Images were acquired using the Zeiss LSM880 Fast Airy scan. For quantification of %YFP within the IBA1+/F4/80+ population in Fig. 4a, 13-14 sections/images were analyzed per region, for each mouse, using Fiji (ImageJ) version 2.1.0/1.53c.

MILAN Multiplex immunohistochemistry of human brain sections

Tissue staining

Multiplex immunohistochemistry was performed according to the previously published method⁶²⁻⁶⁴. Tissue sections of 3 µm were prepared out of formalin-fixed paraffin-embedded (FFPE) human glioblastoma samples (collected at the UZ/KULeuven biobank according to protocols S59804, S61081 and S62248). Following dewaxing, antigen retrieval was performed using PT link (Agilent) using 10mM EDTA in Tris-buffer pH 8. Immunofluorescent staining was performed using the Bond RX Fully Automated Research Stainer (Leica biosystems) with the primary antibodies as indicated in Supplementary Table 21. The sections were incubated during 4 hours with the primary antibodies, washed and visualized with secondary antibodies (30 min; see Supplementary Table 21). To amplify the signal, negative isotype controls were added for 30 min followed by a second incubation with the same secondary antibody for 30 min after three washing steps. A coverslip was placed on the slides with medium containing DAPI and were scanned in the Zeiss Axio Scan Z.1 (Zeiss) with 10X magnification. After completion of the staining procedure, the coverslips were manually removed after 30 min soaking in washing buffer. Consecutive washing steps were thereafter performed using TBS. Stripping of the antibodies was performed in a buffer containing 1% SDS and β-mercaptoethanol during 30 min at 56°C. After this stripping process, the slides were washed in washing buffer for 45 min with frequent changes of the buffer. The staining procedure was thereafter repeated until all markers were stained and scanned on each of the slides.

Image processing

Raw scans were transformed to grayscale 16-bit TIFF images using developer's software (ZEN). Image registration was performed by applying a homomorphic transformation over a set of matched descriptors using a Harris detector. Images were subsequently adjusted for background intensity variations using the rolling ball algorithm, and Autofluorescence (AF) subtraction was performed by subtracting the pre-stained image of the corresponding tissue section from the measured signal (MS). Intensities for both images were normalized using quantile normalization. For cellular segmentation, first a mask was generated in the DAPI channel of the pre-stained image using a local thresholding approach; second, a distance map was calculated using the binary image; and third, the distance map was split in cellular objects using the watershed algorithm. DAPI positive objects were expanded 5 pixels to capture the expression of markers present in the cell cytoplasm. For each cellular object, morphological (nuclear size) and functional (marker intensity) features were extracted. Given that not the whole cell surface expresses all the markers, the expression of every cell was summarized by the 95% quantile of the expression of all its pixels.

Data analysis

Phenotypic identification was implemented as described in Bosisio et al⁶³. Briefly, three different clustering methods were applied (PhenoGraph, FlowSom, and KMeans), from which only those cells that agreed over at least 2 algorithms using a consensus-based approach were retained. Agreeing clusters were then manually annotated by expert pathologists and assigned to the right cell type. Macrophages were further characterized into TMEM119^{high}/low functional groups. To that end, the expression of TMEM119 was z-normalized using the population of all cells and a threshold for high/low discretization was set at $z_tmem119 = 1$. In addition, TMEM119 expression was split in a range in 5 equal bins to define colors as shown in the digital images. Regarding spatial analysis of TMEM119 cells, each tumor section was divided into windows of 100x100 pixels, in which we defined the number of TMEM119^{hi} and TMEM119^{lo} cells based on the above defined criteria, and subsequently calculated and plotted colored boxes on the original image with the following criteria: if TMEM119^{hi}/CD68⁺ >50% of TAMs in window = red; otherwise = cyan).

Statistics

All graphs represent mean $\pm$ s.e.m. The statistical tests that were used are indicated in the figure legends and in Supplementary Table 19. All the statistics performed were done using GraphPad 8.3.0. Differential gene expression was assessed using the Wilcoxon rank sum test (two-sided) as implemented in the Seurat R package. P value adjustment was performed using Bonferroni correction based on the total number of genes in the dataset. Data distribution was assumed to be normal, but this was not formally tested. The sample sizes were determined by means of power analysis, using G*Power v.3.1.9.3. Data collection and analysis were not performed blind to the conditions of the experiments.

Data availability

All human and mouse scRNA-Seq and CITE-Seq datasets described in this article can be accessed via our interactive webserver: www.brainimmuneatlas.org, where all gene-cell count and cell annotation matrices can also be downloaded. Through this tool, it is possible to evaluate the expression of genes or proteins via CITE-Seq antibody staining in our datasets and to download DE gene lists. Additionally, all mouse single-cell RNA-Seq/CITE-Seq raw data, mouse gene-cell count matrices and human gene-cell count matrices are deposited at GEO (NCBI) under accession number GSE163120. Raw sequencing reads of the human scRNA-Seq/CITE-Seq experiments have been deposited in the controlled access public repository European Genome-phenome Archive (EGA), under study accession number EGAS00001004871. Other data that support the findings of this study are available from the corresponding author upon request.

Code availability

The R codes that were used for scRNA-Seq and CITE-Seq analysis can be found at Github: <https://github.com/Movahedilab/Glioblastoma>

Methods-only References

49. Boring, L., *et al.* Impaired monocyte migration and reduced type 1 (Th1) cytokine responses in C-C chemokine receptor 2 knockout mice. *J Clin Invest* **100**, 2552-2561 (1997).
50. Jung, S., *et al.* Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and green fluorescent protein reporter gene insertion. *Mol Cell Biol* **20**, 4106-4114 (2000).
51. Srinivas, S., *et al.* Cre reporter strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. *BMC Dev Biol* **1**, 4 (2001).
52. Lee, S.L., *et al.* Unimpaired thymic and peripheral T cell death in mice lacking the nuclear receptor NGFI-B (Nur77). *Science* **269**, 532-535 (1995).
53. Ausman, J.I., Shapiro, W.R. & Rall, D.P. Studies on the chemotherapy of experimental brain tumors: development of an experimental model. *Cancer Res* **30**, 2394-2400 (1970).
54. Baumann, B.C., Dorsey, J.F., Benci, J.L., Joh, D.Y. & Kao, G.D. Stereotactic intracranial implantation and in vivo bioluminescent imaging of tumor xenografts in a mouse model system of glioblastoma multiforme. *J Vis Exp* (2012).
55. Lun, A.T., McCarthy, D.J. & Marioni, J.C. A step-by-step workflow for low-level analysis of single-cell RNA-seq data with Bioconductor. *F1000Res* **5**, 2122 (2016).
56. Stuart, T., *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821 (2019).
57. Bendall, S.C., *et al.* Single-cell mass cytometry of differential immune and drug responses across a human hematopoietic continuum. *Science* **332**, 687-696 (2011).
58. Huber, W., von Heydebreck, A., Sultmann, H., Poustka, A. & Vingron, M. Variance stabilization applied to microarray data calibration and to the quantification of differential expression. *Bioinformatics* **18 Suppl 1**, S96-104 (2002).
59. Cannoodt, R., *et al.* . SCORPIUS improves trajectory inference and identifies novel modules in dendritic cell development. *Preprint at bioRxiv* <https://www.biorxiv.org/content/10.1101/079509v2> (2016).
60. Zhou, Y., *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun* **10**, 1523 (2019).
61. Movahedi, K., *et al.* Different tumor microenvironments contain functionally distinct subsets of macrophages derived from Ly6C(high) monocytes. *Cancer Res* **70**, 5728-5739 (2010).
62. Bolognesi, M.M., *et al.* Multiplex Staining by Sequential Immunostaining and Antibody Removal on Routine Tissue Sections. *J Histochem Cytochem* **65**, 431-444 (2017).
63. Bosisio, F.M., *et al.* Functional heterogeneity of lymphocytic patterns in primary melanoma dissected through single-cell multiplexing. *Elife* **9** (2020).

1479 64. Cattoretti, G., Bosisio, F.M., Marcelis, L. & Bolognesi, M.M. . Multiple Interactive Labeling by
1480 Antibody Neodeposition (MILAN). *Protoc. Exch.* **1–21** (2018).

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 7

Fig. 8

a

Zoom on monocytes/macrophages:

b **c** **d** **e** **f**

a

Human recurrent Mo-TAMs

b **c**

Human recurrent Mo-TAMs

d