Article type : Original

A novel mouse model of cholangiocarcinoma uncovers a role for Tensin-4 in tumor progression

Short title: Tumor progression in cholangiocarcinoma

Mickaël Di-Luoffo¹, Sophie Pirenne^{1,2,#}, Thoueiba Saandi^{1,#}, Axelle Lorient^{1,#}, Claude Gérard¹, Nicolas Dauguet^{1,3}, Fátima Manzano-Núñez¹, Natália Alves Souza Carvalhais⁴, Florence Lamoline¹, Sabine Cordi¹, Katarzyna Konobrocka¹, Vitaline De Greef¹, Mina Komuta⁵, Georg Halder⁴, Patrick Jacquemin¹, and Frédéric P. Lemaigre¹

¹ de Duve Institute, Université catholique de Louvain, Brussels, Belgium.

² Department of Pathology, Cliniques Universitaires Saint-Luc, Université catholique de Louvain, Brussels, Belgium.

³ CYTF platform, Université catholique de Louvain, Brussels, Belgium.

⁴ VIB Center for Cancer Biology and KU Leuven Department of Oncology, University of Leuven, Belgium.

⁵ Department of Pathology, Keio University School of Medicine, Japan.

# These authors contributed equally

Mickael Di-Luoffo, diluoffo.mickael@gmail.com; Sophie Pirenne, sophie.pirenne@uclouvain.be; Thoueiba Saandi, taanlimi@yahoo.fr; Axelle Lorient, axelle.lorient@uclouvain.be; Claude Gérard, claudegerard1982@gmail.com; Sabine

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/HEP.31834](#)

This article is protected by copyright. All rights reserved

Cordi, sabine.cordi@uclouvain.be; Nicolas Dauguet, nicolas.dauguet@uclouvain.be; Fátima Manzano-Núñez, fatima.manzano@uclouvain.be; Natália Alves Souza Carvalhais, natalia.souza@kuleuven.be; Florence Lamoline, florence.lamoline@student.uclouvain.be; Katarzyna Konobrocka, k.konobrocka@yahoo.com; Vitaline De Greef, vitaline.degreeef@gmail.com; Mina Komuta, mina.komuta@gmail.com; Georg Halder, georg.halder@vib.be; Patrick Jacquemin, patrick.jacquemin@uclouvain.be; Frédéric P Lemaigre, frederic.lemaigre@uclouvain.be.

Keywords: intrahepatic cholangiocarcinoma; intraductal papillary neoplasm of bile ducts; cholangitis; KRAS; SOX17

Footnotes

Corresponding author: Frédéric P. Lemaigre, de Duve Institute, Université catholique de Louvain, Avenue Hippocrate 75/B1-7503, B-1200 Brussels, Belgium. Email: frederic.lemaigre@uclouvain.be. Phone: +32 2 764 7583. Fax +32 2 764 7573.

List of abbreviations: AmotL2, angiomin like-2; BillN, biliary intraepithelial neoplasia; Ctgf, connective tissue growth factor; CK19, cytokeratin 19; Cyr61, cysteine-rich angiogenic protein 61; DDC, 3,5-diethoxycarbonyl-1,4 dihydrocollidine; EGF, epidermal growth factor; EGFR, epidermal growth factor receptor; EpCAM, epithelial cell adhesion molecule; ERK, extracellular-regulated kinase; ERBB2, v-erb-b2 erythroblastic leukemia viral oncogene homolog 2/human epidermal growth factor receptor 2; eYFP, enhanced yellow fluorescent protein; HES1, hairy and enhancer of split-1; H&E, hematoxylin and eosin; IL-33, interleukin-33; iCCA, intrahepatic cholangiocarcinoma; IPNB, intraductal papillary neoplasm of bile ducts; KRAS, v-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog; MAPK, mitogen activated protein kinase PI3K, phosphoinositide-3-kinase; SMAD2, small mothers against decapentaplegic 2; SOX17, SRY-related HMG box transcription factor 17; TCGA, The Cancer Genome Atlas; TGF β , transforming growth factor beta; TNS4, tensin 4.

Financial Support: The work is supported by the National Fund for Scientific Research FRS-FNRS (Belgium, grants J003717 and J.0105.19 to FPL), Télévie (Belgium, grants 7.4544.18 to FPL and MK, and 7.8515.18 to PJ), the Foundation against Cancer (Belgium, grants 2014-125 and 2018-078 to FPL; 2016-089 and 2018-076 to PJ), the D.G. Higher Education and Scientific Research of the French Community of Belgium (grant ARC 15/20-065 to FPL), and the Fonds Joseph Maisin (Belgium, grant 2018-2019) to FPL and PJ. SP holds a PhD fellowship from Télévie (Belgium, grant 7.4544.18). ND is supported by the Belgian National Lottery. PJ is Senior Research Associate at the FRS-FNRS (Belgium). NAS and GH were supported by The Research Foundation – Flanders (FWO, grants G0C2418N and G069321N).

Abstract

Earlier diagnosis and treatment of intrahepatic cholangiocarcinoma (iCCA) are necessary to improve therapy, yet limited information is available about initiation and evolution of iCCA precursor lesions. Therefore, there is a need to identify mechanisms driving formation of precancerous lesions and their progression towards invasive tumors using experimental models that faithfully recapitulate human tumorigenesis. To this end, we generated a new mouse model which combines cholangiocyte-specific expression of *Kras*^{G12D} with 3,5-diethoxycarbonyl-1,4-dihydrocollidine (DDC) diet-induced inflammation to mimic iCCA development in patients with cholangitis. Histological and transcriptomic analyses of the mouse precursor lesions and iCCA were performed and compared with human analyses. The function of genes overexpressed during tumorigenesis was investigated in human cell lines. We found that mice expressing *Kras*^{G12D} in cholangiocytes and fed a DDC diet developed cholangitis, ductular proliferations, intraductal papillary neoplasms of bile ducts (IPNBs) and eventually iCCAs. The histology of mouse and human IPNBs were similar, and mouse iCCAs displayed histological characteristics of human mucin-producing large duct type iCCA. Signaling pathways activated in human iCCA were also activated in mice. The identification of transition zones between IPNB and iCCA on tissue sections, combined with RNA-sequencing analyses of the lesions supported that iCCAs derive from IPNBs. We further provide evidence that Tensin-4 (*TNS4*), which is stimulated by *KRAS*^{G12D} and SRY-related HMG box transcription factor 17 (*SOX17*), promotes tumor progression.

Conclusion. We developed a novel mouse model that faithfully recapitulates human iCCA tumorigenesis and identified a gene cascade which involves *TNS4* and promotes tumor progression.

Introduction

Prognosis of intrahepatic cholangiocarcinoma (iCCA) remains dismal. Since this results in part from delayed diagnosis, there is a need to understand the mechanisms that drive development of precancerous lesions and their progression towards invasive tumors. A range of perturbed signaling pathways and genomic alterations is associated with late-stage cholangiocarcinoma, but there is still limited knowledge about how such pathways and mutations evolve during tumorigenesis.⁽¹⁻³⁾ Moreover, our understanding of this process is further complicated by the observation that iCCA can arise from several cell types.⁽⁴⁻¹⁰⁾

Histopathological studies identified two iCCA subtypes in humans. Small duct subtype is located at the liver periphery and presents as tubular-type adenocarcinoma with cuboidal cells and ductular component without mucin production. The large duct subtype is found near the hilum and exhibits a tubular pattern with mucin-producing columnar cells.⁽¹¹⁾ Histopathological studies classified biliary intraepithelial neoplasia (BillIN), intraductal tubulopapillary neoplasms and intraductal papillary neoplasms of bile ducts (IPNB) as non-invasive precursor lesions.⁽¹²⁻¹⁴⁾ Whereas BillINs develop in large bile ducts and present as flat or micropapillary dysplasia of the bile duct epithelium, IPNBs are macroscopic lesions of the large ducts characterized by exophytic papillary structures with fibrovascular core and prominent intraductal growth.^(15, 16) von Meyenburg complexes are irregular ductal structures surrounded by dense stroma. These complexes are benign but were reported to evolve towards iCCA in rare instances.

Gain-of-function mutations of the small GTPase v-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog (*KRAS*) are detected in iCCA at a frequency of 15-25%, and are associated with a significant decrease of survival.⁽¹⁷⁻¹⁹⁾ Importantly, *KRAS* mutations were detected in BillINs and IPNBs in 25-50% of cases, suggesting that such mutations may contribute to initiate tumorigenesis.⁽²⁰⁻²²⁾ Chronic inflammation is a known risk factor of iCCA. Patients with primary sclerosing cholangitis are at risk for iCCA and molecular classification of late-stage cholangiocarcinomas underscored the importance of inflammatory gene expression in iCCA pathogeny.⁽²³⁻²⁵⁾

Several animal models of intra- and extrahepatic CCA have been designed.^(26, 27) However, limited attention was paid to tumor initiation and progression, and in several instances genetic alterations were induced at the prenatal stage or simultaneously in hepatocytes and cholangiocytes, unlike in humans. Here, we generated an original mouse model of iCCA tumorigenesis in which expression of oncogenic *Kras*^{G12D} is specifically induced in cholangiocytes and is combined with diet-induced inflammation that mimics chronic cholangitis. The mice developed IPNB similar to those in humans, and our data support that those lesions evolve toward iCCA. We also identified a gene regulatory pathway that is activated during tumorigenesis and which promotes cell proliferation and migration.

Experimental Procedures

Animals

Osteopontin-iCreERT² (Opn-Cre^{ER}), loxP-stop-loxP-Kras^{G12D/+} (LSL-Kras^{G12D/+}) and Rosa26R^{eYFP} mice were described⁽²⁸⁻³⁰⁾ and bred in a CD1-enriched background. For Cre-LoxP recombination, 6 week-old mice were injected intraperitoneally three times at two days interval with tamoxifen (T5648; Sigma-Aldrich, Bornem, Belgium) dissolved in corn oil (30 mg/mL, at a dose of 150 mg/kg). Mice were fed a standard diet or a diet containing 3,5-diethoxycarbonyl-1,4 dihydrocollidine 0.1% (DDC; 137030 Sigma-Aldrich) and monitored for signs of illness as described in the Results section and in Supporting Fig. S1.

Animal numbers: Wild-type mice or tamoxifen-treated Opn-Cre^{ER} mice (devoid of mutated *loxP-stop-loxP-Kras*^{G12D} allele) were indistinguishable at the histological level and were used as controls. The number of mice used in the study was as follows: wild-type *Kras* + DDC: 20; *Kras*^{G12D}: 30; *Kras*^{G12D}+DDC: 60.

Laser capture microdissection (LCM), RNA extraction and sequencing

The livers were collected, embedded in OCT, frozen in liquid nitrogen and stored at -80 °C. Microscope support glass, membrane slides and tools were pre-treated to remove RNase (RNaseZAP, Sigma-Aldrich). Ten µm sections were cut using Cryostar NX70

(Thermo Fisher Scientific) at -25°C for knife/blade temperature and -13°C for specimen temperature. Liver sections were deposited on slides adapted for LCM (SuperFrost Plus (25x75x1mm); VWR, Leuven, Belgium) and stored at -80°C inside plastic containers with silica granules to prevent tissue hydration. Within 12h, the liver sections were microdissected using Leica LMD7000 microscope (GIGA, Liège, Belgium) and RNA was extracted with RNAqueous®-Micro total RNA isolation AM1931 Kit (Thermo Fisher Scientific). RNA sequencing was performed by the Next Generation Sequencing Service (NGS Genewiz, South Plainfield, NJ, USA). Quality of samples was controlled by NanoDrop (NanoDrop 2000 (Thermo Fisher Scientific) and Qubit fluorometry (Invitrogen, Carlsbad, CA)) for measuring concentration and calculation of RNA integrity number (RIN) (Agilent 2100 Bioanalyzer System). RNA was amplified using SMARTer® Universal Low Input RNA Kit for sequencing (Takara, Mountain View, CA, USA). Ribosomal RNA was depleted using RiboGone™-Mammalian rRNA removal kit (Takara, Mountain View, CA, USA), and cDNA libraries were generated using the TruSeq total stranded RNA-seq kit (Illumina, San Diego, CA, USA). cDNA libraries were sequenced using Illumina HiSeq 2x150bp configuration with estimated data output ≥350M raw paired-end reads per lane. Data are stored under GEO accession number GSE141511.

Other Materials and Methods are available in the Supporting Information.

Results

Cholangiocyte-specific expression of $Kras^{G12D}$ combined with DDC-induced cholangitis causes development of IPNB and iCCA.

To recapitulate tumorigenesis in human iCCA, we generated a mouse model which combines cholangiocyte-specific $Kras^{G12D}$ expression with chronic cholangitis. Opn-Cre^{ER} mice, which express tamoxifen-inducible Cre^{ER} recombinase specifically in cholangiocytes, were mated with LSL- $Kras^{G12D/+}$ mice, which contain a inducible mutant $Kras^{G12D}$ allele.⁽²⁸⁻³⁰⁾ The mice also contained a $Rosa26R^{eYFP}$ allele. $Kras^{G12D}$ expression was induced by administration of tamoxifen to 6 week-old mice. Mice were then fed a normal diet for 4 weeks to enable wash out of tamoxifen. Thereafter, chronic cholangitis

was induced by feeding the mice a 3,5-diethoxycarbonyl-1,4-dihydrocollidine (DDC) diet.⁽³¹⁾ DDC feeding was interrupted after 2 weeks because of weight loss. Following 1 week of weight recovery under normal diet, DDC diet was resumed, and livers were collected after 8 and 18 weeks of cholangitis, and at several time points between 21 and 31 weeks of cholangitis (Fig. 1A). During that period the mice accepted the DDC diet; the experiments were interrupted when animals showed signs of severe illness and had reached humane endpoints. These conditions did not allow to reveal a significant impact of *Kras*^{G12D} on the survival of DDC-treated animals (Supporting Figs S1A-C).

Expression of *Kras*^{G12D} in the absence of DDC did not affect bile duct morphology. Administration of DDC for 8 weeks to wild-type mice induced accumulation of porphyrin plugs, ductular proliferation, as well as emerging papillary lesions and periductal fibrosis of large ducts. In contrast, expression of *Kras*^{G12D} in mice receiving an 8-week DDC treatment was associated with large duct lesions which displayed papillary proliferation, similar to IPNB (Supporting Fig. S1D).

At 18 weeks of DDC feeding, large duct lesions in mice expressing wild-type *Kras* had evolved to IPNB, and ductular proliferations were more extended (Fig. 1B). Expression of *Kras*^{G12D} in the absence of DDC still did not affect the histology of the biliary epithelium (Figs. 1B and 2A). At 18 weeks of DDC treatment, expression of *Kras*^{G12D} was associated with IPNBs which did not significantly differ from those monitored in mice expressing wild-type *Kras* (Fig. 1B).

Starting at 21 weeks of DDC diet, iCCAs were observed, but only in mice combining DDC-induced cholangitis and cholangiocyte-specific *Kras*^{G12D} expression (Fig. 1C). At that stage iCCAs had developed in one third of cases (12/36 males; 9/24 females) and showed tubular structures with desmoplastic changes typical of human cholangiocarcinoma (Fig. 1C). iCCAs were proliferative as illustrated by Ki67 staining, and expressed cytokeratin 19 (CK19) and epithelial cell adhesion molecule (EpCAM); they also expressed eYFP, supporting their biliary origin (Fig. 2A). IPNBs maintained their biliary fate as evidenced by the expression of SOX9, and lack of HNF4. iCCA cells expressed SOX9. A subset of iCCA cells were HNF4-positive (Fig. 2A). Given the high

cholangiocyte specificity of Opn-Cre^{ER}, it is unlikely that these HNF4-positive cells derive from hepatocytes; rather, HNF4 expression likely reflects dedifferentiation. The epithelium of the iCCA was columnar to cuboidal, and was mucin-positive as shown by alcian blue staining (Fig. 2B). This phenotype is similar to the human mucin-producing large duct type iCCA. Some tumors appeared less desmoplastic (Fig. 2B). Interestingly, adjacent to large ducts, we often found areas of glandular proliferation with expansive growth pattern and cytonuclear atypia, which we qualified as early stage of invasive cancer. Such lesions were detected starting around 18 weeks of DDC treatment (Supporting Fig. S1C). Importantly, focal transitions between IPNB and early invasive cancer were detected, suggesting that IPNB evolved to iCCA (arrows in Fig. 2C).

We concluded that the mice develop IPNB and iCCA which faithfully recapitulate the histology of their human counterparts.

Signaling pathways driving iCCA development in humans are activated in mice.

To investigate the signaling pathways often perturbed in human iCCA, we first evaluated mitogen activated protein kinase (MAPK) activity and looked at phosphorylated extracellular-regulated kinase (p-ERK).⁽³²⁾ p-ERK was detected in cholangiocytes following *Kras*^{G12D} induction or DDC treatment, and in iCCA, but not in untreated normal cholangiocytes (Fig. 3A). Second, phosphoinositide-3-kinase (PI3K) signaling⁽³²⁾ was activated in IPNB cells of DDC-treated mice expressing wild-type *Kras*, and in iCCA of DDC-treated mice expressing mutant *Kras*^{G12D}, as evidenced by staining with anti-phosphorylated AKT serine/threonine kinase (p-AKT) (Fig. 3A). Third, we also detected accumulation of hairy and enhancer of split-1 (HES1)⁽³³⁾ in nuclei of iCCA, indicating that the Notch pathway was stimulated (Fig. 3A). Phospho-smad2 against decapentaplegic 2 (SMAD2), a marker of Transforming Growth Factor beta (TGFβ) activity, was detected at high levels only in IPNB, not in other conditions. Finally, v-erb-b2 erythroblastic leukemia viral oncogene homolog 2/human epidermal growth factor receptor 2 (ERBB2/HER2)⁽³⁴⁾ was detected in all conditions. Microdissection of IPNBs and iCCAs from tissues of mice expressing mutant *Kras*^{G12D} and treated for 25 weeks with DDC, followed by western blotting indicated that the amount of ERBB2 protein was higher in iCCA than in IPNB, and that ERBB2 was active, as evidenced by detection of

phosphorylated ERBB2^{Y1248} (Fig. 3B). Also, RT-qPCR analysis of total RNA from normal purified cholangiocytes and from microdissected IPNB and iCCA showed higher expression of epidermal growth factor receptor (EGFR) in the lesions (Fig. 3B).

We concluded that MAPK, AKT, Notch, TGF β and ERBB signaling was activated during tumorigenesis in our mouse model, like in human iCCA.

IPNB and iCCA display similar gene expression profiles and overexpress TNS4

To gain insight into the mechanisms of tumor progression we investigated the gene expression profile of ductular proliferations, IPNBs and iCCAs in DDC-treated mice expressing mutant *Kras*^{G12D}. The three types of lesions co-existed in livers treated for 25 weeks with DDC and were isolated at that stage. Microdissected iCCAs displayed the histology of early stage of invasive cancer, as illustrated in Fig. 2B. Microdissection was performed by laser capture microdissection (LCM) followed by RNA sequencing. Principal component analysis (PCA) of the transcriptomic data revealed that ductular proliferations clustered separately from iCCAs and IPNBs which clustered more closely (Fig. 4A). Combined with the location of iCCAs near large ducts containing IPNB and the existence of transitional zones between IPNB and iCCA (Fig. 2C), the closely related gene expression pattern suggests that iCCA originate from IPNB. Nineteen genes were found to be differentially expressed between IPNB and iCCA (fold-change >2, adjusted p-value <0.05), out of which 10 were overexpressed (Fig. 4B). Except for *Aspa/Acy2*, these 10 genes were also overexpressed in human iCCA versus non-tumor tissue (The Cancer Genome Atlas cohort (TCGA); n=34; <http://firebrowse.org>) and/or Gene Expression Omnibus cohort GSE119336 (n=15),⁽³⁵⁾ supporting the relevance of the mouse data with human data (Fig. 4C).

The focal adhesion protein Tensin-4 (TNS4; also called C-terminal tensin-like protein CTEN) attracted our attention because it is upregulated by ERK signaling and promotes tumor progression in hepatocellular carcinoma.⁽³⁶⁾ RNA sequencing showed significant overexpression of *Tns4* in iCCA as compared to IPNB (Fig. 4B). RT-PCR showed higher mean values of *Tns4* in iCCA as compared to IPNB, and in *Kras*^{G12D}-expressing cholangiocytes as compared to wild-type, albeit with variability among samples. Western

blotting of microdissected IPNBs and iCCAs confirmed that TNS4 is significantly higher in iCCA compared to IPNB (Fig. 5A). Immunostainings confirmed the expression of TNS4 in mouse bile ducts, IPNB and iCCA (Fig. 5B). Also in humans, TNS4 was upregulated in iCCA as compared to controls (Fig. 5C).

TNS4 expression was also investigated in iCCA derived from hepatocytes. We overexpressed AKT and the intracellular domain of Notch1 (NICD) in hepatocytes of adult mice by hydrodynamic tail vein injection of plasmid DNA as described.⁽⁸⁾ These mice developed iCCAs 6 weeks after injection (Fig. 5D). Similarly, expression of AKT, NICD and KRAS^{G12V} also resulted in development of iCCA 3 weeks after injection. Immunostaining revealed strong expression of TNS4 in the iCCA cells (Fig. 5D), indicating that TNS4 is expressed both in cholangiocyte-derived (Opn-Cre^{ER}/LSL-Kras^{G12D} + DDC model) and in hepatocyte-derived iCCA (AKT+NICD and AKT+NICD+KRAS^{G12V} models).

TNS4 is stimulated by KRAS signaling and stimulates migration and proliferation

We next explored if KRAS activated expression of TNS4. Since EGI-1 cells constitutively express KRAS^{G12D}, we transiently transfected NMC cells with *Kras*^{G12D} to verify if it stimulates *Tns4* in cholangiocytes; this was the case (Fig. 6A). Since TNS4 is known to promote hepatocarcinoma cell migration,⁽³⁶⁾ we verified if this holds true in biliary cells. We generated EGI-1 and NMC clones that stably overexpress doxycycline-inducible TNS4 (Fig. 6B) and found that induction of TNS4 in both cell types accelerated cell migration in a wound healing assay (Fig. 6D). The assay lasted for 8h, *i.e.* below the duration of doubling time. More important, doxycycline-mediated increase of TNS4 stimulated proliferation of EGI-1 cells, as shown by crystal violet staining (Fig. 6E). We also used CRISPR-Cas9 to generate EGI-1 cells knockdown for TNS4 (Fig. 6C and Supporting Fig. S3). *In vitro* knockdown cells displayed reduced proliferation as evidenced by the longer time required to reach confluence (Fig. 6E).

To test whether TNS4 was sufficient to stimulate growth *in vivo*, we overexpressed TNS4 in EGI-1 cells and subcutaneously injected them into NOD-scid gamma (NSG) mice. The results showed that TNS4 accelerates tumor growth *in vivo* (Fig. 6F). Conversely,

subcutaneous injection of 10^6 EGI-1 or TNS4 knockdown EGI-1 cells resulted in strongly-reduced tumor development as compared to control EGI-1 cells (Fig. 6F).

SOX17 controls expression of TNS4

Our data above show that the transcription factor SOX17 is among the 10 genes that are overexpressed in iCCA as compared to IPNB in mice expressing *Kras*^{G12D} and treated for 25 weeks with DDC (Fig. 4B). RT-PCR data showed that the mean expression of *Sox17* was increased in iCCA as compared to IPNB, albeit with variability among samples. Western blotting of microdissected IPNBs and iCCAs confirmed the results and showed significant upregulation of SOX17 protein in iCCA as compared to IPNB (Fig. 5A). By immunostaining on liver sections, SOX17 was not detected in normal cholangiocytes or in *Kras*^{G12D}-expressing cholangiocytes. In IPNBs from DDC-treated animals expressing wild-type *Kras*, a low number of cells expressed SOX17 at levels slightly higher than background. In contrast, in DDC-treated animals expressing *Kras*^{G12D}, patches of epithelial cells in IPNB and iCCA showed significant levels of nuclear SOX17 (Fig. 5B). These immunostaining data were confirmed by Western blotting (Fig. 5A). SOX17 was also detected in endothelial CD31-positive cells, as expected.⁽³⁷⁾ Importantly, human livers displayed a similar profile with no detectable SOX17 in normal cholangiocytes, but with obvious expression in iCCA cells (Fig. 5C).

TNS4 was detected at the cell membrane in all tested conditions. SOX17-positive cells co-expressed TNS4 in iCCA (Fig. 5C). In addition, SOX17 and *TNS4* mRNA expression levels were correlated in human iCCA (Fig. 7A, Supporting Fig. S4). Moreover, since KRAS is known to stimulate SOX17 expression,⁽³⁸⁾ we tested the possibility that SOX17 activates TNS4. We generated EGI-1 clones that stably overexpress doxycycline-inducible SOX17 (Fig. 7B), and found that SOX17 stimulated TNS4 and *KRAS* expression (Fig. 7C). Stimulation of TNS4 is likely indirect since screening 28 kilobase pairs of the *TNS4* gene (-2500 to + 25500) for SOX17 binding sites using PROMO (http://alggen.lsi.upc.es/cgi-bin/promo_v3/promo/promoinit.cgi?dirDB=TF_8.3) or AliBaba2 (<http://gene-regulation.com/pub/programs/alibaba2/index.html>) revealed no SOX17 consensus binding site. We also transfected EGI-1 cells with a TNS4-luciferase construct comprising 2197 bp (-2148 to +49) of the human *TNS4* promoter subcloned

upstream of the firefly luciferase cDNA in pGL3-basic. This construct generated a 22-fold higher luciferase activity than pGL3-basic demonstrating *TNS4* promoter activity. However, it was not stimulated following co-transfection of a *SOX17* expression vector (fold change = 0.79 +/- 0.27 (SD); p = 0.38; n= 3). In addition, *SOX17* expression was activated by *Kras*^{G12D} as shown in NMC cells transiently transfected with a *Kras*^{G12D} expression vector (Fig. 6A). Finally, in EGI-1 cells overexpressing *TNS4* under control of doxycycline, *KRAS* was slightly induced (Fig. 7C). Together, the data suggest the existence of a *KRAS*-*SOX17*-*TNS4* cascade with positive feedbacks (Fig. 7C).

Discussion

Mouse models of cholangiocarcinoma often induce genetic modifications at prenatal stages, or simultaneously in multiple liver cell types, or in several organs.^(26, 27) We selected *Opn-Cre*^{ER} for its high efficiency and cholangiocyte specificity to induce expression of *Kras*^{G12D}. Indeed, earlier work showed that *Opn-Cre*^{ER} triggers recombination of a reporter in up to 99.9% of cholangiocytes, and in only 0.0074% hepatocytes; stellate cells and macrophages did not show *Opn-Cre*^{ER} activity.^(28, 29, 39) *Opn-Cre*^{ER} activity in other organs is illustrated in Supporting Fig. S5. Tumors in extrahepatic ducts, gallbladder and pancreas can show histological features comparable with iCCA. The absence of detectable tumors in these organs excluded that metastatic lesions were misidentified as iCCAs.

In our model, *Kras*^{G12D} is expressed from its endogenous locus. This was combined with chronic cholangitis to mimic tumor development in a context of chronic inflammation, a major risk factor for iCCA. Tumorigenesis was associated with activation of ERK, AKT, NOTCH and ERBB as in many human iCCAs. Also, mouse iCCAs displayed a tubular pattern with mucin-producing columnar cells, which is typical for human large duct type iCCA; intense desmoplasia was also detected. Therefore, our animal model displays many histological and biochemical features of human iCCA. No metastases were detected, yet their development may require prolonged DDC treatment, but the deteriorating health prompted us to euthanize the animals. Moreover, the transcriptomic

profile of IPNB and early stage of invasive iCCA, together with the histological identification of junctions between the two lesions strongly suggest that iCCA derive from IPNB. Interestingly, in humans, mutant *KRAS* is detected already in human low-grade IPNB, and tumor progression is associated with decreased TGF β signaling,⁽²²⁾ as in our model which shows strong nuclear staining of p-SMAD2 in IPNB, but no detectable p-SMAD2 in iCCA. Activation of the Wnt pathway has been reported in a subset of human IPNB,⁽⁴⁰⁾ but we did not detect nuclear accumulation of β -catenin in our mice.

Fickert and coworkers described the inflammation occurring in DDC-treated livers.⁽³¹⁾ We monitored an increase in interleukin-33 (IL-33) expression in IPNB and further in iCCA (Supporting Fig. S6A). IL-33 is a mitogen which induces iCCA when combined with constitutively active AKT and Yes-associated protein (YAP), and with cholestasis⁽⁴¹⁾. Fig. 3A shows that AKT signaling was active during tumorigenesis in IPNB and iCCA, but the expression of Hippo-YAP signaling markers did not correlate with tumor progression (Supporting Fig. S6B). In our model, IL-33 injection to DDC-fed mice expressing *Kras*^{G12D} contributed to growth of the early stage cancer lesions, but did not promote development toward a more advanced stage of iCCA (Supporting Fig. S6C).

Tensins are focal adhesion proteins involved in cell migration by bridging the extracellular matrix receptors and the actin skeleton. In hepatocellular cancer TNS4 is activated by ERK and enhances cell motility.⁽³⁶⁾ In our mouse model we detected TNS4 expression in normal cholangiocytes, and monitored an increase in its expression during tumorigenesis. Moreover, TNS4 and SOX17 are co-localized in IPNB/iCCA SOX17-positive cholangiocytes, and in transfection experiments we positioned TNS4 epistatically downstream of SOX17. TNS4 and SOX17 mRNA expression was positively correlated in human iCCA, as well as in breast carcinoma and prostate cancer but not in other tumor types (Supporting Fig. S4B).

During tumorigenesis we found that SOX17 becomes detectable at low level in IPNB of DDC-treated mice expressing wild-type *Kras*. In iCCA, SOX17 was heterogeneously distributed, with large epithelial cell patches displaying high expression of SOX17, while other tumor cells lacked SOX17. SOX17 was not detected in hepatocyte-derived iCCA

cells (Fig. 5D). SOX17 was also strongly expressed in endothelial cells, both in cholangiocyte- and hepatocyte-derived ICCAs (Fig. 5 B,D), meaning that care is needed to interpret transcriptomic and epigenomic analyses of SOX17 expression from sampled tumor tissues composed of multiple cell types. The higher level of SOX17 in microdissected iCCA as compared to IPNB likely reflects the higher number of cells expressing the gene in iCCA, and possibly the fact that SOX17-positive epithelial cells in iCCA express higher levels of SOX17 than in IPNB cells, as evidenced by more intense immunostaining in iCCA than in IPNB.

Further, SOX17 is not expressed at the protein level in normal human or mouse intrahepatic cholangiocytes, but regulates development of the extrahepatic biliary tract and is detected in adult gallbladder epithelium.⁽⁴²⁾ Earlier work claims that SOX17 is detected in normal intrahepatic cholangiocytes.^(43, 44) However, the immunostainings revealed only non-nuclear SOX17 in intrahepatic ducts, unlike in extrahepatic cholangiocytes where nuclear staining was obvious. In the absence of known nuclear-cytoplasmic shuttling, non-nuclear detection of SOX17 in intrahepatic ducts deserves re-evaluation. Interestingly, *Sox17* mRNA expression in wild-type or *Kras*^{G12D} mutant cholangiocytes does not correlate with SOX17 protein detection, indicating that this gene undergoes post-transcriptional regulation. SOX17 is regarded as a tumor suppressor in iCCA^(44, 45), and we confirmed that EGI-1 cells overexpressing SOX17 proliferate less well *in vivo* than control EGI-1 cells, as shown by Merino-Azpitarte *et al.* Importantly, since TNS4 is expressed in several cells devoid of SOX17, the latter is not strictly required for TNS4 expression. However, our data showing that SOX17 can control TNS4 expression indicate that a global tumor suppressor role of SOX17 does not exclude a contribution to tumor progression. In that context, we note that SOX17 was shown to be induced in precursor lesions of pancreatic ductal adenocarcinoma, and to cooperate with oncogenic KRAS^{G12D} to enhance transformation of normal pancreas.⁽³⁸⁾ Also, SOX17 is overexpressed in high grade malignant ERBB2-transformed cholangiocarcinoma cells as compared to less tumorigenic BDE-1 cells.⁽⁴⁶⁾ In conclusion, we have developed a novel mouse model that recapitulates formation of iCCA precursor lesions and their evolution toward malignancy, much like in humans. Carcinogenesis is associated with upregulation of TNS4 which promotes tumor progression and is regulated by KRAS and SOX17.

Acknowledgements

The authors thank Sandra Ormenese, Mieke Dewerchin, Javed Manesia, Satdarshan Paul Monga and Michael Oertel for advice with microdissection; Noémie Ammar-Khodja, Jean-Nicolas Lodewijckx and Mourad El Kaddouri for help.

Figure Legends

FIG. 1. Biliary lesions develop following diet-induced chronic cholangitis and cholangiocyte-specific expression of *Kras*^{G12D}. (A) Timing of *Kras*^{G12D} expression and of DDC-induced cholangitis. (B) Opn-Cre^{ER}/LSL-*Kras*^{G12D} mice expressing *Kras*^{G12D} following tamoxifen administration (*Kras*^{G12D}) do not show any lesion, whereas mice treated with DDC for 18 weeks in the absence (wild-type *Kras*+DDC) or presence of *KRAS*^{G12D} (*Kras*^{G12D}+DDC) develop ductular proliferations and IPNB with fibrous thickening of the bile duct wall. (C) iCCA are detected only in *Kras*^{G12D}+DDC mice and starting at 21 weeks of DDC treatment. Boxed region in left panel is magnified in right panel. bd, bile duct; H&E, hematoxylin and eosin; pv: portal vein; W, weeks. *, porphyrin plug.

FIG. 2. Histomorphology of biliary lesions in mice expressing *Kras*^{G12D} and fed a cholangitis-inducing diet. (A) In Opn-Cre^{ER}/LSL-*Kras*^{G12D}/Rosa26^{eYFP} mice fed a DDC diet for 21 weeks (*Kras*^{G12D}+DDC), cells of IPNB and iCCA proliferate, as shown by Ki67 staining; they also express CK19 and EpCAM; eYFP expression demonstrates their cholangiocyte origin. IPNB and iCCA express SOX9, but not HNF4. In iCCA, a subset of cells express detectable HNF4. (B) iCCA in *Kras*^{G12D}+DDC mice shows glandular structures displaying columnar to cuboidal morphology and fibroinflammatory stroma. iCCA cells express mucins, as evidenced by alcian blue staining. (C) A transition zone between IPNB and early stage of invasive iCCA (black arrows) is detected in *Kras*^{G12D}+DDC mice. Inset in panel C (right) points to nuclear atypia. Boxed regions in left panels are magnified in right panels. bd, bile duct; H&E, hematoxylin and eosin; pv: portal vein.

FIG. 3. Signaling pathways activated during tumor progression. (A) p-ERK staining is detected in cholangiocytes following *Kras*^{G12D} expression, in DDC-induced IPNB and *Kras*^{G12D}+DDC-induced iCCA. p-AKT is detected in IPNB and iCCA, while HES1 is detected only in iCCA. p-SMAD2 is detected only in IPNB. ERBB2 is detected in biliary cells in all conditions. (B) Levels of ERBB2 are higher in iCCA than in IPNB, and ERBB2 protein is active as evidenced by detection of p-ERBB2^{Y1248}. Expression of *Egfr* and *ErbB2* mRNA was higher in IPNB and iCCA than in normal purified cholangiocytes. Dot plots show means +/- SEM. *, p<0.05.

FIG. 4. Differentially expressed genes between IPNB and iCCA. (A) Principal component analysis of RNA sequencing data of ductular proliferations, IPNB and iCCA. The lesions were microdissected from tamoxifen-treated Opn-Cre^{ER}/LSL-Kras^{G12D}/Rosa26R^{eYFP} mice fed a DDC diet for 25 weeks. (B) Table of the 19 genes differentially expressed between IPNB and iCCA. (C) Genes that are overexpressed in murine iCCA *versus* IPNB are also overexpressed in murine iCCA *versus* control cholangiocytes. Control cholangiocytes are FACS-purified from tamoxifen-treated Opn-Cre^{ER}/Rosa26R^{eYFP} mice. Most of these genes are overexpressed in human iCCA *versus* control tissue. Two human iCCA cohorts are compared with mouse iCCA.

FIG. 5. TNS4 and SOX17 expression in mouse and human iCCA. (A) RT-PCR shows higher mean values of *Tns4* and *Sox17* in iCCA as compared to IPNB, and in *Kras*^{G12D}-expressing cholangiocytes as compared to wild-type, albeit with variability among samples. Western blotting of microdissected IPNBs and iCCAs show significant upregulation of TNS4 and SOX17 proteins in iCCA as compared to IPNB. IPNB and iCCA were microdissected from mice expressing *Kras*^{G12D} and fed a DDC diet for 25 weeks. Cholangiocytes from mice fed a normal diet were FACS-sorted from tamoxifen-treated Opn-Cre^{ER}/Rosa26R^{eYFP} or Opn-Cre^{ER}/LSL-Kras^{G12D}/Rosa26R^{eYFP} mice, as indicated. Data in dot plots are means +/- SEM. **, p<0.01. (B) Expression of TNS4 is detected in biliary cells from normal mouse ducts, IPNB and iCCA. SOX17 is detected in DDC-induced IPNB and iCCA, not in cholangiocytes from untreated mice. (C) TNS4 and SOX17 are detected in human iCCA. Dotted lines delineate bile ducts; white arrowheads,

SOX17-expressing biliary cells; yellow arrowheads, SOX17-expressing CD31-positive endothelial cells. (D) TNS4 is detected in iCCA cells which developed in mice overexpressing AKT and NICD, or AKT, NICD, and Kras^{G12V} in hepatocytes. SOX17 expression was restricted to endothelial cells. bd, bile duct; ns, non-specific labeling; pv, portal vein. Insets represent high magnifications.

FIG. 6. TNS4 is stimulated by KRAS^{G12D} and promotes cell migration and tumor growth. (A) Transient transfection of *Kras*^{G12D} stimulates *Tns4* and *Sox17* expression in NMC cells. (B) Overexpression of TNS4 in EGI-1 cells stably expressing doxycycline-inducible TNS4 (TNS4-EGI-1), as compared to doxycycline-treated EGI-1 stably transfected with pSBtet-GN vector (EGI-1). Data in (B) and (C) represent fold inductions relative to control +/- SEM. *, p<0.05. (C) Reduced expression of TNS4 in EGI-1 cells after CRISPR-Cas9-mediated targeting of the *TNS4* gene. Details are in Supporting Fig. S3. (D) Doxycycline-induced TNS4 accelerates migration of EGI-1 and NMC cells in a wound-healing assay. At each timepoint, the width of the wound is indicated as mean +/- SEM (μm) of 3 widths measured on 6 pictures covering the entire wound. (E) Doxycycline-induced TNS4 stimulates proliferation of EGI-1 cells; 16 x 10⁴ cells were plated at day 1 (upper panels). Cells illustrated were counted after crystal violet staining (lower graphs). Data in graphs are means +/- SEM. (F) Doxycycline-induced TNS4 stimulates tumor growth *in vivo* after subcutaneous injection of 10⁶ EGI-1 or TNS4-EGI-1 cells in NOD-scid gamma (NSG) mice. In contrast, TNS4kd-EGI-1 cells display reduced tumorigenesis as compared to control EGI-1 cells after subcutaneous injection. Tumor growth was measured with a caliper and tumors were collected 5-6 weeks after injection (lower graphs). Data in graphs are means +/- SEM; * p<0.05 and ***, p<0.001.

FIG. 7. A KRAS-stimulated cascade involving TNS4 and SOX17 is active during iCCA tumorigenesis. (A) Expression of *SOX17* and *TNS4* mRNA is correlated in human iCCA. Patient cohorts are from TCGA and Sia *et al.*⁽²⁵⁾ (B) Overexpression of SOX17 in EGI-1 cells stably expressing doxycycline-inducible SOX17 (SOX17-EGI-1), as compared to doxycycline-treated EGI-1 stably transfected with pSBtet-GN vector (EGI-1). (C) Left panel, SOX17 stimulates *KRAS* and *TNS4* in SOX17-EGI-1 cells. Right panel, TNS4

stimulates *KRAS* in TNS4-EGI-1. The structure of the *KRAS*-TNS4-SOX17 cascade is shown.

References

Author names in bold designate shared co-first authorship.

1. Banales JM, Cardinale V, Carpino G, Marzioni M, Andersen JB, Invernizzi P, et al. Expert consensus document: Cholangiocarcinoma: current knowledge and future perspectives consensus statement from the European Network for the Study of Cholangiocarcinoma (ENS-CCA). *Nat Rev Gastroenterol Hepatol* 2016;13:261-280.
2. Sirica AE, Gores GJ, Groopman JD, Selaru FM, Strazzabosco M, Wei Wang X, Zhu AX. Intrahepatic Cholangiocarcinoma: Continuing Challenges and Translational Advances. *Hepatology* 2019;69:1803-1815.
3. Fouassier L, Marzioni M, Afonso MB, Dooley S, Gaston K, Giannelli G, et al. Signalling networks in cholangiocarcinoma: Molecular pathogenesis, targeted therapies and drug resistance. *Liver Int* 2019;39 Suppl 1:43-62.
4. **Hill MA, Alexander WB**, Guo B, Kato Y, Patra K, O'Dell MR, et al. Kras and Tp53 Mutations Cause Cholangiocyte- and Hepatocyte-Derived Cholangiocarcinoma. *Cancer Res* 2018;78:4445-4451.
5. **Nakagawa H, Suzuki N**, Hirata Y, Hikiba Y, Hayakawa Y, Kinoshita H, et al. Biliary epithelial injury-induced regenerative response by IL-33 promotes cholangiocarcinogenesis from peribiliary glands. *Proc Natl Acad Sci U S A* 2017;114:E3806-E3815.
6. Guest RV, Boulter L, Kendall TJ, Minnis-Lyons SE, Walker R, Wigmore SJ, et al. Cell Lineage Tracing Reveals a Biliary Origin of Intrahepatic Cholangiocarcinoma. *Cancer Res* 2014;74:1005-1010.
7. Sekiya S, Suzuki A. Intrahepatic cholangiocarcinoma can arise from Notch-mediated conversion of hepatocytes. *J Clin Invest* 2012;122:3914-3918.
8. **Fan B, Malato Y, Calvisi DF**, Naqvi S, Razumilava N, Ribback S, et al. Cholangiocarcinomas can originate from hepatocytes in mice. *J Clin Invest* 2012;122:2911-2915.

9. **Saha SK, Parachoniak CA**, Ghanta KS, Fitamant J, Ross KN, Najem MS, et al. Mutant IDH inhibits HNF-4alpha to block hepatocyte differentiation and promote biliary cancer. *Nature* 2014;513:110-114.
10. Komuta M, Govaere O, Vandecaveye V, Akiba J, Van Steenberghe W, Verslype C, et al. Histological diversity in cholangiocellular carcinoma reflects the different cholangiocyte phenotypes. *Hepatology* 2012;55:1876-1888.
11. Nakanuma Y, Klimstra DS, Komuta M, Zen Y: Intrahepatic cholangiocarcinoma. In: WHO classification of tumours - 5th edition. Digestive system tumours: WHO, 2019; 254-259.
12. Basturk O, Aishima S, Esposito I: Biliary intraepithelial neoplasia. In: WHO classification of tumours - 5th edition. Digestive system tumours: WHO, 2019; 273-275.
13. Nakanuma Y, Basturk O, Esposito I, Klimstra D, Komuta M, Zen Y: Intraductal papillary neoplasms of the bile ducts. In: WHO classification of tumours - 5th edition. Digestive system tumours: WHO, 2019; 279-282.
14. **Kendall T, Verheij J**, Gaudio E, Evert M, Guido M, Goeppert B, Carpino G. Anatomical, histomorphological and molecular classification of cholangiocarcinoma. *Liver Int* 2019;39 Suppl 1:7-18.
15. Aishima S, Kubo Y, Tanaka Y, Oda Y. Histological features of precancerous and early cancerous lesions of biliary tract carcinoma. *J Hepatobiliary Pancreat Sci* 2014;21:448-452.
16. Ohtsuka M, Shimizu H, Kato A, Yoshitomi H, Furukawa K, Tsuyuguchi T, et al. Intraductal papillary neoplasms of the bile duct. *Int J Hepatol* 2014;2014:459091.
17. **Ong CK, Subimerb C**, Pairajkul C, Wongkham S, Cutcutache I, Yu W, et al. Exome sequencing of liver fluke-associated cholangiocarcinoma. *Nat Genet* 2012;44:690-693.
18. **Zou S, Li J, Zhou H, Frech C**, Jiang X, Chu JS, et al. Mutational landscape of intrahepatic cholangiocarcinoma. *Nat Commun* 2014;5:5696.
19. **Nakamura H, Arai Y, Totoki Y, Shiota T, Elzawahry A**, Kato M, et al. Genomic spectra of biliary tract cancer. *Nat Genet* 2015;47:1003-1010.

20. **Hsu M, Sasaki M**, Igarashi S, Sato Y, Nakanuma Y. KRAS and GNAS mutations and p53 overexpression in biliary intraepithelial neoplasia and intrahepatic cholangiocarcinomas. *Cancer* 2013;119:1669-1674.
21. Sasaki M, Matsubara T, Nitta T, Sato Y, Nakanuma Y. GNAS and KRAS Mutations are Common in Intraductal Papillary Neoplasms of the Bile Duct. *PLoS ONE* 2014;8:e81706.
22. Schlitter AM, Born D, Bettstetter M, Specht K, Kim-Fuchs C, Riener MO, et al. Intraductal papillary neoplasms of the bile duct: stepwise progression to carcinoma involves common molecular pathways. *Mod Pathol* 2014;27:73-86.
23. Khan SA, Tavolari S, Brandi G. Cholangiocarcinoma: Epidemiology and risk factors. *Liver Int* 2019;39 Suppl 1:19-31.
24. Andersen JB, Spee B, Blechacz BR, Avital I, Komuta M, Barbour A, et al. Genomic and genetic characterization of cholangiocarcinoma identifies therapeutic targets for tyrosine kinase inhibitors. *Gastroenterology* 2012;142:1021-1031.
25. Sia D, Hoshida Y, Villanueva A, Roayaie S, Ferrer J, Tabak B, et al. Integrative molecular analysis of intrahepatic cholangiocarcinoma reveals 2 classes that have different outcomes. *Gastroenterology* 2013;144:829-840.
26. Erice O, Vallejo A, Ponz-Sarvisé M, Saborowski M, Vogel A, Calvisi DF, et al. Genetic Mouse Models as In Vivo Tools for Cholangiocarcinoma Research. *Cancers* 2019;11:1868.
27. Pirenne S, Lemaigre F. Genetically-engineered animal models of biliary tract cancers. *Current Opinion in Gastroenterology* 2020;36:90-98.
28. **Espanol-Suner R, Carpentier R**, Van Hul N, Legry V, Achouri Y, Cordi S, et al. Liver progenitor cells yield functional hepatocytes in response to chronic liver injury in mice. *Gastroenterology* 2012;143:1564-1575.
29. **Lesaffer B, Verboven E**, Van Huffel L, Moya IM, van Grunsven LA, Leclercq IA, et al. Comparison of the Opn-CreER and Ck19-CreER Drivers in Bile Ducts of Normal and Injured Mouse Livers. *Cells* 2019;8:380.
30. Hingorani SR, Petricoin EF, Maitra A, Rajapakse V, King C, Jacobetz MA, et al. Preinvasive and invasive ductal pancreatic cancer and its early detection in the mouse. *Cancer Cell* 2003;4:437-450.

31. **Fickert P, Stoger U**, Fuchsbichler A, Moustafa T, Marschall HU, Weiglein AH, et al. A new xenobiotic-induced mouse model of sclerosing cholangitis and biliary fibrosis. *Am J Pathol* 2007;171:525-536.
32. Schmitz KJ, Lang H, Wohlschlaeger J, Sotiropoulos GC, Reis H, Schmid KW, Baba HA. AKT and ERK1/2 signaling in intrahepatic cholangiocarcinoma. *World J Gastroenterol* 2007;13:6470-6477.
33. **El Khatib M, Bozko P**, Palagani V, Malek NP, Wilkens L, Plentz RR. Activation of Notch signaling is required for cholangiocarcinoma progression and is enhanced by inactivation of p53 in vivo. *PLoS One* 2013;8:e77433.
34. Sirica AE. Role of ErbB family receptor tyrosine kinases in intrahepatic cholangiocarcinoma. *World J Gastroenterol* 2008;14:7033-7058.
35. Zhou G, Cao P, Li Y. RNA over-editing leads to aggressiveness of intrahepatic cholangiocarcinoma. *Gene Expression Omnibus*, accession number GSE119336 2019.
36. Chan LK, Chiu YT, Sze KM, Ng IO. Tensin4 is up-regulated by EGF-induced ERK1/2 activity and promotes cell proliferation and migration in hepatocellular carcinoma. *Oncotarget* 2015;6:20964-20976.
37. **Corada M, Orsenigo F**, Morini MF, Pitulescu ME, Bhat G, Nyqvist D, et al. Sox17 is indispensable for acquisition and maintenance of arterial identity. *Nat Commun* 2013;4:2609.
38. Delgiorno KE, Hall JC, Takeuchi KK, Pan FC, Halbrook CJ, Washington MK, et al. Identification and manipulation of biliary metaplasia in pancreatic tumors. *Gastroenterology* 2014;146:233-244
39. Mu X, Espanol-Suner R, Mederacke I, Affo S, Manco R, Sempoux C, et al. Hepatocellular carcinoma originates from hepatocytes and not from the progenitor/biliary compartment. *J Clin Invest* 2015;125:3891-3903.
40. Fujikura K, Akita M, Ajiki T, Fukumoto T, Itoh T, Zen Y. Recurrent Mutations in APC and CTNNB1 and Activated Wnt/beta-catenin Signaling in Intraductal Papillary Neoplasms of the Bile Duct: A Whole Exome Sequencing Study. *Am J Surg Pathol* 2018;42:1674-1685.

41. Li J, Razumilava N, Gores GJ, Walters S, Mizuochi T, Mourya R, et al. Biliary repair and carcinogenesis are mediated by IL-33-dependent cholangiocyte proliferation. *J Clin Invest* 2014;124:3241-3251.
42. **Uemura M, Ozawa A**, Nagata T, Kurasawa K, Tsunekawa N, Nobuhisa I, et al. Sox17 haploinsufficiency results in perinatal biliary atresia and hepatitis in C57BL/6 background mice. *Development* 2013;140:639-648.
43. **Carpino G, Cardinale V**, Onori P, Franchitto A, Berloco PB, Rossi M, et al. Biliary tree stem/progenitor cells in glands of extrahepatic and intrahepatic bile ducts: an anatomical in situ study yielding evidence of maturational lineages. *J Anat* 2012;220:186-199.
44. Merino-Azpitarte M, Lozano E, Perugorria MJ, Esparza-Baquer A, Erice O, Santos-Laso A, et al. SOX17 regulates cholangiocyte differentiation and acts as a tumor suppressor in cholangiocarcinoma. *J Hepatol* 2017;67:72-83.
45. Goeppert B, Konermann C, Schmidt CR, Bogatyrova O, Geiselhart L, Ernst C, et al. Global alterations of DNA methylation in cholangiocarcinoma target the Wnt signaling pathway. *Hepatology* 2014;59:544-554.
46. Dumur CI, Campbell DJ, DeWitt JL, Oyesanya RA, Sirica AE. Differential gene expression profiling of cultured neu-transformed versus spontaneously-transformed rat cholangiocytes and of corresponding cholangiocarcinomas. *Exp Mol Pathol* 2010;89:227-235.

Figure 5