

MONOMIAL GENERATORS OF COMPLETE PLANAR IDEALS

MARIA ALBERICH-CARRAMIÑANA, JOSEP ÀLVAREZ MONTANER, AND GUILLEM BLANCO

ABSTRACT. We provide an algorithm that computes a set of generators for any complete ideal in a smooth complex surface. More interestingly, these generators admit a presentation as monomials in a set of maximal contact elements associated to the minimal log-resolution of the ideal. Furthermore, the monomial expression given by our method is an equisingularity invariant of the ideal. As an outcome, we provide a geometric method to compute the integral closure of a planar ideal and we apply our algorithm to some families of complete ideals.

1. INTRODUCTION

Let (X, O) be a germ of smooth complex surface and $\mathcal{O}_{X,O}$ the ring of germs of holomorphic functions in a neighbourhood of O , which we identify with $\mathbb{C}\{x, y\}$ by taking local coordinates, and let \mathfrak{m} be the maximal ideal at O . Let $\pi : X' \rightarrow X$ be a proper birational morphism that can be achieved as a sequence of blow-ups along a set of points. Given an effective \mathbb{Z} -divisor D in X' we may consider its associated (sheaf) ideal $\pi_*\mathcal{O}_{X'}(-D)$ whose stalk at O we simply denote as H_D . This type of ideals were systematically studied by Zariski in [32]. They are complete ideals of $\mathcal{O}_{X,O}$ and \mathfrak{m} -primary whenever D has exceptional support. Among the class of divisors defining the same complete ideal we may find a unique maximal representative which happens to have the property of being *antinef*. Actually, Zariski [32] showed that that above correspondence is in fact an isomorphism of semigroups between the set of complete \mathfrak{m} -primary ideals and the set of antinef divisors with exceptional support.

The aim of this work is to make computationally explicit this correspondence. Namely, given a proper birational morphism $\pi : X' \rightarrow X$ and an antinef divisor D in X' we provide an algorithm that gives us a system of generators of the ideal H_D . Moreover, our method captures the topological type of D and gives an output that discerns between the equisingularity class of H_D and its analytic type: it produces a set of monomial expressions, which are invariant of the equisingularity class of H_D (or equivalently invariant of the topological type of D); and when the set of variables in these monomial expressions are substituted by a set of maximal contact elements of the resolution π , they give a system of generators of H_D . Our approach uses two main ingredients: the Zariski decomposition of complete ideals into simple ones, and subtle properties in the theory of adjacent ideals, which relay on results obtained in the study of sandwiched singularities and the Nash conjecture of arcs on these singularities.

2010 *Mathematics Subject Classification*. Primary 13B22, 14E15. Secondary 32S15.

Key words and phrases. Complete ideals, log-resolution.

All three authors are supported by Spanish Ministerio de Economía y Competitividad MTM2015-69135-P and Generalitat de Catalunya 2017 SGR-932 project. They are with the Barcelona Graduate School of Mathematics (BGSMath). MAC is also with the Institut de Robòtica i Informàtica Industrial (CSIC-UPC).

Our main result is an algorithm (see Algorithm 3.5) which we briefly describe. We start by fixing a set $\{f_0, \dots, f_g\}$ of *maximal contact elements* associated to π . These are irreducible curves that are geometrically adapted to the proper birational morphism in the sense that they are parameterized by the dead-ends of the dual graph of the divisor D (see Subsection 2.4 for details). Once we have this fixed set we consider the decomposition $D = \rho_1 D_1 + \dots + \rho_r D_r$ of the divisor D corresponding to the Zariski decomposition $H_D = H_{D_1}^{\rho_1} \cdots H_{D_r}^{\rho_r}$, where the ideals H_{D_i} are simple and complete (see [32], [9] for details). For each ideal H_{D_i} we compute in a very specific unique way an *adjacent ideal below* H_{D_i} , i.e. a codimension one ideal $H_{\widehat{D}_i} \subsetneq H_{D_i}$, and we prove in Proposition 4.4 that we may pick a maximal contact element (or a power of it) belonging to H_{D_i} but not belonging to $H_{\widehat{D}_i}$. These adjacent ideals are no longer simple so we may repeat the same procedure. This process finishes after finitely many steps since these adjacent ideals have *smaller support* (see Lemma 4.1). We visualize the steps of our algorithm as a tree of ideals with leaves corresponding to the maximal ideal. From the elements of maximal contact that we pick at each step we construct a system of generators of H_D .

As an outcome of our method we obtain a set of generators of a planar complete ideal given by monomials in this fixed set of maximal contact elements (see Theorem 4.5). That is, we obtain an expression of the form $H_D = (\mathbf{f}^{\alpha_1}, \dots, \mathbf{f}^{\alpha_r})$, where $\mathbf{f}^{\alpha} = f_0^{a_0} \cdots f_g^{a_g}$ for $\alpha = (a_0, \dots, a_g) \in \mathbb{Z}_{\geq 0}^{g+1}$. More interestingly, if we fix another set of maximal contact elements $\{f'_0, \dots, f'_g\}$, the algorithm returns the same monomial expression (in this new set) for the generators of H_D since it only uses the information of the equisingularity types of the maximal contact elements. Actually, the monomial expression that we have for H_D also works for any other ideal $H_{D'} \subseteq \mathcal{O}_{X,O}$ whose associated divisor D' has the same weighted dual graph as D . Hence the monomial expression given by our method is an equisingularity invariant of the ideal.

As a first application of our algorithm we provide a method to compute the integral closure of any ideal $\mathfrak{a} \subseteq \mathcal{O}_{X,O}$. More precisely, given a log-resolution $\pi : X' \rightarrow X$ of the ideal \mathfrak{a} , let D be the effective Cartier divisor such that $\mathfrak{a} \cdot \mathcal{O}_{X'} = \mathcal{O}_{X'}(-D)$. Then, the integral closure $\bar{\mathfrak{a}}$ is the ideal H_D . We should point out that, given a set of generators of the ideal \mathfrak{a} we may compute the divisor D using the algorithm that we developed in [2].

There are some general algorithms to compute the integral closure as those proposed by Vasconcelos [30, 31], de Jong [11], Leonard and Pelikaan [21] or Singh and Swanson [27] which have been implemented in computer algebra systems such as Macaulay2 [17] or Singular [10]. When comparing these algorithms with our method in the case of planar ideals we notice that the system of generators that we produce is not minimal in general. However, our method reflects more closely some geometric properties. For instance, if two different ideals belong to the same *equisingularity class*, meaning that they have the similar weighted cluster of base points [9, 8.3] or equivalently their minimal log-resolution have the same weighted dual graph, their integral closure have the same monomial expression in the respective set of maximal contact elements.

The usefulness of our method becomes still more apparent when dealing with families of complete ideals dominated by the same log-resolution. This is the case of *multiplier ideals* which is an invariant of singularities that has received a lot of attention in recent years. Combining the algorithms given in [2] and [5] with Algorithm 3.5 we develop a method that, given a set of generators of a planar ideal, returns a set of generators of

the corresponding multiplier ideals. Since multiplier ideals are invariant up to integral closure, we obtain a result that resembles a formula given by Howald [18] in the sense that the multiplier ideals of a monomial ideal (in the set of maximal contact elements) is monomial as well.

Another interesting family of complete ideals was considered by Teissier in [28]. These ideals are described by valuative conditions given by the intersection multiplicity of the elements of $\mathcal{O}_{X,O}$ with a fixed germ of plane curve. With the help of Algorithm 3.5 we may provide an explicit system of generators for these ideals.

Finally, we would like to mention that Casas-Alvero [8] proposes a geometrical procedure to obtain a minimal set of generators of the complete ideal H_D based on a construction of a filtration by complete ideals, namely a chain of consecutive adjacent ideals between H_D and $\mathfrak{m}H_D$. The main drawback in order to computationally implement this method is that one does not know a priori which log-resolution of H_D will dominate all the log-resolutions of each complete ideal appearing in the filtration, since it depends on the choices of the adjacent ideal at each step. In contrast, our procedure fixes a proper birational morphism π from the very beginning and we always keep working with complete ideals whose log-resolution is dominated by π . The further points which must be blown up after π in the Casas-Alvero's method can not be chosen following an intrinsic unique way which works for any ideal whose log-resolution is dominated by π . Moreover these choices in Casas-Alvero's method determine the generators of H_D : the generators must go through certain infinitely near points and have to skip some others. Both sets of points are among those blown up by π and by the further blowing-ups. Therefore an hypothetical computational implementation of Casas-Alvero's method would suffer from a lack of robustness and invariance, which are the strengths of our method. For discounted, the fact that the monomial expression given by our method is an equisingularity invariant of the ideal is not achieved by any other previous method.

The algorithms developed in this paper have been implemented in the computer algebra system Magma [7] and are available at

github.com/gblanco92/IntegralClosureDim2

2. PRELIMINARIES

The aim of this section is to introduce all the background that we will use throughout this work.

2.1. Proper birational morphisms and infinitely near points. Let X be a smooth complex surface and $\mathcal{O}_{X,O} \cong \mathbb{C}\{x,y\}$ the ring of germs of holomorphic functions in a neighborhood of a smooth point $O \in X$ and $\mathfrak{m} = \mathfrak{m}_{X,O} \subseteq \mathcal{O}_{X,O}$ the maximal ideal at O . Given a proper ideal $\mathfrak{a} \subseteq \mathcal{O}_{X,O}$ we have a decomposition $\mathfrak{a} = (a) \cdot \mathfrak{a}'$, where $a \in \mathcal{O}_{X,O}$ is the greatest common divisor of the elements of \mathfrak{a} and \mathfrak{a}' is \mathfrak{m} -primary.

Let $\pi : X' \rightarrow X$ a proper birational morphism, with X' a smooth surface. Any such proper birational morphism can be achieved as a sequence of blow-ups

$$\pi : X' = X_{r+1} \rightarrow X_r \rightarrow \cdots \rightarrow X_1 = X$$

with $X_{i+1} = \text{Bl}_{p_i} X_i$ for a point $p_i \in X_i$. Denote by $E = \text{Exc}(\pi)$ the exceptional locus. Since we are in a local framework, we shall assume throughout this work that E

is connected. The set K of points which have been blown up gives an index set for the exceptional components $\{E_p\}_{p \in K}$ in E . We say that these points are a cluster of *infinitely near to the origin* and we may establish a *proximity relation* between them. Namely, we say that a point $q \in K$ is *proximate* to $p \in K$ if and only if q belongs to the exceptional divisor E_p corresponding to p as proper or infinitely near point. We will denote this relation as $q \rightarrow p$ and we collect all these relations by means of the *proximity matrix* $P = (P_{p,q})$ defined as:

$$P_{p,q} := \begin{cases} 1 & \text{if } p = q, \\ -1 & \text{if } q \rightarrow p, \\ 0 & \text{otherwise.} \end{cases}$$

The proximity matrix is related to the negative-definite *intersection matrix* $N = (E_p \cdot E_q)$ by the formula $N = -P^\top P$. An infinitely near point q is proximate to just one or two points. In the former case we say that q is a *free point*, and in the later it is a *satellite point*. Besides, one can establish a *partial ordering* in K . Namely, $q \leq p$ if and only if p is infinitely near to q .

A *log-resolution* of a proper ideal \mathfrak{a} is a proper birational morphism $\pi : X' \rightarrow X$ where X' is smooth such that there is an effective Cartier divisor F satisfying $\mathfrak{a} \cdot \mathcal{O}_{X'} = \mathcal{O}_{X'}(-F)$, and the divisor $F + \text{Exc}(\pi)$ has simple normal crossings. The *weighted dual graph* of π is the dual graph of the divisor $F = \sum_{1 \leq i \leq s} v_i(F) E_i$ weighted by the values $\{v_i(F)\}_{1 \leq i \leq s}$, where $E_i = E_{p_i}$ with $1 \leq i \leq r$ are the exceptional components, and E_i with $r < i \leq s$ are the non-exceptional irreducible components of F .

Definition 2.1. Two proper ideals \mathfrak{a} and \mathfrak{a}' are *equisingular*, or belong to the same *equisingularity class* if one of the following equivalent conditions hold:

- They have respectively minimal log-resolutions with equal weighted dual graph.
- The divisors F and F' associated to their respective minimal log-resolutions are homeomorphic (topologically equivalent).
- The weighted cluster of base points of \mathfrak{a} and \mathfrak{a}' are similar in the sense of [9, 8.3] in case \mathfrak{a} and \mathfrak{a}' are \mathfrak{m} -primary.

Notice that when dealing with principal ideals $\mathfrak{a} = (f)$ and $\mathfrak{a}' = (g)$ the notion of equisingularity of the ideals is equivalent to the classical notion of equisingularity (or topological equivalence) of the germs $f = 0$ and $g = 0$. Equisingular ideals have equisingular generic members. Moreover, equisingular \mathfrak{m} -primary complete ideals have besides equal codimension.

2.2. Divisor basis. Let $\Lambda_\pi := \bigoplus_{p \in K} \mathbb{Z} E_p$ be the lattice of integral divisors in X' with exceptional support. We have two different basis of this \mathbb{Z} -module given by the *strict transforms* and the *total transforms* of the exceptional components. For simplicity we will also denote the strict transforms by E_p and the total transforms by \bar{E}_p . In particular, any divisor $D \in \Lambda_\pi$ can be presented in two different ways

$$D = \sum_{p \in K} v_p(D) E_p = \sum_{p \in K} e_p(D) \bar{E}_p,$$

where the weights $v_p(D)$ (resp. $e_p(D)$) are the *values* (resp. *multiplicities*) of D . The relation between values and multiplicities is given by the proximity relations

$$(2.1) \quad v_q(D) = e_q(D) + \sum_{q \rightarrow p} v_p(D).$$

that provide a base change formula $\mathbf{e}^\top = P \cdot \mathbf{v}^\top$, where we collect the multiplicities and values in the vectors $\mathbf{e} = (e_p(D))_{p \in K}$ and $\mathbf{v} = (v_p(D))_{p \in K}$, respectively.

Aside from the total and strict transform basis $\{\overline{E}_p\}_{p \in K}$ and $\{E_p\}_{p \in K}$ of the lattice Λ_π of exceptional divisors, we may also consider the *branch basis* $\{B_p\}_{p \in K}$ defined as the dual of $\{-E_p\}_{p \in K}$ with respect to the intersection form. Any divisor $D \in \Lambda_\pi$ has a presentation

$$(2.2) \quad D = \sum_{p \in K} \rho_p(D) B_p,$$

where $\rho_p(D) = -D \cdot E_p$ is the *excess* at p and the relation between excesses and multiplicities is given by $\boldsymbol{\rho}^\top = P^\top \mathbf{e}^\top$, where $\boldsymbol{\rho} = (\rho_p(D))_{p \in K}$ denote the vectors of excesses.

The *support* $\text{Supp}(D)$ of a divisor $D \in \Lambda_\pi$ is the union of irreducible divisors of D , i.e. if $D = \sum_\alpha v_\alpha(D) E_\alpha$ is the expression of D in the strict transform basis then $\text{Supp}(D) = \{E_\alpha \mid v_\alpha(D) \neq 0\}$. We will also consider the same construction in the total transform basis. Namely, let $D = \sum_\alpha e_\alpha(D) \overline{E}_\alpha$ be the expression of D in the total transform basis, then we define its *support in the total transform basis* as $\text{Supp}_{\overline{E}}(D) = \{\overline{E}_\alpha \mid e_\alpha(D) \neq 0\}$. To avoid confusion with the usual notion of support, we will always be explicit when referring to the support in the total transform basis.

Remark 2.2. Let $\xi : f = 0$ be the germ of curve defined by an element $f \in \mathcal{O}_{X,O}$ satisfying that π dominates a resolution of the principal ideal (f) . The *total transform* of ξ is the pull-back $\overline{\xi} := \pi^* f$ and its *strict transform* ξ' is the closure of $\pi^{-1}(\xi - \{O\})$. Then we have a presentation

$$\overline{\xi} = \xi' + \sum_{p \in K} v_p(f) E_p = \xi' + \sum_{p \in K} e_p(f) \overline{E}_p = \xi' + \sum_{p \in K} \rho_p(f) B_p,$$

where $v_p(f) := v_p(D)$, $e_p(f) := e_p(D)$ and $\rho_p(f) := \rho_p(D)$ for $D = \text{Div}(\pi^* f)_{\text{exc}}$.

Throughout this work we will be interested in germs of curves associated to branch basis divisors.

Construction 2.3. Given a point $p \in K$, let $f_p \in \mathcal{O}_{X,O}$ be an irreducible element such that its strict transform by the resolution π intersects transversely E_p at a smooth point of E . We have that $B_p = \text{Div}(\pi^* f_p)_{\text{exc}}$. Conversely, any $g \in \mathcal{O}_{X,O}$ with $B_p = \text{Div}(\pi^* g)_{\text{exc}}$ is irreducible and its strict transform by the resolution π intersects transversely E_p at a smooth point of E .

Since the elements $f_p \in \mathcal{O}_{X,O}$ are irreducible, the points $q \in K$ such that $e_q(B_p) \neq 0$ are totally ordered. Furthermore, the resolution of any $f_p \in \mathcal{O}_{X,O}$, $p \in K$ is dominated by π . Moreover, any f whose resolution is dominated by π can be written as a product $f = \prod_p f_p^{\rho_p}$ of suitable elements f_p obtained as in Construction 2.3.

Remark 2.4. Let $\xi : f = 0$, $\eta : g = 0$ be germs of curves such that π dominates their resolutions and set $D = \text{Div}(\pi^* f)_{\text{exc}}$ and $C = \text{Div}(\pi^* g)_{\text{exc}}$. Noether's intersection formula [9, Theorem 3.3.1] gives an expression for their intersection multiplicity at O as $[\xi \cdot \eta]_O = \sum_p e_p(D) \cdot e_p(C) = (\sum_p e_p(D) \overline{E}_p) \cdot (\sum_p e_p(C) \overline{E}_p) = D \cdot C$. Hence, $v_p(f) = [\xi \cdot \eta_p]_O = D \cdot B_p$, where $\eta_p : f_p = 0$, with $f_p \in \mathcal{O}_{X,O}$ as in Construction 2.3.

2.3. Complete ideals and antinef divisors. Given an effective divisor $D = \sum_{p \in K} v_p E_p \in \Lambda_\pi$, we may consider its associated ideal sheaf $\pi_* \mathcal{O}_{X'}(-D)$. Its stalk at O is

$$(2.3) \quad H_D = \{f \in \mathcal{O}_{X,O} \mid v_p(f) \geq v_p \text{ for all } E_p \leq D\},$$

This ideal H_D is *complete*, see [32], and \mathfrak{m} -primary since D has only exceptional support. Complete ideals are closed under all standard operations on ideals, except addition: the intersection, product and quotient of complete ideals is complete.

Recall that an effective divisor $D \in \text{Div}(X')$ is called *antinef* if $\rho_p = -D \cdot E_p \geq 0$, for every exceptional component $E_p, p \in K$. This notion is equivalent, in the total transform basis, to

$$(2.4) \quad e_p(D) \geq \sum_{q \rightarrow p} e_q(D), \quad \text{for all } p \in K.$$

These are usually called *proximity inequalities*, see [9, §4.2]. By means of the relation given in Equation (2.3), Zariski [32] establishes an isomorphism of semigroups between the set of ideals H_D and the set of antinef divisors in Λ_π .

Given a non-antinef divisor D , one can compute an equivalent antinef divisor \tilde{D} , called the *antinef closure*, under the equivalence relation that both divisors define the same ideal, i.e. $\pi_* \mathcal{O}_{X'}(-D) = \pi_* \mathcal{O}_{X'}(-\tilde{D})$, and via the so called *unloading* procedure. This is an inductive procedure which was already described in the work of Enriques [13, IV.II.17] (see [9, §4.6] for more details). The version that we present here is the one considered in [5].

Algorithm 2.5. (*Unloading procedure* [5])

Input: A divisor $D = \sum d_p E_p \in \Lambda_\pi$.

Output: The antinef closure \tilde{D} of D .

Repeat:

- Define $\Theta := \{E_p \in \Lambda_\pi \mid \rho_p = -D \cdot E_p < 0\}$.
- Let $n_p = \left\lceil \frac{\rho_p}{E_p^2} \right\rceil$ for each $E_p \in \Theta$. Notice that $(D + n_p E_p) \cdot E_p \leq 0$.
- Define a new divisor as $\tilde{D} = D + \sum_{E_p \in \Theta} n_p E_p$.

Until the resulting divisor \tilde{D} is antinef.

A fundamental result of Zariski [32] establishes the unique factorization of complete ideals into simple complete ideals, an ideal being simple if it is not the product of ideals different from the unit. A reinterpretation of this result in a more geometrical context is given by Casas-Alvero in [9, §8.4].

Theorem 2.6. [32], [9, §8.4] *Let $D \in \Lambda_\pi$ an antinef divisor expressed as $D = \sum_{p \in K} \rho_p B_p$ in the branch basis. Then,*

$$(2.5) \quad H_D = \prod_{p \in K} H_{B_p}^{\rho_p},$$

with H_{B_p} being simple complete ideals for any $p \in K$.

In the sequel, we will call *simple divisor* the unique antinef divisor defining a simple complete ideal. As a corollary of Theorem 2.6, simple divisors in Λ_π will always be equal to B_p for some $p \in K$.

2.4. Maximal contact elements. Let $\xi : f = 0$ be a germ of an irreducible element such that π dominates the resolution of (f) . Its equisingularity class, or topological equivalence class, is determined by the set of *characteristic exponents* $\{m_1/n, \dots, m_g/n\}$, where $n = e_O(f)$ introduced in [33]. Then, f has a Puiseux series of the form

$$(2.6) \quad s(x) = \sum_{\substack{j \in (n_0) \\ m_0 \leq j < m_1}} a_j x^{j/n} + \sum_{\substack{j \in (n_1) \\ m_1 \leq j < m_2}} a_j x^{j/n} + \cdots + \sum_{\substack{j \in (n_{g-1}) \\ m_{g-1} \leq j < m_g}} a_j x^{j/n} + \sum_{\substack{j \in (n_g) \\ j \geq m_g}} a_j x^{j/n},$$

where $m_0 = 0$, and $n_i = \gcd(n, m_1, \dots, m_i)$ so that, in particular, $n_0 = n$ and $n_g = 1$.

Another way of describing the equisingularity class is using its *semigroup* (see [34]). Consider $v_\xi : \mathcal{O}_{X,O} \rightarrow \mathbb{Z}$ the valuation induced by the intersection multiplicity of ξ with another element $\zeta : h = 0, h \in \mathcal{O}_{X,O}, h \notin (f)$. It is defined by $v_\xi(\zeta) = [\xi \cdot \zeta]_O = \text{ord}_t h(t^n, s(t^n))$, and does not depend on the equation defining the germ ζ or the parameterization of ξ . Then, the semigroup of ξ is

$$\Sigma(\xi) = \{v_\xi(\zeta) \in \mathbb{Z} \mid \zeta : h = 0, h \in \mathcal{O}_{X,O}, h \notin (f)\},$$

and it is finitely generated. Namely, $\Sigma(\xi) = \langle n, \check{m}_1, \dots, \check{m}_g \rangle$ where

$$(2.7) \quad \check{m}_i = \sum_{j=1}^{i-1} \frac{(n_{j-1} - n_j)m_j}{n_{i-1}} + m_i, \quad \text{for } i = 1, \dots, g.$$

Throughout this work we will be interested in those elements $f_i \in \mathcal{O}_{X,O}$ such that $v_\xi(\gamma_i) = [\xi \cdot \gamma_i]_O = \check{m}_i$, with $\gamma_i : f_i = 0$ for $i = 0, \dots, g$. They will be called *maximal contact elements* of ξ . Several choices for each f_i giving raise to different germs can be made; for instance, if ξ is not tangent to the y -axis, then $f_0 = x$ and $f_1 = y + a_1x + a_2x^2 + \cdots$. In general, these elements can be explicitly computed. Namely, if the equation defining ξ is in Weierstrass form, the maximal contact elements correspond to its approximate roots (see [22], [23]). Alternatively, if one has a Puiseux series of ξ as in (2.6), then the equations of the maximal contact elements f_i have Puiseux series:

$$(2.8) \quad s_i(x) = \sum_{\substack{j \in (n_0) \\ m_0 \leq j < m_1}} a_j x^{j/n} + \cdots + \sum_{\substack{j \in (n_{i-1}) \\ m_{i-1} \leq j < m_i}} a_j x^{j/n} + \cdots \quad \text{for } i = 1, \dots, g,$$

where the non-explicit terms are assumed not to increase the polydromy order, [9, §1.2], n/n_{i-1} of s_i , and either $f_0 = x$ or $f_0 = y$ depending on whether ξ is tangent to the x -axis or the y -axis respectively, see [9, §5.8]. Notice that the multiplicity at the origin of these maximal contact elements is $e_O(f_i) = n/n_{i-1}$ for $i = 1, \dots, g$.

We introduce here the notion of *maximal contact elements* for a proper birational morphism $\pi : X' \rightarrow X$. The maximal contact elements of π with exceptional divisor E are those $f_p \in \mathcal{O}_{X,O}$ with $p \in K$ considered in Construction 2.3, such that the vertex p is a dead-end of the dual graph of E_{exc} , i.e. the dual graph remains connected when the vertex is removed. In particular, a dead-end vertex will always be a free point or the origin O , since satellite points are always proximate to two points. A *set of maximal contact elements* $\{f_i\}_{i \in I}$ contains a unique $f_{p_i} = f_i$ for each dead-end vertex $p_i \in K$. Since these elements are determined by a finite number of valuative conditions, the elements f_i can always be chosen to be polynomials instead of power series. Moreover, these polynomials can be chosen in an intrinsic unique way which is valid for any π . This will be a key feature for computational issues.

This definition coincides with the one given for an irreducible element $f \in \mathcal{O}_{X,O}$. The elements $f_p \in \mathcal{O}_{X,O}$ such that p is a dead-end of the dual graph of the resolution of (f) are exactly those with Puiseux series as in (2.8). Conversely, we can construct equations for the maximal contact elements f_p with $p \in K$ of any divisor F parameterized by K : the coordinates, as points in the projective line, $q \in E_{q'} \cong \mathbb{P}_{\mathbb{C}}^1, q \rightarrow q'$, for all $q \leq p, e_q(f_p) \neq 0$ determine, and are determined by, the coefficients of a Puiseux series of any $f_p \in \mathcal{O}_{X,O}$, see [9, §5.7], which will be as in (2.8).

Remark 2.7. From any set of maximal contact elements $\{f_i\}$ one can always recover the morphism $\pi : X' \rightarrow X$ as the minimal resolution dominating the resolutions of $\{f_i\}$.

Example 2.8. Consider the proper birational morphism $\pi : X' \rightarrow X$ consisting in blowing-up four points with the configuration given by the following dual graph:



Take, for instance, $D = 4E_O + 6E_{p_1} + 12E_{p_2} + 13E_{p_3} + 26E_{p_4}$. The dead-end points are precisely O, p_1, p_3 . Therefore, a set of maximal contact elements for π is $\{f_0, f_1, f_2\}$ with

$$\begin{aligned} f_0 &= x + a_{2,0}x^2 + a_{0,2}y^2 + a_{1,1}xy + \cdots, \\ f_1 &= y + b_2x^2 + b_3x^3 + \cdots, \\ f_2 &= y^2 - x^3 + \sum_{3i+2j>6} c_{i,j}x^i y^j, \end{aligned}$$

and different choices of $a_{i,j}, b_i, c_{i,j} \in \mathbb{C}$ will give different sets of maximal contact elements. For instance, if all the coefficients are chosen to be zero, $f_0 = x, f_1 = y, f_2 = y^2 - x^3$, which are polynomials.

3. AN ALGORITHM TO COMPUTE H_D

Let (X, O) be a germ of smooth complex surface and let $\pi : X' \rightarrow X$ be a proper birational transform. In this section we present an algorithm which computes a set of generators for the \mathfrak{m} -primary ideal H_D for any divisor D with exceptional support in X' , i.e. $D \in \Lambda_\pi$.

We briefly describe the main ideas behind Algorithm 3.5. We start with divisor $D \in \Lambda_\pi$ which can be assumed to be antinef. It decomposes into simple divisors $D = \rho_{q_1}B_{q_1} + \cdots + \rho_{q_r}B_{q_r}$ with all $\rho_{q_i} > 0$. For each simple divisor $B_{q_i}, q_i \neq O$, we compute the antinef closure of $B_{q_i} + E_O$ which we denote \widehat{D}_i . This new divisor describes a particular *adjacent ideal* $H_{\widehat{D}_i}$ below $H_{B_{q_i}}$, i.e. an ideal $H_{\widehat{D}_i} \subsetneq H_{B_{q_i}}$ such that $\dim H_{B_{q_i}}/H_{\widehat{D}_i} = 1$ as \mathbb{C} -vector space. Next, we find, among the set of maximal contact elements of π , an element $f \in \mathcal{O}_{X,O}$ belonging to $H_{B_{q_i}}$ but not to $H_{\widehat{D}_i}$. Now, \widehat{D}_i is no longer simple but has smaller support than B_{q_i} in the total transform basis. Therefore we may repeat the same procedure with $D := B_{q_j}, j < i$ until $D = B_O := \text{Div}(\pi^*\mathfrak{m})$.

The first part of the algorithm can be represented by a tree where each vertex is an antinef divisor. The leaves of the tree are all $B_O = \text{Div}(\pi^*\mathfrak{m})$ and the root is the initial

divisor D . The second part of the algorithm traverses the tree bottom-up computing, in each node, the ideal associated to the divisor of that node. Using the notations from the above paragraph, given any node in the tree with divisor D , the ideal H_D is computed multiplying the ideals in child nodes and adding the element f to the resulting generators.

Before giving a more explicit description of the algorithm, let us first state two technical results. The first one presents some properties of adjacent ideals based on results obtained by Fernández-Sánchez (see [14], [15], [16]) in the study of sandwiched singularities and the Nash conjecture of arcs on these singularities.

Proposition 3.1. *Let H_D be the complete ideal defined by an antinef divisor $D \in \Lambda_\pi$. Consider \widehat{D} the antinef closure of $D + E_O$, obtained from $D + E_O$ by unloading on a given¹ subset of points $T \subseteq K$. Then, $H_{\widehat{D}} \not\subsetneq H_D$ are adjacent ideals if and only if $\rho_O(D) = 0$.*

Furthermore, if $H_{\widehat{D}} \not\subsetneq H_D$ are adjacent then,

- i) $\sum_{p \in T} E_p$ is the connected component of $\sum_{p \in K, \rho_p(D)=0} E_p$ containing E_O ;
- ii) $e_O(\widehat{D}) = e_O(D) + 1$, and $e_p(D) - 1 \leq e_p(\widehat{D}) \leq e_p(D)$ for any $p \in K$, $p \neq O$;
moreover $\rho_O(\widehat{D}) > 0$;
- iii) if $p \in K \setminus T$ and p is proximate to some point in T then, $e_p(\widehat{D}) = e_p(D) - 1$.

Proof. Consider the cluster K' obtained from K by adding $r := \rho_O(D) + 1$ free points p_1, \dots, p_r lying on E_O . Let $\pi' : Y' \rightarrow X$ be the composition of π with the sequence of blow-ups of the points p_1, \dots, p_r . Denote by $\overline{G} \in \text{Div}(Y')$ the pullback of any $G \in \text{Div}(X')$. For simplicity, denote the strict and the total transform basis by $\{E_p\}_{p \in K'}$ and $\{\overline{E}_p\}_{p \in K'}$ respectively in the lattice $\Lambda_{\pi'}$.

Clearly, both $\overline{D} + E_O$ and $\overline{D} + E_{p_1} + \dots + E_{p_r}$ are not consistent, whereas $\overline{D} + E_{p_1} + \dots + E_{p_i}$ are consistent for all $1 \leq i < r$. Moreover, when applying the unloading procedure described in Algorithm 2.5, we find that the antinef closures of $\overline{D} + E_O$ and $\overline{D} + E_{p_1} + \dots + E_{p_r}$ are the same, say it \widehat{D}' , and $e_{p_i}(\widehat{D}') = 0$ for all $1 \leq i \leq r$. Indeed, the first step of the unloading procedure applied to $\overline{D} + E_O$ or $\overline{D} + E_{p_1} + \dots + E_{p_r}$ gives the same divisor $\overline{D} + E_O + E_{p_1} + \dots + E_{p_r}$. Furthermore, \widehat{D}' is the pullback of the antinef closure \widehat{D} of $D + E_O$ in $\text{Div}(X')$ and hence they define the same complete ideal $H_{\widehat{D}'} = H_{\widehat{D}}$.

Now, from [9, §4.7], the codimension of a complete ideal H_G defined by a divisor $G \in \Lambda_\pi$ satisfies

$$\dim \mathcal{O}_{X,O}/H_G = \sum_{p \in K} \frac{e_p(\widetilde{G})(e_p(\widetilde{G}) + 1)}{2} \leq \sum_{p \in K} \frac{e_p(G)(e_p(G) + 1)}{2},$$

where \widetilde{G} is the antinef closure of G . Hence,

$$H_{\widehat{D}} = H_{\overline{D} + E_{p_1} + \dots + E_{p_r}} \not\subsetneq H_{\overline{D} + E_{p_1} + \dots + E_{p_{r-1}}} \not\subsetneq \dots \not\subsetneq H_{\overline{D} + E_{p_1}} \not\subsetneq H_D$$

is a chain of adjacent complete ideals, giving $\dim H_{B_{q_i}}/H_{\widehat{D}_i} = r = \rho_O(D) + 1$. Therefore, $H_{\widehat{D}} \not\subsetneq H_D$ are adjacent, if and only if $\rho_O(D) = 0$.

Finally, from [3, 2.1] and [14, 4.6] claim i) follows. Claim ii) and iii) are consequences of [15, 4.2] and [16, 2.2] (see also [4, 3.1]). \square

¹ T is the set of points $p \in K$ that parameterize the support of $\widehat{D} - (D + E_O)$. Notice that T may be empty.

Remark 3.2. Although there may be multiple adjacent ideals $H_{\widehat{D}}$ to a fixed ideal H_D , the adjacent ideal considered in Proposition 3.1 is unique with the property that $e_O(\widehat{D}) = e_O(D) + 1$, so we will refer to it as *the* adjacent ideal to H_D . This property turns out to be crucial for the finiteness of the algorithm.

Remark 3.3. Notice that if D is a simple divisor such that $H_D \neq \mathfrak{m}$, then $\rho_O(D) = 0$ and $H_{\widehat{D}}$ is always adjacent. Furthermore, the unloading step is always required, i.e. T is always non empty in this case.

Lemma 3.4. *Let H_D be the complete ideal defined by an antinef divisor $D \in \Lambda_\pi$. The divisor $D + \overline{E}_O$ is antinef and $H_{D+\overline{E}_O} = \mathfrak{m}H_D$.*

Proof. Clearly $\mathfrak{m} = \{f \in \mathcal{O}_{X,O} \mid e_O(f) \geq 1\}$. Thus, $\mathfrak{m} = H_{\overline{E}_O}$, and since $\overline{E}_O = B_O$ it is antinef and the result follows by the correspondence between antinef divisor and complete ideals in section 2.3. \square

Algorithm 3.5. (*Generators for H_D*)

Input: *A proper birational morphism $\pi : X' \rightarrow X$ and an antinef divisor $D \in \Lambda_\pi$.*

Output: *Generators for the ideal H_D .*

1. *Compute and fix a set of maximal contact elements $\{f_i\}$ with $i \in I$ of π .*
2. *Set $D^{(0)} := D$ and proceed from step (0.1).*

Step (i):

i.1 Decompose $D^{(i)}$ into $d_i := \#\{p \in K \mid \rho_p(D^{(i)}) > 0\}$ simple divisors.

i.2 For each $j = 1, \dots, d_i$, consider $q_j \in \{p \in K \mid \rho_p(D^{(i)}) > 0\}$ and assume $B_{q_j} = \sum_{p \in K} e_p \overline{E}_p$.

*i.j.1 **Stop at the maximal ideal:***

If $B_{q_j} = B_O := \text{Div}(\pi^\mathfrak{m})$, then set $H_{B_{q_j}} = (f_{i_0}, f_{i_1})$ for $i_0, i_1 \in I$ such that they are smooth and transverse at O , then stop. Otherwise, proceed from i.j.2.*

*i.j.2 **Compute the adjacent ideal to $H_{B_{q_j}}$:***

Perform unloading on the divisor $B_{q_j} + E_O$ to get its antinef closure \widehat{D}_j .

*i.j.3 **Select a maximal contact element in $H_{B_{q_j}} \setminus H_{\widehat{D}_j}$:***

Let $p \in K$ be the last free point such that $e_p \neq 0$. Take $\tau_j \in I$ such that $e_p(f_{\tau_j}) = 1$ and $e_O(f_{\tau_j}) \leq e_O$. Define the integer $n_j := e_O/e_O(f_{\tau_j})$.

*i.j.4 **Recursive step:***

Assume that $H_{\widehat{D}_j}$ has been computed after performing step (i + 1) with $D^{(i+1)} := \widehat{D}_j$.

*i.j.5 **Set:***

$$H_{B_{q_j}} = \left(f_{\tau_j}^{n_j} \right) + H_{\widehat{D}_j}.$$

i.3 Apply Zariski's factorization theorem:

Compute the product $H_{D^{(i)}} = \prod_{j=1}^{d_i} H_{B_{q_j}}^{\rho_{q_j}}$, giving generators h_1, \dots, h_{s_i} .

i.4 Set, using Nakayama's lemma:

$$H_{D^{(i)}} = \left(h_k \mid \pi^* h_k \notin \mathcal{O}_{X'}(-D^{(i)} - \bar{E}_O), k = 1, \dots, s_i \right) \mathcal{O}_{X,O}.$$

3. **Return:** $H_D = H_{D^{(0)}}$.

Remark 3.6. In order to clarify some steps of the algorithm we point out the following:

- At step 1 of Algorithm 3.5 a set of maximal contact elements $\{f_i\}_{i \in I}$ of π is fixed. The specific choice of the germs $f_i = 0$ nor of the equations f_i do not affect the output of the algorithm: the monomial expression remains the same for whatever choice, since the algorithm only uses the information of the equisingularity types of the maximal contact elements. However, as mentioned in Section 2.4, the set $\{f_i\}$ can be chosen in an intrinsic unique way, which is convenient for computational issues.
- At steps (i.j.1) and (i.j.3) of Algorithm 3.5 we have to choose maximal contact elements. These choices are not necessary unique as several maximal contact elements may fulfill the required conditions. However, by simply keeping record of the dead-ends indexing these chosen maximal contact elements, our algorithm is able to produce the same output whatever the input equations of the ideal are, and whenever it is used with two different input ideals which have the same integral closure.
- Since the sheaf ideals $\mathcal{O}_{X'}(-D)$, with $D \in \text{Div}(X')$, are defined by valuations, testing whether the pullback of an element f belongs to $\mathcal{O}_{X'}(-D)$ or not is only a matter of comparing the values $v_p(\text{Div}(\pi^* f))$ and $v_p(D)$ for all $p \in K$.
- It is clear from Nakayama's lemma and Lemma 3.4 that a set of elements of $\mathcal{O}_{X,O}$ is a system of generators of H_D if and only if its classes modulo $H_{D+\bar{E}_O}$ are a system of generators of $H_D/H_{D+\bar{E}_O}$ as \mathbb{C} -vector space. Equivalently, any element of $H_{D+\bar{E}_O}$ is redundant in a system of generators of H_D .

Example 3.7. We will compute H_D for the divisor D and the morphism π from Example 2.8. Let us fix the set of maximal contact elements $f_0 = x, f_1 = y, f_2 = y^2 - x^3$. The steps of Algorithm 3.5 applied to $D = 4E_O + 6E_{p_1} + 12E_{p_2} + 13E_{p_3} + 26E_{p_4}$ will be illustrated by means of the tree-shaped graph in Figure 1.

Each vertex of the tree contains an antinef divisor. In this example, we use dual-graphs to represent them. The root node contains the initial divisor D . Dashed arrows connect simple divisors B_{q_j} with its corresponding adjacent \widehat{D}_j from step (i.j.2). The maximal contact elements from step (i.j.3) that belong to $H_{B_{q_j}}$ but not to $H_{\widehat{D}_j}$ are represented next to dashed arrows. Solid arrows connect $\widehat{D}_j =: D^{(i+1)}$ with each of its irreducible components B_p , with $p \in K$. Finally, the weight $\rho_p^{(i)}$ of each divisor $B_p, p \in K$, in $\widehat{D}^{(i)}$ is written next to the solid arrows.

We have added some extra indices to the divisors appearing in the algorithm to highlight at which step we encounter them. Hopefully it does not create any confusion since its




FIGURE 1. Tree of divisors from Algorithm 3.5 in Example 3.7.

meaning should be clear from the context. The generators of the ideals associated to the divisors in each intermediate step are then:

- $H_{B_O} = \mathfrak{m} = (x, y)$.
- $H_{B_{p_1}} = (y) + H_{D_1^{(2)}} = (y) + \mathfrak{m}^2 = (y, x^2, xy, y^2)$.
- $H_{B_{p_2}} = (y^2) + H_{D_2^{(2)}} = (y^2) + \mathfrak{m}^3 = (y^2, x^3, x^2y, xy^2, y^3)$.
- $H_{D^{(1)}} = B_O^2 \cdot B_{p_1} \cdot B_{p_2} = (x, y)^2 \cdot (y, x^2, xy, y^2) \cdot (y^2, x^3, x^2y, xy^2, y^3)$
 $= (x^7, \cancel{x^6y}, \cancel{x^5y^2}, \cancel{x^4y^3}, \dots, x^5y, x^4y^2, \dots, \cancel{x^3y^3}, \cancel{x^2y^4}, x^2y^3, xy^4, y^5)$
 $= (x^7, x^5y, x^4y^2, x^2y^3, xy^4, y^5)$.
- $B_{p_4} = ((y^2 - x^3)^2) + H_{D^{(1)}} = ((y^2 - x^3)^2, x^7, x^5y, x^4y^2, x^2y^3, xy^4, y^5)$.
- $H_D := H_{D^{(0)}} = B_{p_4} = ((y^2 - x^3)^2, x^7, x^5y, x^4y^2, x^2y^3, xy^4, y^5)$.

The crossed out elements are those that are redundant by step (i.4) and Remark 3.6. Observe that, although many crossed out elements are actually multiple of other elements, step (i.4) and Remark 3.6 allows us to remove y^5 which is not multiple of any other element.

Remark 3.8. As an outcome of the algorithm, we see that H_D admits the following monomial expression

$$H_D = (f_2^2, f_0^7, f_0^5 f_1, f_0^4 f_1^2, f_0^2 f_1^3, f_0 f_1^4),$$

in the set of maximal contact elements $f_0 = x, f_1 = y, f_2 = y^2 - x^3$ associated to π that we fixed in the beginning. It is worth remarking that we would get this same monomial expression for any other set of maximal contact elements chosen in the beginning.

However, we might get a different monomial expression depending on the maximal contact elements (or powers of) that we choose in step (i.j.3) of Algorithm 3.5. In this example, when choosing an element in $H_{B_{p_3}}$ that does not belong to $H_{\widehat{D}_2^{(1)}}$ we took $f_1^2 = y^2$, but we could also had chosen $f_2 = y^2 - x^3$. In this later case the final system of generators is

$$H_D = ((y^2 - x^3)^2, x^2y(y^2 - x^3), xy^2(y^2 - x^3), x^7, x^5y, x^4(y^2 - x^3), x^4y^2),$$

so we get the monomial expression

$$H_D = (f_2^2, f_0^2 f_1 f_2, f_0 f_1^2 f_2, f_0^7, f_0^5 f_1, f_0^4 f_2, f_0^4 f_1^2).$$

4. CORRECTNESS OF THE ALGORITHM

In this section we will prove that Algorithm 3.5 developed in Section 3 is correct. First, we need to check that it ends after a finite number of steps. The key point is to prove that the divisor \widehat{D}_j defining the adjacent ideal to the simple ideal $H_{B_{q_j}}$ has smaller support in the total transform basis than B_{q_j} .

Lemma 4.1. *Using the notations in Algorithm 3.5, assume that B_{q_j} is a simple divisor different from $B_O = \text{Div}(\pi^*\mathfrak{m})$. Let \widehat{D}_j be the antinef closure of $B_{q_j} + E_O$ computed in step (i.j.2). Then, \widehat{D}_j has smaller support than B_{q_j} in the total transform basis. That is,*

$$|\text{Supp}_{\overline{E}}(B_{q_j})| > |\text{Supp}_{\overline{E}}(\widehat{D}_j)|.$$

Proof. Since $B_{q_j} \neq B_O$, the excess of B_{q_j} at O is $\rho_O(B_{q_j}) = 0$. Then, according to Proposition 3.1, the ideal defined by $B_{q_j} + E_O$ is an adjacent ideal below $H_{B_{q_j}}$. Let $T \subseteq K$ be the points on which unloading is performed to obtain the antinef closure \widehat{D}_j from $B_{q_j} + E_O$. By Proposition 3.1, T are the points $p \in K$ whose associated exceptional divisor E_p belongs to the same connected component as E_O in $\sum_{O \leq p < q_j} E_p$. Observe that $\sum_{O \leq p < q_j} E_p$ has either one or two components, according if q_j is either free or satellite. In both cases, q_j is proximate to the point $p \in T$ whose exceptional divisor cuts E_{q_j} , i.e. $E_p \cdot E_{q_j} = 1$. Hence, invoking Proposition 3.1 again, the multiplicity at q_j of \widehat{D}_j , after performing unloading on $B_{q_j} + E_O$, decreases by one. Since B_{q_j} is simple, the multiplicity of B_{q_j} at q_j is one. Hence, the multiplicity of \widehat{D}_j at q_j is zero, giving the desired result. \square

In the next proposition we prove that Algorithm 3.5 ends after a finite number of steps. To emphasize the dependence of the divisors on a specific step (i) of the algorithm we will use the notation $B_{q_j}^{(i)}$ and $\widehat{D}_j^{(i)}$.

Proposition 4.2. *Algorithm 3.5 ends after a finite number of steps.*

Proof. As noted earlier, the points $q \in K$ such that $e_q(B_p) \neq 0$ are totally ordered. Then, using Equation (2.4), the sequence of multiplicities of B_p decrease along those points. Hence, we have $|B_p|_{\overline{E}} = 1$ for some $p \in K$ if and only if $p = O$ and then, $B_p = \text{Div}(\pi^*\mathbf{m})$.

Using the notations in Algorithm 3.5, assume that we are in step (i) and we have a simple divisor B_{q_j} in step ($i.j.1$). If $q_j = O$, then $B_O = \text{Div}(\pi^*\mathbf{m})$ and we are done. Otherwise, since $q_j \neq O$, we have that $|\text{Supp}_{\overline{E}}(B_{q_j})| > |\text{Supp}_{\overline{E}}(\widehat{D}_j^{(i)})|$ by Lemma 4.1. Since $D^{(i+1)} := \widehat{D}_j^{(i)}$ admits a decomposition $D^{(i+1)} = \sum_{p \in K} \rho_p^{(i+1)} B_p$, we have $|\text{Supp}_{\overline{E}}(D^{(i+1)})| \geq |\text{Supp}_{\overline{E}}(B_p)|$ for all B_p with $\rho_p^{(i+1)} > 0$. Hence, $|\text{Supp}_{\overline{E}}(B_{q_j})| > |\text{Supp}_{\overline{E}}(B_p)|$, for all p with $\rho_p^{(i+1)} > 0$, and by induction we obtain the desired result. \square

Lemma 4.3. *Let B_q be a branch basis divisor associated to a satellite point $q \in K$. Let $\Sigma(\xi) = \langle n, \check{m}_1, \dots, \check{m}_r \rangle$ be the semigroup of $\xi : f_q = 0$ and take $\gamma : f_r = 0$ such that $[\gamma \cdot \xi]_O = \check{m}_r$. Then, $B_q^2 = [\zeta \cdot \xi]_O$ with $\zeta : f_r^{n_r-1} = 0$. Furthermore, $v_p(f_r^{n_r-1}) \geq v_p(B_q)$, for all $p \leq q$.*

Proof. Assume that $B_q = \sum_{p \in K} e_p \overline{E}_p$. The first claim follows from the following computation:

$$B_q^2 = \sum_{p \in K} e_p^2 = \sum_{i=1}^r (m_i - m_{i-1}) n_{i-1} = \sum_{i=1}^{r-1} (n_{i-1} - n_i) m_i + n_{r-1} m_r = n_{r-1} \check{m}_r = n_{r-1} [\gamma \cdot \xi]_O = [\zeta \cdot \xi]_O.$$

where the second equality is true since $q \in K$ is a satellite point (see [9, §5.10]) and the fourth equality comes from (2.7).

To prove the second claim, in virtue of Remark 2.4, it suffices to check the inequalities $[\zeta \cdot \eta_p]_O \geq [\xi \cdot \eta_p]_O$ for any $p \leq q$ with $\eta_p : f_p = 0$. We will use known properties of the ultrametric $d_{\mathcal{C}}$ distance (introduced in [25]) defined over the space \mathcal{C} of plane branches as $\frac{1}{d_{\mathcal{C}}(C,D)} = \frac{[C,D]_O}{e_O(C)e_O(D)}$ for any $C, D \in \mathcal{C}$. Hence, the inequalities above are equivalent to $d_{\mathcal{C}}(\xi, \eta_p) \geq d_{\mathcal{C}}(\gamma, \eta_p)$ for any $p \leq q$, since in our case

$$\frac{[\zeta \cdot \eta_p]_O}{e_O(\zeta)e_O(\eta_p)} = \frac{n_{g-1} [\gamma \cdot \eta_p]_O}{n_{g-1} e_O(\gamma) e_O(\eta_p)} = \frac{1}{d_{\mathcal{C}}(\gamma, \eta_p)}.$$

Notice that $f_q = f_{q_g}$ for some point q_g , $O \leq q_g \leq q$, which corresponds to a dead-end in the dual graph of B_q . Now we summarize the results on the ultrametric space of plane branches of [1, 3.1, 3.2 and 3.4] adapted to our setting:

- $d_{\mathcal{C}}(\xi, \eta_p) = d_{\mathcal{C}}(\gamma, \eta_p)$, if $O \leq p < q_g$;
- $d_{\mathcal{C}}(\xi, \eta_p) = d_{\mathcal{C}}(\gamma, \eta_p)$, if $q_g < p \leq q$ and in the dual graph of B_q the vertex of p lies on the segment joining the vertexes q and q_g ;
- $d_{\mathcal{C}}(\xi, \eta_p) > d_{\mathcal{C}}(\gamma, \eta_p)$, otherwise.

Hence, the second claim follows. \square

Proposition 4.4. *Using the notations in Algorithm 3.5, at any step (i) of the algorithm, there exists a power of a maximal contact element $f_{\tau_j}^{n_j} \in \mathcal{O}_{X,O}$ as required at step (i.j.3) and such element belongs to $H_{B_{q_j}}$ but not to $H_{\widehat{D}_j}$.*

Proof. We are going to break the proof of the first statement in two cases depending on whether the point $q_j \in K$ is free or satellite. With the notations from step (i.j.3), $p \in K$ will be the last free point such that $e_p(B_{q_j}) \neq 0$.

Assume first that q_j is free, i.e. $p = q_j$. If, in addition, the vertex of q_j is a dead-end of the dual graph of F we are done, since $f_{\tau_j} = f_{q_j}$ and $f_{q_j} \in H_{B_{q_j}}$. If q_j is not a dead-end of the dual graph, there is a dead-end $q \in K$ and a totally ordered sequence $q_j \leq p_1 \leq \dots \leq p_r \leq q$ of free points such that $e_{q_j}(B_q) = e_{p_1}(B_q) = \dots = e_{p_r}(B_q) = e_q(B_q) = 1$, by (2.4). Therefore, $f_{\tau_j} = f_q$ with $e_{q_j}(f_q) = 1$ and $B_q = B_p + \overline{E}_{p_1} + \dots + \overline{E}_{p_r} + \overline{E}_q$, which implies, by (2.1), that $f_q \in H_{B_q} \subsetneq H_{B_{q_j}}$. In either case we have that $n_j = e_O(B_{q_j})/e_O(f_{\tau_j}) = 1$.

Now, assume that q_j is satellite and hence $p < q_j$. Let $\Sigma(\xi) = \langle n, \check{m}_1, \dots, \check{m}_g \rangle$ the semigroup of $\xi : f_{q_j} = 0$. By [9, §5.8], p has the property that any $\gamma : f_p = 0, f_p \in \mathcal{O}_{X,O}$ satisfies $[\gamma.\xi]_O = B_p \cdot B_{q_j} = \check{m}_g$. However, it may happen that $p \in K$ is not a dead-end. In this case, using the same argument as before, there is a dead-end $q \in K$ and a totally ordered sequence $p \leq p_1 \leq \dots \leq p_r \leq q$ of free points and $B_q = B_p + \overline{E}_{p_1} + \dots + \overline{E}_{p_r} + \overline{E}_q$. Since p is the last free point of B_{q_j} , $B_{q_j} \cdot \overline{E}_{p_i} = 0$ for $i = 1, \dots, r$ and also $B_{q_j} \cdot \overline{E}_q = 0$. Hence, $[\tilde{\gamma}.\xi]_O = B_q \cdot B_{q_j} = \check{m}_g$ with $\tilde{\gamma} : f_q = 0$, i.e. we can take $f_{\tau_j} = f_q$. We can then apply Lemma 4.3 to B_{q_j} with $f_g = f_{\tau_j}$ yielding that $f_{\tau_j}^{n_j} \in H_{B_{q_j}}$ with $n_j = e_O(B_{q_j})/e_O(f_{\tau_j})$.

Finally, if $f_{\tau_j}^{n_j} \in \mathcal{O}_{X,O}$ fulfills the requirements of step (i.j.3), then $e_O(f_{\tau_j}^{n_j}) = e_O(B_{q_j})$, but $e_O(\widehat{D}_j) > e_O(B_{q_j})$ by Proposition 3.1, therefore we have that $f_{\tau_j}^{n_j} \notin H_{\widehat{D}_j}$. \square

Theorem 4.5. *Let $\pi : X' \rightarrow X$ be a proper birational morphism and let $D \in \Lambda_{\pi}$. Then, Algorithm 3.5 computes a set of generators for H_D that are monomial in any given set of maximal contact elements of π .*

Proof. Let us prove that the i -th step of the algorithm returns a system of generators of $H_{D^{(i)}}$ which has the desired properties. By Zariski's factorization Theorem 2.6, it is enough to focus on computing generators for each simple ideal $H_{B_{q_j}}, j = 1, \dots, d_i$, in the decomposition of $D^{(i)}$. Fixing B_{q_j} at step (i.2), we will make induction on the order of the neighbourhood that $q_j \in K$ belongs to, and we will show that Algorithm 3.5 computes generators for $H_{B_{q_j}}$ which are monomials in the set of maximal contact elements.

If $q_j = O$, then $B_O = \text{Div}(\pi^*\mathfrak{m})$ and step (i.j.1) returns $H_{D^{(i)}} = \mathfrak{m}$, since a pair of smooth transverse elements generate \mathfrak{m} . By construction, any set of maximal contact elements contain such a pair of elements.

Assume now that $q_j \neq O$ and that the algorithm computes the generators of the ideals associated to B_p for $p < q_j$. By Proposition 3.1, $H_{\widehat{D}_j} \subsetneq H_{B_{q_j}}$ are adjacent ideals. Since $\widehat{D}_j = \sum_{p < q_j} \rho_p^{(i)} B_p$, we can apply the induction hypothesis to the simple divisors $B_p, p < q_j$ such that $\rho_p^{(i)} \neq 0$ and apply Theorem 2.6 to get

$$H_{\widehat{D}_j} = \prod_{p < q_j} H_{B_p}^{\rho_p^{(i)}} \subsetneq H_{D^{(i)}}.$$

At this point is it enough to add any element that belongs to $H_{B_{q_j}}$ but not to $H_{\widehat{D}_j}$ to get a system of generators of $H_{B_{q_j}}$. By Proposition 4.4, the element chosen at step (i.j.3) has the desired properties, namely, it is a power of a maximal contact element. Finally, we can remove unnecessary elements from the system of generators of $H_{D^{(i)}}$ using Lemma 3.4.

The dependency on the set of maximal contact elements $\{f_i\}_{i \in I}$ is only used in step (i.j.3). The conditions required to $\{f_p\}_{i \in I}$ depend only on a finite number of valuations associated to the exceptional divisors of π . These conditions are fulfilled by an infinite number of elements which can be part of a set of maximal contact elements. \square

Remark 4.6. We would like to stress the generality of the monomial generators in Theorem 4.5. Consider the monomials $\mathbf{z}^\alpha = \prod_{i \in I} z_i^{\alpha_i}, \alpha = (\alpha_i)_{i \in I}$ in the variables $z_i, i \in I$. Each variable z_i formally represents all possible elements f_p for a fixed dead-end of the dual graph of π . They all have the same value for the valuations associated to the exceptional divisors. Take now any set of maximal contact elements $\mathbf{f} = \{f_{p_i}\}_{i \in I}$ and denote $\mathbf{z}_{\mathbf{f}}^\alpha = \prod_{i \in I} f_{p_i}^{\alpha_i}$ the specialization $z_i \mapsto f_{p_i}$.

The result of Theorem 4.5 is that Algorithm 3.5 returns formally $(\mathbf{z}^{\alpha_1}, \dots, \mathbf{z}^{\alpha_r})$ and that for any two sets of maximal contact elements $\mathbf{f} = \{f_{p_i}\}_{i \in I}$ and $\mathbf{g} = \{g_{p_i}\}_{i \in I}$, both specializations $H_{D, \mathbf{f}} = (\mathbf{z}_{\mathbf{f}}^{\alpha_1}, \dots, \mathbf{z}_{\mathbf{f}}^{\alpha_r})$ and $H_{D, \mathbf{g}} = (\mathbf{z}_{\mathbf{g}}^{\alpha_1}, \dots, \mathbf{z}_{\mathbf{g}}^{\alpha_r})$ are equal, $H_D = H_{D, \mathbf{f}} = H_{D, \mathbf{g}}$, and are wanted the complete ideal H_D .

Corollary 4.7. *The set of monomial expressions returned by Algorithm 3.5 is a topological invariant of D , i.e. it is an invariant of the weighted dual graph of D .*

Proof. From the proof of Theorem 4.5 it follows that the algorithm only uses the information of the equisingularity types of the maximal contact elements and of the dual graph of D weighted by the natural numbers $v_p(D)$ for $p \in K$. \square

5. APPLICATION TO SOME FAMILIES OF COMPLETE IDEALS

Let (X, O) be a germ of smooth complex surface and let $\pi : X' \rightarrow X$ be a proper birational transform. The sheaf ideals $\mathcal{O}_{X'}(D)$ and its pushforward, for some $D \in \Lambda_\pi$, arise in many different contexts. The goal of this sections is two show how Algorithm 3.5 and Theorem 4.5 applies to different problems.

Our approach is specially useful when studying families of divisor $\{D_i\}$ in Λ_π since all the generators for all the ideals H_{D_i} will be given as monomials in any set of maximal contact elements.

5.1. Integral closure. Let $\mathfrak{a} \subseteq \mathcal{O}_{X, O}$ be an ideal which can be assumed to be \mathfrak{m} -primary after considering the decomposition $\mathfrak{a} = (a) \cdot \mathfrak{a}'$ with $a = \gcd(\mathfrak{a})$. Let $\pi : X' \rightarrow X$ be a

log-resolution of the ideal \mathfrak{a} , i.e., a proper birational morphism such that there exists F an effective Cartier divisor such that $\mathfrak{a} \cdot \mathcal{O}_{X'} = \mathcal{O}_{X'}(-F)$. Then, the integral closure $\bar{\mathfrak{a}}$ of \mathfrak{a} is just the ideal H_F , see [9, §8.3].

Therefore, we have a very simple method to compute the integral closure of any planar ideal that boils down to the following steps:

- Compute the divisor F associated to the log-resolution of \mathfrak{a} by using the algorithm from [2].
- Compute a set of generators for the ideal H_F using Algorithm 3.5.

Let us illustrate this situation with an small example.

Example 5.1. Let $\mathfrak{a} = ((y^2 - x^3)^2, x^2y^3) \subseteq \mathcal{O}_{X,O}$ an ideal. One can compute the log-resolution divisor of \mathfrak{a} using the algorithm from [2]. The log-resolution and its associated divisor F of \mathfrak{a} are precisely the proper birational morphism π and the divisor D from Example 2.8.

Namely, $F = 4E_O + 6E_{p_1} + 12E_{p_2} + 13E_{p_3} + 26E_{p_4}$ and we can take $f_0 = x, f_1 = y, f_2 = y^2 - x^3$ as a set of maximal contact elements of the log-resolution π . Thus, from the computation in Example 3.7, one deduces that

$$\bar{\mathfrak{a}} = H_F = \pi_* \mathcal{O}_{X'}(F) = ((y^2 - x^3)^2, x^7, x^5y, x^4y^2, x^2y^3, xy^4).$$

The following results follows directly from Theorem 4.5 and its corollaries.

Theorem 5.2. *Let $\mathfrak{a} \subseteq \mathcal{O}_{X,O}$ be an ideal. There exists a set of generators of the integral closure $\bar{\mathfrak{a}}$ that are monomial in any given set of maximal contact elements of the log-resolution of \mathfrak{a} .*

Corollary 5.3. *Let $\pi : X' \rightarrow X$ be a proper birational morphism. Any complete ideal $\mathfrak{a} \subseteq \mathcal{O}_{X,O}$ whose log-resolution is dominated by π admits a system of generators given by monomials in any set of maximal contact elements associated to π .*

Corollary 5.4. *The set of monomial expressions returned by Algorithm 3.5 for the integral closure of an ideal \mathfrak{a} is an equisingular invariant of \mathfrak{a} .*

5.2. Multiplier ideals. Let $\pi : X' \rightarrow X$ be a log-resolution of an ideal $\mathfrak{a} \subseteq \mathcal{O}_X$ and let F be the divisor such that $\mathfrak{a} \cdot \mathcal{O}_{X'} = \mathcal{O}_{X'}(-F)$. The *multiplier ideal* associated to \mathfrak{a} and some rational number $\lambda \in \mathbb{Q}_{\geq 0}$ is defined as

$$\mathcal{J}(\mathfrak{a}^\lambda) = \pi_* \mathcal{O}_{X'}(\lceil K_\pi - \lambda F \rceil),$$

where K_π is the so-called *relative canonical divisor* (see [20] for details). Multiplier ideals form a discrete nested sequence of ideals

$$\mathcal{O}_X \supsetneq \mathcal{J}(\mathfrak{a}^{\lambda_0}) \supsetneq \mathcal{J}(\mathfrak{a}^{\lambda_1}) \supsetneq \mathcal{J}(\mathfrak{a}^{\lambda_2}) \supsetneq \cdots \supsetneq \mathcal{J}(\mathfrak{a}^{\lambda_i}) \supsetneq \cdots$$

and the rational numbers $0 < \lambda_0 < \lambda_1 < \cdots$ where an strict inclusion of ideals is achieved are called the *jumping numbers* associated to \mathfrak{a} .

There are general algorithms as those developed by Shibuta [26] and Berkersch and Leykin [6] that, given a set of generators of \mathfrak{a} , return the list of jumping numbers and a minimal set of generators of the corresponding multiplier ideals. These algorithms use the theory of Bernstein-Sato polynomials and have been implemented in Macaulay2. However, it is difficult to compute large examples due to the complexity of these algorithms.


In the case of planar ideals, there are methods given by Järvilletho [19], Naie [24] and Tucker [29] to compute the set of jumping numbers. The first two authors of this manuscript and Dachs-Cadefau [5] gave an algorithm that computes sequentially the list of jumping numbers of a planar ideal and the antinef divisor associated to the corresponding multiplier ideal.

Combining the algorithms developed in [2] and [5] with Algorithm 3.5 we may provide a method that, given a set of generators of a planar ideal \mathfrak{a} , returns the set of jumping numbers and a set of generators of the corresponding multiplier ideals. Namely, we have to perform the following steps:

- Compute the divisor F associated to the log-resolution of \mathfrak{a} by using the algorithm from [2].
- Compute the sequence of jumping numbers $\{\lambda_j\}_{j \in \mathbb{Z}_{\geq 0}}$ and the divisor corresponding to the associated multiplier ideals $\{\mathcal{J}(\mathfrak{a}^{\lambda_j})\}_{j \in \mathbb{Z}_{\geq 0}}$, i.e. the antinef closures D_{λ_j} of $\lfloor \lambda_j F - K_\pi \rfloor$, using the main algorithm of [5].
- Compute a set of generators for the ideals $H_{D_{\lambda_j}}$ using Algorithm 3.5.

This method is illustrated with the following:

Example 5.5. Consider the ideal $\mathfrak{a} = ((y^2 - x^3)^3, x^3(y^2 - x^3)^2, x^6y^3) \subseteq \mathcal{O}_{X,O}$. The log-resolution of \mathfrak{a} can be computed using the algorithm from [2] and is represented by means of the following dual-graph:



The divisor F such that $\mathfrak{a} \cdot \mathcal{O}'_X = \mathcal{O}_{X'}(-F)$ is $F = 6E_O + 9E_1 + 18E_2 + 20E_3 + 21E_4 + 42E_5$. A set of maximal contact elements for the log-resolution of \mathfrak{a} is, for instance, $f_0 = x, f_1 = y, f_2 = y^2 - x^3$.

The jumping numbers smaller than 1 computed using the algorithm from [5] and generators for the associated multiplier ideals computed using Algorithm 3.5 can be found in Table 1.

It is a known result that the multiplier ideals $\mathcal{J}(\mathfrak{a}^\lambda)$ associated to \mathfrak{a} are the same when taking the completion $\bar{\mathfrak{a}}$ of \mathfrak{a} , i.e. $\mathcal{J}(\mathfrak{a}^\lambda) = \mathcal{J}(\bar{\mathfrak{a}}^\lambda)$. As a corollary of Theorem 4.5 and Corollary 5.3 we obtain the following result which resembles Howald's theorem [18] on the fact that multiplier ideals of monomial ideals are also monomial.

Theorem 5.6. *Let $\mathfrak{a} \subseteq \mathcal{O}_{X,O}$ be an ideal and consider its completion $\bar{\mathfrak{a}}$, that can be generated by monomials in any given set of maximal contact elements. Then, the multiplier ideals $\mathcal{J}(\mathfrak{a}^\lambda)$ are also generated by monomials in the same set of maximal contact elements of the log-resolution of \mathfrak{a} .*

λ_i	$\mathcal{J}(\mathfrak{a}^{\lambda_i})$
$\frac{5}{18}$	x, y
$\frac{7}{18}$	y, x^2
$\frac{4}{9}$	x^2, xy, y^2
$\frac{1}{2}$	xy, y^2, x^3
$\frac{23}{42}$	y^2, x^3, x^2y
$\frac{25}{42}$	$y^2 - x^3, x^2y, xy^2, x^4$
$\frac{11}{18}$	x^2y, xy^2, y^3, x^4
$\frac{9}{14}$	xy^2, y^3, x^4, x^3y
$\frac{29}{42}$	$x(y^2 - x^3), y(y^2 - x^3), x^3y, x^2y^2, x^5$
$\frac{13}{18}$	y^3, x^3y, x^2y^2, x^5
$\frac{31}{42}$	$y(y^2 - x^3), x^2y^2, xy^3, x^2(y^2 - x^3), x^4y$
$\frac{7}{9}$	$x^2y^2, xy^3, y^4, x^5, x^4y$
$\frac{11}{14}$	$x^2(y^2 - x^3), xy(y^2 - x^3), y^2(y^2 - x^3), x^4y, x^3y^2, x^6$
$\frac{5}{6}$	$xy(y^2 - x^3), y^2(y^2 - x^3), x^3(y^2 - x^3), x^3y^2, x^2y^3, x^5y$
$\frac{37}{42}$	$xy(y^2 - x^3), y^2(y^2 - x^3), x^3(y^2 - x^3), x^2y^3, xy^4, x^5y, x^4y^2, x^7$
$\frac{8}{9}$	$y^2(y^2 - x^3), x^5y, x^3(y^2 - x^3), x^2y^3, x^2y(y^2 - x^3), xy^4, x^4y^2, x^7$
$\frac{13}{14}$	$y^2(y^2 - x^3), x^3(y^2 - x^3), x^2y(y^2 - x^3), xy^4, y^5, x^4y^2, x^3y^3, x^7, x^6y$
$\frac{17}{18}$	$x^2y(y^2 - x^3), xy^2(y^2 - x^3), y^3(y^2 - x^3), x^4y^2, x^4(y^2 - x^3), x^3y^3, x^6y$
$\frac{41}{42}$	$x^2y(y^2 - x^3), xy^2(y^2 - x^3), y^3(y^2 - x^3), x^4(y^2 - x^3), x^3y^3, x^2y^4, x^6y, x^5y^2, x^8$

TABLE 1. The jumping numbers smaller than 1 and generators of the associated multiplier ideal for $\mathfrak{a} = ((y^2 - x^3)^3, x^3(y^2 - x^3)^2, x^6y^3)$.

5.3. Valuation filtration. Throughout this subsection we will use the notation from Section 2.4. Let $\xi : f = 0, f \in \mathcal{O}_{X,O}$ be a plane branch and let v_ξ be the valuation induced by the intersection multiplicity. Let \mathfrak{V}_i denote the ideal of all the elements in $\mathcal{O}_{X,O}$ with valuation greater or equal to i :

$$\mathfrak{V}_i := \{\zeta \in \mathcal{O}_{X,O} \mid v_\xi(\zeta) \geq i\}.$$

These ideals form a filtration in $\mathcal{O}_{X,O}$:

$$(5.1) \quad \mathcal{O}_{X,O} = \mathfrak{V}_0 \supseteq \mathfrak{V}_1 \supseteq \cdots \supseteq \mathfrak{V}_i \supseteq \mathfrak{V}_{i+1} \supseteq \cdots$$

such that $\mathfrak{V}_i \cdot \mathfrak{V}_j \subset \mathfrak{V}_{i+j}$ and $\bigcap_{i \in \mathbb{Z}_{\geq 0}} \mathfrak{V}_i = (f)$. This type of filtration was considered by Teissier in [28]. Since, the ideals \mathfrak{V}_i are defined by valuations, they are complete, and hence, have the form $\pi_* \mathcal{O}_{X'_i}(D_i)$ for some effective divisor D_i in some surface X'_i .

Example 5.7. Consider the germ of plane curve $\xi : f = 0, f = (y^2 - x^3)^2 - x^5y \in \mathcal{O}_{X,O}$. A resolution morphism $\pi : X' \rightarrow X$ of ξ is the same as the one given for the ideal in

Example 2.8. Using Algorithm 3.5 we obtain the generators of the filtration (5.1) until the last \mathfrak{V}_i dominated by the log-resolution of (f) , which we collect in Table 2.

i	\mathfrak{V}_i
1, 2, 3, 4	x, y
5, 6	y, x^2
7, 8	xy, x^2, y^2
9, 10	xy, y^2, x^3
11, 12	x^2y, y^2, x^3
13	$y^2 - x^3, x^2y, xy^2, x^4$
14	x^2y, xy^2, y^3, x^4
15, 16	xy^2, y^3, x^4, x^3y
17	$x(y^2 - x^3), y(y^2 - x^3), x^3y, x^2y^2, x^5$
18	y^3, x^3y, x^2y^2, x^5
19	$y(y^2 - x^3), x^2(y^2 - x^3), x^4y, x^2y^2, xy^3$
20	$x^4y, x^2y^2, xy^3, y^4, x^5$
21	$x^2(y^2 - x^3), xy(y^2 - x^3), y^2(y^2 - x^3), x^4y, x^3y^2, x^6$
22	$xy^3, y^4, x^4y, x^3y^2, x^6$
23	$xy(y^2 - x^3), y^2(y^2 - x^3), x^3y^2, x^3(y^2 - x^3), x^2y^3, x^5y$
24	$y^4, x^3y^2, x^2y^3, x^6, x^5y$
25	$(y^2 - x^3)^2, x^3(y^2 - x^3), x^2y(y^2 - x^3), x^5y, x^4y^2, x^7$
26	$(y^2 - x^3)^2, x^2y^3, xy^4, x^5y, x^4y^2, x^7$

TABLE 2. The ideals \mathfrak{V}_i of the filtration associated to the plane branch $f = (y^2 - x^3)^2 - x^5y$ for $i = 1, 2, \dots, 26$.

REFERENCES

- [1] I. Abío, M. Alberich-Carramiñana, V. González-Alonso, *The ultrametric space of plane branches*, *Comm. Algebra*, 39: 4206–4220, 2011. [15](#)
- [2] M. Alberich-Carramiñana, J. Álvarez Montaner and G. Blanco, *Effective computation of base points of two-dimensional ideals*, preprint available at arXiv:1605.05665. [2](#), [17](#), [18](#)
- [3] M. Alberich-Carramiñana, J. Fernández-Sánchez, *Equisingularity classes of birational projections of normal singularities to a plane*, *Adv. Math.*, 216: 753–770, 2007. [9](#)
- [4] M. Alberich-Carramiñana, J. Fernández-Sánchez, *On adjacent complete ideals above a given complete ideal*, *Proc. Royal Soc. Edinburgh*, 140 A: 225–238, 2010. [9](#)
- [5] M. Alberich-Carramiñana, J. Álvarez Montaner and F. Dachs-Cadefau, *Multiplier ideals in two-dimensional local rings with rational singularities*, *Mich. Math. J.*, 65: 287–320, 2016. [2](#), [6](#), [18](#)

- [6] C. Berkersch and A. Leykin, *Algorithms for Bernstein-Sato polynomials and multiplier ideals.*, ISSAC 2010—Proceedings of the 2010 International Symposium on Symbolic and Algebraic Computation, 99–106, 2010. [17](#)
- [7] W. Bosma, J. Cannon and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput., 24: 235–265, 1997. [3](#)
- [8] E. Casas-Alvero, *Filtrations by complete ideals and applications*, Collect. Math. 49: 265–272, 1998. [3](#)
- [9] E. Casas-Alvero, *Singularities of Plane Curves*. Number 276 in London Math. Soc. Lecture Note Series. Cambridge University Press, Cambridge, UK, first edition, 2000. [2](#), [4](#), [5](#), [6](#), [7](#), [8](#), [9](#), [14](#), [15](#), [17](#)
- [10] W. Decker, G.-M. Greuel, G. Pfister and H. Schönemann, SINGULAR – A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de>. [2](#)
- [11] T. de Jong, *An algorithm for computing the integral closure*, J. Symb. Comput., 26(3): 273–277, 1998. [2](#)
- [12] F. Delgado, C. Galindo and A. Núñez, *Generating sequences and Poincaré series for a finite set of plane divisorial valuations*, Adv. Math., 219(5): 1632–1655, 2008.
- [13] F. Enriques and O. Chisini, *Lezioni sulla teoria geometrica delle equazioni e delle funzioni algebriche*, N. Zanichelli, Bologna, (1915). [6](#)
- [14] J. Fernández-Sánchez, *On sandwiched singularities and complete ideals*, J. Pure Appl. Algebra, 185: 165–175, 2003. [9](#)
- [15] J. Fernández-Sánchez, *Nash families of smooth arcs on a sandwiched singularity*, Math. Proc. Camb. Phil. Soc., 138: 117–128, 2005. [9](#)
- [16] J. Fernández-Sánchez, *Some results relating to adjacent ideals in dimension two*, J. Pure Appl. Algebra, 207: 387–395, 2006. [9](#)
- [17] D. Grayson and M. Stillman, *Macaulay2*. Available at: <http://www.math.uiuc.edu/Macaulay2>. [2](#)
- [18] J. A. Howald, *Multiplier ideals of monomial ideals*, Transactions of the American Mathematical Society, 353(7), 2665–2671, 2001. [3](#), [18](#)
- [19] T. Järvilehto, *Jumping numbers of a simple complete ideal in a two-dimensional regular local ring*, Mem. Amer. Math. Soc. **214** (2011), no. 1009, viii+78 pp. [18](#)
- [20] R. Lazarsfeld, *Positivity in algebraic geometry II*, volume 49, (2004), Springer-Verlag, xviii+385. [17](#)
- [21] D. A. Leonard and R. Pellikaan, *Integral closures and weight functions over finite fields*, Finite Fields Appl., 9(4): 479–504, 2003. [2](#)
- [22] T. Moh and S. S. Abhyankar, *Newton-Puiseux expansion and generalized Tschirnhausen transformation. I*. J. Reine Angew. Math., 260: 47–83, 1973. [7](#)
- [23] T. Moh and S. S. Abhyankar, *Newton-Puiseux expansion and generalized Tschirnhausen transformation. II*. J. Reine Angew. Math., 261: 29–54, 1973. [7](#)
- [24] D. Naie, *Jumping numbers of a unibranch curve on a smooth surface*, Manuscripta Math. **128** (2009), 33–49. [18](#)
- [25] A. Płoski, *Remarque sur la multiplicité d’intersection des branches planes*. Bull. Pol. Acad. Sci. Math., 33: 601–605, 1985. [14](#)
- [26] T. Shibuata, *Algorithms for computing multiplier ideals.*, J. Pure Appl. Algebra, 215: 2829–2842, 2011. [17](#)
- [27] A. Singh and I. Swanson, *An algorithm for computing the integral closure*. Algebra Number Theory, 3(5): 587–595, 2009. [2](#)
- [28] B. Teissier *Appendix in [34]* [3](#), [19](#)
- [29] K. Tucker, *Jumping numbers on algebraic surfaces with rational singularities*, Trans. Amer. Math. Soc. **362** (2010), 3223–3241. [18](#)
- [30] W. V. Vasconcelos, *Computing the integral closure of an affine domain*, Proc. Amer. Math. Soc., 113(3): 633–638, 1991. [2](#)
- [31] W. V. Vasconcelos, *Divisorial extensions and the computation of integral closures*, J. Symbolic Comput., 30(5): 595–604, 2000. [2](#)
- [32] O. Zariski, *Polynomial ideals defined by infinitely near points*. Amer. J. Math., 60(1): 151–204, 1938. [1](#), [2](#), [6](#)
- [33] O. Zariski, *Studies in equisingularity III: saturation of local rings and equisingularity*, Amer. J. Math., 90(3): 961–1023, 1968. [7](#)

- [34] O. Zariski, *Le problème des modules pour les branches planes*. Hermann, Paris, 1986. [7](#), [21](#)

DEPARTAMENT DE MATEMÀTIQUES, UNIV. POLITÈCNICA DE CATALUNYA, AV. DIAGONAL 647,
BARCELONA 08028, SPAIN

E-mail address: Maria.Alberich@upc.edu, Josep.Alvarez@upc.edu, Guillem.Blanco@upc.edu