ORIGINAL ARTICLE

Response of the autonomic activity to stress provocation in females with cervicogenic headache compared to asymptomatic controls: a cross-sectional study

Sarah MINGELS ^{1, 2 *}, Marita GRANITZER ¹

¹REVAL Rehabilitation Research Center, Faculty of Rehabilitation Sciences, Hasselt University, Hasselt, Belgium; ²Musculoskeletal Research Unit, Department of Rehabilitation Sciences, Faculty of Kinesiology and Rehabilitation Sciences, Catholic University of Leuven, Leuven, Belgium

*Corresponding author: Sarah Mingels, Faculty of Rehabilitation Sciences, Hasselt University, Agoralaan, 3590, Diepenbeek, Belgium.
Email: sarah.mingels@uhasselt.be

ABSTRACT

BACKGROUND: Because abnormal activity of the autonomic nervous system is associated with chronification of pain, early detection of such dysfunction is important.

AIM: Although several studies highlight autonomic dysfunction in the chronification of headache, no study discussed its role in episodic cervicogenic headache.

DESIGN: Case-controlled cross-sectional single-blind comparative study between women with episodic cervicogenic headache and matched controls.

SETTING: Outpatient setting, Hasselt University.

POPULATION: Autonomic activity of 17 females with episodic cervicogenic headache (26.6±11.6 years) was compared with 17 age, gender and socio-economic matched asymptomatic controls (26.8±11.9 years).

METHODS: Autonomic activity was compared via repeated measures of the activity of the dermal sweat glands (μmho), peripheral circulation (%), electrical activity of the bilateral upper trapezius (μV) before, during and after cognitive stress provocation.

RESULTS: Whereas the autonomic parameters of the control group behaved as expected, participants in the headache group showed: 1) to stress provocation a significant lower dermal sweat gland activity (3.03±0.44 vs. 4.19±0.91 μmho, $P<0.0001$), higher vasodilatation (-5.56±1.45% vs. -5.61±1.85%, $P=0.03$), lower activity of the left upper trapezius (0.21±0.44 vs. 0.89±0.59 μV, $P=0.03$), significant less recuperation of the dermal sweat gland activity (-2.57±0.40 vs. -3.29±0.84 μmho, $P<0.0001$); 2) no recuperation of the activity (μV) of the left ($P=0.83$) and right ($P=0.99$) upper trapezius; 3) from stress provocation to recuperation a significant negative correlation ($\rho=0.69$, $P=0.04$) between dermal sweat gland and right upper trapezius activity.

CONCLUSIONS: Females with episodic cervicogenic headache reacted less to cognitive stress provocation. Recuperation after such provocation was absent. More research is needed to associate autonomous responses with a possible chronification process.

CLINICAL REHABILITATION IMPACT: A dysfunctional reaction to cognitive stress could be a threat to allostasis.

(Cite this article as: Mingels S, Granitzer M. Response of the autonomic activity to stress provocation in females with cervicogenic headache compared to asymptomatic controls: a cross-sectional study. Eur J Phys Rehabil Med 2020;56:175-83. DOI: 10.23736/S1973-9087.20.05836-0)

KEY WORDS: Post-traumatic headache; Autonomic nervous system; Physiological stress.

Headache ranks in the global top-ten of the most disabling conditions. The prevalence of current headache in adults is estimated to amount 50%. Among those individuals, 1.7% to 4% evolve into a chronic state.^{1, 2} Long-term coping with chronic headache (*i.e.* headache occurring on 15 or more days per month for at least 3 months) is a predisposition to develop other illnesses, and thereby

imposes a substantial burden on personal and social life.² Therefore, factors contributing to the chronification process should be early recognized.

Chronic pain is a multidimensional interaction between physiological, psychological, and social factors.³ Identifying those factors in an acute phase contributes to the prevention of chronification. Autonomic dysregulation for

instance can be such a chronification-promoting factor.⁴⁻⁶ Lack of physiological recuperation after stress, a typical feature of autonomic dysregulation, might be a risk factor that reduces the capacity to respond to new stressors.^{4, 5} In addition, an increased sympathetic activity facilitates the sensitization of headache by feeding the nociceptive circuits at the primary nociceptive source and corresponding dorsal horn neurons.^{7, 8} The development of cervicogenic headache (CeH), *i.e.* headache caused by a disorder of the cervical spine and its bony, disc and/or soft tissue elements, usually but not invariably accompanied by neck pain, is suggested to be associated with a spread of rostral neuraxial sensitization of the caudal trigemino-cervical nucleus.⁹⁻¹²

Since a dysfunction in the activity of the ANS might be a catalyst for the progression from episodic to chronic headache by sensitizing the caudal trigemino-cervical nucleus, its role should be further analysed in patients with CeH.^{13, 14} Although multiple pathways induce (central) sensitization, the specific objective of the current study is to explore if differences in autonomic responses can be detected between participants with episodic CeH and asymptomatic controls during cognitive stress provocation.

It is expected that a normal autonomic response is characterized by decreased sympathetic activity during relaxation, which increases during stress provocation and recovers after the omission of stress.^{5, 10, 15-17} Participants with episodic CeH are hypothesized to show a maladaptive autonomic response.^{4, 15, 17}

Materials and methods

The study is registered as an ‘Observational Study’ at ‘ClinicalTrials.gov (NCT02887638)’ on September 1st, 2016. The Medical Ethical Committee of the Oost-Limburg Hospital (Chairperson: dr. P. Noyens) granted approval on 21/05/2015 (B371201423025). All participants signed the written informed consent in which information was given concerning confidentiality of the data. The protection of personal data is legally determined by the Belgian law of December 8th, 1992 on the protection of privacy.

Design

A case-controlled cross-sectional single-blind design was used to compare autonomic responses before (relaxation phase), during (cognitive stress phase) and after (recuperation phase) a cognitive stress provocation between women with episodic CeH and matched asymptomatic controls (control group).

Participants

Sixty-one potential candidates for the episodic CeH group responded to a general call launched at the Hasselt University campus between October 2015 and January 2016. Inclusion criteria for the episodic CeH group were: females, Dutch-speaking, between 18 and 45 years, diagnosed with episodic CeH based on the International Classification of Headache Disorders 3, 2018 (Table I) by a neurologist.

Inclusion criteria for the Control group were: asymptomatic females, Dutch-speaking, between 18 and 45 years. Exclusion criteria for the episodic CeH and Control group were: smoking, pregnancy, physiotherapy for head- or neck-related disorders, comorbid pathology (neurological, cardiovascular, endocrine, musculoskeletal, autoimmune, psychiatric), medication use (neuroleptics, anti-epileptic’s, Ca²⁺-blockers, beta-blockers, anti-depressants, hormonal supplements), medication overuse (ergotamine, NSAID, opioid, acetylsalicylic acid, triptans, simple analgesics >10 days a month during >3 months), substance abuse, history of neck/head trauma, headaches with autonomic features (photo- or phonophobia, nausea).

Forty-four participants were excluded from the episodic CeH group due to: no CeH diagnosis (N.=27), male gender

TABLE I.—Diagnostic criteria of cervicogenic headache by the International Classification of Headache Disorders 3.

Description	Headache caused by a disorder of the cervical spine and its component bony, disc and/or soft tissue elements, usually but not invariably accompanied by neck pain
Diagnostic criteria	A. Any headache fulfilling criterion C B. Clinical and/or imaging evidence of a disorder or lesion within the cervical spine or soft tissues of the neck, known to be able to cause headache C. Evidence of causation demonstrated by at least two of the following: 1. headache has developed in temporal relation to the onset of the cervical disorder or appearance of the lesion 2. headache has significantly improved or resolved in parallel with improvement in or resolution of the cervical disorder or lesion 3. cervical range of motion is reduced and headache is made significantly worse by provocative maneuvers 4. headache is abolished following diagnostic blockade of a cervical structure or its nerve supply D. Not better accounted for by another ICHD-3 diagnosis
Provocation	Headache provoked by at least one of the following: 1. Posture (<i>e.g.</i> forward head posture) 2. Repetitive cervical movement
Autonomous	1. No nausea or vomiting 2. No photophobia or no phonophobia

(N.=7), trauma (N.=2), neurological comorbidity (N.=3), cardiovascular comorbidity (N.=3), metabolic comorbidity (N.=1), and pregnancy (N.=1). Eventually, 17 participants met the criteria for the episodic CeH group. Seventeen asymptomatic female participants were selected after a general call (Hasselt University campus) to compose the control group. Purposive sampling was applied to match the control group for age and socioeconomic status (level of education, job).

Outcomes, measurements and instruments

Primary outcome was the activity of the ANS. Autonomic activity was estimated by following the psychophysiological profile of: 1) the activity of the dermal sweat glands (skin conductance [SC], in μmho); 2) the peripheral circulation, characterized by vasoconstriction and vasodilatation (blood volume pulse [BVP], in %); 3) surface electrical activity of the left and right upper trapezius (surface electromyography [sEMG], in μV) evaluated with the Procomp+ and Biograph 2.1 software (Thought Technology Ltd, 2016, Montreal, Canada).¹⁸ sEMG-sensors (pre-amplified MyoScan-Pro™ Sensor, filter 0-400 μV) with AgCl-electrodes (Triode™, reference electrode included) were used. The Muscular Activity Index (MAI) was cal-

culated to estimate muscular activity (μV) during samples of the last 10 seconds in each phase (except in the stress phase data of 10 seconds peak stress were used). Electromyographically signals were automatically converted into root mean squares (RMS) (Table II). An asymmetry index was calculated to estimate a possible asymmetry between the left and right upper trapezius using the following formula: $\text{Asymmetry Index} = (\text{RMS}_{\text{right}} - \text{RMS}_{\text{left}}) / (\text{RMS}_{\text{right}} + \text{RMS}_{\text{left}}) \times 100$.¹⁹

Secondary outcomes, namely headache intensity (100-mm Visual Analogue Scale for pain intensity; VAS in mm per attack),²⁰ duration (hours per attack) and frequency (days per month), were extracted from the Belgian Headache Society diary which was completed by the episodic CeH group four weeks before the start of the study.

Test procedure

A VAS for headache intensity and stress of <30 mm on the test day were prerequisites to be tested. The intake of painkillers, alcohol, and caffeine (coffee, energy drinks, etc.) was prohibited 24 hours before testing. The experiment was carried out by a blinded physiotherapist in a quiet, temperature-controlled and dimly lit room at the Hasselt University (Belgium). The non-invasive psychophysiological testing consisted of four consecutive phases: baseline (1 min), relaxation phase (3 min), cognitive stress test phase (3 min) and recuperation phase (= relax after omission of the stressor) (3 min). To relax and recuperate abdominal breathing was taught by a physiotherapist before the start of the measurements (Supplementary Digital Material 1: Supplementary Text File 1). Sensors for the measurement of SC, BVP and sEMG were applied (Table II) while the participant was seated on an ergonomic chair (Gymna). The chair could be individually adjusted for comfort.²¹ Autonomic activity was evaluated via a baseline measurement in which no instructions were given, followed by a relaxation where abdominal breathing was performed. Next, stress was provoked using a cognitive stress test: the participant was instructed to count down from 500 to 0 in steps of 7 during 3 minutes.²² Hesitations or incorrect answers were penalized by restarting the test.²³ To recuperate 3 minutes of abdominal breathing was performed. In the episodic CeH group a 100-mm VAS for headache intensity was questioned before and after testing.

Statistical analysis

Analysis was done by using SAS JMP Pro 13 (Wittington House, Henley Road, Marlow, UK). Descriptive statistics were provided by presenting the absolute mean group val-

TABLE II.—Summary of the sensor and electrode placement and interpretation of the psychophysiological measurements

Measurement	Sensor and electrode placement - Interpretation
SC, μmho	<p><i>Sensor placement:</i> Two sensors attached to the distal phalanges of the ring and middle finger of the non-dominant hand</p> <p><i>Interpretation:</i> Evaluation of electrical SC depends on the dermal sweat gland activity. SNS activity increases conductance by elevating the dermal sweat gland activity</p>
BVP, %	<p><i>Sensor placement:</i> Sensor attached to the palmar thumb side of the non-dominant hand</p> <p><i>Interpretation:</i> The BVP-sensor evaluates peripheral circulation through photoplethysmography. The amount of reflected light is measured and presented as a bandwidth amplitude. Stress reduces the amplitude of the bandwidth through peripheral capillary constriction. An increased bandwidth indicates a vasodilatation of the peripheral capillaries</p>
sEMG, μV	<p><i>Electrode placement:</i> Triodes were placed parallel with the fibers of the upper trapezius, 10 mm lateral of the center of the line between the acromion and spinous process of C7 [17,21]</p> <p><i>Interpretation:</i> The sEMG measures the electrical activity of the muscle. During stress the sEMG-signal increases</p>
SNS: sympathetic nervous system; SC: skin conductance; BVP: blood volume pulse; sEMG: surface electromyography.

Figure 1.—Visualization of the mean (SEM) autonomic activity in the episodic CeH group and Control group from baseline to recuperation. Cg: control group.

ues of every phase (Figure 1) and the group characteristics (Table III).

Two- or one-tailed P values were reported with a 95% confidence level ($P < 0.05$). Equality of groups was tested by an unpaired t -test. For the primary outcome, activity

of the ANS, measures of the SC, BVP and sEMG were averaged from the different phases. Results were independent of the baseline (linear regression model). Mixed models with random and fixed effects were used to analyze the repeated measures of the mean SC, BVP and sEMG

TABLE III.—Summary of the mean group characteristics.

Characteristics	Episodic CeH (N.=17)	Control (N.=17)	P value
Mean age, years	26.6±11.6	26.8±11.9	0.19 †
Side dominance, right/left	16/1	15/2	1 ‡
Mean headache VAS score per attack, mm	40.2±22	NA	NA
Mean headache frequency, days/month	13.6±7.3	NA	NA
Mean headache duration, h per attack	6.8±3.2	NA	NA
SC, µmho (SE)	2.7 (0.4)	2.5 (0.4)	0.64
BVP, % (SE)	10.4 (1.7)	9.4 (2.4)	0.85
sEMG left, µV (SE)	2.3 (0.6)	1.8 (0.6)	0.40
sEMG right, µV (SE)	1.3 (0.3)	1.7 (0.8)	0.82

Results concerning the headache characteristics for the episodic CeH group were retrieved from the headache diary.

CeH: cervicogenic headache; SE: standard error; SC: skin conductance; BVP: blood volume pulse; sEMG: surface electromyography; VAS: 100-mm Visual Analogue Scale; NA: not applicable.

† Unpaired t -test; ‡ Fisher Exact test. Statistical significance is set at $P < 0.05$.

from baseline to recuperation within and between groups (Tukey-Kramer test). Fixed effects were the phases: baseline, relaxation, cognitive stress and recuperation. The group or the participants were random effects. Dependent variables were the mean SC, BVP and sEMG. Conditions to apply linear models were met (*i.e.* normal distribution (Shapiro-Wilk $P>0.05$, linearity, homoscedasticity [Levene's tests $P>0.05$] of the residuals).

Conditions to apply parametric statistics to assess correlations were violated. Therefore, Spearman's rho was used to estimate associations between autonomic parameters (SC, BVP, sEMG) and headache-characteristics (intensity, duration, frequency), and between autonomic parameters mutually.

The Minimal Detectable Change (MDC), which is the minimal amount of change a measurement must show to be larger than the within-subject variability and measurement error, was calculated based on the formula: $1.96 \times \sqrt{2} \times \text{SEM}$. Effect sizes (ES) (Cohen's d) were calculated in case of statistical significance ($P<0.05$) (0.2=small, 0.5=medium, 0.8=large ES). The sample size ($N=17$) (power of 80%; .05 probability of a type-I error) was estimated *a priori*.¹²

Results

Group characteristics

Autonomic activity at baseline did not differ between the episodic CeH group and Control group (Table III).

TABLE IV.—Comparisons of the autonomic activity from baseline to recuperation within the episodic CeH ($N=17$) and the control group ($N=17$).

Parameter		Baseline	Relaxation	P value [#]	Relaxation	Stress provocation	P value [‡]	Stress provocation	Recuperation	P value [†]
Skin conductance μmho (SE) (95% CI)	CeH	2.7 (0.4) (1.5-3.2)	1.4 (0.3) (0.7-2.1)	0.02*	1.4 (0.3) (0.7-2.1)	4.4 (0.7) (3-5.9)	<0.0001*	4.4 (0.7) (3-5.9)	1.9 (0.4) (1-2.8)	<0.0001*
	Control	2.5 (0.4) (1.6-3.5)	1.6 (0.4) (0.8-2.3)	0.71	1.6 (0.4) (0.8-2.3)	5.8 (1.1) (3.5-8)	<0.0001*	5.8 (1.1) (3.5-8)	2.5 (1.8) (1.6-3.4)	<0.0001*
Blood volume pulse, % (SE) (95% CI)	CeH	10.4 (1.7) (6.8-14)	12.3 (1.8) (8.5-16.2)	0.66	12.3 (1.8) (8.5-16.2)	6.8 (1.3) (4-9.6)	0.002*	6.8 (1.3) (4-9.6)	7.6 (1.3) (4.8-10.4)	0.96
	Control	9.4 (2.4) (4.2-14.5)	11.8 (2.5) (6.5-17.2)	0.41	11.8 (2.5) (6.5-17.2)	6.2 (1.5) (3.1-9.3)	0.0001*	6.2 (1.5) (3.1-9.3)	8.9 (1.9) (4.9-12.9)	0.34
sEMG left, μV (SE) (95% CI)	CeH	2.3 (0.6) (1.1-3.5)	2.2 (0.4) (1.3-3.1)	0.99	2.2 (0.4) (1.3-3.1)	2.4 (0.4) (1.5-3.3)	0.97	2.4 (0.4) (1.5-3.3)	2.8 (0.6) (1.5-4.1)	0.83
	Control	1.8 (0.6) (0.5-3.2)	1.7 (0.6) (0.5-2.9)	0.99	1.7 (0.6) (0.5-2.9)	2.6 (0.7) (1.1-4.1)	0.002*	2.6 (0.7) (1.1-4.1)	1.5 (0.4) (0.8-2.3)	0.003*
sEMG right, μV (SE) (95% CI)	CeH	1.3 (0.3) (0.5-2)	1.8 (0.4) (0.8-2.7)	0.88	1.8 (0.4) (0.8-2.7)	3.3 (1) (1.1-5.4)	0.03*	3.3 (1) (1.1-5.4)	3.1 (1.1) (0.9-5.3)	0.99
	Control	1.7 (0.8) (0-3.5)	1.8 (0.7) (0.3-3.4)	0.91	1.8 (0.7) (0.3-3.4)	4.1 (1) (2.1-6.2)	0.01*	4.1 (1) (2.1-6.2)	1.9 (0.4) (0.9-2.8)	0.02*

CeH: cervicogenic headache; SE: standard error.

[#] P value (mixed model with random and fixed effects) from baseline to relaxation; [‡] P value (mixed model with random and fixed effects) from relaxation to stress provocation; [†] P value (mixed model with random and fixed effects) from stress provocation to recuperation; *statistically significant change ($P<0.05$).

Descriptive statistics

Figure 1 illustrates the mean profiles of the autonomic parameters during the different phases for both groups. Deviating patterns were observed for the sEMG of the upper trapezius in the episodic CeH group: sEMG right did not respond to relaxation and sEMG left did not recover during the recuperation phase after stress provocation.

Primary outcomes

Responses from baseline to relaxation

SC within the episodic CeH group decreased significantly ($P=0.02$). This drop was larger than the MDC (0.38) (Table IV). The SC within the Control group did not change significantly. In addition, changes in BVP and sEMG, observed within both groups, were not significant (Table IV). Changes in SC, BVP and sEMG differed not significantly between groups (Table V).

Responses from relaxation to stress provocation

Within both the episodic CeH group and Control group SC increased significantly ($P<0.0001$) and BVP decreased significantly ($P<0.01$) (Table IV). Changes for the SC and BVP were larger than the MDCs (CeH: 1.22 and 4.02, respectively; control 2.52 and 5.13, respectively) (Table IV). Response of the sEMG right increased significantly ($P=0.03$) within the episodic CeH group, whereas sEMG left increased significantly ($P=0.002$) within the Control group (Table IV).

TABLE V.—Comparisons of the changes in autonomic activity from baseline to recuperation between the episodic CeH (N.=17) and the control group (N.=17).

Parameter	Baseline to relaxation (Δ)			Relaxation to stress (Δ)			Stress to recuperation (Δ)		
	CeH	Control	P value	CeH	Control	P value (ES)	CeH	Control	P value (ES)
Skin conductance, μmho (SE)	-0.96 (0.14)	-0.96 (0.04)	0.32	3.03 (0.44)	4.19 (0.91)	<0.0001	-2.57 (0.4)	-3.29 (0.84)	<0.0001
MDC	0.38	0.11		1.22	2.52	(1.56)*	1.11	2.33	(1.26)*
Blood volume pulse, % (SE)	1.92 (1.79)	2.46 (1.24)	0.63	-5.56 (1.45)	-5.61 (1.85)	0.003	0.84 (1.84)	2.68 (1.2)	0.77
MDC	4.96	3.43		4.02	5.13	(0.73)*	5.1	3.33	
sEMG left, μV (SE)	0.13 (0.06)	-0.13 (0.14)	0.99	.21 (0.44)	.89 (0.59)	0.03	0.43 (0.41)	-1.01 (0.62)	0.56
MDC	0.16	0.38		1.22	1.64	(0.54)*	1.14	1.72	
sEMG right, μV (SE)	0.49 (0.05)	0.11 (0.17)	0.98	1.5 (0.61)	2.31 (0.88)	0.07	-0.14 (0.51)	-2.28 (0.96)	0.39
MDC	0.14	0.47		1.7	2.44		1.41	2.66	

Negative values are indicative for a decrease, positive values for an increase.

CeH: cervicogenic headache; SE: standard error; MDC: minimal detectable change; Δ : differences between phases; P value: mixed model with random and fixed effects; ES: effect size (Cohen's d).

*Statistically significant change ($P<0.05$).

The episodic CeH group differed from the Control group by showing a significantly lower SC ($P<0.0001$), higher BVP ($P=0.02$) and lower sEMG left ($P=0.03$) (Table V).

Responses from stress provocation to recuperation

The episodic CeH and Control group responded with a significant drop in SC ($P<0.0001$) after stress provocation. The observed decrease of the SC in both groups was larger than the MDC (episodic CeH: -2.57 vs. 1.11; Control: -3.29 vs. 2.33) (Table IV). Whereas sEMG left and right decreased significantly ($P=0.003$ respectively $P=0.02$) in the Control group, sEMG left ($P=0.83$) and right ($P=0.20$) did not change significantly in the episodic CeH group (Table IV). The SC recuperated significantly ($P<0.0001$) less in the episodic CeH group compared to the Control group. No significant between-group differences can be presented for BVP ($P=0.77$), sEMG left ($P=0.56$) and sEMG right ($P=0.39$) (Table V).

Correlations

Positive correlations between sEMG left and right during relaxation ($\rho=0.59$, $P=0.01$) and stress ($\rho=0.69$, $P=0.04$) were seen in the episodic CeH group. Contrarily, a negative correlation between SC and sEMG right ($\rho=-0.51$) ($P=0.03$) was calculated during recuperation.

Secondary outcomes

Pain

The mean 100-mm VAS for headache intensity (26 mm) before the psychophysiological measurement did not differ significantly from the mean VAS for headache intensity (30 mm) after the measurements in the episodic CeH group.

Discussion

Although several studies highlight an involvement of autonomic dysfunction in the chronification of headache, no study discussed its role in episodic CeH.^{6, 24} Whereas the autonomic parameters of our Control group responded as expected, participants in the episodic CeH group showed: 1) to relaxation no response of the right upper trapezius; 2) to stress provocation a significant lower dermal sweat gland activity, higher vasodilatation, and lower activity of the left upper trapezius; 3) significant less recuperation of the dermal sweat gland activity and no recuperation of the activity of the left and right upper trapezius after stress provocation; 4) from stress provocation to recuperation a significant negative correlation between dermal sweat gland activity and activity of the right upper trapezius.

Dermal sweat gland activity

Cutaneous sweat glands are exclusively innervated by cholinergic nerves of the sympathetic nervous system (SNS). Electrodermal activity therefore reflects the SNS's arousal.²⁵ Orthostatic, physical and cognitive stressors are used to assess such arousal.²⁶ An increased sympathetic response to cognitive stress might contribute to an increased sensitivity to painful stimuli.²⁷ However, such findings are in contrast with the significant lower electrodermal reactivity in our episodic CeH group during the stress response, which could indicate a lower sympathetic activity.^{5, 6} The dampened electrodermal reactivity might reflect a depletion of the activity of the SNS. Sympathetic hypofunction has been mooted as a trait of depression, and as a potential patho-mechanism leading to higher risks for negative health outcomes.²⁸

In healthy young controls (mean 23.3 years) electrodermal activity recovers after a cognitive (arithmetic) test.²² Similar results were found in our Control group (26.8 years). However, in the episodic CeH group, dermal sweat gland activity recovered significantly less. A delayed inhibitory process might be associated with such results.²² In addition, a persistent augmented activity of the SNS facilitates sensitization processes by decreasing sensory and nociceptive thresholds, and increasing muscular activity.⁸ The latter was observed in the episodic CeH group.

The negative correlation between the SC and sEMG right during recuperation in the episodic CeH group, opposing the normal expectation, might be an indication of a dissociated pattern of sympathetic activity.^{29, 30}

Peripheral vascular response

The peripheral vascular response to stress provocation, *i.e.* vasoconstriction, was significantly lower in the episodic CeH group compared to the Control group. Analogously, previous analysis of Mayer (vasomotor) waves in patients with chronic tension-type headache revealed a sympathetic down-regulation. Such reaction was hypothesized to be the result of an habituation process.³¹ The ability to habituate to nociceptive stimuli is a trait of chronic headache.³² Yet, in our study participants in the episodic CeH group responded to a lesser extent to stress provocation. It could be hypothesized that these participants are already evolving into a chronic headache state. However, results from the current study cannot accept nor refute such hypothesis.

Muscular activity

Participants in the episodic CeH group seem to have difficulties to relax the bilateral upper trapezius after stress provocation: sEMG of the left upper trapezius was 27%, and sEMG of the right upper trapezius 58% higher after recuperation compared to at the end of the relaxation (Figure 1). Based on the asymmetry index, activity of the right upper trapezius was 36% higher compared to the left after recuperation. Such lacking muscular recuperation might be indicative for a dysregulated ANS.^{15, 17} Sustained sEMG-activity is suggested to be the result of a defective central inhibition, in which low threshold motor units remain activated. This fits the concept of “sustained arousal.”^{33, 34} The observed left-right difference in sEMG activity is hypothesized to be related to two mechanisms. Firstly, hand-dominance might support such difference since dominant upper trapezius muscles demonstrate higher amplitudes of activity and less muscular rest compared to non-dominant

muscles.³⁵ Secondly, pain-related increased activation of local muscle sympathetic nerve activity might reinforce muscular activity.³⁶ The higher sEMG activity of the right upper trapezius might be explained by the fact that most participants in the episodic CeH group were right-handed (94%) and suffered from right-sided CeH (right 53% vs. left 23.5%).

However, stress provocation could not provoke headache in the episodic CeH group. Bansevicius *et al.* concluded that the mean level of phasic sEMG responses during stressful conditions are of little importance to develop muscular pain.³⁷ It can nonetheless be argued that other physiological responses, such as prolonged activity of low-threshold motor units, might be important for such development. Since chronic pain conditions are caused by multimodal interactions, muscular responses might only partially contribute to the development of chronic pain.^{17, 38}

Pain provocation

Cathcart *et al.* supported the hypothesis in which mental stress contributes to chronic headache through sensitization of nociceptive afferents.³⁹ The cognitive stressor in our study did not provoke instant headache. The duration of the stressor could have been too short, or a delayed headache might have developed.

Limitations of the study

The authors acknowledge that case-controlled studies and purposive sampling are sensitive to selection bias. The generalization of the results is therefore limited to females with episodic CeH between 18 and 45 years.^{40, 41} More research is needed to provide normative data, and determine clinically significant changes (minimal clinically important difference, MCID). No causative statements can be made based on the current cross-sectional study. Longitudinal designs (follow-up) are needed to determine if an autonomic dysregulation can contribute to the chronification of episodic CeH. Further, a 24-hour VAS should be questioned.

This study used a cognitive stress provocation with a limited recuperation phase (3 minutes). More research is needed to assess the contribution of an incomplete recovery to the chronification of pain.³⁷ Based on the current study we advise to measure the time needed to recuperate completely (=baseline) since the prolonged activity of low-threshold motor units might be important for the development of pain originating from a muscle.³⁷ In addition, although outside the scope of the current study, we

found a dependency (linear regression model) between the magnitude of the SC, BVP, sEMG left and right in the recuperation phase (dependent variable) and the stress phase (independent variable); meaning that higher responses to stress were predictive for a stronger recuperation (Supplementary Digital Material 2: Supplementary Table I).

Future research should question psychosocial (anxiety, depression, cognition, personality) and lifestyle factors (sleep quality, physical activity) to accentuate the biopsychosocial character of stress.

Finally, no Bonferroni corrections were applied given the explorative nature of the study. The authors realize the consequence (Type I (α) error).

Conclusions

The Control group responded as was expected from previous studies. However, females with episodic CeH reacted with a lower activity of the dermal sweat glands, a more relaxed peripheral circulation and less activity of the left upper trapezius to cognitive stress provocation. Recuperation after such provocation was contrarily absent.

References

- Blume HK. Childhood Headache: A Brief Review. *Pediatr Ann* 2017;46:e155–65.
- World Health Organization. Lifting The Burden. Atlas of headache disorders and resources in the world; 2011 [Internet]. Available from: www.who.int/mental_health/management/atlas_headache_disorders/en/ [cited 2020, Jan 22].
- Jensen MP, Turner JA, Romano JM. Changes after multidisciplinary pain treatment in patient beliefs and coping are associated with concurrent changes in patient functioning. *Pain* 2007;131:38–47.
- Koenig J, Williams DP, Kemp AH, Thayer JF. Vagally mediated heart rate variability in headache patients—a systematic review and meta-analysis. *Cephalalgia* 2016;36:265–78.
- Leistad RB, Nilsen KB, Stovner LJ, Westgaard RH, Rø M, Sand T. Similarities in stress physiology among patients with chronic pain and headache disorders: evidence for a common pathophysiological mechanism? *J Headache Pain* 2008;9:165–75.
- Gass JJ, Glaros AG. Autonomic dysregulation in headache patients. *Appl Psychophysiol Biofeedback* 2013;38:257–63.
- Cortelli P, Pierangeli G. Chronic pain-autonomic interactions. *Neurol Sci* 2003;24(Suppl 2):S68–70.
- Drummond PD. Sensory disturbances in complex regional pain syndrome: clinical observations, autonomic interactions, and possible mechanisms. *Pain Med* 2010;11:1257–66.
- Castien R, De Hertogh W. A Neuroscience Perspective of Physical Treatment of Headache and Neck Pain. *Front Neurol* 2019;10:276.
- Bogduk N. The neck and headaches. *Neurol Clin* 2014;32:471–87.
- Christensen MB, Bendtsen L, Ashina M, Jensen R. Experimental induction of muscle tenderness and headache in tension-type headache patients. *Cephalalgia* 2005;25:1061–7.
- Chua NH, van Suijlekom HA, Vissers KC, Arendt-Nielsen L, Wilder-Smith OH. Differences in sensory processing between chronic cervical zygapophyseal joint pain patients with and without cervicogenic headache. *Cephalalgia* 2011;31:953–63.
- Fishbain DA, Cutler R, Cole B, Rosomoff HL, Rosomoff RS. International Headache Society headache diagnostic patterns in pain facility patients. *Clin J Pain* 2001;17:78–93.
- Jacobs SC, Friedman R, Parker JD, Tofler GH, Jimenez AH, Muller JE, *et al.* Use of skin conductance changes during mental stress testing as an index of autonomic arousal in cardiovascular research. *Am Heart J* 1994;128:1170–7.
- Bansevicius D, Westgaard RH, Sjaastad OM. Tension-type headache: pain, fatigue, tension, and EMG responses to mental activation. *Headache* 1999;39:417–25.
- DiPietro JA, Costigan KA, Gurewitsch ED. Fetal response to induced maternal stress. *Early Hum Dev* 2003;74:125–38.
- Leistad RB, Sand T, Westgaard RH, Nilsen KB, Stovner LJ. Stress-induced pain and muscle activity in patients with migraine and tension-type headache. *Cephalalgia* 2006;26:64–73.
- Thought Technology Ltd. FAQ - ProComp/FlexComp Infinity; 2016 [Internet]. Available from: http://thoughttechnology.com/index.php/faq_profex/ [cited 2020, Jan 22].
- Wieczorek A, Loster JE. Activity of the masticatory muscles and occlusal contacts in young adults with and without orthodontic treatment. *BMC Oral Health* 2015;15:116.
- Karcioglu O, Topacoglu H, Dikme O, Dikme O. A systematic review of the pain scales in adults: which to use? *Am J Emerg Med* 2018;36:707–14.
- Vincent MB. Cervicogenic headache: a review comparison with migraine, tension-type headache, and whiplash. *Curr Pain Headache Rep* 2010;14:238–43.
- Visnovcova Z, Mestanik M, Gala M, Mestanikova A, Tonhajzerova I. The complexity of electrodermal activity is altered in mental cognitive stressors. *Comput Biol Med* 2016;79:123–9.
- Hassinger HJ, Semenchuk EM, O'Brien WH. Cardiovascular responses to pain and stress in migraine. *Headache* 1999;39:605–15.
- Evers S, Voss H, Bauer B, Sörös P, Husstedt IW. Peripheral autonomic potentials in primary headache and drug-induced headache. *Cephalalgia* 1998;18:216–21.
- Critchley HD. Electrodermal responses: what happens in the brain. *Neuroscientist* 2002;8:132–42.
- Posada-Quintero HF, Florian JP, Orjuela-Cañón AD, Aljama-Corrales T, Charleston-Villalobos S, Chon KH. Power spectral density analysis of electrodermal activity for sympathetic function assessment. *Ann Biomed Eng* 2016;44:3124–35.
- Crofford LJ. Chronic Pain: Where the Body Meets the Brain. *Trans Am Clin Climatol Assoc* 2015;126:167–83.
- Mestanikova A, Ondrejka I, Mestanik M, Hrtanek I, Snircova E, Tonhajzerova I. Electrodermal activity in adolescent depression. *Adv Exp Med Biol* 2016;935:83–8.
- Vissing SF. Differential activation of sympathetic discharge to skin and skeletal muscle in humans. *Acta Physiol Scand Suppl* 1997;639:1–32.
- Ruiz-Robledillo N, Moya-Albiol L. Lower electrodermal activity to acute stress in caregivers of people with autism spectrum disorder: an adaptive habituation to stress. *J Autism Dev Disord* 2015;45:576–88.
- Sliwka U, Harscher S, Diehl RR, van Schayck R, Niesen WD, Weiller C. Spontaneous oscillations in cerebral blood flow velocity give evidence of different autonomic dysfunctions in various types of headache. *Headache* 2001;41:157–63.
- Ozkul Y, Ay H. Habituation of sympathetic skin response in migraine and tension type headache. *Auton Neurosci* 2007;134:81–4.
- Lundberg U. [Psychological stress and musculoskeletal disorders: psychobiological mechanisms. Lack of rest and recovery greater problem than workload]. *Lakartidningen* 2003;100:1892–5. Swedish.

34. Logan JG, Barksdale DJ. Allostasis and allostatic load: expanding the discourse on stress and cardiovascular disease. *J Clin Nurs* 2008;17(7B):201–8.
35. Wixted F, O’Riordan C, O’Sullivan L. Inhibiting the Physiological Stress Effects of a Sustained Attention Task on Shoulder Muscle Activity. *Int J Environ Res Public Health* 2018;15:E115.
36. Burton AR, Birzniece I, Bolton PS, Henderson LA, Macefield VG. Effects of deep and superficial experimentally induced acute pain on muscle sympathetic nerve activity in human subjects. *J Physiol* 2009;587:183–93.
37. Lovallo WR. Can exaggerated stress reactivity and prolonged recovery predict negative health outcomes? The case of cardiovascular disease. *Psychosom Med* 2015;77:212–4.
38. McEwen BS. Protective and damaging effects of stress mediators. *N Engl J Med* 1998;338:171–9.
39. Cathcart S, Winefield AH, Lushington K, Rolan P. Effect of mental stress on cold pain in chronic tension-type headache sufferers. *J Headache Pain* 2009;10:367–73.
40. Pothineni NV, Shirazi LF, Mehta JL. Gender Differences in Autonomic Control of the Cardiovascular System. *Curr Pharm Des* 2016;22:3829–34.
41. Russo MA, Santarelli DM, O’Rourke D. The physiological effects of slow breathing in the healthy human. *Breathe (Sheff)* 2017;13:298–309.

Conflicts of interest.—The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

Authors’ contributions.—Sarah Mingels drafted the manuscript. Marita Granitzer provided essential comments on the structure, spelling and content of the manuscript. Both authors read and approved the final manuscript.

Acknowledgements.—The authors acknowledge Heleen Debrabandere and Shilly van Gelder for providing assistance in the data collection.

History.—Article first published online: January 14, 2020. - Manuscript accepted: January 7, 2020. - Manuscript revised: November 29, 2019. - Manuscript received: May 4, 2019.

Supplementary data.—For supplementary materials, please see the HTML version of this article at www.minervamedica.it