UNDER PRESSURE

Effects of standardisation of compression therapy in venous leg ulcers in community nursing

UNDER PRESSURE

Catholic University College Ghent

Campus Dirk Martens

Research unit Nursing

Kwalestraat 154

B-9320 Aalst

Belgium

www.kahosl.be

Hilde Boucqué

MSN, RN

Course coordinator nursing KAHO

dr. Jos Van der Veken

MD, PhD

Management staff member
Department of Educational
Affairs Ghent University

Caroline Goeman

MSN, RN

Teaching staff and researcher in nursing KAHO

Contact: caroline.goeman@kahosl.be



BACKGROUND

Venous ulcers are the most common chronic wounds in the aging population. Compression therapy forms the cornerstone of venous leg ulcer treatment. There is strong evidence that it enhances leg ulcer healing rates and prevents recurrence.

Nevertheless a discrepancy between evidence and practice exists. Literature reveals a lack of knowledge and skills of nurses regarding compression therapy as one of the key reasons.





AIM

To examine if a standard education package for community nursing regarding venous ulcer treatment, will lead to a more adequate subbandage pressure, a better performance of the ulcer, a pain reduction and a longer wearing of compression bandages.

METHOD

Based on literature and international guidelines an evidence-based education package was developed. Experts in wound care, doctors and community nurses were contacted to validate the education package trough a Delphi-procedure.

The training focuses both on knowledge and skills related to compression therapy. A group of community nurses will be trained using the standard education package. An other group of home care nurses will receive no additional education. The subbandage pressure, wound evolution, pain and compression wearing time of venous ulcer patients will be compared between both groups.

RELEVANCE

Enhancing the knowledge and skills of nurses may result in a better quality of life of patients with a venous ulcer and may reduce the costs of long-term health care treatment.



