

Safety of Everolimus Plus Exemestane In Patients With Hormone-Receptor-Positive, HER2-Negative Locally Advanced or Metastatic Breast Cancer Progressing on Prior Non-Steroidal Aromatase Inhibitors: Primary Results of a Phase 3b, Open-Label, Single-Arm, Expanded-Access Multicenter Trial (BALLET)

G. Jerusalem¹, G. Mariani², E. M. Ciruelos³, M. Martin⁴, V. C. G. Tjan-Heijnen⁵, P. Neven⁶, J. G. Gavila⁷, A. Michelotti⁸, F. Montemurro⁹, D. Generali¹⁰, E. Simoncini¹¹, I. Lang¹², J. Mardiak¹³, B. Naume^{14,15}, M. Camozzi¹⁶, K. Lorizzo¹⁶, S. Bianchetti¹⁶, P. Conte^{17,18}

¹CHU Sart Tilman Liege and Liege University, Department of Medical Oncology, Domaine Universitaire du Sart Tilman, Liege, Belgium

²Fondazione IRCCS Istituto Nazionale Dei Tumori, Milan, Italy

³Medical Oncology Department, Breast Cancer Unit, University Hospital 12 de Octubre, Madrid, Spain

⁴Instituto de Investigación Sanitaria Gregorio Marañón, Univesidad Complutense, Madrid, Spain

⁵Maastricht University Medical Centre, Department of Medical Oncology, GROW, Maastricht, The Netherlands

⁶KULeuven (University of Leuven), Department of Oncology, Multidisciplinary Breast Center, University Hospitals Leuven, Belgium

⁷Medical Oncology Unit of Fundacion Instituto Valenciano De Oncologia, Valencia, Spain

⁸UO Oncologia Medica I, Azienda Ospedaliera Universitaria Pisana, Santa Chiara Hospital, Pisa, Italy

⁹Unit of Investigative Clinical Oncology (INCO), Fondazione del Piemonte per l'Oncologia, Institute of Candiolo Cancer Center (IRCCs), Candiolo (Torino), Italy

¹⁰Department of Medical, Surgery and Health Sciences, University of Trieste, Piazza Ospitale 1, 34129 Trieste, Italy

¹¹Breast Unit, Azienda Ospedaliera Spedali Civili, Brescia, Italy

¹²National Institute of Oncology, Medical Oncology and Clinical Pharmacology, Budapest, Hungary

¹³Narodny Onkologicky Ustav Klenova 1, Bratislava, Slovakia

¹⁴Department of Oncology, Oslo University Hospital, Oslo, Norway

¹⁵K.G. Jebsen Center for Breast Cancer Research, Institute for Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway

¹⁶Novartis Farma S.p.A., Origgio/VA, Italy

¹⁷Department of Surgery, Oncology and Gastroenterology, University of Padova, Padova, Italy

¹⁸Medical Oncology 2, Istituto Oncologico Veneto, Padova, Italy

Correspondence to: Prof. Guy Jerusalem, CHU Sart Tilman Liege and Liege University, Department of Medical Oncology, Domaine Universitaire du Sart Tilman, Liege, Belgium, Phone: 003243668414; Fax: 003243667688, E-mail: g.jerusalem@chu.ulg.ac.be

Part of these data has been presented previously:

- Presented as a poster at the 2014 San Antonio Breast Cancer Conference; December 9-13, 2014; San Antonio, Texas, USA
- Presented as a poster at the 2015 European Society for Medical Oncology; September 25-29, 2015; Vienna, Austria

Key Message: "This is the largest ever reported safety dataset on a general patient population presenting ABC treated with EVE plus EXE and included a sizeable elderly subset. This is also the first trial reporting the impact of BMI on safety. These data reinforce observations from the BOLERO-2 trial that the safety profile of everolimus plus exemestane was manageable; there were no new safety signals."

ABSTRACT

Background: This European phase 3b, expanded-access multicenter trial evaluated the safety of EVE plus EXE in a patient population similar to BOLERO-2.

Patients and methods: Postmenopausal women aged ≥ 18 years with hormone receptor-positive, human epidermal growth factor-receptor-2–negative advanced breast cancer (ABC) recurring/progressing during/after prior non-steroidal aromatase inhibitors (NSAIs) were enrolled. Primary objective was safety of EVE plus EXE based on frequency of adverse events (AEs), and serious AEs (SAEs). Secondary objective was to evaluate AEs of grade 3/4 severity.

Results: Median treatment duration was 5.1 months (95% CI, 4.8-5.6) for EVE and 5.3 months (95% CI, 4.8-5.6) for EXE. Overall, 2131 patients were included in the analysis; 81.8% of patients experienced EVE- or EXE-related or EVE/EXE-related AEs (investigator assessed); 27.2% were of grade 3/4 severity. The most frequently reported non-hematologic AEs were (overall %, % EVE-related) stomatitis (52.8%; 50.8%) and asthenia (22.8%; 14.6%). The most frequently reported hematologic AE were (overall %, % EVE-related) anemia (14.4%; 8.1%) and thrombocytopenia (5.9%; 4.6%). AE-related treatment discontinuations were higher in elderly (≥ 70 years) versus non-elderly patients (23.8% vs. 13.0%). The incidence of EVE-related AEs in both elderly and non-elderly patients appeared to be lower in first-line ABC versus later lines. The incidence of AEs (including stomatitis/pneumonitis) was independent of BMI status (post-hoc analysis). Overall, 8.5% of patients experienced at least one EVE-related SAE. Of the 121 on-treatment deaths (5.7%), 66 (3.1%) deaths were due to disease progression and 46 (2.2%) due to AEs; 4 deaths were suspected to be EVE-related.

Conclusions: This is the largest ever reported safety dataset on a general patient population presenting ABC treated with EVE plus EXE and included a sizeable elderly subset. Although the patients were more heavily pretreated, the safety profile of EVE plus EXE in BALLET was consistent with BOLERO-2.

Clinical trial registration: EudraCT Number: 2012-000073-23

Key words: advanced breast cancer, BMI, elderly, everolimus, hormone-receptor positive, stomatitis

Introduction

Endocrine therapy (ET) is the treatment of choice for patients with HR+, human epidermal growth factor receptor 2-negative (HER2-) advanced breast cancer (ABC) in both adjuvant and advanced settings [1-5]. However, despite the effectiveness of ET many women experience disease progression, either de novo or acquired [6]. Hence, identification of effective targeted therapies, which may enhance or prolong endocrine sensitivity in these patients continues to be of clinical importance. Two different targeted agents, everolimus (a mammalian target of rapamycin [mTOR] inhibitor) and palbociclib (a cyclin-dependent kinase 4/6 inhibitor) have each shown efficacy in this patient population [7, 8].

Extensive cross-talk between the phosphatidylinositol 3-kinase (PI3K)/protein kinase B (AKT)/mTOR pathway and estrogen receptor signaling has been implicated as one of the key mechanisms of endocrine resistance [9-11]. Preclinical studies have shown that co-targeting both signaling pathways can synergistically inhibit tumor progression [9, 13, 14]. The pivotal BOLERO-2 trial showed that dual-blockade with everolimus (EVE), an mTOR inhibitor, plus EXE more than doubled median progression free survival (PFS) vs EXE alone in patients with HR+, HER2- ABC recurring/progressing on prior NSAIs (7.8 months vs 3.2 months) [14]. The present European phase 3b, expanded-access multicenter trial, BALLET, further evaluated the safety of EVE plus EXE in patients with HR+, HER2- ABC recurring/progressing on prior NSAIs.

Methods

Patients

Postmenopausal women aged ≥ 18 years with histologically/cytologically confirmed HR+ ABC unamenable to curative resection or radiotherapy, recurring or progressing on prior NSAIs were enrolled. NSAIs were not necessarily the last treatment prior to enrollment. There was no restriction on the number of prior lines of chemotherapy. Patients were excluded if they were HER2+ by local testing, were hypersensitive to mTOR inhibitors or EXE, had received radiotherapy within four weeks prior to enrollment, had symptomatic visceral disease, brain or central nervous system metastasis, or had

inadequate liver, renal, cardiac or bone marrow functions. Hormone replacement therapy had to be discontinued prior to enrollment.

Written informed consent was obtained from all patients. The study was performed in accordance with the Good Clinical Practice guidelines and the Declaration of Helsinki. The study protocol was approved by an independent ethics committee or institutional review board at each site. Dose modifications for specific treatment-emergent toxicities were advised in the protocol.

Study design and treatment

BALLET was a European, open-label, single-arm, multicenter phase 3b, expanded access trial (EudraCT Number: 2012-000073-23). Enrolled patients self-administered EVE on Day-1 and continued daily doses of EVE (either 2x5 mg or 1x10 mg) plus EXE (25 mg/day) in 28-day cycles. Study treatment continued until disease progression, intolerable toxicity, discontinuation from study due to other reasons, local reimbursement of EVE, or death. All patients also received best supportive care for pre-existing medical conditions or adverse events (AEs), as per standard local practice. Dose adjustments were allowed for AEs that were suspected to be related to EVE (additional information in Supplementary material). Permissible dose adjustments for EVE were 5 mg daily; 5 mg every other day. Relative dose intensity was defined as dose intensity (dosing unit/unit of time)/planned dose intensity. Permanent discontinuation involved discontinuation of either EVE or both EVE and EXE. Treatment once interrupted due to unacceptable toxicity was resumed only after recovery to grade ≤ 1 was achieved; reintroduction was at the initial/lower dose level according to study protocol. Patients were withdrawn from the study if the interruption was >28 days.

Safety assessment

The primary objective was to assess the safety of EVE plus EXE based on the frequency of AEs, serious AEs (SAEs) and number of laboratory abnormalities. The secondary objective was to evaluate AEs of grade 3 and 4 severity in routine clinical practice. Exploratory analysis included safety assessments in a subset of patients aged ≥ 70 years. Post hoc exploratory analyses evaluated the impact of body mass index (BMI) on the safety profile of EVE plus EXE.

Safety assessments included recording of all AEs and SAEs with their severity and relationship to study treatments, and deaths which occurred throughout the study and up to 28 days after the last treatment. The severity of AEs was graded according to National Cancer Institute Common Terminology Criteria for AEs v4.03 or on a scale of grade 1–4. Eastern Cooperative Oncology Group Performance Status (ECOG-PS) was assessed at baseline and at each study visit. Patients who dropped out of the study but continued on EVE through reimbursement, were not followed up after their exit.

Statistical analysis

The study did not formally test any hypotheses; all analyses were descriptive. The safety analysis included all patients who received at least one dose of EVE or EXE and were evaluable at least one time point. No formal sample size calculation was performed; the estimated sample size of 2500 patients was chosen based on the expected average accrual rate and duration of the trial based on the expected date of reimbursement in each participating country. Kaplan-Meier method was used for the exploratory assessment of the median treatment durations in the full population and when the study treatment was administered in the first-line setting, after censoring patients who discontinued treatment due to reimbursement/switch to other EVE programs.

Results

Demographics

Between 16-May-2012 and 31-December-2013, 2133 patients were enrolled at 267 centers in 14 countries; 2131 patients were included in the safety analysis; two patients were excluded due to missing baseline safety assessment.

The baseline demographic characteristics were comparable between the two treatment arms (Table 1; comparison with BOLERO-2 in Supplementary Table S1). The majority of patients (65.1%) received EVE plus EXE as a third-line of therapy or beyond in the advanced setting. Baseline patient characteristics of elderly patients were comparable to the non-elderly except for ECOG-PS 0/1/2 and less frequent prior chemotherapy in metastatic setting (additional information in Supplementary materials).

Treatment exposure

The median duration of follow-up was 4.6 months (range: <1–24.2 months). The median treatment duration was 3.7 months each for both EVE and EXE and appeared to be longer in patients who received EVE plus EXE as first-line therapy (4.4 months for EVE and 4.6 months for EXE). After censoring patients who discontinued treatment due to reimbursement/switch to other everolimus programs the median treatment duration was 5.1 months for EVE and 5.3 months for EXE and as first-line therapy, 6.1 months for both EVE and EXE; Figure 1A). Most common reasons for treatment discontinuation were disease progression (36.8%), local reimbursement of EVE (33.4%) and AEs (15.8%). Those patients who dropped out of the study continued on reimbursed EVE, however, the study did not plan to follow-up on them until treatment discontinuation. The overall median RDI was 0.98 (range: 0.1–1.1) for EVE and 1.0 (range: 0.4–1.0) for EXE.

In the elderly subset, the median duration of exposure was 3.2 months for EVE and 3.5 months for EXE. After censoring elderly patients who discontinued treatment due to reimbursement or crossover to other EVE programs, median duration of exposure was 3.8 months for EVE and 4.1 months for EXE, which was shorter compared to that in non-elderly patients, 5.0 months for EVE and 5.2 months for EXE. The median RDI for EVE was lower (0.95) in the elderly in comparison to non-elderly patients (1.0).

Dose interruptions, reductions and discontinuations

Overall, 29.6% and 1.2% of patients required dose reduction for EVE and EXE, respectively, the most common reasons being AEs (14.1% vs 0%) and medical decision (10.3% vs 0.1%). Dose interruptions for EVE and EXE were required for 55.9% and 19.3% of patients, respectively. More than half (52.3%) of the patients who needed a dose interruption were able to restart EVE at full dose. The median duration of dose interruption was 17 days. The median time to first dose modification was 32 days for EVE and 45 days for EXE. Permanent discontinuation of EVE was avoided through dose reduction in most patients; only 18.1% of these patients were unable to continue therapy thereafter for AEs. On the other hand, 15.1% of patients who never had a dose reduction stopped EVE permanently due to AEs.

Dose reductions and interruptions were higher in the elderly compared to non-elderly patients (37.7% and 60.5%, respectively vs. 26.7 % and 54.2%, respectively) (Supplementary Table S2). The most common AEs requiring dose adjustment in the elderly versus non-elderly patients were stomatitis (27.9% vs. 21.2%), asthenia (8.9% vs. 3.6%), anaemia (6.2% vs. 3.1%) and non-infectious pneumonitis (NIP, 5.9% vs. 4.8%), respectively. The median time to first dose modification was 30 days in the elderly and 33 days in non-elderly patients.

AEs led to permanent treatment discontinuation in 17.1% of patients (grade 3/4, 10%); the majority of these occurred within the first three months from treatment initiation. The most frequently reported AEs that led to permanent treatment discontinuation were (all grade; grade 3/4) NIP (2.4%; 0.8%), stomatitis (1.9%; 1.0%), asthenia (1.5%; 0.9%), and dyspnea (1.1%; 0.5%). AE-related treatment discontinuations were reported for 18.9% of the elderly versus 10.6% of non-elderly patients.

AE-related discontinuations were higher in the elderly versus non-elderly patients (23.8% vs. 13.0%). In the elderly, the most frequently reported AEs leading to permanent discontinuation of EVE or EVE plus EXE in the elderly versus non-elderly patients were stomatitis (3.9% vs. 1.2%) and NIP (3.4% vs. 2%).

Safety

Overall, 94.7% of patients experienced at least one any grade AE and 42.7% of patients experienced at least one grade 3/4 AE irrespective of relationship to EVE. Of these, 81.8% and 15.1% of AEs were assessed by the investigators to be EVE- or EXE-related, respectively; 27.2% and 1.6%, respectively were of grade 3/4 severity. The most frequently reported nonhematologic (all grade) AEs were stomatitis (52.8%; 50.8% were EVE-related), and asthenia (22.8%; 14.6% were EVE-related). Anemia (14.4%; 8.1% were EVE-related), and thrombocytopenia (5.9%; 4.6% were EVE-related) were the most frequently reported hematologic AEs (Table 2). The majority of EVE-related AEs were of grade 1/2 severity; the incidences of grade 3 or 4 stomatitis or NIP were low (Table 2; comparison with BOLERO-2 in Supplementary Table S3).

In the elderly subset, 95.2% of patients experienced at least one AE. The most common any grade AEs in elderly versus non-elderly patients were stomatitis (55.5% vs. 51.9%), asthenia (28.5% vs. 20.7%) and

decreased appetite (22.4% vs. 13.7%); the most frequent grade 3 or 4 AEs were stomatitis (12.3% vs. 8.3%), asthenia (5.7% vs. 2.9%) and hyperglycaemia (4.6% vs. 2.3%). NIP was reported in 11.2% of elderly versus 8.9% of non-elderly patients (Supplementary Table S4).

Post hoc analysis showed that the incidences of any grade AEs following EVE, including grade 3/4 AEs of special interest like stomatitis (range: 5.4-9.8%), NIP (range: 1.3-2.7%), asthenia/fatigue (range: 4.1-8.1%) were independent of the patients' BMI status (Supplementary Table S5).

The median time to onset for stomatitis events and NIP events were 29 days (range: 1–396) and 87 days (range: 1–231), respectively (Figure 1B). The median duration of a stomatitis event was 16 days and of a NIP event was 19 days. The incidence of EVE-related AEs appeared to be lower in the first-line setting for ABC vs later lines, with a numerically lower incidence observed for stomatitis (45.6% vs 51.4%), rash (11.4% vs 15.1%), asthenia (10.9% vs 15.1%) and diarrhea (9.1% vs 10.7%). Grade 3/4 stomatitis (7.7% vs 9.4%), diarrhea (0.5% vs 0.9%), rash (0.5% vs 1.0%) and NIP (0.9% vs 1.9%) were also reported less frequently in the first-line setting vs later lines.

Overall, 21.2% of patients experienced at least one SAE regardless of the relationship to any study treatment [Supplementary Table S6]; 8.5% of patients experienced at least one EVE-related SAE. The most frequent SAEs were dyspnea (2.4%), NIP (2.2%), pyrexia (1.6%), anemia (1.3%) and pleural effusion (1.2%). Treatment-related SAEs were reported in 10.3% of the elderly versus 7.8% of non-elderly patients.

Survival and follow-up status

At the time of analysis, 121 (5.7%) on-treatment deaths were recorded in the full study population. Deaths were attributable to disease progression (66 [3.1%]), AEs (46 [2.2%]) and unknown reasons (9 [0.4%]).

On-treatment deaths were reported in 39 (6.9%) elderly patients; the deaths were attributable to progressive disease (18 [3.2%]), AEs (16 [2.8%]) (additional data in Supplementary Table S7), and others reasons (5 [0.9%]). AEs suspected to be EVE-related leading to death were reported in 4 patients: NIP (2 patients), general physical health deterioration, and cardio-respiratory arrest (1 patient each).

Overall, 69.2% of patients received at least one anti-neoplastic medication since study treatment discontinuation [Supplementary Table S8].

Discussion

To our knowledge, this is the largest ever reported safety dataset on a general patient population presenting HR+, HER2– ABC progressing on prior NSAIs, treated with EVE plus EXE. Further, the exploratory safety analysis in the elderly subset in BALLEET is by far the largest safety dataset in this age group.

Overall, the patient population in BALLEET was more heavily pretreated compared with BOLERO-2 [14]. Patients in BALLEET were treated in later lines (65% vs 54% in BOLERO-2 in third-line and beyond) [14], and with more prior chemotherapy (60% vs 26% in BOLERO-2; 20.8% of patients in BALLEET received ≥ 3 lines of chemotherapy in the metastatic setting) because the trial did not impose limitations in term of number of prior chemotherapy [14]. This may be one of the reasons why fewer patients in BALLEET received any post-treatment therapy than those in BOLERO-2 (69.2% vs 84%) [15]. The short follow-up period could be another reason for this observation.

The incidence of stomatitis (52.8% vs 59%) and NIP (9.5% vs 16%) were lower in BALLEET compared with BOLERO-2 [14]; this difference may be attributed to the shorter median follow-up in BALLEET compared with BOLERO-2 (4.6 months vs 17.7 months). This could be because long-term safety profile of EVE could not be evaluated in those patients in BALLEET who dropped out of the study but continued on EVE under reimbursement. Another plausible reason for this difference could be the variability across study sites in the reporting of low-grade stomatitis. Consistent with BOLERO-2, the most common AEs leading to treatment discontinuation in BALLEET were NIP (2.4% vs 5.6% in BOLERO-2) and stomatitis (1.9% vs 2.7% in BOLERO-2) [14]. In BALLEET, fewer treatment-emergent AEs were reported in the first-line setting compared with later lines.

Median duration of treatment with EVE in BALLEET was lower compared with BOLERO-2 (16 weeks vs 23.9 weeks), even though the median RDI was higher [14]. This may suggest an improvement in

treatment optimization. Indeed, fewer patients in BALLEET discontinued study drug due to AEs (17.1% compared with patients in BOLERO-2 [26.3%]) [14] and clinicians preferred a temporary treatment interruption over permanently discontinuation.

The trends in the safety profile of EVE plus EXE in elderly patients in BALLEET were consistent with the ≥ 70 years subset (n=121) in the BOLERO-2 trial [16]. The median duration of exposure to EVE and EXE was shorter and the corresponding RDIs were lower in the elderly compared to non-elderly patients. Dose reductions, interruptions and AE-related study treatment discontinuations, grade 3/4 AEs, treatment-related SAEs and on-treatment deaths were higher in the elderly compared with non-elderly patients.

BALLEET is also the first trial reporting the impact of BMI on safety. Post hoc analysis showed that the incidence of AEs was independent of BMI status. This observation bears clinical implications as it appears that patients with lower BMI do not require a lower EVE starting dose.

Given that globally relevant clinical practice guidelines for breast cancer recommend EVE plus AIs including EXE as a treatment option for patients with HR+, HER2- ABC recurring or progressing on/after prior NSAIs [1, 17], these data add safety information to the clinical relevance of the efficacy data. Given the early incidence of AE and dose interruption/modification observed in BALLEET, close follow-up in the first months of therapy is indicated. We recommend a first visit 2 weeks after starting EVE (only one month in BALLEET) in order to further reduce EVE-related discontinuation rate.

The rate of study discontinuation due to EVE reimbursement (33.4%) was high. Although these patients continued on reimbursed EVE, they could not be followed-up until definitive treatment discontinuation. This might have influenced the long-term safety profile of the study treatments and treatment duration. An exploratory analysis of treatment duration censoring patients who discontinued treatment due to reimbursement showed the median duration of treatment to be 5.1 months and 5.3 months for EVE and EXE, respectively but a similar analysis for safety was not feasible from a statistical point of view due to lack of sufficient power. Despite these limitations, the inherent strength of BALLEET lies in the sizeable study population that was evaluated to provide meaningful safety data that support the utilization of the dual inhibition strategy with EVE plus EXE in this patient population.

Given that BALLEET was an expanded access program, the methodology did not allow for the assessment of PFS; however, given that the duration of exposure is surrogate marker for PFS, this has allowed cross-trial comparisons, in particular with the BOLERO-2 trial. As is inherent to expanded access trial designs, this design of BALLEET allowed for inclusion of patients unrestricted by the demographic profile of the BOLERO-2 patient population. This allowed for meaningful evaluation of the safety profile of EVE plus EXE in an a larger patient population that mimicked the real-world setting.

In conclusion, this is the largest ever reported safety dataset on a general patient population including a sizeable elderly subset, in patients with HR+, HER2– ABC progressing on prior NSAIs. The safety profile of EVE plus EXE in BALLEET is consistent with previous observations from BOLERO-2. There were no new safety signals. Diligent monitoring, proactive communication, early detection and implementation of appropriate AE-management strategies can ensure better treatment optimization.

Acknowledgments:

We thank the patients who participated in the BALLEET trial; the investigators, study nurses, and clinical research associates from the individual trial centers who provided ongoing support; and Avishek Pal (Novartis Healthcare Pvt. Ltd.) for providing medical editorial assistance with this manuscript.

Funding

This study was sponsored by Novartis Pharmaceuticals Corporation. No grant number is applicable.

Declarations of interests

Guy Jerusalem: during the conduct of the study: grants, personal fees and non-financial support from Novartis; outside of submitted work: grants, personal fees and non-financial support from Novartis and Roche, personal fees from Celgene, grants from MSD, non-financial support from GSK.

Gabriella Mariani: nothing to disclose.

Eva M Ciruelos: nothing to disclose.

Miguel Martin: speakers honoraria and advisory boards, and research funding from Novartis.

Vivianne CG Tjan-Heijnen: grants from Novartis , outside the submitted work.

Patrick Neven: nothing to disclose.

Joaquin G Gavila: nothing to disclose.

Andrea Michelotti: nothing to disclose.

Filippo Montemurro: outside the submitted work: speaker's bureau fees from Hoffmann La Roche and Astra Zeneca S.P.A and consultation fees from GlaxoSmithKline.

Daniele Generali: nothing to disclose.

Edda Simoncini: nothing to disclose.

Istvan Lang: nothing to disclose.

Jozef Mardiak: nothing to disclose.

Bjorn Naume: speakers honoraria from GlaxoSmithKline and Roche; institutional research funding from Sanofi

Maura Camozzi: employee of Novartis.

Katia Lorizzo: employee of Novartis.

Sara Bianchetti: employee of Novartis.

Pierfranco Conte: nothing to disclose.


References

- 1 Cardoso F, Costa A, Norton L, et al. ESO-ESMO 2nd international consensus guidelines for advanced breast cancer (ABC2). *Annals of Oncology* 2014; 25(10): 1871–1888.
- 2 Nabholz JM, Buzdar A, Pollak M, et al. Anastrozole is superior to tamoxifen as first-line therapy for advanced breast cancer in postmenopausal women: results of a North American multicenter randomized trial. *J Clin Oncol.* 2000; 18(22): 3758–3767.
- 3 Mouridsen H, Gershanovich M, Sun Y, et al. Phase III study of letrozole versus tamoxifen as first-line therapy of advanced breast cancer in postmenopausal women: analysis of survival and update of efficacy from the International Letrozole Breast Cancer Group. *J Clin Oncol.* 2003; 21(1): 2101–2109.
- 4 Bergh J, Jonsson P-E, Lidbrink EK, et al. FACT: An open-label randomized phase III study of fulvestrant and anastrozole in combination compared with anastrozole alone as first-line therapy for patients with receptor-positive postmenopausal breast cancer. *J Clin Oncol.* 2012; 30(16): 1919-1925.
- 5 Chia S, Gradishar W, Mauriac L, et al. Double-blind, randomized placebo controlled trial of fulvestrant compared with exemestane after prior nonsteroidal aromatase inhibitor therapy in postmenopausal women with hormone receptor-positive, advanced breast cancer: results from EFECT. *J Clin Oncol.* 2008; 26(10): 1664–1670.
- 6 Osborne KC, Schiff R. Mechanisms of endocrine resistance in breast cancer. *Annu Rev Med.* 2011; 62: 233–247.
- 7 Baselga J, Campone M, Piccart M, et al. Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer. *N Engl J Med.* 2012; 366(6): 520–529.
- 8 Turner NC, Ro J, Andre F, et al. Palbociclib in Hormone-Receptor-Positive Advanced Breast Cancer. *N Engl J Med.* 2015; 373(3): 209–19.
- 9 Gnant M. The role of mammalian target of rapamycin (mTOR) inhibition in the treatment of advanced breast cancer. *Curr Oncol Rep.* 2013; 15(1): 14–23.
- 10 Jerusalem G, Rorive A, Collignon J. Use of mTOR inhibitors in the treatment of breast cancer: an evaluation of factors that influence patient outcomes. *Breast Cancer: Targets and Therapy.* 2014; 6: 43–57.


- 11 Jerusalem G, Bachelot T, Neven P, et al. A new era of improving progression-free survival with dual blockade in postmenopausal HR(+), HER2(-) advanced breast cancer. *Cancer Treat Rev.* 2015; 4(2): 94–104.
- 12 Gnant M. Overcoming endocrine resistance in breast cancer: importance of mTOR inhibition. *Expert Rev Anticancer Ther.* 2012; 12(12): 1579–1589.
- 13 Buolay A, Rudloff J, Ye J, et al. Dual inhibition of mTOR and estrogen receptor signaling in vitro induces cell death in models of breast cancer. *Clin Cancer Res.* 2005; 11(14): 5319–28.
- 14 Yardley DA, Noguchi S, Burris HA, et al. Everolimus plus exemestane in postmenopausal patients with HR(+) breast cancer: BOLERO-2 final progression-free survival analysis. *Adv Ther.* 2013; 30(10): 870–84.
- 15 Piccart M, Hortobagyi GN, Campone M, et al. Everolimus plus exemestane for hormone-receptor-positive, human epidermal growth factor receptor-2-negative advanced breast cancer: overall survival results from BOLERO-2. *Ann Oncol.* 2014; 25(12): 2357–2362.
- 16 Pritchard KI, Burris HA, Ito Y, et al. Safety and efficacy of everolimus with exemestane vs. exemestane alone in elderly patients with HER2-negative, hormone receptor-positive breast cancer in BOLERO-2. *Clin Breast Cancer.* 2013; 13(6): 421–432.e8.
- 17 National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Breast Cancer. V1.2015.

Figures Legends

- Figure 1A** Kaplan Meier curve for time to treatment discontinuation after censoring patients who discontinued treatment due to reimbursement or natural death (Safety set)
- Figure 1B** Median time to onset for stomatitis events and noninfectious pneumonitis (NIP) events (Safety set)



	Number of subjects still at risk												
	0	2	4	6	8	10	12	14	16	18	20	22	24
Everolimus	2129	1666	978	602	303	170	85	39	13	5	1	1	0
Exemestane	2128	1732	1017	620	307	173	84	37	12	5	1	1	0



Number of subjects still at risk

Stomatitis	2131	1062	369	355	194	110	53	23	5	0	0	0	0	0	0
NIP	2131	1908	1231	735	401	220	123	58	22	9	3	1	1	0	0

Table 1 Baseline and treatment characteristics (including data for prior therapies by number of prior chemotherapy lines) (Full analysis set)

Characteristic	N = 2133
Median age (range), years	63.0 (28-90)
Age categories	
< 70 years, <i>n</i> (%)	1570 (73.6%)
≥ 70 years, <i>n</i> (%)	563 (26.4%)
Median BMI (range)	25.7 (14.69-54.57)
Race, <i>n</i> (%)	
Caucasian	2100 (98.5%)
Black	5 (0.2%)
Asian	1 (0.0%)
Native American	1 (0.0%)
Others	26 (1.2%)
Ethnicity, <i>n</i> (%)	
Hispanic/Latino	314 (14.7%)
Indian	1 (0.0%)
Other	1818 (85.2%)
ECOG PS	
0	1383 (64.8%)
1	670 (31.4%)
2	56 (2.6%)
Missing	24 (1.1%)
Characteristic	N = 2131
Current disease status, <i>n</i> (%)	
Metastatic	1762 (82.7%)
Locally advanced	369 (17.3%)
Metastatic site, <i>n</i> (%)	
Bone only	543 (25.5%)
Visceral	1277 (59.9%)
Visceral only	208 (9.8%)
Bone and visceral	924 (43.4%)
Others	775 (36.4%)
Number of metastatic site, <i>n</i> (%)	
≥5	469 (22.0%)
4	281 (13.2%)
3	437 (20.5%)
2	509 (23.9%)
1	435 (20.4%)
Median time to first diagnosis (range), days	2908 (-16218- -76)
Key comorbidities, <i>n</i> (%)	
Vascular	808 (37.9%)
Metabolic and nutritional	676 (31.7%)
Musculoskeletal and connective tissue	599 (28.1%)
Psychiatric	410 (19.2%)
Gastrointestinal	259 (12.2%)
Number of lines of prior antineoplastic therapy in metastatic setting, <i>n</i> (%)	
None (adjuvant therapy only)	222 (10.4)
1	518 (24.3%)
2	483 (22.7%)
3	345 (16.2%)
4	209 (9.8%)
≥5	350 (16.4%)

Key prior antineoplastic therapies, <i>n</i> (%)	2131 (100%)
Aromatase inhibitors	2122 (99.6%)
Anti-estrogens	1579 (74.1%)
Chemotherapy in metastatic setting	1284 (60.3%)
Immunosuppressant	564 (26.5%)
Monoclonal antibodies	272 (12.8%)

Table 2 Adverse events of $\geq 10\%$ incidence in either treatment group or grade 3, 4 AEs of $\geq 0.5\%$ incidence in either treatment group (Safety set)

Adverse event (N = 2131), n (%)	All grades	Grade 3	Grade 4
Stomatitis	1126 (52.8%)	198 (9.3%)	2 (0.1%)
Asthenia	485 (22.8%)	75 (3.5%)	2 (0.1%)
Diarrhea	359 (16.8%)	26 (1.2%)	1 (0.0%)
Rash	351 (16.5%)	20 (0.9%)	0
Decreased appetite	341 (16.0%)	24 (1.1%)	0
Anemia	306 (14.4%)	0	0
Pyrexia	299 (14.0%)	7 (0.3%)	3 (0.1%)
Fatigue	298 (14.0%)	26 (1.2%)	1 (0.0%)
Hyperglycemia	265 (12.4%)	60 (2.8%)	3 (0.1%)
Peripheral edema	259 (12.2%)	13 (0.6%)	0
Nausea	255 (12.0%)	13 (0.6%)	0
Cough	254 (11.9%)	8 (0.4%)	0
Dyspnea	220 (10.3%)	39 (1.8%)	4 (0.2%)
Decreased weight	217 (10.2%)	2 (0.1%)	0
Hypercholesterolemia	216 (10.1%)	1 (0.0%)	1 (0.0%)
NIP	203 (9.5%)	35 (1.6%)	6 (0.3%)

NIP, noninfectious pneumonitis