

Identification and characterisation of Nanobodies targeting the EphA4 receptor

Lies Schoonaert<sup>1,2\*</sup>, Laura Rué<sup>1,2\*</sup>, Bart Roucourt<sup>3</sup>, Mieke Timmers<sup>1,2</sup>, Susan Little<sup>4</sup>, Lucía Chávez Gutiérrez<sup>2,5</sup>, Maarten Dewilde<sup>2,5</sup>, Peter Joyce<sup>4</sup>, Adam Curnock<sup>4</sup>, Peter Weber<sup>4</sup>, Jurgen Haustreaete<sup>6</sup>, Gholamreza Hassanzadeh-Ghassabeh<sup>7</sup>, Bart De Strooper<sup>2,5,8</sup>, Ludo Van Den Bosch<sup>1,2</sup>, Philip Van Damme<sup>1,2,9</sup>, Robin Lemmens<sup>1,2,9</sup>, Wim Robberecht<sup>1,9</sup>

<sup>1</sup> KU Leuven - University of Leuven, Department of Neurosciences, Laboratory of Neurobiology & Experimental Neurology and Leuven Research Institute for Neuroscience and Disease (LIND), B-3000 Leuven, Belgium

<sup>2</sup> VIB, Center for Brain & Disease, Leuven, Belgium

<sup>3</sup> Laboratory for Signal Integration in Cell Fate Decision, University of Leuven, Leuven, Belgium

<sup>4</sup> Vertex Pharmaceuticals Incorporated (Europe) Ltd., Biology Dept., Abingdon, OX14 4RW, UK

<sup>5</sup> KU Leuven, Center for Human Genetics, and Leuven Research Institute for Neuroscience and Disease (LIND), Leuven, Belgium

<sup>6</sup> Protein Service Facility, Inflammation Research Center, VIB, Ghent University, Ghent, Belgium

<sup>7</sup> Nanobody Service Facility, VIB, Vrije Universiteit Brussel, Brussels, Belgium

<sup>8</sup> Institute of Neurology, University College London, London, UK

<sup>9</sup> University Hospitals Leuven, Department of Neurology, Leuven, Belgium

\* Authors contributed equally to the paper

Running title: *EphA4 inhibiting Nanobodies*

To whom correspondence should be addressed: Wim Robberecht, MD, PhD, Laboratory of Neurobiology, KU Leuven, Campus Gasthuisberg, O&N4 Herestraat 49 box 602, Leuven, Belgium, Telephone: (0032)16373188; FAX: (0032)16330212; E-mail: Wim.Robberecht@kuleuven.be

**KEYWORDS:**

Single-domain antibody (sdAb,nanobody), VHH, EphA4, ligand-binding domain, receptor tyrosine kinase, phosphorylation, protein-protein interaction, inhibition, regeneration

**ABSTRACT**

The ephrin receptor A4 (EphA4) is one of the receptors in the ephrin system that plays a pivotal role in a variety of cell-cell interactions, mostly studied during development. In addition, EphA4 has been found to play a role in cancer biology as well as in the pathogenesis of several neurological disorders such as stroke, spinal cord injury, multiple sclerosis (MS), amyotrophic lateral sclerosis (ALS) and Alzheimer's disease (AD). Pharmacological blocking of EphA4 has been suggested to be a therapeutic strategy for these disorders. Therefore, the aim of our study was to generate potent and selective Nanobodies against the ligand-binding domain of the human EphA4 receptor. We identified two Nanobodies, Nb 39 and Nb 53, that bind EphA4 with affinities in the nanomolar range. These Nanobodies were most selective for EphA4, with residual binding to EphA7 only. Using Alphascreen technology, we found that both Nanobodies displaced all known EphA4-binding ephrins from the receptor. Furthermore, Nb 39 and Nb 53 inhibited ephrin-induced phosphorylation of the EphA4 protein in a cell-based assay. Finally, in a cortical neuron primary culture, both Nanobodies were able to inhibit endogenous EphA4-mediated growth cone collapse induced by ephrin-B3. Our results demonstrate the potential of Nanobodies to target the ligand-binding domain of EphA4. These Nanobodies may deserve further evaluation as potential therapeutics in disorders in which EphA4-mediated signalling plays a role.

Ephrin receptors, the largest family of receptor tyrosine kinases, are subdivided in A- and B- class receptors. In mammals there are nine EphA receptors (EphA1-EphA8, EphA10) that interact with five ephrin-A ligands (ephrin-A1- ephrin-A5) and five EphB receptors (EphB1-EphB4, EphB6) that interact with three ephrin-B ligands (ephrin-B1-ephrin-B3) (1). There is some interclass promiscuity as EphA4 can also interact with ephrin-B ligands while EphB2 also interacts with ephrin-A5 (2-4).

The ephrin system plays a major role in a variety of cell-cell interactions. In the developing nervous system it is pivotal as an axonal guidance system, while in the adult brain, it is involved in synaptic plasticity and long-term potentiation (LTP) (5,6). Importantly, the ephrin system also plays a role in cancer biology and in

the pathogenesis of several neurological disorders (5). EphA4 is upregulated in spinal cord injury, traumatic brain injury and stroke, and blocking of the receptor increases functional recovery in models for these conditions (7-9). Interestingly, antagonism of EphA4 improves LTP defects in a mouse model for Alzheimer's Disease (AD) and improves outcome in animal models for Amyotrophic Lateral Sclerosis (ALS) and stroke (9-12). EphA4 expression is inversely correlated with survival time in ALS patients, which suggests that EphA4 is also involved in ALS human pathology (11). These findings suggest that EphA4 inhibition may have potential for the treatment of neurological disorders.

Inhibition of EphA4 signaling can be accomplished by targeting the ATP-binding pocket in the kinase domain or by blocking the interaction with ephrin ligands (13,14). Because the ATP-binding pocket is highly conserved among tyrosine kinases it is very difficult to develop selective inhibitors (13). Alternatively targeting the ligand-binding domain (LBD) certainly allows the development of more selective compounds, but this poses several other problems. On one hand, the protein interaction surface to be covered is large; 900 Å<sup>2</sup> in the case of EphA4 (15,16). On the other hand, the LBD is dynamic by nature, as EphA4 can adopt a conformation similar to other EphA receptors upon interaction with ephrin-A ligands, or characteristics of EphB receptors when interacting with ephrin-B ligands (2,4,17).

Nevertheless several peptides and small molecules have been identified that bind the EphA4 LBD and block its interaction with ephrin ligands (18-22). One of these EphA4 antagonists is the KYL-peptide, which has been extensively characterised and shown to be effective in several *in vitro* assays as well as in *in vivo* spinal cord injury and ALS models (11,19,23,24), suggesting the potential of an EphA4-based therapeutic approach.

The aim of this study was to develop highly selective and potent EphA4 inhibitors. In order to achieve this goal, we took advantage of the Nanobody technology (25-27). Nanobodies (Nbs) or VHHs are small antigen-binding fragments derived from camelid heavy-chain only antibodies that are devoid of light chains. They are superior to conventional antibodies in terms of stability, solubility and immunogenicity (27). Furthermore, they are much smaller than

conventional antibodies (12-15 kDa versus 150-160 kDa) and can penetrate small clefts and cavities (28).

We were able to generate Nbs against the LBD of the EphA4 receptor. Two of these Nbs specifically bind the EphA4 receptor with nanomolar affinities and block ephrin-induced EphA4 phosphorylation and EphA4-mediated actin remodelling in a growth cone collapse assay. These results demonstrate the potential of Nbs to selectively target the LBD of the EphA4 receptor. These Nbs may be useful as a therapeutic strategy in disorders in which EphA4 plays a pathogenic role.

## Results

### **Generation of anti-EphA4 LBD Nbs**

An alpaca was immunized with recombinant human EphA4 LBD according to standard procedures (29). In addition to conventional antibodies, alpacas also produce heavy-chain only antibodies in response to an immunogen. A phagemid library displaying Nbs was constructed from the RNA extracted from peripheral blood lymphocytes and transformed in *E.coli* TG1 cells. A library of about  $2 \times 10^8$  independent transformants was obtained and about 87% of transformants harboured the vector with the right insert size. Phage particles were generated and subjected to panning. After four consecutive rounds of panning, we obtained 41 colonies that expressed antigen-specific Nbs in their periplasmic extracts, as determined by ELISA. Sequencing of the Nb genes from these 41 positive colonies resulted in 15 different Nbs. Characteristically Nbs contain three complementary determining regions (CDRs), which contribute to antigen binding specificity. Based on sequence homology they belonged to nine different clonally-unrelated B-cell clones (Figure 1). Nbs belonging to the same group showed very high sequence similarity suggesting that they are originating from clonally-related B-cells that underwent hypermutation. Nbs 39, 16, 71, 28 and 22 most likely belong to unrelated B-cell clones.

### **Expression of the anti-EphA4 LBD Nbs**

Phagemids encoding the Nbs of interest were transformed into WK6 *E.coli* cells to allow expression of the Nbs without the pIII fusion.

The expressed Nbs were recovered from the periplasmic extracts and purified. Their purity and was analysed by Coomassie-stained SDS-polyacrylamide gels and their identity was determined with Western Blot detection using His<sub>6</sub>-tag (data not shown). All Nbs could be detected at a position corresponding to about 15 kDa. However, there were obvious differences in expression levels of the Nbs (data not shown). Concentration of poorly expressed Nbs 90, 16, 71, 28 and 10 resulted in protein aggregation. Therefore, we did not include these Nbs in further analyses.

### **Analysis of EphA4 binding**

We next determined whether the Nbs recognised EphA4 under reducing, non-native conditions. None of them bound recombinant EphA4 (data not shown) on Western Blot. An EphA4 monoclonal antibody was used as a positive control and readily bound the denatured recombinant protein. These data suggested that the Nbs target a conformational epitope on the EphA4 LBD. To test whether the Nbs were indeed able to bind native EphA4, we performed immunoprecipitation experiments (Figure 2). Nbs 31, 39, 50, 53 and 57 precipitated EphA4 protein and EphA4 in turn precipitated these Nbs (data shown only for Nbs 53 and 39 in figure 2A, B). In contrast to the other Nbs, Nb 22 showed lower binding to mouse EphA4 compared to human EphA4 (Figure 2C). Nbs 19, 34 and 47 revealed non-specific binding to the beads and to recombinant ephrin-B2 (data shown only for Nb 19 and 47 in Figure 2D, E). Nb 60 did not show non-specific binding to the beads, but precipitated with recombinant ephrin-B2 indicating cross-reactivity with ephrin-B2 (Figure 2F).

To identify the Nbs with the highest affinity, we determined their binding kinetics to EphA4 using Surface Plasmon Resonance (SPR). The EphA4 LBD was immobilized onto the chip and Nbs (1 nM to 300 nM) were used as analyte. SPR analyses revealed that Nbs 34 and 47 do not bind to the EphA4 LBD (Table 1). Nbs 22, 31, 39, 50, 53, 57 and 60 bound in the low nanomolar range, while Nb 19 only interacted with EphA4 LBD in the higher nanomolar range (Table 1).

We next investigated the affinity of the Nbs for full-length recombinant human and mouse

EphA4 receptor. Most Nbs showed similar binding to the full-length receptor as to the EphA4 LBD. Nb 22 showed a two-fold difference in binding affinity between human and mouse EphA4, in agreement with what we found in immunoprecipitation experiments (Figure 2C).

Based on these results, we selected seven Nbs (60, 57, 39, 22, 31, 50 and 53) with affinities in the nanomolar range for both the EphA4 LBD and the full-length receptor.

### ***Cross-reactivity with other Eph receptors***

Since the homology between the different Eph receptors is very high, we used Alphascreen technology to study the specificity of the Nbs generated. All Eph receptors tested showed binding with ephrin-A5 and/or ephrin-B2 indicating that all receptors adopted their correct conformation. Ephrin-A5 interacted with all EphA receptors and with EphB2 as has been previously shown (data not shown) (3). Ephrin-B2 interacted with all EphB receptors and EphA4, but not with EphB6 for which no ligands have been identified so far (data not shown) (4,30). The control Nb did not bind to any of the Eph receptors as expected (data not shown). Alphascreen analyses showed that Nb 60 interacted with several Eph receptors other than EphA4 (Figure 3A), in agreement with the immunoprecipitation results obtained with this Nb (Figure 2F). Nb 57 bound EphA4, but also EphA3 and EphA7, and Nb 50 bound EphA4 but also EphA3, EphA6, EphA7 and EphA8 (Figure 3B,F). Nbs 22, 31, 39 and 53 were almost completely selective for EphA4, although some binding to EphA7 was detected (Figure 3C, D, E, G). We therefore selected the latter four for further screening.

### ***Competition with ephrin ligands for the interaction with EphA4***

EphA4 interacts with most of the ephrins (2,4). We therefore investigated whether the selected Nbs were able to displace the ligands from the EphA4 receptor. We determined the optimal concentration for the different ephrin ligands using a fixed concentration of the EphA4 LBD to avoid the hooking effect and we added

increasing concentrations of the Nbs. The control Nb did not compete with any ephrin ligand for the interaction with EphA4, as expected (data not shown). The KYL peptide, a known EphA4 inhibitor, was used as a positive control.

Nb 39 and Nb 53 completely displaced all ephrin ligands from EphA4 binding in a concentration range lower than the KYL-peptide (Figure 4A-H). Nbs 22 and 31 were less potent. Nb 22 completely displaced ephrin-A3 (Figure 4C) and almost completely displaced ephrin-B ligands (Figure 4F, G, H) from EphA4 binding, but no full displacement of the other ephrin-A ligands was obtained with the concentrations tested (Figure 4A, B, D, E). Its potency was comparable to that of the KYL peptide. Nb 31 completely displaced ephrin-A1, ephrin-B1 and ephrin-B2 at lower concentrations than the KYL-peptide (Figure 4A, F, G), and completely displaced ephrin-A3 with concentrations similar to the KYL-peptide (Figure 4C). However, ephrin-A2, ephrin-A4, ephrin-A5 and ephrin-B3 could not be completely displaced with the concentrations tested (Figure 4B, D, E). These data show that Nbs 39 and 53 were able to block the interaction of all ephrin ligands with the EphA4 LBD.

To examine the potency of Nb 39 and 53, in inhibiting another Eph receptor, EphA7, compared to EphA4, we determined the capacity of these two Nbs to modify the interaction between EphA7 and ephrin-A5 in a competition assay. We determined the optimal binding concentrations of EphA7 and ephrin-A5. At a concentration in which Nb 39 and Nb 53 were able to inhibit binding of ephrin-A5 to EphA4 (Figure 4E) only partial inhibition of the interaction between EphA7 and ephrin-A5 could be detected (Figure 5). A control Nb did not compete with ephrin-A5-binding to EphA7 (data not shown).

### ***Inhibition of ephrin-induced EphA4 activation***

To assess the antagonistic properties of the Nbs, we tested their effect on ephrin-induced EphA4 phosphorylation (31). Upon ephrin binding, Eph receptors are dimerised and activated by autophosphorylation (phosphorylation of tyrosine residues in the kinase domain and juxtamembrane region) (31). After clustering, the phosphorylated tyrosine residues form binding

sites for cytoplasmic targets with Src Homology 2 (SH2) or Phosphotyrosine Binding (PTB) domains (31,32). In general, EphA4 phosphorylation results in cell-cell segregation in this way regulating cytoskeleton dynamics and morphology (1). Ephrin-induced phosphorylation was examined using the PathHunter assay in U2OS cells. Cells were stimulated with ephrin-A1 with or without the presence of an EphA4 antagonist, and the resulting EphA4 phosphorylation was measured. The control Nb did not have any effect on receptor phosphorylation (data not shown). The KYL-peptide achieved complete inhibition of ephrin-A1-induced phosphorylation with an  $IC_{50}$  value of 53  $\mu$ M (Figure 6). Nbs 39 and 53 also completely inhibited ephrin-A1-induced phosphorylation, at lower concentrations than the KYL-peptide, with an  $IC_{50}$  of 170 nM and 261 nM respectively (Figure 6). These data indicate that Nbs 39 and 53 are 200 to 300 times more potent than the KYL-peptide in inhibiting ephrin-A1-induced phosphorylation of the EphA4 receptor.

### ***Inhibition of ephrin-induced growth cone collapse***

A common cellular response downstream of Eph receptor activation is actin cytoskeleton reorganization, leading to cell morphology changes (1,33). Ephrin-B3 treatment of cortical neurons *in vitro* induces collapse of growth cones at the tip of the neurites, by reorganization of actin filaments (34,35). Cortical neurons lacking functional EphA4 show insensitivity to the ephrin-B3-mediated growth cone collapse, suggesting that this effect is EphA4-mediated (36). We next determined whether the Nbs are also able to antagonize a cellular response such as growth cone collapse induced by an EphA4 receptor agonist. To this end, we investigated their effect on the ephrin-B3 induced inhibition of neurite elongation in E17.5 embryonic cortical neurons. As described before, stimulation with preclustured recombinant ephrin-B3-Fc resulted in growth cone collapse as compared to an unstimulated condition and a condition stimulated with Fc only (Figure 7). As expected, KYL peptide antagonized this effect on growth cone collapse. Nb 53 as well as Nb 39 also inhibited the growth cone collapse effect

mediated by ephrin-B3-Fc, at the concentration of 1  $\mu$ M, a concentration at which the KYL peptide was ineffective (Figure 7).

### ***Nb stability in plasma***

We incubated Nb 39 and Nb 53 in heparinized mouse plasma at 37°C for up to 168h (7 days) to study the stability in plasma and obtained aliquots at different time-points. We determined the stability of Nbs with Western Blot using an anti-HA antibody. Some cleavage products could be observed from 24 hours onwards for both Nb 39 and 53 (Figure 8A). Whereas full-length Nb 39 remained stable across the timecourse, full-length Nb 53 levels diminished with time from 72 hours onwards, suggesting that Nb 53 is more prone to degradation in plasma than Nb 39 (Figure 8A, B). No degradation products were detected for any of the Nbs after incubation in PBS at 37°C for the same time-course length (Figure 8C). In addition we studied if the binding capacity of Nbs to EphA4 was altered after incubation in plasma by Alphascreen technology. After 72h of incubation in plasma at 37°C, Nbs were still capable of binding EphA4 LBD (Figure 8D).

### **Discussion**

In view of EphA4's role in normal physiology as well as in cancer and neurodegeneration, the generation of potent and selective antagonists against this receptor is of great interest. The EphA4 LBD may be a better target than its tyrosine kinase moiety, because of the difficulty to obtain selective Eph tyrosine kinase inhibitors, and because of the need for such molecules to enter the cell. Here we show for the first time the generation of single domain antibodies targeting the LBD of EphA4.

Nbs are in many ways superior to small molecules when targeting the LBD of Eph receptor proteins, as high affinity binding of the LBD requires the coverage of the protein interaction surface (900  $\text{\AA}^2$  for EphA4) in order to block the interaction with EphA4 ligands. Both Nb 39 and Nb 53 were able to block the interaction of EphA4 with all ephrin ligands and to inhibit ephrin-induced phosphorylation of

EphA4 and growth cone collapse triggered by ephrin-B3 binding to endogenous EphA4.

In previous studies, Nbs were used to fix receptors in one conformation (28,37). As a consequence Nbs are an ideal probe to target dynamic structures such as the EphA4 LBD (2,4,17). Four different conformations have been reported for the EphA4 LBD: a conformation similar to other EphA receptors bound with ephrin-A ligands (2,38), a conformation similar to EphB receptors bound with ephrin-B ligands (2,4,15) and two unbound conformations when free in solution (17). Fixing the EphA4 LBD in a specific conformation might limit the accessibility for all or a specific class of ephrin ligands, thereby blocking the interaction of those ligands with the LBD.

Nb 39 and Nb 53 were most selective for EphA4, but still showed residual binding to EphA7, suggesting that they target a region that is not highly conserved in all Eph receptors. However concentrations of Nb 39 and Nb 53 that strongly inhibit EphA4 binding of ephrin ligands, the interaction between EphA7 and ephrin-A5 was only partially reduced. Interestingly, EphA7 KO mice as well as rats treated with EphA7 antisense oligonucleotides showed enhanced recovery after Spinal Cord Injury (SCI) compared to control mice (39), similar to the effects described for EphA4 antagonists (24). Therefore, if some binding to EphA7 would occur in addition to EphA4, this could potentially be of benefit rather than detrimental.

In the last years, extensive work has been performed aiming to develop EphA4 antagonists with high affinity, specificity and with good pharmacokinetic profiles (14). One EphA4 inhibitor, the KYL-peptide, has been studied extensively in models of neurological disorders, such as acute injuries including spinal cord injury and stroke, and neurodegenerative disorders such as ALS and AD (10,11). Unfortunately, this peptide has a  $K_D$  value of approximately 1  $\mu$ M as determined with Isothermal Titration Calorimetry (ITC) and a very short half-life in serum (11 min in mouse serum) (19,24). Interestingly, a recent study reported the generation of highly selective EphA4 antagonists with higher potency, the cyclic peptide APY-derivatives, APY-d3 and APY-d4 (16,22). APY-d3 and APY-d4 have  $K_D$

values of 30 nM and 20 nM as determined with ITC, and inhibit ephrin-A5-induced phosphorylation of EphA4 with an  $IC_{50}$  of 240 nM and 310 nM. Both cyclic peptides are stable in mouse plasma for more than 72 hours and are currently the best available EphA4 inhibitors (22). Nb 39 and 53 are more potent than the KYL-peptide, but similarly potent as APY-d3 and APY-d4, with  $K_D$  values in the nanomolar range, as measured with SPR. Moreover, both Nbs could inhibit EphA4 phosphorylation with an  $IC_{50}$  of 170 nM and 261 nM respectively. Similar to APY-d3 and APY-d4, Nb 39 and Nb 53 were stable in mouse serum for more than 72 hours and maintained their full capability of binding EphA4 LBD. The affinity of Nbs could still be further increased through error-prone PCR mutagenesis and/or making bispecific Nbs (40,41). Bivalent or bispecific Nbs can be obtained by connecting two identical or different Nbs with a linker thereby improving the avidity. However, this strategy requires caution in the setting of developing EphA4 antagonists, as dimerisation of Eph receptors may induce clustering and subsequently activation, similar to what is obtained with a preclustered antibody (42).

EphA4 has been found to play a role in cancer biology and in the pathogenesis of several neurological disorders (5). In the central nervous system, EphA4 is highly expressed in cell bodies, dendritic spines and axons of neurons in various brain regions (43,44). EphA4 expression is upregulated in axonal stumps after injury (45) and in sprouting neurons of aged mice after stroke, which contributes to reduced recovery (46). Furthermore, blocking EphA4 induced more sprouting after spinal cord injury (8). EphA4's negative effect on axonal regeneration is mediated through the interaction of the receptor with ephrin ligands on surrounding cells such as muscle cells, astrocytes, microglia and oligodendrocytes (47). Neuronal EphA4 may interact with ephrin-A5 and ephrin-B2 of which the expression is highly upregulated in reactive astrocytes after injury (48-50). This high astrocytic ephrin-A5 expression inhibited axonal sprouting and motor recovery after stroke (49). Also, ephrin-B2 expression decreased axonal sprouting as deleting ephrin-B2 from reactive astrocytes reduced glial scar formation and improved recovery after spinal cord injury and this was correlated with an increased

regenerative capacity of sprouting spinal cord axons (45,48,50). Oligodendrocytes precursor cells are glial cells responsible for remyelination during neuronal injury or degeneration, thereby inhibiting axonal outgrowth (51). Neuronal EphA4 may interact with ephrin-A1, ephrin-A3, ephrin-A5, ephrin-B1 and ephrin-B2, which have been identified in oligodendrocyte precursor cells and in mature oligodendrocytes (52-54). In addition, enhanced functional recovery was observed in ephrin-B3 KO mice after spinal cord injury and blocking ephrin-B3 promotes remyelination *in vivo* in a rat model of remyelination (55,56). The relative importance of the different surrounding cells and different ligands is not yet known. However, as EphA4 is a promiscuous receptor, targeting EphA4 by blocking the interaction with all ephrin ligands might be the most efficient therapeutic strategy against neurological diseases.

Treating neurological disorders by targeting EphA4 implies using substances that are able to penetrate the Blood Brain Barrier (BBB) in order to reach their target. In acute injuries, this barrier is impaired. In neurodegenerative diseases such as ALS, getting drugs across the BBB remains a major challenge, despite of the fact that the BBB is abnormal in this disease (57-59). Several strategies to get biologics across the BBB have been developed, such as transcytosis through clathrin vesicles and receptor-mediated transcytosis by targeting the Low-Density Lipoprotein receptor, the transferrin receptor or the insulin receptor (60-65). Interestingly, one study showed that Nbs with a high isoelectric point can cross the BBB spontaneously (66).

Nbs also have some drawbacks. Although Nb 39 and Nb 53 remained stable in mouse plasma for more than 72 hours, the small size of Nbs limits their half-life to about 1.5 h after *in vivo* administration (67). Due to their small size they are rapidly cleared from blood via the kidney (27). This low half-life can be overcome by linking the Nb to serum albumin, which can increase the half-life to 20-30 h in mice (68). In humans, this approach extended the half-life of Nbs to 19 days (68,69). Another strategy to increase the half-life of Nbs is coupling polyethylene glycol (PEG) groups to the Nbs (70). The addition of PEG groups increases the apparent molecular weight above the glomerular

filtration limit avoiding renal clearance and/or evades cellular clearance mechanisms (71).

In summary, we describe for the first time the development of Nbs that target the EphA4 LBD. We identified two Nbs that were specific for EphA4 with residual EphA7 binding, and with  $K_D$  and  $IC_{50}$  values in the nanomolar range. Both Nbs were able to block the interaction of EphA4 with all ephrin ligands, inhibit EphA4 phosphorylation, and the growth cone collapse mediated by EphA4 activation upon ephrin-B3 interaction. These Nbs may be useful tools to study the role of the EphA4 receptor in preclinical cancer or neurodegeneration models. Furthermore they may represent a starting point for an EphA4-based therapeutic approach using Nbs.

## EXPERIMENTAL PROCEDURES

**Expression and purification of the EphA4 ligand-binding domain** - The EphA4 LBD (AA 22 – AA 203 (17)) was cloned from the EphA4 Human cDNA ORF clone (Origene, US) and expressed in the *E. coli* strain BL21 codon + pICA2 transformed with the pLH36Epha plasmid. Expression was induced by isopropyl b-D-1-thiogalactopyranoside under control of a pL-promotor developed by the Protein Service Facility of VIB. The pLH36 plasmid was provided with a His<sub>6</sub>-tag followed by a murine caspase-3 site. The murine caspase-3 site can be used to remove the His<sub>6</sub>-tag attached at the N-terminus of the protein of interest during purification. The transformed bacteria were grown in Luria Bertani medium supplemented with ampicillin (100 µg/mL) and kanamycin (50 µg/mL) overnight at 28°C inoculation (1/100) in a 20 liter fermenter provided with Luria Bertani medium supplemented with ampicillin (100 µg/mL) and 1% glycerol. The initial stirring and airflow was 200 rpm and 1.5 L/min, respectively. Further, this was automatically adapted to keep the pO<sub>2</sub> at 30 %. The temperature was kept at 28°C. The cells were grown to an optical density of  $A_{600nm} = 1.0$ , transferred at 20°C, and expression was induced by addition of 1 mM isopropyl b-D-1-thiogalactopyranoside overnight. Cells were then harvested and frozen at -20°C. After thawing, the cells were resuspended at 3 mL/g in 20 mM NaH<sub>2</sub>PO<sub>4</sub> pH 7.5, 500 mM NaCl, 20 mM

imidazole and 1 mM phenylmethylsulfonyl fluoride. The cytoplasmic fraction was prepared by sonication of the cells followed by centrifugation at 18,000g for 30 min. All steps were conducted at 4°C. The clear supernatant was applied to a 74 mL Ni-Sepharose 6 FF column (GE Healthcare, Belgium), equilibrated with 20 mM NaH<sub>2</sub>PO<sub>4</sub> pH 7.5, 500 mM NaCl, 20 mM imidazole and 0.1% CHAPS. The column was eluted with 20 mM NaH<sub>2</sub>PO<sub>4</sub> pH7.4, 20mM NaCl, 400mM imidazole and 0.1% CHAPS after an extra wash step with 50 mM imidazole. The elution fraction was diluted 1/20 with 20 mM Tris pH 8.0 and 0.1% CHAPS and loaded on a 20-mL Source 15Q column (GE Healthcare, Belgium) to remove contaminants. After equilibration, the protein of interest was eluted by a linear gradient over 20 column volumes of NaCl from 0 to 1 M in 20 mM Tris pH 8.0 and 0.1% CHAPS. To the EphA4-containing fractions, activated murine caspase-3 (1/100 % m/m murine caspase-3/Epha4) with 10 mM DTT was added to remove the His<sub>6</sub>-tag. After 1 h incubation at 37°C, the reaction solution was injected on a HiLoad 26/60 Superdex 75 prep grade with PBS as running solution for formulation and to remove minor contaminants, His<sub>6</sub>-tag and murine caspase-3. The obtained fractions were analysed by SDS-PAGE and the concentration was determined using the Micro-BCA assay (ThermoFisher Scientific, Belgium).

**Construction of a VHH library** - Nbs targeting the EphA4 LBD were obtained in collaboration with the VIB Nanobody Service Facility. An alpaca was injected subcutaneously on days 0, 7, 14, 21, 28 and 35, each time with 250 µg of human EphA4 LBD. On day 39 anticoagulated blood was collected for lymphocyte preparation. A VHH library was constructed as previously described (72-74) and screened for the presence of antigen-specific Nbs. In brief, total RNA from peripheral blood lymphocytes was used as template for first strand cDNA synthesis with oligo(dT) primer. Using this cDNA, the VHH encoding sequences were amplified by PCR, digested with PstI and NotI, and cloned into the PstI and NotI sites of the phagemid vector pMECS. A VHH library of about 2 × 10<sup>8</sup> independent transformants was obtained.

#### **Isolation of hEphA4 LBD-specific Nbs -**

To screen for the presence of EphA4-specific Nbs, the library was subjected to 4 consecutive rounds of panning, performed on solid-phase coated EphA4 LBD (concentration: ~200 mg/ml, ~20 mg/well). The enrichment for antigen-specific phages after each round of panning was assessed by comparing the number of phages eluted from antigen-coated wells with the number of phages eluted from negative control (only-blocked) wells. These experiments suggested that the phage population was enriched for antigen-specific phages after the 3<sup>rd</sup> and 4<sup>th</sup> round of panning. In total, 190 individual colonies from the 3<sup>rd</sup> and 4<sup>th</sup> round (95 from each round) were randomly selected and analysed by ELISA for the presence of antigen-specific Nbs in their periplasmic extracts (ELISA using crude periplasmic extracts including soluble Nbs). Of 190 colonies, 41 colonies (14 and 27 from the 3<sup>rd</sup> and 4<sup>th</sup> round, respectively) scored positive in this assay. The VHH genes of the selected genes were sequenced to identify the different Nbs.

**Expression and purification of recombinant Nbs** - pMECS vectors harbouring Nb genes were transformed into WK6 *E. coli* cells and the transformed cells were grown in Luria Bertani medium supplemented with ampicillin (100 µg/mL) and 0.1 % glucose at 37°C overnight. Subsequently, the cultures were inoculated 1/100 to have 1 liter productions in Terrific Broth medium supplemented with ampicillin (100 µg/mL) and 0.1 % glucose in baffles shake flasks. The temperature was kept at 37°C. The cells were grown to an optical density of A<sub>600nm</sub> = 1.0, transferred at 28°C, and expression was induced by addition of 1 mM isopropyl β-D-1-thiogalactopyranoside overnight. Cells were then harvested and frozen at -20°C. The expressed Nbs were extracted from the periplasm by osmotic shock and purified using His GraviTrap (GE Healthcare, Belgium) in parallel, equilibrated with 20 mM NaH<sub>2</sub>PO<sub>4</sub>, pH 7.5, 300 mM NaCl, 20 mM imidazole, and 1 mM PMSF. After loading, the columns were washed with 20 column volumes of the same buffer. The Nbs were first eluted with 20 mM NaH<sub>2</sub>PO<sub>4</sub>, pH 7.5, 20 mM NaCl, 50 mM imidazole, 1 mM PMSF and then with 400 mM imidazole in the same buffer. Finally, the Nbs were desalted to PBS on sephadex G25 (GE Healthcare, Belgium). The obtained fractions were analysed with Coomassie-stained SDS-polyacrylamide gels. Protein concentration was



measured by the Micro-BCA assay (ThermoFisher Scientific, Belgium).

**Western Blot** – For immunoprecipitation experiments, 1  $\mu\text{g}$  of EphA4 LBD, hEphA4-Fc (R&D, UK) and mEphA4-Fc (R&D, UK) were boiled for 10 min in reducing sample buffer (ThermoFisher Scientific, Belgium), and proteins were separated in a 4-12% Bis-Tris SDS-PAGE gel (ThermoFisher Scientific, Belgium). After SDS-PAGE the gel was transferred to Immobilon-P membrane (Merck Millipore, Belgium) and subsequently blocked with 10% Blotting-grade blocker (Biorad, Belgium) for 1 h at room temperature. One  $\mu\text{g}$  Nb and mouse anti-HA antibody 1/1000 (clone 16B12, MMS-101P-200, lot D13FF01646, Covance, US) were used to detect EphA4. To determine Nb stability, 10 ng of Nbs were denatured with reducing sample buffer (ThermoFisher Scientific, Belgium) and separated in a 12% Tris-glycine SDS-PAGE gel. The gel was transferred to Immobilon-P membrane, which was afterwards blocked as described above. To detect the Nbs, 1/1000 anti-HA antibody (clone C29F4, 3724S, lot 8, Cell Signaling, US) was used.

**Immunoprecipitation experiments** - One and a half mg protein G magnetic Dynabeads (ThermoFisher Scientific, Belgium) was preblocked with 1% BSA for 1 h at room temperature, washed four times with PBS and incubated with 2.5  $\mu\text{g}$  of recombinant mouse or human EphA4 protein (R&D, UK) for 10 min at room temperature. After washing four times with PBS the beads were incubated with 1  $\mu\text{g}$  Nb overnight at 4°C. The beads were washed four times with PBS, boiled for 10 min in reducing sample buffer (ThermoFisher Scientific, Belgium), and proteins were separated in a 4-12% Bis-Tris SDS-PAGE gel (ThermoFisher Scientific, Belgium). After SDS-PAGE the gel was transferred to Immobilon-P membrane (Merck Millipore, Belgium) and subsequently blocked with 10% Blotting-grade blocker (Biorad, Belgium) for 1 h at room temperature. Mouse N-terminal anti-EphA4 antibody 1/1000 (EM2801, lot 1, ECM Biosciences, US) and mouse anti-HA antibody 1/1000 (clone 16B12, MMS-101P-200, lot D13FF01646, Covance, US) were used to detect EphA4 and Nb respectively. To capture EphA4 with Nb anti-HA magnetic Dynabeads (ThermoFisher Scientific,

Belgium) were used and all following steps were performed as described above.

**Stability experiments** - To determine the Nb stability in plasma, Nbs were incubated at 37°C at a concentration of 10 ng/ $\mu\text{l}$  in sodium heparin not-filtered C57BL/6 mouse plasma (BioreclamationIVT, US). At every time-point an aliquot was collected and diluted 1/10 in PBS for further Western Blot analysis. As a control, Nbs were incubated for the same time and at the same concentration at 37°C in PBS. Nb stability in all samples was next determined with Western Blot as described above. Statistical analysis was performed using the two-way ANOVA followed by Sidak's multiple comparisons posthoc test as indicated in the figure legends. A 95 % confidence interval was used and values of  $P < 0.05$  were considered as statistically significant.

To determine functional stability of Nbs, they were first incubated at 37°C in sodium heparin not-filtered C57BL/6 mouse plasma at 500 nM concentration. At every time-point and for a maximum of 72 hours, an aliquot was collected and stored for further analysis with Alphascreen technology.

**Surface Plasmon Resonance** - The equilibrium dissociation constant ( $K_D$ ) and the association ( $k_{a1}$  and  $k_{a2}$ ) and dissociation rates ( $k_{d1}$  and  $k_{d2}$ ) were determined using surface plasmon resonance detection on a BIAcore T200 (GE Healthcare, Belgium). Two approaches were used. First, the extracellular N-terminal domain of human EphA4 was immobilized directly onto a CM5 S series sensor chip (GE Healthcare, Belgium) using standard amine coupling. After activation of the carboxyl moieties on the matrix on the chip surface with a 7-min injection of a 1:1 mixture of 0.4 M EDC and 0.1 M NHS, the N-terminal domain of human EphA4 (50  $\mu\text{g}/\text{ml}$  in 10 mM acetate, pH 4.5) was immobilized to a predefined level of 300 response units (RU). A flow channel activated with EDC/NHS and immediately afterwards blocked by ethanolamine served as reference channel. Throughout the analysis 10 mM Hepes, 150 mM NaCl, 3 mM EDTA, 0.01 mM Surfactant P-20 pH 7.35 was used as running buffer. 20 mM glycine pH 2.5 was used to regenerate the channels (remove all bound proteins). The second approach used the anti-human Fc capture kit as described by the manufacturer. Briefly, approximately 8000 RU

anti-human Fc antibody was captured on the surface of the channels using amine coupling as described above. In a next step, EphA4-hFc (both human and mouse EphA4) (R&D, UK) fusion protein (2 µg/ml diluted in running buffer) was injected (10 µl/min for 6 min) over the channel resulting in the capture of approximately 500 RU of EphA4-hFc fusion protein on the surface of the channel. In this approach, a channel with only anti-human Fc antibody served as reference. 3M MgCl<sub>2</sub> was used as regeneration buffer. The Nbs were diluted to the indicated concentrations in running buffer and injected (60 µl/min) over the channel with immobilized EphA4 and the reference channel. After correction of the response using the responses from the reference channel and a blank injection of running buffer over the EphA4-immobilized channel (double referencing), kinetic parameters were determined using Biacore T200 evaluation software (GE Healthcare, Belgium). Interactions of Nbs with different EphA4 recombinant proteins were calculated with a 1:1 binding model or a two-state model. The latter model was used to calculate all Nb interactions with EphA4 LBD and interactions of Nb 34, 31 and 50 with human and mouse EphA4.

**AlphaScreen** - To test the specificity of the Nbs for the different Eph receptors all Nbs were biotinylated with a five times molar excess of EZ link NHS biotin (ThermoFisher Scientific, Belgium). The Nbs were incubated with the biotin for two hours on ice allowing the interaction of the biotin with the primary amines on the surface of the protein. To remove unbound biotin, dialysis was performed with PBS in the Slide-A-Lyzer mini dialysis device (10 kDa cutoff, ThermoFisher Scientific, Belgium). Ten microliter of biotinylated Nbs (100 nM) were incubated with 10 µl of a subhooking concentration of mouse recombinant Eph receptor (Fc-tagged; R&D, UK) for one hour in standard buffer (50 mM Hepes, 100 mM NaCl, 0.1% Triton and 0.1% BSA) in white opaque 384-well microplates (Perkin Elmer, Belgium) to avoid the hooking effect (oversaturation of donor and acceptor beads inhibiting their association). The determined subhooking concentrations were 10 nM for EphA2, A3, A4, A6, B2 and B6, 30 nM for EphA7 and B3, B4 and 100 nM for EphA8. Subsequently we incubated first one hour with

10 µl of anti-IgG AlphaLISA acceptor beads (20 µg/ml, Perkin Elmer, Belgium) and then incubated an additional 30 minutes with 10 µl streptavidin donor beads (20 µg/ml, Perkin Elmer, Belgium). Incubation steps were performed at room temperature and protected from light.

To test the inhibition of interaction between EphA4 and its different ligands, the EphA4 LBD (2 mg/ml) was biotinylated with a two and a half times molar excess of EZ link NHS biotin (ThermoFisher Scientific, Belgium) for two hours on ice allowing the interaction of the biotin with primary amines on the surface of the protein. To remove unbound biotin, dialysis was performed with PBS in the Slide-A-Lyzer mini dialysis device (10 kDa cutoff, ThermoFisher Scientific, Belgium). Five microliter of a subhooking concentration of biotinylated EphA4 LBD (10 nM) was incubated with 5 µl of different concentrations of Nb for one hour in standard buffer (50 mM Hepes, 100 mM NaCl, 0.1% Triton and 0.1% BSA). Subsequently, 5 µl of subhooking concentration of recombinant ephrin ligand (Fc-tagged; R&D, UK) was added and incubated for one hour at room temperature. The determined subhooking concentrations were 3 nM for ephrin-A1 and ephrin-A4, and 10 nM for ephrin-A2/3, ephrin-A5 and ephrin-B1-3. Next, 5 µl of anti-IgG AlphaLISA acceptor beads (20 µg/ml; Perkin Elmer, Belgium) and five microliter streptavidin donor beads (10 µg/ml; Perkin Elmer, Belgium) were added and incubated for one hour and 30 minutes respectively at room temperature, protected from light. A Nb targeting SOD1 was used as negative control in all Eph binding assays.

To test the inhibition of EphA7 and ephrin-A5 interaction, His-tagged human ephrin-A5 (Sino Biological, China) was biotinylated with a five times molar excess of EZ link NHS biotin (ThermoFisher Scientific, Belgium), as it is described for the Nbs. Briefly, ephrin-A5 was incubated with the biotin for two hours on ice, and dialysed in PBS with the Slide-A-Lyzer mini dialysis device (10 kDa cutoff, ThermoFisher Scientific, Belgium), to remove unbound biotin. microliter of a subhooking concentration of human EphA7 (3 nM) was incubated with 5 µl of different concentrations of Nb for one hour in standard buffer (50 mM Hepes, 100 mM NaCl, 0.1% Triton and 0.1% BSA). Next, 5 µl of subhooking concentration of

biotinylated ephrin-A5 ligand (30 nM) was added and incubated for one hour at room temperature. Finally, 5  $\mu$ l of anti-IgG AlphaLISA acceptor beads (20  $\mu$ g/ml; Perkin Elmer, Belgium) and 5  $\mu$ l streptavidin donor beads (20  $\mu$ g/ml; Perkin Elmer, Belgium) were added and incubated for one hour and 30 minutes respectively at room temperature, protected from light. A control Nb targeting chicken lysozyme was used as negative control, and untagged ephrin-A5 was used as positive control.

In order to determine the capability of Nbs to bind EphA4 after plasma incubation, 500 nM Nbs pre-incubated in plasma were first diluted in PBS at the time of the Alphascreen assay, to reach a final subhooking Nb concentration of 10 nM. Five microliter of a subhooking concentration of biotinylated EphA4 LBD (10 nM) was incubated with 5  $\mu$ l of 10 nM Nb 39 and Nb 53 for one hour in standard buffer (50 mM Hepes, 100 mM NaCl, 0.1% Triton and 0.1% BSA). Next, 5  $\mu$ l of anti-HA AlphaLISA acceptor beads (20  $\mu$ g/ml; Perkin Elmer, Belgium) and five microliter streptavidin donor beads (10  $\mu$ g/ml; Perkin Elmer, Belgium) were added and incubated for one hour and 30 minutes respectively at room temperature, protected from light. As a control to assess the specificity of the binding, 300 nM of ephrin-B2 ligand ((Fc-tagged; R&D, UK) was incubated with EphA4 LBD for one hour prior to the addition of the Nbs.

Plates were read on the Envision Multilabel Reader (Perkin Elmer, Belgium).

**Phosphorylation assay** - The amount of phosphorylation of EphA4 was determined using the PathHunter assay (DiscoverX Corporation, UK) with U2OS cells adapted for the EphA4 receptor according to manufacturer's instructions. In short, a small peptide epitope is expressed recombinantly on the intracellular C-terminus of the EphA4 receptor tyrosine kinase. An interaction partner containing SH2 domains is co-expressed with a larger sequence, termed enzyme activator (EA). Activation with recombinant human ephrin-A1-Fc causes EphA4 receptor dimerisation, leading to cross-phosphorylation of tyrosine residues on the cytoplasmic domain of the receptor. The SH2-EA fusion protein binds the phosphorylated receptor enabling the complementation of EA and the peptide epitope, yielding an active  $\beta$ -

galactosidase enzyme. This interaction can be visualised with a chemiluminescent substrate. Increasing concentrations of the Nbs were added to the medium before ephrin-A1-Fc stimulation. 20  $\mu$ l of 5000 EphA4-expressing U2OS cells were plated in 384-well plates and incubated for 24 h at 37°C 5% CO<sub>2</sub>. Five  $\mu$ l Nb dilutions or vehicle were added per well followed by incubation for 1 h at 37°C. Five  $\mu$ l of ephrin-A1-Fc (1.2  $\mu$ g/ml) or vehicle were added to each well followed by incubation for 3 h at room temperature. Twelve  $\mu$ l of detection reagent (Galacton Star, Emerald II solution, PathHunter cell assay buffer in relative volumes of 1:5:19 respectively) was added, incubated for 1 h at room temperature and read on a Pherastar (BMG Labtech, Germany). Percentage activity was calculated as (signal - non-stimulated control (no ephrin added)) / (ephrin stimulated condition - non-stimulated control) \* 100. Positive and negative controls were Dasatinib and a Nb targeting Superoxide Dismutase 1 (SOD1) respectively.

**Growth cone collapse** - The cortex of 7.5-day-old mice was dissociated by trypsinisation and trituration. Cortical neurons were cultured on 13 mm-diameter cover-slips coated with poly-L-Lysine, at a density of 15000 cells/cover-slip, in MEM medium with Earle's salt and L-glutamine and supplemented with 10% heat-inactivated horse serum and penicillin and streptomycin (ThermoFisher Scientific, Belgium). Cultures were kept in a 5% CO<sub>2</sub> humidified incubator at 37°C. Medium was replaced for Neurobasal medium supplemented with B27, 500  $\mu$ M L-glutamine and penicillin and streptomycin (ThermoFisher Scientific), 4 hours after plating. 24 hours after plating the cultures were incubated for 30 min with KYL peptide at a concentration of 1  $\mu$ M or 30  $\mu$ M, or Nb 53 or Nb 39 at a concentration of 1  $\mu$ M. Cultures were next stimulated with 1  $\mu$ g/ml mouse pre-clustered ephrin-B3-Fc, or Fc (R&D, UK) as a control, for 30 min in the presence of the KYL peptide or the Nbs. Cortical neurons were fixed for 20 min in 4% paraformaldehyde (ThermoFisher Scientific, Belgium), permeabilized in 0.2% Triton X-100 in PBS and stained with Alexa Fluor 555-conjugated phalloidin (ThermoFisher Scientific, Belgium). Growth cone collapse was scored for each condition under a Zeiss Axioimager M1 epifluorescence and brightfield upright

microscope with a Zeiss A-Plan 40X/0.65  $\infty$ /0.17 objective (Carl Zeiss, Germany). Growth cone collapse was considered when no lamellipodia or filopodia were present at the tip of the longest neurite of every scored neuron. Growth cone collapse was scored in a blinded manner for 40-160 neurons in every condition and experiment. Images were obtained with AxioCam MRm monochrome digital camera and Zeiss

AxioVision V 4.8.2.0 software (Carl Zeiss, Germany) at the same magnification. Statistical analysis was performed using the one-way ANOVA followed by Tukey's post hoc test as indicated in the figure legends. A 95 % confidence interval was used and values of  $P < 0.05$  were considered as statistically significant.

## **Acknowledgments**

This work has been supported by grants from the Flanders Institute for Biotechnology (VIB), the University of Leuven (GOA/11/014), Vertex Pharmaceuticals Incorporated, Research Foundation Flanders (FWO-Vlaanderen; G.0996.14N), the ALS League Belgium, the Thierry Latran Foundation, Geneeskundige Stichting Koningin Elisabeth (G.S.K.E.), Laevers Fund for ALS Research, Fund ‘Een Hart voor ALS’. The research leading to these results has received funding from the European Community’s Health Seventh Framework Programme (FP7/2007-2013) under grant agreement number 259867, the European Research Council and the European’s Seventh Framework Programme (FP7/2007-2013)/ ERC grant agreement n° 340429 and structural funding for medium-scale research infrastructure (HERCULES, AUGE/09/040). LS and BD have received funding from the Agency for Innovation by Science and Technology in Flanders (IWT-Vlaanderen; projects 101318 and IWT-VIND 135043 respectively). RL and PVD hold a clinical investigatorship of FWO-Vlaanderen. RL is supported by Research Fund KU Leuven. WR, PVD, RL and LVDB are supported by the “Opening the Future” Fund and WR by the E. von Behring Chair for Neuromuscular and Neurodegenerative Disorders, both from the University of Leuven. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health. The authors gratefully acknowledge the assistance of An Snellinx, Katrien Horr  and Kathleen Craessaerts for the preparation of the mouse cortical neuron cultures.

## **Conflict of interest**

The authors declare that they have no conflicts of interest with the contents of this article.

## **Author contributions**

LS and LR designed, performed and analysed most of the experiments and wrote the manuscript. BR, MT, SL, AC, PJ and PW designed, performed and analysed some experiments. LCG and MD assisted with the analysis of results. JH expressed, purified and isolated the N-terminal EpA4 protein and the Nbs. GHZG generated the anti-EphA4 LBD Nbs. BD, PVD, LVDB, RL supervised the experiments and contributed to the analysis. WR conceived the idea for the project, supervised the experiments and wrote the manuscript. All authors contributed to the final manuscript.

1. Pitulescu, M. E., and Adams, R. H. (2010) Eph/ephrin molecules--a hub for signaling and endocytosis. *Genes & development* **24**, 2480-2492
2. Bowden, T. A., Aricescu, A. R., Nettleship, J. E., Siebold, C., Rahman-Huq, N., Owens, R. J., Stuart, D. I., and Jones, E. Y. (2009) Structural plasticity of eph receptor A4 facilitates cross-class ephrin signaling. *Structure* **17**, 1386-1397
3. Himanen, J. P., Chumley, M. J., Lackmann, M., Li, C., Barton, W. A., Jeffrey, P. D., Vearing, C., Geleick, D., Feldheim, D. A., Boyd, A. W., Henkemeyer, M., and Nikolov, D. B. (2004) Repelling class discrimination: ephrin-A5 binds to and activates EphB2 receptor signaling. *Nat Neurosci* **7**, 501-509
4. Qin, H., Noberini, R., Huan, X., Shi, J., Pasquale, E. B., and Song, J. (2010) Structural characterization of the EphA4-Ephrin-B2 complex reveals new features enabling Eph-ephrin binding promiscuity. *J Biol Chem* **285**, 644-654
5. Boyd, A. W., Bartlett, P. F., and Lackmann, M. (2014) Therapeutic targeting of EPH receptors and their ligands. *Nature Reviews Drug Discovery* **13**, 39-62
6. Pasquale, E. B. (2008) Eph-ephrin bidirectional signaling in physiology and disease. *Cell* **133**, 38-52
7. Frugier, T., Conquest, A., McLean, C., Currie, P., Moses, D., and Goldshmit, Y. (2012) Expression and activation of EphA4 in the human brain after traumatic injury. *J Neuropathol Exp Neurol* **71**, 242-250
8. Goldshmit, Y., Galea, M. P., Wise, G., Bartlett, P. F., and Turnley, A. M. (2004) Axonal regeneration and lack of astrocytic gliosis in EphA4-deficient mice. *J Neurosci* **24**, 10064-10073
9. Lemmens, R., Jaspers, T., Robberecht, W., and Thijs, V. N. (2013) Modifying expression of EphA4 and its downstream targets improves functional recovery after stroke. *Hum Mol Genet* **22**, 2214-2220
10. Fu, A. K. Y., Hung, K. W., Huang, H. Q., Gu, S., Shen, Y., Cheng, E. Y. L., Ip, F. C. F., Huang, X. H., Fu, W. Y., and Ip, N. Y. (2014) Blockade of EphA4 signaling ameliorates hippocampal synaptic dysfunctions in mouse models of Alzheimer's disease. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 9959-9964
11. Van Hoecke, A., Schoonaert, L., Lemmens, R., Timmers, M., Staats, K. A., Laird, A. S., Peeters, E., Philips, T., Goris, A., Dubois, B., Andersen, P. M., Al-Chalabi, A., Thijs, V., Turnley, A. M., van Vught, P. W., Veldink, J. H., Hardiman, O., Van Den Bosch, L., Gonzalez-Perez, P., Van Damme, P., Brown, R. H., van den Berg, L. H., and Robberecht, W. (2012) EPHA4 is a disease modifier of amyotrophic lateral sclerosis in animal models and in humans. *Nature Medicine* **18**, 1418+
12. Vargas, L. M., Leal, N., Estrada, L. D., Gonzalez, A., Serrano, F., Araya, K., Gysling, K., Inestrosa, N. C., Pasquale, E. B., and Alvarez, A. R. (2014) EphA4 activation of c-Abl mediates synaptic loss and LTP blockade caused by amyloid-beta oligomers. *PLoS One* **9**, e92309
13. Noberini, R., Lamberto, I., and Pasquale, E. B. (2012) Targeting Eph receptors with peptides and small molecules: progress and challenges. *Semin Cell Dev Biol* **23**, 51-57
14. Tognolini, M., Hassan-Mohamed, I., Giorgio, C., Zanotti, I., and Lodola, A. (2014) Therapeutic perspectives of Eph-ephrin system modulation. *Drug Discov Today* **19**, 661-669
15. Guo, F. Y., and Lesk, A. M. (2014) Sizes of interface residues account for cross-class binding affinity patterns in Eph receptor-ephrin families. *Proteins* **82**, 349-353
16. Lamberto, I., Lechtenberg, B. C., Olson, E. J., Mace, P. D., Dawson, P. E., Riedl, S. J., and Pasquale, E. B. (2014) Development and structural analysis of a nanomolar cyclic peptide antagonist for the EphA4 receptor. *ACS Chem Biol* **9**, 2787-2795
17. Singla, N., Goldgur, Y., Xu, K., Paavilainen, S., Nikolov, D. B., and Himanen, J. P. (2010) Crystal structure of the ligand-binding domain of the promiscuous EphA4 receptor reveals two distinct conformations. *Biochem Biophys Res Commun* **399**, 555-559

18. Han, X., Xu, Y., Yang, Y., Xi, J., Tian, W., Duggineni, S., Huang, Z., and An, J. (2013) Discovery and characterization of a novel cyclic peptide that effectively inhibits ephrin binding to the EphA4 receptor and displays anti-angiogenesis activity. *PLoS One* **8**, e80183
19. Lamberto, I., Qin, H., Noberini, R., Premkumar, L., Bourgin, C., Riedl, S. J., Song, J., and Pasquale, E. B. (2012) Distinctive binding of three antagonistic peptides to the ephrin-binding pocket of the EphA4 receptor. *Biochem J* **445**, 47-56
20. Noberini, R., Koolpe, M., Peddibhotla, S., Dahl, R., Su, Y., Cosford, N. D., Roth, G. P., and Pasquale, E. B. (2008) Small molecules can selectively inhibit ephrin binding to the EphA4 and EphA2 receptors. *J Biol Chem* **283**, 29461-29472
21. Wu, B., Zhang, Z., Noberini, R., Barile, E., Giulianotti, M., Pinilla, C., Houghten, R. A., Pasquale, E. B., and Pellecchia, M. (2013) HTS by NMR of combinatorial libraries: a fragment-based approach to ligand discovery. *Chem Biol* **20**, 19-33
22. Olson, E. J., Lechtenberg, B. C., Zhao, C., Rubio de la Torre, E., Lamberto, I., Riedl, S. J., Dawson, P. E., and Pasquale, E. B. (2016) Modifications of a Nanomolar Cyclic Peptide Antagonist for the EphA4 Receptor To Achieve High Plasma Stability. *ACS Med Chem Lett* **7**, 841-846
23. Fabes, J., Anderson, P., Brennan, C., and Bolsover, S. (2007) Regeneration-enhancing effects of EphA4 blocking peptide following corticospinal tract injury in adult rat spinal cord. *Eur J Neurosci* **26**, 2496-2505
24. Murai, K. K., Nguyen, L. N., Koolpe, M., McLennan, R., Krull, C. E., and Pasquale, E. B. (2003) Targeting the EphA4 receptor in the nervous system with biologically active peptides. *Mol Cell Neurosci* **24**, 1000-1011
25. De Meyer, T., Muyldermans, S., and Depicker, A. (2014) Nanobody-based products as research and diagnostic tools. *Trends Biotechnol* **32**, 263-270
26. Hamers-Casterman, C., Atarhouch, T., Muyldermans, S., Robinson, G., Hamers, C., Songa, E. B., Bendahman, N., and Hamers, R. (1993) Naturally occurring antibodies devoid of light chains. *Nature* **363**, 446-448
27. Muyldermans, S. (2013) Nanobodies: natural single-domain antibodies. *Annu Rev Biochem* **82**, 775-797
28. Steyaert, J., and Kobilka, B. K. (2011) Nanobody stabilization of G protein-coupled receptor conformational states. *Curr Opin Struct Biol* **21**, 567-572
29. Harmsen, M. M., and De Haard, H. J. (2007) Properties, production, and applications of camelid single-domain antibody fragments. *Appl Microbiol Biotechnol* **77**, 13-22
30. Truitt, L., and Freywald, A. (2011) Dancing with the dead: Eph receptors and their kinase-null partners. *Biochem Cell Biol* **89**, 115-129
31. Binns, K. L., Taylor, P. P., Sicheri, F., Pawson, T., and Holland, S. J. (2000) Phosphorylation of tyrosine residues in the kinase domain and juxtamembrane region regulates the biological and catalytic activities of Eph receptors. *Molecular and cellular biology* **20**, 4791-4805
32. Lisabeth, E. M., Falivelli, G., and Pasquale, E. B. (2013) Eph receptor signaling and ephrins. *Cold Spring Harbor perspectives in biology* **5**
33. Kania, A., and Klein, R. (2016) Mechanisms of ephrin-Eph signalling in development, physiology and disease. *Nature reviews. Molecular cell biology* **17**, 240-256
34. Dent, E. W., Gupton, S. L., and Gertler, F. B. (2011) The growth cone cytoskeleton in axon outgrowth and guidance. *Cold Spring Harbor perspectives in biology* **3**
35. Kullander, K., Croll, S. D., Zimmer, M., Pan, L., McClain, J., Hughes, V., Zabski, S., DeChiara, T. M., Klein, R., Yancopoulos, G. D., and Gale, N. W. (2001) Ephrin-B3 is the midline barrier that prevents corticospinal tract axons from recrossing, allowing for unilateral motor control. *Genes & development* **15**, 877-888
36. Egea, J., Nissen, U. V., Dufour, A., Sahin, M., Greer, P., Kullander, K., Mrcic-Flogel, T. D., Greenberg, M. E., Kiehn, O., Vanderhaeghen, P., and Klein, R. (2005) Regulation of EphA 4

- kinase activity is required for a subset of axon guidance decisions suggesting a key role for receptor clustering in Eph function. *Neuron* **47**, 515-528
37. Rasmussen, S. G., Choi, H. J., Fung, J. J., Pardon, E., Casarosa, P., Chae, P. S., Devree, B. T., Rosenbaum, D. M., Thian, F. S., Kobilka, T. S., Schnapp, A., Konetzki, I., Sunahara, R. K., Gellman, S. H., Pautsch, A., Steyaert, J., Weis, W. I., and Kobilka, B. K. (2011) Structure of a nanobody-stabilized active state of the beta(2) adrenoceptor. *Nature* **469**, 175-180
  38. Xu, K., Tzvetkova-Robev, D., Xu, Y., Goldgur, Y., Chan, Y. P., Himanen, J. P., and Nikolov, D. B. (2013) Insights into Eph receptor tyrosine kinase activation from crystal structures of the EphA4 ectodomain and its complex with ephrin-A5. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 14634-14639
  39. Figueroa, J. D., Benton, R. L., Velazquez, I., Torrado, A. I., Ortiz, C. M., Hernandez, C. M., Diaz, J. J., Magnuson, D. S., Whittemore, S. R., and Miranda, J. D. (2006) Inhibition of EphA7 up-regulation after spinal cord injury reduces apoptosis and promotes locomotor recovery. *Journal of Neuroscience Research* **84**, 1438-1451
  40. Saerens, D., Ghassabeh, G. H., and Muyldermans, S. (2008) Single-domain antibodies as building blocks for novel therapeutics. *Curr Opin Pharmacol* **8**, 600-608
  41. Sheedy, C., MacKenzie, C. R., and Hall, J. C. (2007) Isolation and affinity maturation of hapten-specific antibodies. *Biotechnol Adv* **25**, 333-352
  42. Vearing, C., Lee, F. T., Wimmer-Kleikamp, S., Spirkoska, V., To, C., Stylianou, C., Spanevello, M., Brechbiel, M., Boyd, A. W., Scott, A. M., and Lackmann, M. (2005) Concurrent binding of anti-EphA3 antibody and ephrin-A5 amplifies EphA3 signaling and downstream responses: potential as EphA3-specific tumor-targeting reagents. *Cancer Res* **65**, 6745-6754
  43. Martone, M. E., Holash, J. A., Bayardo, A., Pasquale, E. B., and Ellisman, M. H. (1997) Immunolocalization of the receptor tyrosine kinase EphA4 in the adult rat central nervous system. *Brain Res* **771**, 238-250
  44. Tremblay, M. E., Riad, M., Bouvier, D., Murai, K. K., Pasquale, E. B., Descarries, L., and Doucet, G. (2007) Localization of EphA4 in axon terminals and dendritic spines of adult rat hippocampus. *The Journal of comparative neurology* **501**, 691-702
  45. Fabes, J., Anderson, P., Yanez-Munoz, R. J., Thrasher, A., Brennan, C., and Bolsover, S. (2006) Accumulation of the inhibitory receptor EphA4 may prevent regeneration of corticospinal tract axons following lesion. *The European journal of neuroscience* **23**, 1721-1730
  46. Li, S., Overman, J. J., Katsman, D., Kozlov, S. V., Donnelly, C. J., Twiss, J. L., Giger, R. J., Coppola, G., Geschwind, D. H., and Carmichael, S. T. (2010) An age-related sprouting transcriptome provides molecular control of axonal sprouting after stroke. *Nat Neurosci* **13**, 1496-1504
  47. Ilieva, H., Polymenidou, M., and Cleveland, D. W. (2009) Non-cell autonomous toxicity in neurodegenerative disorders: ALS and beyond. *J Cell Biol* **187**, 761-772
  48. Bundesen, L. Q., Scheel, T. A., Bregman, B. S., and Kromer, L. F. (2003) Ephrin-B2 and EphB2 regulation of astrocyte-meningeal fibroblast interactions in response to spinal cord lesions in adult rats. *Journal of Neuroscience* **23**, 7789-7800
  49. Overman, J. J., Clarkson, A. N., Wanner, I. B., Overman, W. T., Eckstein, I., Maguire, J. L., Dinov, I. D., Toga, A. W., and Carmichael, S. T. (2012) A role for ephrin-A5 in axonal sprouting, recovery, and activity-dependent plasticity after stroke. *Proceedings of the National Academy of Sciences of the United States of America* **109**, E2230-2239
  50. Ren, Z., Chen, X., Yang, J., Kress, B. T., Tong, J., Liu, H., Takano, T., Zhao, Y., and Nedergaard, M. (2013) Improved axonal regeneration after spinal cord injury in mice with conditional deletion of ephrin B2 under the GFAP promoter. *Neuroscience* **241**, 89-99
  51. Bradl, M., and Lassmann, H. (2010) Oligodendrocytes: biology and pathology. *Acta Neuropathol* **119**, 37-53



52. Benson, M. D., Romero, M. I., Lush, M. E., Lu, Q. R., Henkemeyer, M., and Parada, L. F. (2005) Ephrin-B3 is a myelin-based inhibitor of neurite outgrowth. *Proc Natl Acad Sci U S A* **102**, 10694-10699
53. Linneberg, C., Harboe, M., and Laursen, L. S. (2015) Axo-Glia Interaction Preceding CNS Myelination Is Regulated by Bidirectional Eph-Ephrin Signaling. *ASN Neuro* **7**
54. Prestoz, L., Chatzopoulou, E., Lemkine, G., Spassky, N., Lebras, B., Kagawa, T., Ikenaka, K., Zalc, B., and Thomas, J. L. (2004) Control of axonophilic migration of oligodendrocyte precursor cells by Eph-ephrin interaction. *Neuron Glia Biol* **1**, 73-83
55. Duffy, P., Wang, X., Siegel, C. S., Tu, N., Henkemeyer, M., Cafferty, W. B., and Strittmatter, S. M. (2012) Myelin-derived ephrinB3 restricts axonal regeneration and recovery after adult CNS injury. *Proc Natl Acad Sci U S A* **109**, 5063-5068
56. Syed, Y. A., Zhao, C., Mahad, D., Mobius, W., Altmann, F., Foss, F., Senturk, A., Acker-Palmer, A., Lubec, G., Lilley, K., Franklin, R. J., Nave, K. A., and Kotter, M. R. (2016) Antibody-mediated neutralization of myelin-associated EphrinB3 accelerates CNS remyelination. *Acta Neuropathol* **131**, 281-298
57. Garbuzova-Davis, S., and Sanberg, P. R. (2014) Blood-CNS Barrier Impairment in ALS patients versus an animal model. *Front Cell Neurosci* **8**, 21
58. Garbuzova-Davis, S., Saporta, S., Haller, E., Kolomey, I., Bennett, S. P., Potter, H., and Sanberg, P. R. (2007) Evidence of compromised blood-spinal cord barrier in early and late symptomatic SOD1 mice modeling ALS. *PLoS one* **2**, e1205
59. Miyazaki, K., Ohta, Y., Nagai, M., Morimoto, N., Kurata, T., Takehisa, Y., Ikeda, Y., Matsuura, T., and Abe, K. (2011) Disruption of neurovascular unit prior to motor neuron degeneration in amyotrophic lateral sclerosis. *J Neurosci Res* **89**, 718-728
60. Abulrob, A., Sprong, H., Van Bergen en Henegouwen, P., and Stanimirovic, D. (2005) The blood-brain barrier transmutating single domain antibody: mechanisms of transport and antigenic epitopes in human brain endothelial cells. *Journal of Neurochemistry* **95**, 1201-1214
61. Boado, R. J., Hui, E. K., Lu, J. Z., and Pardridge, W. M. (2012) Glycemic control and chronic dosing of rhesus monkeys with a fusion protein of iduronidase and a monoclonal antibody against the human insulin receptor. *Drug Metabolism and Disposition: The Biological Fate of Chemicals* **40**, 2021-2025
62. Farrington, G. K., Caram-Salas, N., Haqqani, A. S., Brunette, E., Eldredge, J., Pepinsky, B., Antognetti, G., Baumann, E., Ding, W., Garber, E., Jiang, S., Delaney, C., Boileau, E., Sisk, W. P., and Stanimirovic, D. B. (2014) A novel platform for engineering blood-brain barrier-crossing bispecific biologics. *FASEB Journal* **28**, 4764-4778
63. Rissiek, B., Koch-Nolte, F., and Magnus, T. (2014) Nanobodies as modulators of inflammation: potential applications for acute brain injury. *Front Cell Neurosci* **8**, 344
64. Wang, D., El-Amouri, S. S., Dai, M., Kuan, C. Y., Hui, D. Y., Brady, R. O., and Pan, D. (2013) Engineering a lysosomal enzyme with a derivative of receptor-binding domain of apoE enables delivery across the blood-brain barrier. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 2999-3004
65. Xiao, G., and Gan, L. S. (2013) Receptor-mediated endocytosis and brain delivery of therapeutic biologics. *Int J Cell Biol* **2013**, 703545
66. Li, T., Bourgeois, J. P., Celli, S., Glacial, F., Le Sourd, A. M., Mecheri, S., Weksler, B., Romero, I., Couraud, P. O., Rougeon, F., and Lafaye, P. (2012) Cell-penetrating anti-GFAP VHH and corresponding fluorescent fusion protein VHH-GFP spontaneously cross the blood-brain barrier and specifically recognize astrocytes: application to brain imaging. *FASEB Journal* **26**, 3969-3979
67. Coppieters, K., Dreier, T., Silence, K., de Haard, H., Lauwereys, M., Casteels, P., Beirnaert, E., Jonckheere, H., Van de Wiele, C., Staelens, L., Hostens, J., Revets, H., Remaut, E., Elewaut, D., and Rottiers, P. (2006) Formatted anti-tumor necrosis factor alpha VHH proteins derived

- from camelids show superior potency and targeting to inflamed joints in a murine model of collagen-induced arthritis. *Arthritis Rheum* **54**, 1856-1866
68. Dennis, M. S., Zhang, M., Meng, Y. G., Kadkhodayan, M., Kirchofer, D., Combs, D., and Damico, L. A. (2002) Albumin binding as a general strategy for improving the pharmacokinetics of proteins. *J Biol Chem* **277**, 35035-35043
  69. Dixon, F. J., Maurer, P. H., and Deichmiller, M. P. (1953) Half-lives of homologous serum albumins in several species. *Proc Soc Exp Biol Med* **83**, 287-288
  70. Veronese, F. M., and Mero, A. (2008) The impact of PEGylation on biological therapies. *Biodrugs* **22**, 315-329
  71. Chapman, A. P. (2002) PEGylated antibodies and antibody fragments for improved therapy: a review. *Adv Drug Deliv Rev* **54**, 531-545
  72. Els Conrath, K., Lauwereys, M., Wyns, L., and Muyldermans, S. (2001) Camel single-domain antibodies as modular building units in bispecific and bivalent antibody constructs. *J Biol Chem* **276**, 7346-7350
  73. Saerens, D., Kinne, J., Bosmans, E., Wernery, U., Muyldermans, S., and Conrath, K. (2004) Single domain antibodies derived from dromedary lymph node and peripheral blood lymphocytes sensing conformational variants of prostate-specific antigen. *J Biol Chem* **279**, 51965-51972
  74. Steeland, S., Puimege, L., Vandenbroucke, R. E., Van Hauwermeiren, F., Hastraete, J., Devoogdt, N., Hulpiau, P., Leroux-Roels, G., Laukens, D., Meuleman, P., De Vos, M., and Libert, C. (2015) Generation and characterization of small single domain antibodies inhibiting human tumor necrosis factor receptor 1. *J Biol Chem* **290**, 4022-4037

**Table 1. Overview of the EphA4-Nanobody binding constants.** Association constants ( $k_{a1}$  and  $k_{a2}$ ), dissociation constants ( $k_{d1}$  and  $k_{d2}$ ) and equilibration constants ( $K_D$ ) were determined by Surface Plasmon Resonance (SPR) through kinetics analysis. SPR analyses were performed in duplicate. Alternating red and black colour indicates different Nb groups, based on their clonal origin.

	EphA4 LBD					Human EphA4					Mouse EphA4				
	$k_{a1}$ (1/Ms)	$k_{d1}$ (1/s)	$k_{a2}$ (1/s)	$k_{d2}$ (1/s)	$K_D$ (nM)	$k_{a1}$ (1/Ms)	$k_{d1}$ (1/s)	$k_{a2}$ (1/s)	$k_{d2}$ (1/s)	$K_D$ (nM)	$k_{a1}$ (1/Ms)	$k_{d1}$ (1/s)	$k_{a2}$ (1/s)	$k_{d2}$ (1/s)	$K_D$ (nM)
Nb 60	1.73 x 10 <sup>6</sup>	0.11484	0.00014	0.00018	36.96	1.23 x 10 <sup>6</sup>	0.04885			39.58	2.70 x 10 <sup>9</sup>	74.05041			27.41
Nb 19	1.82 x 10 <sup>5</sup>	0.10771	0.00480	0.02886	506.93	1.07 x 10 <sup>5</sup>	0.09097			851.56	1.18 x 10 <sup>5</sup>	0.13456			1143
Nb 57	3.85 x 10 <sup>6</sup>	0.25686	0.00021	0.00133	57.52	5.97 x 10 <sup>6</sup>	0.22869			38.31	7.63 x 10 <sup>6</sup>	0.28821			37.79
Nb 39	1.01 x 10 <sup>6</sup>	0.06067	0.00011	0.00186	56.56	9.65 x 10 <sup>5</sup>	0.02906			30.13	6.71 x 10 <sup>5</sup>	0.01582			23.59
Nb 34	/	/	/	/	/	5807.55	0.16824	0.00734	0.01209	18030	8245.13	0.34532	0.01260	0.02262	26900
Nb 47	Hardly any signal above background														
Nb 22	2.32 x 10 <sup>5</sup>	0.01433	0.00036	0.00255	54.10	3.14 x 10 <sup>5</sup>	0.01477			47.08	2.21 x 10 <sup>5</sup>	0.02299			104.08
Nb 31	3.47 x 10 <sup>5</sup>	0.15483	0.00355	0.00676	293.04	7.08 x 10 <sup>5</sup>	0.20028	0.00647	0.01046	174.95	4.22 x 10 <sup>5</sup>	0.15132	0.00447	0.01155	258.45
Nb 50	1.61 x 10 <sup>6</sup>	0.19233	0.00023	0.00062	87.33	2.41 x 10 <sup>6</sup>	0.16758	0.00018	0.00220	64.21	1.10 x 10 <sup>6</sup>	0.11397			103.68
Nb 53	1.24 x 10 <sup>6</sup>	0.05942	0.00011	0.00213	45.74	6.91 x 10 <sup>5</sup>	0.01563			22.62	1.07 x 10 <sup>6</sup>	0.02445			22.82

## FIGURE LEGENDS

**Figure 1. Amino acid sequence of 15 different anti-EphA4 LBD Nbs.** Nbs were numbered according to Kabat numbering with reference sequence VH3-23/JH5 (human germline sequence on average most closely related to Nanobody sequences) on top. The CDRs were assigned according to the AbM definitions and are shown with a grey background. The Nbs belonged to nine clonally-unrelated B-cell clones and the different groups are shown alternatively in black and purple. According to the amino acid sequences Nb 90 and 60; Nb 19 and 57; Nb 34 and 47; Nb 31, 50, 10 and 53 belonged to the same group. Nb 39, 16, 71, 28 and 22 belonged to unrelated B-cell clones.

**Figure 2. Nbs bind native EphA4 LBD.** Nbs and EphA4 co-immunoprecipitated together, both when immunoprecipitating different Nbs (left panel) or when immunoprecipitating EphA4 (right panel). (A, B) Nb 53 and Nb 39 could capture EphA4 and could be captured by EphA4. (C) Nb 22 showed lower affinity for mouse EphA4 compared to human EphA4. (D, E) Nb 19 and 47 showed unspecific binding to the beads. (F) Nb 60 showed cross-reactivity with ephrin-B2, which was performed as a negative control experiment.

**Figure 3. Selectivity for EphA4 and cross-reactivity with other Eph receptors.** Using AlphaScreen technology we determined the selectivity of the Nbs for EphA4 receptor and the cross-reactivity with other Eph receptors. (A) Next to EphA4, Nb 60 also interacted with many other different Eph receptors. (B-F) Nb 57, 22, 39, 31 and 53 presented the highest selectivity to the EphA4 receptor. (F) Nb 50 had the lowest selectivity for EphA4 as it also interacted with EphA3, EphA6, EphA7 and EphA8. All assays were done in duplicate and data are represented as mean  $\pm$  SD.

**Figure 4. Nbs compete with ephrin ligands in binding EphA4.** With AlphaScreen technology we determined the capability of the Nbs in inhibiting the binding to EphA4 LBD. (A, D, E, H) Nb 22 and 31 could not completely inhibit ephrin-ligand binding to EphA4 LBD, whereas Nb 39 and Nb 53 succeeded in doing so even at a lower concentration than the KYL-peptide, which was used as a positive control. (B, F, G) Nb 22 did not show complete inhibition Nb 39, Nb 53 and Nb 31 showed complete inhibition of ligand-binding at a lower concentration than the KYL-peptide. (C) Nb 39 and 53 showed complete inhibition of ligand-binding at a lower concentration than Nb 22, Nb 31 and the KYL-peptide. Trendlines are only shown when complete inhibition was reached. All assays were done in duplicate and data are represented as mean  $\pm$  SD.

**Figure 5. Nb 39 and Nb 53 do not completely inhibit ephrin-A5 binding to EphA7.** With alphascreen technology we determined whether Nb 39 and Nb 53 could inhibit ephrin-A5 binding to EphA7. These Nbs did not completely inhibit the interaction between ephrin-A5 and EphA7 at the highest concentrations tested. This assay was done in triplicate and data are represented as mean  $\pm$  SD.

**Figure 6. Nbs inhibit ephrin-induced EphA4 phosphorylation.** With the PathHunter assay we could determine that Nb 39 and Nb 53 completely inhibited EphA4 phosphorylation triggered by ephrin-A1 binding, even at a lower concentration than the KYL-peptide. All assays were performed in triplicate and data are represented as mean  $\pm$  SD.

**Figure 7. Nbs inhibit ephrin-B3-EphA4 mediated growth cone collapse.** Nb 53 and Nb 39 (1  $\mu$ M) reversed ephrin-B3-mediated growth cone collapse. (A) E17.5 cortical neurons were first treated at 24h after culture with 1  $\mu$ M or 30  $\mu$ M KYL peptide or 1  $\mu$ M Nb for 30 min, and then stimulated with 1  $\mu$ g/ml preclustured ephrin-B3-Fc or Fc as a control for 30 min in the presence of the Nb. A control condition without any kind of treatment was also scored (Control). Cells were stained with Alexa Fluor 555-conjugated phalloidin and DAPI. Growth cone collapse was determined for every neuron by analysing the tip of only the longest neurite (white arrowheads). Scale bar, 10  $\mu$ m. (B) Percentage of growth cone collapse was determined in every experimental condition by assessing 40-160 neurons

under a Zeiss fluorescence microscope. Percentage of collapse is expressed as the mean of collapse of 6 independent experiments  $\pm$  SD, and was compared between different conditions by one-way ANOVA followed by Tukey's multiple comparisons posthoc test. \*\*  $p < 0.01$  and \*\*\*  $p < 0.0001$ , as compared with control and Fc conditions. #  $p < 0.05$ , as compared with the ephrin-B3 condition.

**Figure 8. Nbs are stable in plasma for at least 72 hours.** Nbs were incubated in heparinized mouse plasma at 37°C until a maximum of 168 hours. Western Blot against HA tag was performed to detect Nb 39 and Nb 53 full-length and cleavage products (A). Quantifications of full-length Nb band (B) were referred to the full-length band at 0h for each Nb respectively. Means of 3-5 independent experiments  $\pm$  SD are represented. Time factor was analysed by two-way ANOVA followed by Sidak's multiple comparisons posthoc test. \*  $p < 0.05$  and \*\*\*  $p < 0.0001$ , as compared with 0 h time-point. Differences between the two Nbs were determined by two-way ANOVA, but not identified (n.s.). Nb incubation in PBS did not give rise to degradation products as detected by Western Blot (C). Incubation of Nbs in plasma at 37°C until a maximum of 72 hours did not affect the capability of those in binding EphA4 LBD (D). All values represent the percentage of Nb-EphA4 LBD binding respect to the 0 h time-point. This assay was done in triplicate and data are represented as mean  $\pm$  SD.



**Figure 2.**





Figure 3.





**Figure 4.**




Figure 5.







Figure 6.



**Figure 7.**



**Figure 8.**



## Identification and Characterisation of Nanobodies targeting the EphA4 receptor

Lies Schoonaert, Laura Ru&eacute;acute, Bart Roucourt, Mieke Timmers, Susan Little, Luc&iacute;acutea Ch&aacute;acutevez Guti&eacute;acuterrez, Maarten Dewilde, Peter Joyce, Adam Curnock, Peter Weber, Jurgen Haustraete, Gholamreza Hassanzadeh-Ghassabeh, Bart De Strooper, Ludo Van Den Bosch, Philip Van Damme, Robin Lemmens and Wim Robberecht

*J. Biol. Chem.* published online May 19, 2017

---

Access the most updated version of this article at doi: [10.1074/jbc.M116.774141](https://doi.org/10.1074/jbc.M116.774141)

### Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 0 references, 0 of which can be accessed free at <http://www.jbc.org/content/early/2017/05/19/jbc.M116.774141.full.html#ref-list-1>