microbial biotechnology

Open Access

Expression of fluorescent proteins in *Lactobacillus rhamnosus* to study host–microbe and microbe– microbe interactions

Irina Spacova,^{1,2} Elke Lievens,^{1,2} Tine Verhoeven,¹ Hans Steenackers,¹ Jos Vanderleyden,¹

Sarah Lebeer^{1,2,*} and Mariya I. Petrova^{1,2,*}

¹Centre of Microbial and Plant Genetics, KU Leuven, Heverlee, Belgium.

²Department of Bioscience Engineering, University of Antwerp, Antwerp, Belgium.

Summary

Probiotic Lactobacillus strains are widely used to benefit human and animal health, although the exact mechanisms behind their interactions with the host and the microbiota are largely unknown. Fluorescent tagging of live probiotic cells is an important tool to unravel their modes of action. In this study, the implementation of different heterologously expressed fluorescent proteins for the labelling of the model probiotic strains Lactobacillus rhamnosus GG (gastrointestinal) and Lactobacillus rhamnosus GR-1 (vaginal) was explored. Heterologous expression of mTagBFP2 and mCherry resulted in long-lasting fluorescence of L. rhamnosus GG and GR-1 cells, using the nisin-controlled expression (NICE) system. These novel fluorescent strains were then used to study in vitro aspects of their microbe-microbe and microbe-host interactions. Lactobacillus rhamnosus GG and L. rhamnosus GR-1 expressing mTagBFP2 and mCherry could be visualized in mixed-species biofilms, where they inhibited biofilm formation by Salmonella Typhimurium-gfpmut3 expressing the

Received 22 May, 2017; revised 12 August, 2017; accepted 25 August, 2017.

*For correspondence. E-mail sarah.lebeer@uantwerpen.be (S.L.); Tel. (+32) 32653285; Fax (+32) 32653225.

and

E-mail mariyapetrova3@gmail.com (M.P.); Tel. (+32) 16376647; Fax (+32)16321963.

Microbial Biotechnology (2017) **0**(0), 000–000 doi:10.1111/1751-7915.12872

Funding information

Irina Spacova and Elke Lievens were supported by IWT-SB-Vlaanderen through research projects at the time of this study. Sarah Lebeer was supported by the Fund for Scientific Research (FWO) Vlaanderen with a previous postdoctoral grant and the research grant KaN 28960. Mariya Petrova holds a postdoctoral grant from FWO Vlaanderen. Work at KU Leuven was supported by the BOF programme financing from the KU Leuven (PF/10/018). green fluorescent protein. Likewise, fluorescent *L. rhamnosus* GG and *L. rhamnosus* GR-1 were implemented for the visualization of their adhesion patterns to intestinal epithelial cell cultures. The fluorescent *L. rhamnosus* strains developed in this study can therefore serve as novel tools for the study of probiotic interactions with their environment.

Introduction

The ability to specifically label live bacteria is a powerful advantage for the *in vivo* and *in vitro* tracking of their behaviour, survival and interactions with other microor-ganisms, as well as their potential hosts. Heterologous expression of fluorescent proteins remains one of the most useful and common tools for bacterial imaging (Olenych *et al.*, 2007; Dean and Palmer, 2014). This approach results in higher labelling stability and specificity of live single cells compared with externally added dyes, allows prolonged time-lapse imaging and circumvents the need for exogenous dye internalization.

Currently, a broad selection of fluorescent proteins with varving fluorescence emission spectra and properties is available due to continuous efforts aimed at expanding the spectral range and improving their brightness and photostability (Olenych et al., 2007). For example, the green fluorescent protein (GFP) isolated from the Aequorea victoria jellyfish was the first fluorescent protein to be discovered and subsequently cloned in various live bacterial cells (Chalfie et al., 1994). Improved genetically modified GFP variants in the green (GFPuv) (Crameri et al., 1996), cyan (enhanced cyan fluorescent protein, ECFP) (Cubitt et al., 1995) and yellow (mVenus) (Nagai et al., 2002) spectra have also been developed. At the same time, a new group of fluorescent proteins with other fluorescence emission spectra and efficient maturation at 37°C has been discovered in reef corals and sea anemones, including the monomeric red spectrum protein mCherry (Shaner et al., 2004) and the blue spectrum protein mTagBFP2 (Subach et al., 2011). These developments have significantly broadened the application potential of fluorescent proteins to study various aspects of bacterial physiology.

A wide range of applications for fluorescent protein expression are available for common laboratory strains and pathogens, including *Escherichia coli* (Chalfie *et al.*,

© 2017 The Authors. *Microbial Biotechnology* published by John Wiley & Sons Ltd and Society for Applied Microbiology. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

1994; Ma et al., 2011), Salmonella spp. (Ma et al., 2011; Robijns et al., 2014) and Listeria spp. (Ma et al., 2011; Van der Veen and Abee, 2011). However, fluorescent labelling of probiotic bacteria, known as 'live microorganisms that, when administered in adequate amounts, confer a health benefit on the host' (Hill et al., 2014), is less widespread and has long remained limited to traditional fluorescent proteins, such as GFP and its derivatives (Geoffroy et al., 2000; De Keersmaecker et al., 2006a; Mota et al., 2006; Van der Veen and Abee, 2011). Nevertheless, a wide variety of novel fluorescent proteins with advanced properties that have not yet been tested in probiotic strains, such as mTagBFP2, are currently available for heterologous expression in bacteria, which can offer additional benefits for the study of microbe-microbe and host-microbe interactions (Olenych et al., 2007; Dean and Palmer, 2014).

One important feature for many probiotics is the ability to prevent or treat the occurrence of pathogens in various human body niches and restore microbial homeostasis (Bron et al., 2011; Reid et al., 2011; Ritchie and Romanuk, 2012). For example, it has previously been shown that specific probiotic Lactobacillus strains and their molecular products are capable of inhibiting biofilms of Salmonella spp. (De Keersmaecker et al., 2006b; Petrova et al., 2016a), Listeria monocytogenes, Pseudomonas aeruginosa (Alexandre et al., 2014), E. coli O157:H7 (Gómez et al., 2016) and uropathogenic E. coli (UPEC) (Petrova et al., 2016a). In addition, certain probiotic strains are capable of affecting host immune responses (Karlsson et al., 2012; Lebeer et al., 2012), potentially promoting pathogen recognition and elimination and/or a decrease in inflammatory signalling. The ability of probiotic bacterial cells to adhere to the lining of their respective niche might enhance these beneficial effects. This can be explained by an increased retention time and possibly closer and more prolonged contact with the host epithelial and immune cells, although adherence is not an absolute prerequisite (Lebeer et al., 2008).

Prominent examples of widely used and well-studied probiotic strains with demonstrated beneficial effects on the host are *L. rhamnosus* GG (a gastrointestinal isolate) (Segers and Lebeer, 2014) and *L. rhamnosus* GR-1 (an urogenital isolate) (Reid *et al.*, 1987). Nevertheless, understanding their mechanisms of action by which they improve and/or maintain human health remains a challenge. Therefore, in this study, we aimed to express various fluorescent proteins in the model strains *L. rhamnosus* GG and *L. rhamnosus* GR-1. The recombinant strains that showed the highest count of fluorescent cells and strongest fluorescence compared to the wild-type control strains were used to study their behaviour in mixed-species biofilms with *Salmonella* Typhimurium–*gfpmut3*, as well as for interactions with human epithelial cells *in vitro*.

Results

Expression of mTagBFP2 and mCherry in Lactobacillus rhamnosus GG and Lactobacillus rhamnosus GR-1 results in a fluorescent phenotype

The NICE system was implemented for the expression of fluorescent proteins in *L. rhamnosus* GG and *L. rhamnosus* GR-1 under the control of the inducible *nisA* promoter, as it was previously optimized and successfully used for expression of *gfp* in *L. rhamnosus* GG (De Keersmaecker *et al.*, 2006a). The pMEC45-derived plasmids pCMPG11260, pCMPG11261, pCMPG11262 and pCMPG11263 containing respectively *mTagBFP2, mCherry, mVenus* or *ecfp* downstream of the *nisA* promoter (Fig. S1) were introduced by electroporation into the *nisRK*-containing FAJ1905 strain of *L. rhamnosus* GR-1 (Table 1).

One of the colonies showing the presence of the correct corresponding plasmid (as confirmed by PCR and sequencing) was selected for detection of fluorescent signals. Following induction with commercial nisin from Lactococcus lactis, a fluorescent phenotype could be microscopy observed with epifluorescence in CMPG11260 (L. rhamnosus GG carrying mTagBFP2), CMPG11261 (L. rhamnosus GG carrying mCherry), CMPG11264 (L. rhamnosus GR-1 carrying mTagBFP2) and CMPG11265 (L. rhamnosus GR-1 carrying *mCherry*) compared to the corresponding wild-type L. rhamnosus GG or L. rhamnosus GR-1 used as negative control (Fig. 1A and B). For the strains CMPG11262 (L. rhamnosus GG carrying mVenus), CMPG11263 (L. rhamnosus GG carrying ecfp), FAJ1905/pMEC45 (L. rhamnosus GG carrying gfp), CMPG11266 (L. rhamnosus GR-1 carrying mVenus), CMPG11267 (L. rhamno-GR-1 carrying ecfp) and CMPG11268 SUS (L. rhamnosus GR-1 carrying gfp), no clear distinction at the expected wavelengths with the intrinsic fluorescence produced by the wild-type L. rhamnosus GG and L. rhamnosus GR-1 could be observed (Fig. 1C,D and E).

Subsequently, the fluorescence signals were quantified using flow cytometry. Fluorescent signals for the induced CMPG11260 (*L. rhamnosus* GG carrying *mTagBFP2*), CMPG11261 (*L. rhamnosus* GG carrying *mTagBFP2*) and CMPG11265 (*L. rhamnosus* GR-1 carrying *mTagBFP2*) and CMPG11265 (*L. rhamnosus* GR-1 carrying *mCherry*) strains were detected at the expected wavelength as compared to the corresponding wild-type *L. rhamnosus* GG or *L. rhamnosus* GR-1, which confirmed the microscopy results (Fig. 1A and B). CMPG11260 and CMPG11264 showed the most optimal detection of fluorescence when compared to the fluorescence signal of the wild-type control strains, with respectively 98.2% and

Table 1. Bacterial strains and p	lasmids used in this study.
----------------------------------	-----------------------------

Strain or plasmid	Genotype and/or phenotype	Source and/or references
Strain		
E. coli MC1061	araD139; ∆(ara, leu)7697; ∆lacX74; galU-; gal; hsr–; hsm+; strA	Casadaban and Cohen (1980)
Salmonella enterica serovar Typhimurium (ATCC 14028) – afomut3	<i>S</i> . Typhimurium (ATCC 14028) variant carrying pFPV25.1 with <i>gfpmut3</i> for constitutive GFP production	Robijns <i>et al.</i> (2014)
L. rhamnosus	Wild-type human isolate	Kankainen <i>et al.</i>
<i>L. rhamnosus</i> GR-1 (ATCC 5582)	Wild-type human isolate	Reid <i>et al.</i> (2001)
FAJ1905	<i>L. rhamnosus</i> GG with chromosomal insertion of pMEC10 in <i>attB</i> (phage mv4); Ery ^R	De Keersmaecker <i>et al.</i> (2006a)
CMPG11259 CMPG5357 CMPG11260 CMPG11261 CMPG11262 CMPG11263 FAJ1905/pMEC45	<i>L. rhamnosus</i> GR-1 with chromosomal insertion of pMEC10 in <i>attB</i> (phage mv4); Ery ^R <i>L. rhamnosus</i> GG ∆spaCBA::Tc ^R knockout mutant FAJ1905 strain of <i>L. rhamnosus</i> GG carrying pCMPG11260 with <i>mTagBFP2</i> ; Ery ^R , Cm ^R FAJ1905 strain of <i>L. rhamnosus</i> GG carrying pCMPG11261 with <i>mCherry</i> ; Ery ^R , Cm ^R FAJ1905 strain of <i>L. rhamnosus</i> GG carrying pCMPG11262 with <i>mVenus</i> ; Ery ^R , Cm ^R FAJ1905 strain of <i>L. rhamnosus</i> GG carrying pCMPG11263 with <i>ecfp</i> ; Ery ^R , Cm ^R FAJ1905 strain of <i>L. rhamnosus</i> GG carrying pCMPG11263 with <i>ecfp</i> ; Ery ^R , Cm ^R	This study Lebeer <i>et al.</i> (2012) This study This study This study This study De Keersmaecker <i>et al.</i> (2006a)
CMPG11269	CMPG5357 strain of <i>L. rhamnosus</i> GG carrying pCMPG11260 with <i>mTagBFP2</i> ; Ery ^P , Cm ^R	This study
CMPG11270 CMPG11264	CMPG5357 strain of <i>L. rhamnosus</i> GG carrying pCMPG11261 with <i>mCherry</i> ; Ery ^R , Cm ^R CMPG11259 strain of <i>L. rhamnosus</i> GR-1 carrying pCMPG11260 with <i>mTagBFP2</i> ; Ery ^R , Cm ^R	This study This study
CMPG11265	CMPG11259 strain of <i>L. rhamnosus</i> GR-1 carrying pCMPG11261 with <i>mCherry</i> ; Ery ^R , Cm ^R	This study
CMPG11266	CMPG11259 strain of <i>L. rhamnosus</i> GR-1 carrying pCMPG11262 with <i>mVenus</i> ; Ery ^R , Cm ^R	This study
CMPG11267 CMPG11268 Plasmids	CMPG11259 strain of <i>L. rhamnosus</i> GR-1 carrying pCMPG11263 with <i>ecfp</i> ; Ery ^R , Cm ^R CMPG11259 strain of <i>L. rhamnosus</i> GR-1 carrying pMEC45 with <i>gfp</i> ; Ery ^R , Cm ^R	This study
pMEC10	Integration plasmid in attB at 3' end of tRNAser locus (phage mv4); pNZ950-derivative containing the 3' end of <i>nisP</i> and <i>nisRK</i> (2.7 kb; <i>L. lactis</i> NZ9700) expressed by ery readthrough; Erv ^R	Pavan <i>et al.</i> (2000)
pMEC45	L. lactis pSH71 replicon; pNZ8037 derivative containing gfp _{uv} downstream of nisA promoter (L. lactis NZ9800); Cm ^R	Geoffroy et al. (2000)
pCMPG11260	Derivative of pMEC45 containing <i>mTagBFP2</i> cloned from pBAD- <i>mTagBFP2</i> downstream of <i>nisA</i> promoter; Cm ^R	This study
pBAD- <i>mTagBFP2</i> pCMPG11261	Plasmid used for <i>mTagBFP2</i> amplification Derivative of pMEC45 containing <i>mCherry</i> cloned from pRSETb- <i>mCherry</i> downstream of <i>nisA</i> promoter: Cm ^R	Subach <i>et al.</i> (2011) This study
pRSETb <i>–mCherry</i> pCMPG11262	Plasmid used for <i>mCherry</i> amplification Derivative of pMEC45 containing <i>mVenus</i> cloned from pCMPG13918 downstream of <i>nisA</i> promoter: Cm ^R	Shaner <i>et al.</i> (2004) This study
pCMPG13918	Plasmid used for <i>mVenus</i> amplification	Verstraeten <i>et al.,</i> unpublished
pCMPG11263	Derivative of pMEC45 containing <i>ecfp</i> cloned from pECFP (Clontech) downstream of <i>nisA</i> promoter; Cm ^R	This study
pECFP	Plasmid used for <i>ecfp</i> amplification	Clontech

Cm^R, chloramphenicol resistance; Ery^R, erythromycin resistance; Tc^R, tetracyclin resistance.

99.4% of the analysed events classified as fluorescent (Fig. 1A). For CMPG11261 and CMPG11265, only 25.6% and 59.3% of cells respectively showed no overlap with the fluorescence signal of the respective wild-type control strains when analysed with flow cytometry (Fig. 1B). No clear distinction in fluorescence compared with the respective wild-type controls was observed for the strains CMPG11262 and CMPG11266 carrying *mVenus*, CMPG11263 and CMPG11267 carrying *ecfp*, and

FAJ1905/pMEC45 and CMPG11268 carrying *gfp* (Fig. 1C,D and E).

To explore whether the lack of fluorescence in the recombinant *L. rhamnosus* GG and *L. rhamnosus* GR-1 strains containing *mVenus, ecfp* and *gfp* was due to inefficient gene expression, a real-time quantitative PCR (qRT-PCR) analysis was performed on all the strains designed to express *mTagBFP2, mCherry, mVenus, ecfp* and *gfp* using the NICE system. qRT-PCR analysis



Fig. 1. Detection of possible fluorescent phenotypes by epifluorescence microscopy and flow cytometric analysis of nisin-induced *Lactobacillus rhamnosus* GG and *L. rhamnosus* GR-1 bacterial cells implementing the NICE system to produce (A) mTagBFP2 (CMPG11260 and CMPG11264, in blue), (B) mCherry (CMPG11261 and CMPG11265, in red), (C) mVenus (CMPG11262 and CMPG11266, in yellow), (D) ECFP (CMPG11263 and CMPG11267, in cyan) and (E) GFP (FAJ1905/pMEC45 and CMPG11268, in green). In each panel, top photographs represent phase contrast, and bottom photographs represent fluorescent microscopy of the same field, with the white bars being equal to 10 μm. For each histogram, the fluorescence intensities at the corresponding wavelengths are indicated on the *x*-axis, and the number of events/cell counts is indicated on the *y*-axis. Wild-type *L. rhamnosus* GG and *L. rhamnosus* GR-1 were used as negative controls (depicted in grey).

for mRNA levels of expressed genes demonstrated that all of the fluorescent protein-encoding genes were expressed, although the amount of detected mRNA varied between the tested strains and the respective fluorescent protein genes (Fig. 2). Even though no fluorescence was observed for the strains designed to express *ecfp* and *mVenus* compared with the wild-type negative controls, high levels of both *ecfp* and *mVenus* mRNA could be detected in CMPG11262 and CMPG11266, and CMPG11263 and CMPG11267 respectively (Fig. 2).

The NICE system was likewise implemented to fluorescently label the CMPG5357 strain, a L. rhamnosus GG ∆*spaCBA*::Tc^R knockout mutant previously developed by our group to study the functional role of SpaCBA pili in interactions with the host (Lebeer et al., 2012). The pCMPG11260 and pCMPG11261 plasmids for the expression of respectively mTagBFP2 and mCherry under the control of the nisA promoter were introduced into CMPG5357 by electroporation, resulting in CMPG11269 and CMPG11270. The presence of plasmids with the correct insert was confirmed by PCR and plating out as described above for the wild-type recombinant strains. A fluorescent phenotype following induction with nisin could be observed in both CMPG11269 (CMPG5357 carrying mTagBFP2) and CMPG11270 (CMPG5357 carrying mCherry). The fluorescent signals at the expected wavelengths were visually confirmed by epifluorescence microscopy and quantified using flow cytometric analysis that classified 96.5% of CMPG11269



L. rhamnosus GG-derived strains

L. rhamnosus GR-1-derived strains

Fig. 2. Fluorescent protein-encoding gene expression in nisininduced *Lactobacillus rhamnosus* GG and *L. rhamnosus* GR-1 strains quantified by qRT-PCR. Data are presented as means with standard deviations indicating a ratio of the mRNA level for the fluorescent protein-encoding genes over that of a housekeeping *srtA* gene. cells (Fig. 3, in blue) and 93.1% of CMPG11270 cells as fluorescent (Fig. 3, in red).

In addition to the *nisA* promoter of the NICE system, the use of constitutive or internally regulated promoters for the *msp1*, *msp2*, *spaCBA*, *welE* genes and *dlt* operon from *L. rhamnosus* combined with an integrating pEM40 plasmid has been analysed for fluorescent protein expression in the context of this study. However, up to now, none of the constitutive promoters resulted in a detectable fluorescence signal in *L. rhamnosus* under the tested conditions.

Visualization of mixed-species biofilms of fluorescent Lactobacillus rhamnosus and Salmonella Typhimurium strains

Biofilm formation by single strains (Fig. 4) and mixed strains (Fig. 5) of fluorescent L. rhamnosus was explored using visualization with epifluorescence microscopy. CMPG11260 and CMPG11261, both L. rhamnosus GG wild-type-derived strains (Fig. 4A, CMPG11260 in blue and CMPG11261 in red), showed formation of thick biofilms in AOAC. Biofilm thickness was clearly reduced for CMPG11269 and CMPG11270, derived from the pili knockout mutant of L. rhamnosus GG (Fig. 4A, CMPG11269 in blue and CMPG11270 in red), in agreement with our previous work showing the key role of pili in biofilm formation (Lebeer et al., 2012). Lower levels of biofilm formation in AOAC medium were also observed for CMPG11264 and CMPG11265, both L. rhamnosus GR-1-derived strains (Fig. 4A, CMPG11264 in blue and CMPG11265 in red). These epifluorescence microscopy observations correspond with the results obtained from the Calgary biofilm formation experiments of the wildtype L. rhamnosus GG, L. rhamnosus GR-1 and the CMPG5357 pili knockout mutant of L. rhamnosus GG (Fig. 4B).

Fluorescent *L. rhamnosus* cells can also be distinguished from each other in mixed-strain lactobacilli biofilm set-ups in AOAC or TSB, such as those of CMPG11260 (Fig. 5A, in blue) with CMPG11265 (Fig. 5A, in red), as well as CMPG11261 (Fig. 5B, in red) with CMPG11264 (Fig. 5B, in blue). The *L. rhamnosus* GG-derived strains CMPG11260 and CMPG11261 demonstrate a more prominent biofilm formation capacity compared to CMPG11264 and CMPG11265 (Fig. 5), confirming the results from the single-strain biofilm formation experiment (Fig. 4). However, intriguingly, the *L. rhamnosus* GR-1-derived strains CMPG11264 and CMPG11265 also showed biofilm formation when co-cultured with the *L. rhamnosus* GG-derived CMPG11260 and CMPG11261 strains (Fig. 5).

To validate the functionality of the *L. rhamnosus* strains successfully expressing *mCherry* and



Fig. 3. Detection of a fluorescent phenotype by epifluorescence microscopy and flow cytometric analysis of nisin-induced CMPG5357, a *Lactobacillus rhamnosus* GG pili knockout mutant, implementing the NICE system to produce mTagBFP2 (strain CMPG11269, A, in blue) and mCherry (strain CMPG11270, B, in red). In each panel, top photographs represent phase contrast, and bottom photographs represent fluorescent microscopy of the same field. The white bar is equal to 10 μm. For each histogram, the fluorescence intensities at the corresponding wavelengths are indicated on the *x*-axis, and the number of events/cell counts is indicated on the *y*-axis. CMPG5357 was used as negative control (depicted in grey).

mTagBFP2, microbe-microbe interactions in mixed-species biofilms with fluorescent S. Typhimurium-afpmut3 were studied. A clear visual distinction could be made between the gfp-expressing S. Typhimurium-gfpmut3 cells (Fig. 6A-E, in green) and the induced recombinant L. rhamnosus cells CMPG11260 (Fig. 6B, in blue), CMPG11261 (Fig. 6C, in red), CMPG11264 (Fig. 6D, in blue) and CMPG11265 (Fig. 6E, in red). Fluorescence of induced recombinant L. rhamnosus cells could still be observed after 24 h (Fig. 6, left set of panels) and after 48 h (Fig. 6, right set of panels) of co-incubation without additional induction with nisin. In these co-interaction experiments, biofilm formation by S. Typhimurium-gfpmut3 was reduced by the lactobacilli compared to the negative control S. Typhimurium-gfpmut3 alone in all tested conditions (Fig. 6B-E compared to A).

Fluorescent Lactobacillus rhamnosus can be used to detect adhesion to human epithelial cells in vitro

To further validate the potential use of fluorescent L. rhamnosus strains in unravelling probiotic interactions with the host, an adhesion assay to human intestinal epithelial Caco-2 cells was conducted. The adhesion capacities of L. rhamnosus GG expressing mCherry (CMPG11261, Fig. 7B), L. rhamnosus GR-1 expressing mCherry (CMPG11265, Fig. 7C) and the CMPG5357 spaCBA pili knockout mutant of L. rhamnosus GG expressing mCherry (CMPG11270, Fig. 7D) to Caco-2 cells were compared. Only the mCherry-expressing strains could be visualized by epifluorescence microscopy with Caco-2 cell cultures, as the fluorescence produced by the mTagBFP2-expressing strains could not be distinguished from the intrinsic background fluorescence of the cell cultures at the corresponding wavelength. The L. rhamnosus GG-derived strain CMPG11261 showed a significant level of interaction with the cells and could be visualized in large numbers on top of the Caco-2 cell layer (Fig. 7B). In comparison, the *L. rhamnosus* GR-1-derived strain CMPG11265 (Fig. 7C) demonstrated significantly less binding to the Caco-2 cell layer. Likewise, the *L. rhamnosus* GG pili knockout mutant-derived strain CMPG11270 (Fig. 7D) demonstrated a drastically diminished adhesion capacity to Caco-2 cells compared with the *L. rhamnosus* GG-derived strain CMPG11261 (Fig. 7B), as it has previously been reported that SpaCBA is the key adhesin promoting the good adhesion capacities of *L. rhamnosus* GG to human intestinal epithelial cells (Lebeer *et al.*, 2012).

Discussion

In this study, we developed novel fluorescently labelled probiotic strains to study their interactions with other microorganisms, as well as the host. These strains can be used to better understand probiotic mechanisms of action, as bacterial fluorescence is a powerful tool to visualize and quantify the probiotic cells. The repertoire of fluorescent proteins expressed in probiotic lactobacilli is currently limited, and its expansion remains potentially challenging due to the microbiological properties of lactic acid bacteria, such as the low pH of the bacterial cytoplasm and culture medium, anaerobic growth conditions and the specific environments of the niche in which the strain is studied.

Here, we demonstrate successful labelling of wild-type and mutant strains of the commercially available probiotics *L. rhamnosus* GG and *L. rhamnosus* GR-1 by heterologous expression of the mTagBFP2 and mCherry fluorescent proteins. To our knowledge, this is the first report of the monomeric blue protein mTagBFP2 being



Fig. 4. Single-strain biofilms of nisin-induced fluorescent *Lactobacillus rhamnosus* strains after 48 h incubation in AOAC medium. (A) Biofilms of CMPG11260, an *L. rhamnosus* GG strain producing mTagBFP2 (phase contrast in grey and fluorescent in blue), CMPG11269, the CMPG5357 strain producing mTagBFP2 (phase contrast in grey and fluorescent in blue), CMPG11261, an *L. rhamnosus* GG strain producing mTagBFP2 (phase contrast in grey and fluorescent in blue), CMPG11261, an *L. rhamnosus* GG strain producing mCherry (phase contrast in grey and fluorescent in red), CMPG11265, an *L. rhamnosus* GG strain producing mCherry (phase contrast in grey and fluorescent in red), CMPG11265, an *L. rhamnosus* GR-1 strain producing mCherry (phase contrast in grey and fluorescent in red), CMPG11265, an *L. rhamnosus* GR-1 strain producing mCherry (phase contrast in grey and fluorescent in red) and CMPG11270, the CMPG5357 strain producing mCherry (phase contrast in grey and fluorescent in red) are shown. White bars are equal to $20 \mu m$. (B) Bar graph depicting biofilm formation on pegs by wild-type *L. rhamnosus* GG, *L. rhamnosus* GR-1 and the CMPG5357 pili knockout mutant of *L. rhamnosus* GG and *L. rhamnosus* GR-1 or CMPG5357 are depicted with double asterisks.

used to label any probiotic *Lactobacillus* strain. Furthermore, no heterologous expression of fluorescent proteins in the model vaginal probiotic strain *L. rhamnosus* GR-1 has been described to date. Significant fluorescence could be demonstrated in the novel recombinant *L. rhamnosus* strains when the mTagBFP2 and mCherry proteins were expressed using the nisin-inducible expression system (NICE) system, while this was not observed for the strains designed to express the mVenus or ECFP proteins. The maturation of traditionally used fluorescent proteins, such as GFP, and some of its derivatives, such as GFPuv, mVenus and ECFP, can be



Fig. 5. Mixed *Lactobacillus* strains biofilms of nisin-induced fluorescent *Lactobacillus rhamnosus* strains after 48 h co-incubation in AOAC or TSB media. (A) Biofilms of *L. rhamnosus* GG-derived CMPG11260 producing mTagBFP2 (phase contrast in grey and fluorescent in blue) with *L. rhamnosus* GR-1-derived CMPG11265 producing mCherry (phase contrast in grey and fluorescent in red). (B) Biofilms of *L. rhamnosus* GG-derived CMPG11261 producing mCherry (phase contrast in grey and fluorescent in red). (B) Biofilms of *L. rhamnosus* GG-derived CMPG11261 producing mCherry (phase contrast in grey and fluorescent in red) with *L. rhamnosus* GR-1-derived CMPG11264 producing mTagBFP2 (phase contrast in grey and fluorescent in blue). White bars are equal to 20 μm.

especially sensitive to changes in pH, low oxygen availability and growth temperatures (Scott et al., 2000; Olenych et al., 2007). For example, it has been demonstrated that increased aeration and artificial buffering of the growth medium pH are required for the optimal production and fluorescence of GFP in L. casei (Pérez-Arellano and Pérez-Martínez, 2003). Certain drawbacks resulting from the combination of fluorescent proteins with the properties of specific strains can be compensated for using a superior gene expression system that results in higher protein yield, such as the NICE system (Geoffroy et al., 2000; De Keersmaecker et al., 2006a). The NICE system is potentially food grade and offers high protein expression levels, although it presents a number of limitations, including the requirement for nisin induction and a dual plasmid composition (Geoffroy et al., 2000). To overcome these limitations, the use of constitutive or internally regulated promoters might be considered. However, until now, we were unable to obtain a detectable fluorescence signal when using constitutive promoters in L. rhamnosus under the tested conditions; therefore, the NICE system was chosen for further experiments. Nevertheless, even when the heterologous protein expression system is effective, the resulting fluorescence will still largely depend on the choice of a specific fluorescent protein and its maturation properties. For example, even under the optimized conditions, only 55% of the GFP produced in L. casei was in fact fluorescent (Pérez-Arellano and Pérez-Martínez,

2003). As demonstrated in this study for the strains carrying mVenus and ecfp, efficient fluorescent protein gene expression indeed does not always result in significant fluorescence in lactobacilli, even when corresponding mRNA production can be demonstrated. This might be the result of potential protein aggregation or impaired folding or maturation of proteins such as mVenus and ECFP in the cytoplasmic conditions of L. rhamnosus, which could impair the development of fluorescence even when sufficient mRNA production and translation occur. In addition, codon optimization of fluorescent protein genes could be considered (Van der Veen and Abee, 2011; Karimi et al., 2016) to potentially improve fluorescent protein translation and folding and the resulting fluorescence in lactobacilli. Although fluorescent protein mRNA production for each strain could be demonstrated, comparative analysis of mRNA levels corresponding to different fluorescent proteins is complicated due to the use of different gRT-PCR primer couples, which could affect amplification efficiency.

A number of recent studies have shown that fluorescent proteins other than GFP and its derivatives can indeed be useful for investigating *Lactobacillus* and *Bifidobacterium* strains in certain *in vivo* and *in vitro* assays (Russo *et al.*, 2015; Karimi *et al.*, 2016). In our study, we used mTagBFP2 and mCherry, as they are among the most promising candidates for labelling probiotic bacteria due to their fast and efficient maturation at 37°C, high photostability and lower pH sensitivity compared to



Fig. 6. Mixed-species biofilms of *gfp*-expressing *S*. Typhimurium–*gfpmut3* and nisin-induced fluorescent *Lactobacillus rhamnosus* strains after 24 h (left set of panels) and 48 h (right set of panels) co-incubation. *S*. Typhimurium–*gfpmut3* was incubated alone (A) (phase contrast in grey and fluorescent in green in all panels) or co-incubated with CMPG11260 producing mTagBFP2 (B) (phase contrast in grey and fluorescent in blue), CMPG11261 producing mCherry (C) (phase contrast in grey and fluorescent in red), CMPG11264 producing mTagBFP2 (D) (phase contrast in grey and fluorescent in blue) or CMPG11265 producing mCherry (E) (phase contrast in grey and fluorescent in red). White bars are equal to 20 μm.

GFP and its derivatives (Olenych *et al.*, 2007). mCherry has recently been used with success for labelling other probiotic *Lactobacillus* strains, such as *L. reuteri* (Karimi *et al.*, 2016), *L. plantarum* and *L. fermentum* (Russo et al., 2015), and it also performed well in this study. Indeed, the majority of induced cells designed to express the mCherry protein demonstrated a fluorescent phenotype when analysed by flow cytometry. This was even



Fig. 7. Adhesion pattern of *Lactobacillus rhamnosus* strains expressing *mCherry* to intestinal epithelial Caco-2 cells visualized by epifluorescence microscopy. (A) Caco-2 cells alone used as negative control; (B) Caco-2 cells incubated with CMPG11261 (*L. rhamnosus* GG expressing mCherry, in red); (C) Caco-2 cells incubated with CMPG11265 (*L. rhamnosus* GR-1 expressing mCherry, in red); (D) Caco-2 cells incubated with CMPG11270 (CMPG5357 *spaCBA* knockout expressing mCherry, in red). In each set of panels, left panel represent phase contrast, middle panel represents fluorescence microscopy and right panel represents a combination of phase contrast with fluorescent microscopy. White bars are equal to 20 μm.

more pronounced for the *L. rhamnosus* strains producing mTagBFP2, as 95% or more cells designed to express *mTagBFP2* were classified as fluorescent. Furthermore, for both mTagBFP2 and mCherry, we could show a persistence of fluorescence in recombinant *L. rhamnosus* strains even 48 h after induction with nisin, which makes these strains especially useful for long-term *in vitro* and potentially *in vivo* studies.

Tagging probiotic strains not only with the commonly used GFP, but also with the more recently developed variants of fluorescent proteins, such as mCherry and mTagBFP2, allows for an expanded spectral range and improved bacterial visualization. In this study, we demonstrate that fluorescent *L. rhamnosus* strains can be visualized in mixed-species biofilms, as bacterial cells of GFP-labelled *S.* Typhimurium and different mCherryand mTagBFP2-producing *L. rhamnosus* strains can be distinguished from each other based on their fluorescent tagging. This is especially useful for unravelling probiotic mechanisms of action against pathogenic microorganisms, as many bacterial pathogens, including *L. monocytogenes* (Gómez *et al.*, 2016), uropathogenic *E. coli*

(Petrova et al., 2016a) and P. aeruginosa (Alexandre et al., 2014), are capable of forming biofilms that contribute to their enhanced survival, tolerance and spreading (Olson et al., 2002; Hobley et al., 2015; Steenackers et al., 2016). In this study, visualization of mixed-species biofilms by epifluorescence microscopy demonstrated an inhibitory effect of L. rhamnosus GG and L. rhamnosus GR-1 on biofilm formation by S. Typhimurium. This is in line with previous studies describing a strong antimicrobial effect of L. rhamnosus GG and its molecular products on S. Typhimurium viability (Hutt et al., 2006; Marianelli et al., 2010) and biofilm formation (De Keersmaecker et al., 2006b; Petrova et al., 2016a). Therefore, these probiotic strains can potentially be used for prevention of S. Typhimurium infections and biofilm formation in human and animal disease and food applications. In future studies, a similar approach can be used for testing probiotic interactions with other biofilm-forming microorganisms, such as the clinically relevant strains of E. coli, Clostridium difficile or other pathogens commonly found in the gastrointestinal and vaginal niches.

In addition, visual analysis of single strain and mixedstrain biofilms of different fluorescent strains of L. rhamnosus has demonstrated a diminished capacity for biofilm formation by L. rhamnosus GR-1 and the CMPG5357 spaCBA knockout mutant strain of L. rhamnosus GG compared to L. rhamnosus GG. It has previously been demonstrated that L. rhamnosus GG is capable of efficient biofilm formation in vitro (Lebeer et al., 2007). Our results support the previously developed notion that the SpaCBA pili play a crucial role in this process (Lebeer et al., 2012), as both L. rhamnosus GR-1 and the spaCBA knockout mutant strain lack the pili structures. Surprisingly, when L. rhamnosus GR-1 is co-incubated with L. rhamnosus GG, its biofilm formation properties are enhanced compared to its single-strain biofilms, possibly due to the embedding of L. rhamnosus GR-1 into the L. rhamnosus GG biofilm matrix. Therefore, expression of fluorescent proteins in both pathogens (Ma et al., 2011; Robijns et al., 2014) and probiotic bacteria (Van der Veen and Abee, 2011) provides a non-destructive labelling method for real-time observation of biofilm formation dynamics and mixed-species biofilm interactions.

Furthermore, in this study, we show that the fluorescent *L. rhamnosus* strains, in particular those expressing the red mCherry fluorescent protein, can serve as a useful tool in studying probiotic interactions with human host cells. The adhesion pattern to intestinal epithelial Caco-2 cells visualized with epifluorescence microscopy confirms high binding capacity of *L. rhamnosus* GG to the intestinal epithelium, which is one of the hallmark characteristics of this probiotic strain (Segers and Lebeer, 2014). Results previously obtained by our group suggest that SpaCBA pili present on the outer surface of *L. rhamnosus* GG cells

play a key role in its adhesion to both intestinal epithelial cells (Lebeer *et al.*, 2012) and murine macrophages (Vargas García *et al.*, 2015). Therefore, SpaCBA pili might contribute to the longer retention time of *L. rhamnosus* GG in the human gastrointestinal tract compared with other *Lactobacillus* strains that lack pili-like structures (Kankainen *et al.*, 2009). Indeed, in our adhesion experiment, the non-piliated *L. rhamnosus* GR-1 strain expressing *mCherry* failed to demonstrate significant adhesion to Caco-2 cells in the tested conditions. Likewise, a *spaCBA* knockout mutant of *L. rhamnosus* GG expressing *mCherry* showed a drastically diminished binding to Caco-2 cells, further confirming our previous data (Lebeer *et al.*, 2012).

In the present work, we demonstrate the implementation and feasibility of mTagBFP2 and mCherry fluorescent tagging of probiotic lactobacilli, thus further expanding the tools for studying their molecular and cellular probiotic interactions with other microorganisms, as well as the host. Due to the easily detectable fluorescence of individual cells that persists long after induction, these novel fluorescent strains demonstrate a strong application potential for both *in vitro* and animal *in vivo* probiotic experiments.

Experimental procedures

Bacterial strains, plasmids and growth conditions

The bacterial strains and plasmids used in this study are listed in Table 1. All L. rhamnosus strains were grown in Man-Rogosa-Sharpe (MRS) medium (Difco) at 37°C in static conditions resulting in a limited supply of oxygen. Escherichia coli MC1061 and S. Typhimurium-gfpmut3 strains were grown at 37°C in Luria-Bertani broth in shaking conditions at 200 r.p.m. When appropriate erythromycin and chloramphenicol were supplied at respectively 5 and 10 μ g ml⁻¹ for the *L. rhamnosus* strains, chloramphenicol was supplied at 10 μ g ml⁻¹ for *E. coli* MC1061 and ampicillin was supplied at 100 μ g ml⁻¹ for S. Typhimurium-gfpmut3. Production of fluorescent proteins by L. rhamnosus strains was induced with nisin as previously described with minor modifications (De Keersmaecker et al., 2006a). Briefly, an overnight culture was diluted 1:50 in prewarmed MRS and incubated for 30 min. The diluted culture was subsequently induced with commercial nisin from Lactococcus lactis (Sigma-Aldrich, Diegem, Belgium, N5764) at a final concentration of 500 ng ml⁻¹ and further incubated for 24 h.

Construction of expression vectors and recombinant Lactobacillus rhamnosus strains

Standard molecular techniques were used for the construction of overexpression plasmids (Table 1),

preparation of competent E. coli cells and transformation (Sambrook et al., 1989). Gibson assembly (Gibson et al., 2009) was additionally used for the construction of pCMPG11261 due to a Ncol restriction site present in the coding sequence of *mCherry*, which complicated its insertion in the Ncol cloning site of pMEC45 (Fig. S1) using traditional cloning methods. PCR primers were purchased from Integrated DNA Technologies (IDT) (Belgium) (Table S1). Enzymes used for PCR, restriction digests, dephosphorylation and ligation of constructs were purchased from New England Biolabs (Bioké, Leiden, the Netherlands) and used according to the manufacturer's instructions. Plasmid DNA was isolated from E. coli using QIAGEN miniprep kits (Qiagen Benelux, Antwerp, Belgium) and transferred to highly competent L. rhamnosus GG, L. rhamnosus GR-1, CMPG5357, FAJ1905 or CMPG11259 cells by electroporation as described before (De Keersmaecker et al., 2006a; Petrova et al., 2016b). Electroporated L. rhamnosus cells were incubated at 37°C on MRS agar with corresponding antibiotics, i.e. erythromycin and chloramphenicol at respectively 5 and 10 µg ml⁻¹ for strains electroporated with pMEC45-derived constructs. The presence of plasmids with the correct insert in the resulting recombinant L. rhamnosus strains was confirmed by PCR analysis of colonies grown on erythromycin- and chloramphenicolcontaining MRS agar using the Pro-352 forward primer and the respective reverse primers for the corresponding fluorescent protein genes, being S&P-01367 for mTagBFP2, S&P-01303 for mCherry, S&P-01364 for mVenus and S&P-01365 for ecfp (Table S1).

Visualization of cell fluorescence with epifluorescence microscopy

Epifluorescence microscopy was performed using the Zeiss Axio Imager Z1 microscope with an EC Plan Neofluar (×40 magnification/0.3 numerical aperture) objective using the settings corresponding to the excitation and emission wavelengths of each expressed fluorescent protein. Images were obtained in aerobic conditions with an AXIOCAM MRM and the AXIOVISION software. Each visualization experiment was repeated at least twice.

Quantification of bacterial cell fluorescence using flow cytometry

Flow cytometric analysis of *L. rhamnosus* strains carrying genes for fluorescent protein expression was performed using 24 h cultures incubated at 37°C with appropriate antibiotics and when necessary induced with nisin as previously described (De Keersmaecker *et al.*, 2006a). Cultures were washed once and resuspended in PBS. Bacterial cell fluorescence was analysed in aerobic conditions by flow cytometry with a BD Influx cell sorter (Becton, Dickinson, Erembodegem, Belgium). Data were analysed using FLOWJO software (7.6.4), (FlowJo, LCC, Ashland, OR, USA), and histogram plots were obtained based on cell counts versus fluorescent signals at the expected wavelengths. Each flow cytometry quantification experiment was repeated at least twice.

Quantification of fluorescent protein gene expression by qRT-PCR

Total bacterial RNA was isolated using the SV Total RNA Isolation System Kit (Promega, Leiden, Netherlands) from 24 h cultures of L. rhamnosus strains incubated at 37°C with appropriate antibiotics and induced with nisin as previously described (De Keersmaecker et al., 2006a). To determine fluorescent protein gene expression, reverse transcription (RevertAid First Strand cDNA Synthesis Kit; Life Technologies Europe, Invitrogen, Gent, Belgium) with subsequent real-time quantitative PCR (qRT-PCR) (Power SYBR Green PCR Master Mix; Applied Biosystems Europe, Halle, Belgium) was performed. Primers were designed and synthetized by Integrated DNA Technologies (IDT) (Belgium) and are listed in Table S1. Relative abundance of mRNA corresponding to each fluorescent protein gene in relation to the housekeeping SrtA sortase mRNA was quantified after 40 amplification cycles of 15 s at 94°C and 1 min at 60°C. Expression plasmids containing each fluorescent protein-encoding gene were used as cDNA plasmid controls, and mRNA from wild-type L. rhamnosus GG and L. rhamnosus GR-1 served as negative controls for each pair of primers.

Co-incubation in mixed-species biofilm with Salmonella Typhimurium–gfpmut3

For the visualization of mixed-species biofilms, a *S.* Typhimurium ATCC14028 strain constitutively expressing the *gfpmut3* gene was used, combined with the CMPG11260, CMPG11261, CMPG11264 or CMPG11265 fluorescent strains of *L. rhamnosus.*

To prepare fluorescent *L. rhamnosus* cells for the mixed-species biofilm visualization experiments, overnight cultures of CMPG11260, CMPG11261, CMPG11264 or CMPG11265 were first induced with nisin as previously described (De Keersmaecker *et al.*, 2006a) and incubated for 24 h before being diluted for biofilm formation. For the formation of 24 and 48 h mixed-species biofilms with *S.* Typhimurium–*gfpmut3*, overnight cultures of planktonic *S.* Typhimurium–*gfpmut3* were diluted 1:100 in 1/20 TSB in plastic Petri dishes. Then, the previously induced 24 h cultures of CMPG11260, CMPG11261, CMPG11264 or CMPG11265 were immediately added at a 1:100 dilution

to the Petri dishes already containing diluted *S*. Typhimurium–*gfpmut3* in 1/20 TSB. After co-incubation of *S*. Typhimurium–*gfpmut3* with CMPG11260, CMPG1-1261, CMPG11264 or CMPG11265 for 24 or 48 h at 25°C in static aerobic conditions, the mixed-species biofilms were visualized with epifluorescence microscopy.

For the biofilm visualization experiments of single strain or mixed strains fluorescent *L. rhamnosus* biofilm formation, overnight cultures of CMPG11260, CMPG11261, CMPG11264, CMPG11265, CMPG11269 or CMPG11270 were induced with nisin as previously described (De Keersmaecker *et al.*, 2006a) and incubated for 24 h in MRS before being used for biofilm formation. The cultures were subsequently diluted 1:100 in 1/20 TSB or AOAC (Difco) medium in Petri dishes, and incubated at 25°C for 48 h in static aerobic conditions. The biofilms were visualized with epifluorescence microscopy.

Epifluorescence microscopy was performed using the Zeiss Axio Imager Z1 microscope with an EC Plan Neofluar (×40 magnification/0.3 numerical aperture). Images were obtained with an AXIOCAM MRm and the AXIOVISION software (ZEISS, Jena, Germany). Each visualization experiment was repeated at least twice.

For the quantification of biofilm formation, biofilms of *L. rhamnosus* GG, *L. rhamnosus* GR-1 or CMPG5357 strains were grown on pegs in 96-well plates for 72 h at 37°C in AOAC or TSB medium, and biofilm formation was evaluated using crystal violet staining as previously described (Lebeer *et al.*, 2007). Sterile growth medium was included as negative control. Analysis for each of the strains was performed eight times. Statistical analysis of the data using the Kruskal–Wallis test and Dunn's multiple comparison test was performed in GraphPad Prism 5.

In vitro assay for adhesion to human intestinal epithelial Caco-2 cells

The human intestinal epithelial Caco-2 cell line (American Type Culture Collection; Rockville, MD, USA; ATCC HTB-37TM) was maintained in Dulbecco's modified Eagle's medium (DMEM)-F-12 (Invitrogen) supplemented with 10% foetal calf serum (FCS) at 37°C, 5% CO₂ and 90% relative humidity. For the visualization of L. rhamnosus adhesion to Caco-2 cells, the cell cultures that had reached 70%-80% confluence were treated with trypsin and distributed over glass coverslips as previously described (Petrova et al., 2016a). Briefly, after 24 h of incubation in DMEM-F-12-FCS at 37°C, 5% CO₂ and 90% relative humidity, the cells were fixed on 13 mm coverslips in fixation buffer containing 0.05% glutaraldehyde and 2.5% formaldehyde for 1 h at 37°C, 5% CO₂ and 90% relative humidity. Simultaneously, 10⁷ CFU of fluorescent L. rhamnosus cultures previously induced with nisin as described above were collected by centrifugation for 10 min at 2000 *g*, washed once and dissolved in $1 \times PBS$, which was then added to the Caco-2 cells. After 1 h co-incubation at $37^{\circ}C$ in aerobic conditions, the coverslips were washed twice in PBS to remove unattached bacterial cells and subsequently visualized by epifluorescence microscopy using the Zeiss Axio Imager Z1 microscope with an EC Plan Neofluar (×40 magnification/0.3 numerical aperture). Images were obtained with an AxioCam MRm and the AxioVision software. Each visualization experiment was repeated at least twice.

Acknowledgements

We acknowledge the valuable help of Kenny Appermans, David De Coster, Laura Dell Erba and Rob Dockx during the course of this study. Irina Spacova and Elke Lievens were supported by IWT-SB-Vlaanderen through research projects at the time of this study. Sarah Lebeer was supported by the Fund for Scientific Research (FWO) Vlaanderen with a previous postdoctoral grant and the research grant KaN 28960. Mariya Petrova holds a postdoctoral grant from FWO Vlaanderen. Work at KU Leuven was supported by the BOF programme financing from the KU Leuven (PF/10/018).

Author's contribution

Irina Spacova, Sarah Lebeer and Mariya Petrova designed the experiments and wrote the manuscript. Irina Spacova, Mariya Petrova, Tine Verhoeven and Elke Lievens performed the experimental work. All authors reviewed and corrected the manuscript.

Conflict of interest

None declared.

References

- Alexandre, Y., Le Berre, R., Barbier, G., and Le Blay, G. (2014) Screening of *Lactobacillus* spp. for the prevention of *Pseudomonas aeruginosa* pulmonary infections. *BMC Microbiol* **14:** 107.
- Bron, P.A., van Baarlen, P., and Kleerebezem, M. (2011) Emerging molecular insights into the interaction between probiotics and the host intestinal mucosa. *Nat Rev Microbiol* **10**: 66–78.
- Casadaban, M.J., and Cohen, S.N. (1980) Analysis of gene control signals by DNA fusion and cloning in *Escherichia coli. J Mol Biol* **138**: 179–207.
- Chalfie, M., Tu, Y., Euskirchen, G., Ward, W.W., and Prasher, D.C. (1994) Green fluorescent protein as a marker for gene expression. *Science* **263**: 802–805.

- Crameri, A., Whitehorn, E.A., Tate, E., and Stemmer, W.P. (1996) Improved green fluorescent protein by molecular evolution using DNA shuffling. *Nat Biotechnol* **14:** 315–319.
- Cubitt, A.B., Heim, R., Adams, S.R., Boyd, A.E., Gross, L.A., and Tsien, R.Y. (1995) Understanding, improving and using green fluorescent proteins. *Trends Biochem Sci* **20**: 448–455.
- De Keersmaecker, S.C., Braeken, K., Verhoeven, T.L., Perea Vélez, M., Lebeer, S., Vanderleyden, J., *et al.* (2006a) Flow cytometric testing of green fluorescent protein-tagged *Lactobacillus rhamnosus* GG for response to defensins. *Appl Environ Microbiol* **72:** 4923–4930.
- De Keersmaecker, S.C., Verhoeven, T.L., Desair, J., Marchal, K., Vanderleyden, J., and Nagy, I. (2006b) Strong antimicrobial activity of *Lactobacillus rhamnosus* GG against *Salmonella typhimurium* is due to accumulation of lactic acid. *FEMS Microbiol Lett* **259**: 89–96.
- Dean, K.M., and Palmer, A.E. (2014) Advances in fluorescence labeling strategies for dynamic cellular imaging. *Nat Chem Biol* **10**: 512–523.
- Geoffroy, M.-C., Guyard, C., Quatannens, B., Pavan, S., Lange, M., and Mercenier, A. (2000) Use of green fluorescent protein to tag lactic acid bacterium strains under development as live vaccine vectors. *Appl Environ Microbiol* **66**: 383–391.
- Gibson, D.G., Young, L., Chuang, R.Y., Venter, J.C., Hutchison, C.A. 3rd, and Smith, H.O. (2009) Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* **6**: 343–345.
- Gómez, N.C., Ramiro, J.M.P., Quecan, B.X.V., and de Melo Franco, B.D.G. (2016) Use of potential probiotic Lactic Acid Bacteria (LAB) biofilms for the control of *Listeria monocytogenes*, *Salmonella typhimurium*, and *Escherichia coli* O157:H7 biofilms formation. *Front Microbiol* **7**: 863.
- Hill, C., Guarner, F., Reid, G., Gibson, G.R., Merenstein, D.J., Pot, B., *et al.* (2014) The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol* **11**: 506–514.
- Hobley, L., Harkins, C., MacPhee, C.E., and Stanley-Wall, N.R. (2015) Giving structure to the biofilm matrix: an overview of individual strategies and emerging common themes. *FEMS Microbiol Rev* **39**: 649–669.
- Hutt, P., Shchepetova, J., Loivukene, K., Kullisaar, T., and Mikelsaar, M. (2006) Antagonistic activity of probiotic lactobacilli and bifidobacteria against entero- and uropathogens. J Appl Microbiol 100: 1324–1332.
- Kankainen, M., Paulin, L., Tynkkynen, S., von Ossowski, I., Reunanen, J., Partanen, P., *et al.* (2009) Comparative genomic analysis of *Lactobacillus rhamnosus* GG reveals pili containing a human- mucus binding protein. *Proc Natl Acad Sci USA* **106**: 17193–17198.
- Karimi, S., Ahl, D., Vågesjö, E., Holm, L., Phillipson, M., Jonsson, H., and Roos, S. (2016) In vivo and in vitro detection of luminescent and fluorescent *Lactobacillus reuteri* and application of red fluorescent mCherry for assessing plasmid persistence. *PLoS One* **11**: e0151969.
- Karlsson, M., Scherbak, N., Reid, G., and Jass, J. (2012) Lactobacillus rhamnosus GR-1 enhances NF-kappaB

activation in *Escherichia coli*-stimulated urinary bladder cells through TLR4. *BMC Microbiol* **12:** 15.

- Lebeer, S., Verhoeven, T.L., Perea Vélez, M., Vanderleyden, J., and De Keersmaecker, S.C.J. (2007) Impact of environmental and genetic factors on biofilm formation by the probiotic strain *Lactobacillus rhamnosus* GG. *Appl Environ Microbiol* **73:** 6768–6775.
- Lebeer, S., Vanderleyden, J., and De Keersmaecker, S.C.J. (2008) Genes and molecules of lactobacilli supporting probiotic action. *Microbiol Mol Biol Rev* **72**: 728–764.
- Lebeer, S., Claes, I., Tytgat, H.L., Verhoeven, T.L., Marien, E., von Ossowski, I., *et al.* (2012) Functional analysis of *Lactobacillus rhamnosus* GG pili in relation to adhesion and immunomodulatory interactions with intestinal epithelial cells. *Appl Environ Microbiol* **78**: 185–193.
- Ma, L., Zhang, G., and Doyle, M.P. (2011) Green fluorescent protein labeling of *Listeria*, *Salmonella*, and *Escherichia coli* O157:H7 for Safety-Related Studies. *PLoS One* 6: e18083.
- Marianelli, C., Cifani, N., and Pasquali, P. (2010) Evaluation of antimicrobial activity of probiotic bacteria against *Salmonella enterica* subsp. *enterica* serovar typhimurium 1344 in a common medium under different environmental conditions. *Res Microbiol* **161**: 673–680.
- Mota, R.M., Moreira, J.L.S., Souza, L.R., Horta, M.F., Teixeira, S.M.R., Neumann, E., *et al.* (2006) Genetic transformation of novel isolates of chicken *Lactobacillus* bearing probiotic features for expression of heterologous proteins: a tool to develop live oral vaccines. *BMC Biotechnol* 6: 2.
- Nagai, T., Ibata, K., Park, E.S., Kubota, M., Mikoshiba, K., and Miyawaki, A. (2002) A variant of yellow fluorescent protein with fast and efficient maturation for cell-biological applications. *Nat Biotechnol* **20:** 87–90.
- Olenych, S.G., Claxton, N.S., Ottenberg, G.K., and Davidson, M.W. (2007) The fluorescent protein color palette. *Curr Protoc Cell Biol* **36:** 5: 21.5.1–21.5.34.
- Olson, M.E., Ceri, H., Morck, D.W., Buret, A.G., and Read, R.R. (2002) Biofilm bacteria: formation and comparative susceptibility to antibiotics. *Can J Vet Res* **66**: 86–92.
- Pavan, S., Hols, P., Delcour, J., Geoffroy, M.-C., Grangette, C., Kleerebezem, M., and Mercenier, A. (2000) Adaptation of the nisin-controlled expression system in *Lactobacillus plantarum*: a tool to study in vivo biological effects. *Appl Environ Microbiol* 66: 4427–4432.
- Pérez-Arellano, I., and Pérez-Martínez, G. (2003) Optimization of the green fluorescent protein (GFP) expression from a lactose-inducible promoter in *Lactobacillus casei*. *FEMS Microbiol Lett* **222**: 123–127.
- Petrova, M.I., Imholz, N.C.E., Verhoeven, T.L., Balzarini, J., Van Damme, E.J.M., Schols, D., *et al.* (2016a) Lectin-like molecules of *Lactobacillus rhamnosus* GG inhibit pathogenic *Escherichia coli* and *Salmonella* biofilm formation. *PLoS One* **11**: e0161337.
- Petrova, M.I., Lievens, E., Verhoeven, T.L., Macklaim, J.M., Gloor, G., Schols, D., *et al.* (2016b) The lectin-like protein 1 in *Lactobacillus rhamnosus* GR-1 mediates tissue-specific adherence to vaginal epithelium and inhibits urogenital pathogens. *Sci Rep* **6**: 37437.
- Reid, G., Cook, R.L., and Bruce, A.W. (1987) Examination of strains of lactobacilli for properties that may influence bacterial interference in the urinary tract. *J Urol* **138**: 330–335.

- Reid, G., Bruce, A.W., Fraser, N., Heinemann, C., Owen, J., and Henning, B. (2001) Oral probiotics can resolve urogenital infections. *FEMS Immunol Med Microbiol* **30**: 49– 52.
- Reid, G., Younes, J.A., Van der Mei, H.C., Gloor, G.B., Knight, R., and Busscher, H.J. (2011) Microbiota restoration: natural and supplemented recovery of human microbial communities. *Nat Rev Microbiol* **9**: 27–38.
- Ritchie, M.L., and Romanuk, T.N. (2012) A meta-analysis of probiotic efficacy for gastrointestinal diseases. *PLoS One* 7: e34938.
- Robijns, S.C., Roberfroid, S., Van Puyvelde, S., De Pauw, B., Uceda Santamaría, E., De Weerdt, A., *et al.* (2014) A GFP promoter fusion library for the study of *Salmonella* biofilm formation and the mode of action of biofilm inhibitors. *Biofouling* **30**: 605–625.
- Russo, P., Iturria, I., Mohedano, M.L., Caggianiello, G., Rainieri, S., Fiocco, D., *et al.* (2015) Zebrafish gut colonization by mCherry-labelled lactic acid bacteria. *Appl Microbiol Biotechnol* **99:** 3479–3490.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual.* Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- Scott, K.P., Mercer, D.K., Richardson, A.J., Melville, C.M., Glover, L.A., and Flint, H.J. (2000) Chromosomal integration of the green fluorescent protein gene in lactic acid bacteria and the survival of marked strains in human gut simulations. *FEMS Microbiol Lett* **182**: 23–27.
- Segers, M.E., and Lebeer, S. (2014) Towards a better understanding of *Lactobacillus rhamnosus* GG – host interactions. *Microb Cell Fact* **13(Suppl.** 1): S7.
- Shaner, N.C., Campbell, R.E., Steinbach, P.A., Giepmans, B.N.G., Palmer, A.E., and Tsien, R.Y. (2004) Improved monomeric red, orange and yellow fluorescent proteins derived from *Discosoma* sp. red fluorescent protein. *Nat Biotechnol* 22: 1567–1572.

- Steenackers, H.P., Parijs, I., Dubey, A., Foster, K.R., and Vanderleyden, J. (2016) Experimental evolution in biofilm populations. *FEMS Microbiol Rev* **40**: 980.
- Subach, O.M., Cranfill, P.J., Davidson, M.W., and Verkhusha, V.V. (2011) An enhanced monomeric blue fluorescent protein with the high chemical stability of the chromophore. *PLoS One* **6:** e28674.
- Van der Veen, S., and Abee, T. (2011) Mixed-species biofilms of *Listeria monocytogenes* and *Lactobacillus plantarum* show enhanced resistance to benzalkonium chloride and peracetic acid. *Int J Food Microbiol* **144**: 421–431.
- Vargas García, C.E., Petrova, M.I., Claes, I.J., De Boeck, I., Verhoeven, T.L., Dilissen, E., *et al.* (2015) Piliation of *Lactobacillus rhamnosus* GG promotes adhesion, phagocytosis, and cytokine modulation in macrophages. *Appl Environ Microbiol* 81: 2050–2062.

Supporting information

Additional Supporting Information may be found online in the supporting information tab for this article:

Fig. S1. Map of pMEC45 carrying *gfp* (green arrow) and pMEC45-derived plasmids carrying *mTagBFP2* (dark blue arrow, in pCMPG11260), *mCherry* (red arrow, in pCMPG11261), *mVenus* (yellow arrow, in pCMPG11262) or *ecfp* (cyan arrow, in pCMPG11263) under the control of the *L. lactis* inducible *nisA* promoter (P*nisA*, black arrow). The plasmids contain the *L. lactis* pSH71 replicon (*repA* and *repC* genes) and the chloramphenicol resistance cassette (Cm^R) depicted by grey arrows (adapted from Geoffroy *et al.*, 2000).

Table S1. Primers used in this study.