

TITLE

Competitive inter-species interactions underlie the increased antimicrobial tolerance in multispecies
brewery biofilms

RUNNING TITLE

Competition underlies antimicrobial tolerance

AUTHORS

Ilse Parijs¹, Hans P. Steenackers^{1#}

1 Centre of Microbial and Plant Genetics (CMPG), Department of Microbial and Molecular Systems,
KU Leuven, Kasteelpark Arenberg 20 - box 2460, B-3001 Leuven, Belgium.

Corresponding author. Mailing address: CMPG, KU Leuven, Kasteelpark Arenberg 20 – box 2460, B-
3001 Leuven, Belgium. Phone: +32 16 32 16 31. Fax: +32 16 32 19 66. E-mail:
hans.steenackers@kuleuven.be

The authors declare no conflict of interest.

ABSTRACT

Genetic diversity often enhances the tolerance of microbial communities against antimicrobial treatment. However the sociobiology underlying this antimicrobial tolerance remains largely unexplored. Here we analyze how inter-species interactions can increase antimicrobial tolerance. We apply our approach to 17 industrially-relevant multispecies biofilm models, based on species isolated from 58 contaminating biofilms in three breweries. Sulfathiazole is used as antimicrobial agent because it shows the highest activity out of 22 biofilm inhibitors tested. Our analysis reveals that competitive interactions dominate among species within brewery biofilms. We show that antimicrobial treatment can reduce the level of competition and therefore cause a subset of species to bloom. The result is a lower percentage inhibition of these species and increased tolerance. In addition, we show that the presence of competing species can also directly enhance the inherent tolerance of microbes to antimicrobial treatment, either because species protect each other or because they induce specific tolerance phenotypes as a response to competitors (i.e. competition sensing). Overall, our study emphasizes that the dominance of competitive interactions is central to the enhanced antimicrobial tolerance of the multispecies biofilms and that the activity of antimicrobials against multispecies biofilms cannot be predicted based on their effect against mono-cultures.

INTRODUCTION

Microbes commonly live in surface-attached communities embedded in a self-produced matrix, known as biofilms, which cause major problems and economic losses within industrial and medical sectors (Hall-Stoodley *et al.*, 2004). The majority of natural biofilms contain multiple species and harbor different functions and abilities compared to their monospecies counterparts (Stoodley *et al.*, 2002; Elias and Banin, 2012). One hallmark of multispecies biofilms is their increased tolerance against antimicrobial agents (Baffone *et al.*, 2011; Simões *et al.*, 2010; Shakeri *et al.*, 2007; Kumar and Peng, 2015; Jagmann *et al.*, 2015; Adam *et al.*, 2002; Lopes *et al.*, 2012; Leriche *et al.*, 2003; Whiteley *et al.*, 2001; Luppens *et al.*, 2008; Wang *et al.*, 2013; Schwering *et al.*, 2013; Van der Veen and Abee, 2011; Simões *et al.*, 2009; Harriott and Noverr, 2009; Lee *et al.*, 2014). Although different species within biofilms are closely associated and are expected to strongly interact with each other (Elias and Banin, 2012), little is known about how these interactions affect antimicrobial tolerance. Indeed, most previous studies focused on the overall tolerance of multispecies biofilms, without looking at the contributions of individual species. In the limited cases that multispecies composition before and after treatment was determined, the types of interactions and their interdependency with antimicrobial treatment and tolerance were generally not investigated (Harriott and Noverr, 2009; Van der Veen and Abee, 2011; Simões *et al.*, 2009; Whiteley *et al.*, 2001; Leriche *et al.*, 2003; Luppens *et al.*, 2008; Wang *et al.*, 2013; Schwering *et al.*, 2013).

Interactions can be cooperative or competitive in nature. Cooperative interactions for example involve the secretion of enzymes (Rakoff-Nahoum *et al.*, 2016) or metabolic cross-feeding (Harcombe, 2010). Social evolution theory defines a cooperative adaptation in one species as a phenotype that increases the fitness of another species and that evolved at least in part because of this effect (Mitri and Foster, 2013; Foster and Bell, 2012; West *et al.*, 2007). This implies that both species are benefitting from the interaction, since it is difficult to see how an adaptation that helps another species can evolve when it has a fitness cost to the helping species. The *cooperation criterion*

therefore states that an observed interaction is only consistent with a cooperative adaptation if the total productivity in co-culture is higher than the sum of the mono-culture productivities (which is the null for no interaction) and both species increase their cellular productivity in co-culture vs. mono-culture (Mitri and Foster, 2013). Higher order cooperative interactions could potentially occur when sets of three, four or more species engage in loops of mutually beneficial interactions. In addition to evolved cooperation, also accidental positive effects can occur (no positive feedback loop), for example when waste products in a focal species can be used as resource by another species. This can result in a commensal interaction, in which there is no fitness effect in one direction, or can happen within a competitive association (*vide infra*).

Whenever one or both species experience a disadvantage in co-culture, competition is dominant. Competition is expected to be favoured when coexisting species have overlapping metabolic niches, are spatially mixed and when cell density is high relative to the available resources (Ghoul and Mitri, 2016). Phenotypes involved in microbial competition can be accidental in nature or have evolved for this purpose. The latter phenotypes are called competitive adaptations and include (i) strategies to take away resources from competitors (exploitative competition), for example by fast-but-wasteful-growth (Pfeiffer *et al.*, 2001), production of nutrient-scavenging molecules (Scholz and Greenberg, 2015), or superior positioning within the niche (Kim *et al.*, 2014), and (ii) strategies to directly fight with competing species (interference competition), for example by production of antimicrobials (Riley and Gordon, 1999) or contact dependent inhibition (Russell *et al.*, 2014; Lories *et al.*, 2017). Competition can be further characterized by comparing the observed productivity in co-culture with the weighted average productivity of the constituent species in mono-culture. This allows to determine to which extent inter-species competition differs from intraspecific competition. As described below, this difference is called the *biodiversity effect* and is constituted of a *selection effect* and *complementarity effect*. Previously, this concept has been frequently applied in plant ecology (Cardinale *et al.*, 2007; Morin *et al.*, 2011; Polley *et al.*, 2003; Spehn *et al.*, 2005; Loreau and Hector, 2001).

Here we combine the *cooperation criterion* and *biodiversity effect* to investigate how inter-species interactions underlie the increased tolerance of multispecies biofilms. We apply our approach to industrially-relevant multispecies biofilm models consisting of combinations of species isolated from contaminating biofilms in breweries. In the brewing industry, undesired biofilms can be associated with spoilage organisms, cause corrosion and reduce process efficiency. Several stages of the brewing process, including the pasteurization, the storage and bottling of the beer, are known to be affected by biofilms and the eradication of these biofilms remains highly challenging (Storgards and Tapani, 2006; Mamvura and Iyuke, 2011; Maifreni *et al.*, 2015). Improved understanding of antimicrobial tolerance is thus a prerequisite for designing more effective brewery sanitation procedures.

Our analysis reveals that a complex interplay between antimicrobial treatment and genetic diversity underlies the commonly-observed increased tolerance of multispecies biofilms. Consistent with previous work in other microbial communities (Foster and Bell, 2012; Rivett *et al.*, 2016; Oliveira *et al.*, 2015), we show that competitive interactions dominate among species within the brewery biofilms. We then show that this dominance of competitive interactions is central to the enhanced antimicrobial tolerance of the multispecies biofilms. Antimicrobial treatment, if incomplete, can reduce the level of competition and therefore cause a subset of species to bloom. The result is a lower percentage inhibition of these species in the multispecies biofilm compared to the mono-culture biofilms, which appears -per definition- as increased tolerance. Complete inhibition of all species in the mixture would avoid this effect. However, our results further indicate that the presence of competing species can also directly enhance the inherent tolerance of microbes to antimicrobial treatment. Antimicrobials that are completely effective against mono-culture biofilms are thus not necessarily effective against the same species in co-culture. Overall this emphasizes that the activity of antimicrobials against multispecies biofilms cannot be predicted based on their effect against mono-cultures.

MATERIAL AND METHODS

Sampling

A total of 58 samples were collected from four breweries in Belgium between August and December 2014, before and after cleaning in place (CIP). Sterile cotton swabs (Deltalab) were used for collection of biofilm material from approximately 25 cm² on different surfaces in the bottling plant, filtration room and storage room. After sampling, swabs were submerged in phosphate buffered saline (PBS), which consists of 8.8 g/L NaCl, 1.24 g/L K₂HPO₄, and 0.39 g/L KH₂PO₄ (pH 7.4). Biofilm material was removed from the swabs by 3x 30 s vortexing and sonication at 45 kHz, 80 W. 900 µl of the PBS solution with the biofilm material was frozen at -80°C in 50% glycerol. 100 µl was plated out in dilution 10⁰ to 10⁻⁸ on plate count agar (PCA), which is composed of 5 g/L peptone, 2.5 g/L yeast extract, 1 g/L glucose and 15 g/L agar. The plates were incubated at 25 °C for 7 days. The total microbial load was determined by counting the colonies on the plates and determining the CFU/cm².

Identification of culturable species

Colonies growing on the PCA plates were identified by partial 16S rRNA or ITS gene sequencing. For bacteria, colony PCR using Taq DNA polymerase (Life Technologies) was performed on the 16S rRNA gene, which was targeted by primers BSF8/20 and BSR1541/20 (Cai *et al.*, 2003). The following PCR program was used: 96°C for 6 min, 35 cycles of (i) denaturation at 96°C for 1 min, (ii) annealing at 47.5°C for 1 min, and (iii) elongation at 72°C for 90s, and a final elongation at 72°C for 6 min. Afterwards, the PCR products were loaded on a 1X agarose gel, which was run for 1 hour at 125V and 400A. The band at 1500 bp was cut out and the DNA was extracted by using the GenElute™ Gel Extraction Kit (Sigma-Aldrich). Sanger sequencing was performed on the extracted DNA using primer BSF8/20 (GATC Biotech). The resulting sequence was blasted against the NCBI gene database to identify the closest relative of each colony. For yeast identification, the same protocol was followed, using primers ITS1 and ITS4 (White *et al.*, 1990), with an annealing temperature of 60°C and an elongation time of 1 min. The resulting PCR fragment was 330 bp.

Defined multispecies biofilm models

Defined multispecies biofilm models combined a fixed number of culturable species that were isolated from the same sample. Hereto, each species was grown in liquid PCA culture for 48 hours at 25 °C under shaking conditions. These species were combined with a starting density of 1000 CFU/ml for each species and grown in 1/20 Trypticase Soy Broth in 96 well plates. Similarly, monospecies biofilms were set up with a starting density of 1000 CFU/ml. After incubation for 4 days at 25°C, the amount of living cells in the biofilm formed on the bottom of the 96 well plates was quantified by plate counting. First, the free-living cells were removed from the wells. Second, the biofilm on the bottom of the wells was scraped off in 200 µl PBS and diluted appropriately. Finally, the diluted solutions with biofilm cells were plated out on PCA plates and incubated at 25°C for 2-5 days. Colonies on the plates were counted and the CFU/cm² was determined. In the multispecies biofilms, a distinction was made between the different species based on colony morphology.

Study of inter-species interactions: cooperation criterion and biodiversity effect

15 different defined multispecies biofilms were grown as described above. To determine if inter-species interactions are cooperative or competitive, the cooperation criterion was applied. This criterion requires that the biofilm growth for all species in co-culture is higher than their respective biofilm growth in mono-culture (Mitri and Foster, 2013).

To further characterize the ecological influences on interactions, the biodiversity effect was calculated according to the formula below, in which $N\overline{\Delta RY\overline{M}}$ measures the complementarity effect and $N cov(\Delta RY, M)$ measures the selection effect (Loreau and Hector, 2001).

$$\Delta Y = Y_O - Y_E = N\overline{\Delta RY\overline{M}} + N cov(\Delta RY, M)$$

M_i = biofilm growth of species i in mono-culture

$Y_{O,i}$ = observed biofilm growth of species i in co-culture

$Y_O = \sum_i Y_{O,i}$ = total observed biofilm growth in co-culture

$RY_{E,i}$ = expected relative biofilm growth of species i in co-culture, which is its proportion inoculated

$RY_{O,i} = Y_{O,i}/M_i$ = observed relative biofilm growth of species i in co-culture

$Y_{E,i} = RY_{E,i}M_i$ = expected biofilm growth of species i in co-culture

$Y_E = \sum_i Y_{E,i}$ = total expected biofilm growth in co-culture

$\Delta Y = Y_O - Y_E$ = deviation from the total expected biofilm growth in the co-culture (= biodiversity effect)

$\Delta RY_i = RY_{O,i} - RY_{E,i}$ = deviation from expected relative biofilm growth of species i in co-culture

N = number of species in co-culture

The cooperation criterion requires the total multispecies inoculation density to be equal to the sum of the inoculation densities of the mono-culture biofilm. In contrast, the definition of the biodiversity effect imposes that the inoculation density of each species in the multispecies biofilms should be its inoculation density in mono-culture, divided by the number of species that are present in the multispecies biofilm. All multispecies biofilm models in this study were grown in both set-ups and no differences in final growth were observed. Inoculation densities are indeed not expected to have large effects when biofilms are grown to stationary phase (Foster and Bell, 2012).

Biofilm inhibitor screening

A library of 96 inhibitors that were reported in literature or *in house* developed and that are known to affect biofilm specific processes was composed. Based on commercial availability, low cost and toxicity, 22 compounds were selected for a one-replicate preventive screening against 17 undefined multispecies biofilm, grown directly from the frozen samples. Inhibitors were dissolved in dimethyl sulfoxide (DMSO) at 100 μ M and their activity was tested as described in the next paragraph. In a next step, 12 of the 22 inhibitors were selected based on their activity and chemical properties and were screened against 9 of the 17 frozen sample multispecies biofilms using a range of twofold serial diluted concentrations between 800 and 0.4 μ M. This allows calculating the BIC50, which is defined as the concentration (μ M) of inhibitor needed to inhibit 50% of the biofilm growth.

The undefined multispecies biofilms were set up with a 1000 CFU/ml start inoculum taken directly from the frozen isolated biofilm sample and grown in 1/20 Trypticase Soy Broth on the pegs of the Calgary biofilm device for 4 days at 25 °C. Quantification of biofilm biomass was done by crystal violet staining (Ceri *et al.*, 1999). Briefly, the pegs were first washed in 200 µl PBS and then stained for 30 minutes with 200 µl 0.1% (w/v) crystal violet in an isopropanol/methanol/PBS solution (v/v 1:1:18). Next, the excess stain was washed off the pegs in 200 µl distilled water and the pegs were left to dry for 30 minutes. Finally, the pegs were destained in 200 µl 30 % glacial acetic acid and biofilm matrix was quantified by measuring the OD₅₇₀ of each well using a Synergy MX multimode reader (Biotek, Winooski, VT). The BIC50 was determined for each inhibitor that was tested in a range of concentrations by nonlinear curve fitting (GraphPad Prism software, version 6).

Multispecies biofilm inhibition by sulfathiazole

Defined mono- and multispecies biofilms were set up in 96 well plates as previously described and at the start of the incubation, sulfathiazole (100 µM in DMSO) or an equal amount of DMSO was added. Biofilms were grown and quantified by plating out as described previously. Inter-species interactions were determined as described above. Tolerance to sulfathiazole was calculated in mono- and multispecies biofilms using the following formulas:

$$\% \text{ Tolerance } (species\ i, mono) = \frac{\text{Treated biofilm growth } (species\ i, mono)}{\text{Untreated biofilm growth } (species\ i, mono)} * 100$$

$$\% \text{ Tolerance } (species\ i, multi) = \frac{\text{Treated biofilm growth } (species\ i, multi)}{\text{Untreated biofilm growth } (species\ i, multi)} * 100$$

RESULTS

Construction of industrially relevant multispecies biofilm models

We started by setting up a series of *in vitro* multispecies biofilm models with relevance for the brewing-industry, that were further used throughout this study. Hereto, 103 biofilm samples isolated from different locations in several breweries were microbiologically characterized. The total bacterial load (CFU/cm²) varied between 10² and 10⁸ before cleaning in place (CIP) and between 10¹ and 10⁹ after CIP. As shown in Figure 1, the microbial contamination after CIP was reduced with less than 75% in 52% of the samples and was even increased in 24% of the samples, indicating that CIP is insufficient and that improved antimicrobial treatments are highly needed.




Figure 1: % tolerance to CIP for 33 biofilms sampled from different locations in different breweries.

The genera of the closest known relatives of the culturable microbes were determined by partial 16S rRNA gene sequencing to characterize the microbial diversity (Table S1). The biofilm samples were mainly composed of *Pseudomonas* and *Raoultella* ssp. and also two beer spoiling organisms, *Pediococcus* and *Lactococcus*, were identified. Multispecies biofilm models were then constructed by combining species isolated from the same sample. Seventeen ‘undefined’ biofilm models were set up by directly inoculating part of the frozen isolated biofilm samples. These models were used to screen for broad-spectrum biofilm inhibitors. Because these biofilms likely contain unculturable species, for the study of inter-species interactions an additional 12 ‘defined’ multispecies biofilm models were constructed by inoculating equal ratios of 3 to 6 well-identified, culturable

species (originating from 8 samples taken before CIP and 4 samples taken after CIP). Several biofilms containing *Pseudomonas* and *Raoultella* spp. were included.

Screening of biofilm inhibitors

To study the tolerance of multispecies brewery biofilms, we sought to use a broad-spectrum antimicrobial with a large potential for application against brewery biofilms. Hereto, a library of 22 biofilm inhibitors -with previously reported activity against mono-culture biofilms- was composed that target biofilm specific processes such as adhesion (Opperman *et al.*, 2009), dispersion (Barraud *et al.*, 2006), EPS-production (Nithya *et al.*, 2011) and several others (Lynch and Abbanat, 2010). After an initial screening against 17 undefined multispecies biofilm models using a fixed concentration of 100 μ M (data not shown), we selected 12 inhibitors, which were tested more thoroughly using multiple concentrations. Specifically, we performed a preventive screening against 9 undefined multispecies biofilm models directly grown from the frozen brewery biofilm samples. Crystal violet staining was used to measure the amount of biofilm formed and the 50% inhibitory concentrations (BIC50) were calculated for each biofilm model (Figure 2). Sulfathiazole was found to have the broadest activity-spectrum against the brewery biofilms and was therefore selected for further study. This inhibitor has been described previously to interfere with c-di-GMP biosynthesis in *E. coli* biofilms (Antoniani *et al.*, 2010). C-di-GMP has been reported to play a crucial role in biofilm formation by a wide range of bacterial species, which might explain the broad-spectrum activity of this compound (Cotter and Stibitz, 2007; Hengge, 2009; Romling *et al.*, 2013).




Figure 2: BIC50 values of 12 biofilm inhibitors against 9 undefined biofilms (shown as mean with standard deviation of three biological repeats). BIC50 is defined as the concentration (μM) of inhibitor needed to prevent biofilm growth with 50%. Compounds with BIC50 values over 600 μM are considered ineffective and are not shown.

Effect of inter-species interactions on biofilm growth and composition

We first aimed to determine the role of inter-species interactions in multispecies biofilms, irrespective of antimicrobial treatment. Hereto, we performed a systematic classification of the interactions in 12 defined biofilm models, each consisting of 3 to 6 culturable species, by using two complementary approaches: (i) *cooperation criterion* and (ii) *biodiversity effect*.

The *cooperation criterion* was used to classify interactions as cooperative or competitive. For all biofilm models, the number of biofilm cells of each species (CFU/cm²) in mono-culture was compared to the cell count of each species in co-culture (Figure 3). Most species performed worse in co-culture than in mono-culture indicating that competitive interactions are dominant. The increased cellular productivity observed for a subset of the species in few of the multispecies biofilms (e.g. for 3

out of 4 species in model 5) could be due to exploitation of the remaining, suppressed species, however, cooperation between these species cannot be ruled out. In summary, all models were characterized by competitive interactions that cause some or all species to perform worse than in mono-culture.




Figure 3: Mono-culture growth (Mono), expected (Exp) and observed (Obs) multispecies composition and the biodiversity, selection and complementarity effect for 1 representative repeat of 12 defined multispecies biofilm models.

To further characterize competition, the *biodiversity effect* was measured (Loreau and Hector, 2001). When inter-species competition is equal to intra-specific competition, the observed productivity in co-culture is expected to be equal to the average productivity of the constituent species in mono-culture, weighted by the inoculation frequencies. The biodiversity effect is defined as the difference between the observed and expected multispecies biofilm productivity and is thus a measure for the extent to which inter-species interactions deviate from intra-specific interactions. The

observed productivity of the 12 model biofilms (CFU/cm²) was compared to the expected biofilm growth. In the majority of the multispecies biofilm models (75%), the total amount of biofilm formed was higher than expected, as indicated by a positive biodiversity effect, while the remaining 25% of the cases were characterized by a negative biodiversity effect.

A positive biodiversity effect can either be caused by selection of the best biofilm former or by a (partial) niche separation alleviating competition; conversely a negative biodiversity effect can be caused by selection of the worse biofilm former or by strong interference competition. To distinguish between both possibilities, Loreau & Hector (2001) partitioned the biodiversity effect into a selection and complementarity effect (Material and Methods). Selection occurs when the extent to which the relative productivity in co-culture vs. mono-culture deviates from expected is non-randomly related to the productivity in mono-culture and is measured by a covariance function. Positive selection is indicative of the dominance of the best mono-culture biofilm formers and occurred in 33,3% of the multispecies biofilm models. Negative selection suggests the opposite and appeared in the remaining 67,7% of the multispecies biofilm models. If only selection effects take place, the total relative productivity (sum of relative productivities of all species) is 1, meaning that an increase in productivity in one species is compensated by a decrease in productivity of another species. However, if the total relative productivity is higher or lower than 1 over- or underyielding occurs, which is defined as the complementarity effect. This effect measures whether the relative amount of biofilm formed in co-culture vs. mono-culture is on average higher or lower than expected based on the initial relative abundance and biofilm growth in mono-culture and is thus also a measure for the strength of competition. Complementarity is positive if some degree of niche separation occurs, for example if two species can grow on different resources or if one species is able to use a waste product of another species as a resource. Consequently, the strength of competition decreases and the productivity increases due to a more optimal use of the available niches. Positive complementarity was observed in 91,7% of the multispecies biofilm models. On the other hand, negative complementarity effects

occurred in the remaining 8,3% of the multispecies biofilm models and indicate the occurrence of strong chemical or physical interference competition (Fox, 2005; Turnbull *et al.*, 2013; Loreau, 2000).

The combination of complementarity and selection effects then gives an indication as to which ecological processes are the cause of the total positive or negative biodiversity effect. In our multispecies biofilm models positive biodiversity (75%), could be explained by resource partitioning or facilitation between the different species for 66,7% of the biofilms (only positive complementarity), by dominance of the best biofilm formers for 11,1% of the biofilms (only positive selection) or by a combination of both positive complementarity and selection for 22,2% of the biofilms. Conversely, negative biodiversity effects (25%), were caused by exploitation or interference competition for 33,3% of the biofilms (only negative complementarity), by dominance of poor biofilm formers for 33,3% of the biofilms (only negative selection) or by a combination of negative complementarity and selection for 33,3% of the biofilms (Loreau and de Mazancourt, 2013). Overall, the mainly positive complementarity effects indicate that the competitive interactions in the multispecies biofilm models are in most cases alleviated by partial niche separation.

Link between reduced competition and antimicrobial tolerance in multispecies biofilms

The results above show that competitive inter-species interactions, although in general alleviated by partial niche separation, strongly influence the productivity of each species in the multispecies biofilms. In a next step, we sought to investigate the interplay of these competitive interactions with antimicrobial treatment and their effect on antimicrobial tolerance. Hereto, sulfathiazole was added preventively to three multispecies biofilm models (Table 1). Tolerance to sulfathiazole is defined as the ratio between the amount of biofilm formed in the presence and absence of treatment and was determined for each species in mono- and co-culture conditions (Figure 4-6). In all three models the tolerance of each species was equal or higher in the multispecies biofilm than in the mono-culture. The result is an overall increase in tolerance in each multispecies biofilm, which can be seen by comparing the expected and observed amount of biofilm after treatment. Here the expected amount

is calculated based on the composition before treatment and the percentage of reduction of each species in mono-culture. These results are in line with the increased tolerance generally observed in multispecies biofilms (Mozina *et al.*, 2013; Burmølle *et al.*, 2014). Specifically, our results are consistent with a previous study on sulfathiazole treatment, in which multispecies biofilms isolated from cooling water systems were found to be more tolerant compared to their mono-culture counterparts (Shakeri *et al.*, 2007).

Table 1: Closest known relative genus for each of the species present in the three multispecies biofilm models that were used to study the inhibition by sulfathiazole

	Species 1	Species 2	Species 3	Species 4
Model 1	<i>Epilithonimonas</i>	<i>Aeromonas</i>		
Model 2	<i>Pseudomonas</i>	<i>Pseudoclavibacter</i>	<i>Raoultella</i>	<i>Serratia</i>
Model 3	<i>Pseudomonas</i>	<i>Raoultella</i>		

Analyzing biofilm compositions before and after treatment revealed that the above described dominance of competitive interactions in untreated biofilms is central to the observed enhanced tolerance to antimicrobial treatment. In two out of three biofilm models we found that antimicrobial treatment reduced the level of competition and therefore caused a subset of species to bloom. The result was a lower percentage inhibition of these species in the multispecies biofilm compared to the mono-culture biofilms, which -per definition- appears as increased tolerance.

In duo-species model 1 (Figure 4), species 2 is sensitive to the inhibitor both in the mono- and co-culture biofilm (4 A&B). However, species 1, which is insensitive to the inhibitor in mono-culture, shows a 50-fold increase in growth upon addition of the inhibitor in the duo-species biofilm (4 A&B), resulting in an overall higher tolerance of the duo-species biofilm (4A). In the untreated duo-species biofilm, species 1 is strongly suppressed by species 2 as reflected in the strong competition (4C), large positive selection effect (4D) and negative complementarity (4D). The increased growth of species 1 upon treatment is therefore consistent with an abrogation of the competitive interactions of sensitive species 2 against species 1, which then blooms and shows a net increase in antimicrobial tolerance. This is reflected in a reduced competition (4C), associated with a positive complementarity (4D) in the

treated biofilm. In summary, inhibition of the best competitor results in a bloom of the worse competitor and overall increased tolerance.

A similar mechanism plays in tetra-species model 2 (Figure 5). Three out of four species are completely inhibited both in the mono- and multispecies biofilm (5 A&B). Species 3, however, which is insensitive to the inhibitor in the monospecies biofilm, shows a 1.5-fold increase in growth upon addition of the inhibitor in the multispecies biofilm (5 A&B), resulting in an overall increase in tolerance of the mixed species biofilm (5A). Species 3 experiences competition by the other species in the untreated multispecies biofilm (5 C&D), explaining why inhibition of these other species increases the growth -and tolerance- of species 3 in the treated biofilm. Since there is only one species left after treatment, competition (5C) and biodiversity effect (5D) are zero.

It should be noted that this mechanism of ‘increased tolerance due to reduced competition’ does not involve an increase in absolute cell numbers of the different species in co-culture compared to mono-culture, nor an expression of specific tolerance phenotypes. Nevertheless, the proposed mechanism is of significance. Indeed, similar to our study, antimicrobial tolerance in previous studies was generally measured by calculating the reduction in cell numbers before and after treatment, not by directly comparing the absolute cell numbers between co- and mono-culture conditions (Chorianopoulos *et al.*, 2008; Van der Veen & Abee, 2011; Kostaki *et al.*, 2012; Giaouris *et al.*, 2013; Wang *et al.*, 2013). Therefore, the increased tolerance observed in these studies might as well be explained by decreased competition and should not necessarily be accompanied by any changes in specific tolerance phenotypes.




Figure 4 Model 1: **A:** Number of cells of each species in mono- and co-culture biofilms, grown in the absence and presence of sulfathiazole treatment: suppressed species 1 is able to grow after treatment. **B:** Tolerance = ratio between the number of biofilm cells with and without sulfathiazole treatment, determined for each species in mono- and co-culture conditions. For each species the tolerance is equal or higher within the co-culture biofilm. Significant differences were examined using a two-way anova and Bonferroni correction (* $P < 0.05$). **C:** Cooperation: both in the absence and presence of treatment the criterion for cooperation is not met. **D:** Biodiversity effects in the absence and presence of treatment of the duo-species biofilm: dominating positive selection is replaced by positive complementarity. **E:** Overview: inhibition of species 2 leads to a reduction in the competitive interactions against species 1, which allows species 1 to bloom. Results show the average of 3 biological repeats, except for A, which shows the average of 3 technical repeats of one representative biological repeat.




Figure 5: Model 2: **A:** Number of cells of each species in mono- and co-culture biofilms, grown in the absence and presence of sulfathiazole treatment: suppressed species 3 shows an increased growth upon treatment. **B:** Tolerance = ratio between the number of biofilm cells with and without sulfathiazole treatment, determined for each species in mono- and co-culture conditions. Species 3 shows an increased the tolerance within the multispecies biofilm, while species 1, 2 and 4 are completely inhibited in mono- and co-culture. Significant differences were examined using a two-way anova and Bonferroni correction (**** $P < 0.0001$). **C:** Cooperation: both in the absence and presence of treatment the criterion for cooperation is not met. **D:** Biodiversity effects in the absence and presence of treatment of the multispecies biofilm: negative complementarity becomes positive. **E:** Overview: complete inhibition of species 1, 2 and 4 leads to the abrogation of competitive interactions against species 3, which allows species 3 to bloom. Results show the average of 3 biological repeats, except A, which shows the average of 3 technical repeats of one representative biological repeat.

Direct effect of competitors on antimicrobial tolerance

The findings above indicate that incomplete antimicrobial treatment of multispecies biofilms can reduce the levels of competition and therefore cause a subset of species to bloom, which ultimately results in increased antimicrobial tolerance. Complete inhibition of all species in the mixture would solve this problem. However, our analysis of duo-species model 3 (Figure 6) indicates that the presence of competing species can also directly enhance the inherent tolerance of other species by driving specific tolerance phenotypes. This means that antimicrobials that are completely effective against mono-culture biofilms are not necessarily effective against the same species in co-culture and thus precludes any prediction on multispecies tolerance.

In this duo-species model (Figure 6), both species respond to sulfathiazole treatment in the mono- and co-culture biofilms (6 A&B). However, species 1 shows a 11,1-fold reduction in sensitivity in co-culture, resulting in an overall increased tolerance of the co-culture biofilm (6A). In contrast to the previous model systems, this tolerance of species 1 is associated with an increase in cell number above the mono-culture levels (6A, right panel). These results cannot be explained by a decrease in competition alone (6C-D),and should be attributed to the presence of specific tolerance phenotypes within the multispecies biofilm. These could either be related to a protective effect of species 2 on species 1 or to a direct change in tolerance phenotype of species 1 as a response to species 2.




Figure 6: Model 3: **A:** Number of cells of each species in mono- and co-culture biofilms, grown in the absence and presence of sulfathiazole treatment. The right part of the graph zooms in on the amount of biofilm formed by species 1 in treated mono- and duo-culture: after treatment the growth of species 1 in duo-culture exceeds its growth in mono-culture. **B:** Tolerance = ratio between the number of biofilm cells with and without sulfathiazole treatment, determined for each species in mono- and co-culture conditions. Species 1 shows a higher tolerance within the co-culture biofilm, while there is no difference for species 2. **C:** Cooperation: after treatment species 1 grows better in co-culture than in mono-culture, while there is no difference for species 2. This is consistent with commensalism. **D:** Biodiversity effects in the absence and presence of treatment of the co-culture biofilm: negative complementarity before treatment becomes positive. **E:** Overview: species 1 becomes more tolerant in the presence of competing species 2. The growth of species 1 in the treated co-culture biofilm even exceeds its mono-culture growth, suggesting induction of specific tolerance phenotypes. Results show the average of 3 biological repeats, except A, which shows one representative biological.

DISCUSSION

Functional properties like antimicrobial tolerance strongly differ between multispecies and monospecies biofilm communities (Burmølle *et al.*, 2014; Røder *et al.*, 2016). Although inter-species interactions are expected to be both intense and important within dense communities (Elias and Banin, 2012), little is known about how they affect antimicrobial tolerance. Previous work either focused on microbial interactions in untreated biofilms (Tan *et al.*, 2016; Røder *et al.*, 2016; Ghoul and Mitri, 2016) or on the overall tolerance of multispecies biofilms, without taking contributions of individual species into account (Adam *et al.*, 2002; Burmølle *et al.*, 2006; Baffone *et al.*, 2011; Simões *et al.*, 2010; Lopes *et al.*, 2012). We have bridged the gap and shown that a complex interplay between antimicrobial treatment and inter-species interactions underlies the commonly-observed increased tolerance of multispecies biofilms. We have shown that competitive interactions dominate within industrially relevant multispecies biofilm models and that antimicrobial treatment, if incomplete, can reduce the level of competition and therefore cause subsets of species to bloom, ultimately leading to enhanced overall tolerance. In addition, we have shown that the presence of competitors can also directly enhance the inherent tolerance to antimicrobials by driving specific tolerance phenotypes. Overall, our results emphasize that the increasingly-recognized dominance of competition in multispecies biofilms is central to the enhanced antimicrobial tolerance and that antimicrobial activities against mono-culture biofilms cannot predict efficacy against multispecies biofilms.

Our data indicate that competitive interactions dominate among species within brewery biofilms, although inter-species competition is generally weaker than intraspecific competition. These data fit with a growing body of recent theoretic and experimental work motivating that competition, not cooperation, dominates interactions among microbial species. The genotypic view of social interactions predicts a low chance of evolution of cooperation between species, because this requires both a high within-genotype relatedness and sufficient niche separation to reduce ecological competition. Increased niche separation, however, often implies a decreased exchange of resources,

which counteracts interactions, and further complicates the evolution of cooperation (Mitri and Foster, 2013). These predictions are confirmed by recent systematic screenings of inter-species interactions based on the cooperation criterion (Foster and Bell, 2012; Rivett *et al.*, 2016; Fiegna *et al.*, 2015). Also in these studies inter-species competition was found to be weaker than intraspecific competition (Foster and Bell, 2012; Rivett *et al.*, 2016; Oliveira *et al.*, 2015; Fiegna *et al.*, 2015). Moreover, Rivett *et al.* (2016) showed that initially strong competitive interactions can weaken over time by divergence in resource use and increased niche complementarity. It should be noted that a number of studies did report a prevalence of positive interactions, however, these studies made use of alternative definitions. In a recent study, synergistic interactions were defined as the total amount of multispecies biofilm being higher than the sum of all mono-cultures and synergy was observed in 13% of the biofilms (Madsen *et al.*, 2016). This definition is similar to the cooperation criterion, but since the effect of growth in co-culture on the individual species is not included, the presence of cooperative interactions cannot be confirmed. In earlier studies, synergy required the total amount of multispecies biofilm to be higher than that of the best mono-culture biofilm former and synergistic interactions were reported in respectively 11%, 63% and 30% of the biofilms (Burmølle *et al.*, 2007; Ren *et al.*, 2015; Røder *et al.*, 2015). Also here information on composition of the multispecies biofilm is needed to determine whether the described synergistic interactions are competitive or cooperative. It can however be deduced that these synergistic interactions are associated with a positive biodiversity effect, since both definitions imply the total amount of multispecies biofilm to be higher than the weighted average of the mono-cultures. Notably, this positive biodiversity effect does imply niche complementarity, but can also partly be caused by positive selection effects. In conclusion, the importance of competition among species over cooperation is increasingly recognized and our data are consistent with this. However, an important note is that all studies described above, including ours, are based on culturable species, which might exclude species that are only able to grow in the presence of other species. Therefore, the prevalence of cooperation might be underestimated (Foster and Bell, 2012; Røder *et al.*, 2016).

The enhanced overall antimicrobial tolerance against sulfathiazole that we observed for each multispecies biofilm model compared to the mono-culture biofilms is consistent with the enhanced resistance found in the majority of multispecies biofilm studies (Baffone *et al.*, 2011; Simões *et al.*, 2010; Shakeri *et al.*, 2007; Kumar and Peng, 2015; Jagmann *et al.*, 2015; Adam *et al.*, 2002; Lopes *et al.*, 2012; Leriche *et al.*, 2003; Whiteley *et al.*, 2001; Luppens *et al.*, 2008; Wang *et al.*, 2013; Schwering *et al.*, 2013; Van der Veen and Abee, 2011; Simões *et al.*, 2009; Harriott and Noverr, 2009; Lee *et al.*, 2014; Hoffman *et al.*, 2006). In most of these studies the enhanced tolerance was attributed to protective effects of the species on each other, however, generally without unraveling the mechanism of tolerance. In contrast, a minority of studies did not observe an effect of multispecies conditions on antimicrobial tolerance (Gkana *et al.*, 2017) or did even measure a decrease in tolerance in multispecies conditions (Lindsay *et al.*, 2002; Chorianopoulos *et al.*, 2008; Kart *et al.*, 2014; Yassin *et al.*, 2016; Feldman *et al.*, 2016).

Our data indicate that the commonly-observed enhanced antimicrobial tolerance of multispecies biofilms is associated with a reduction in the level of competition upon treatment, causing a subset of species to bloom. The dominance of competition among species over cooperation in untreated biofilms is therefore central to the enhanced antimicrobial tolerance. Indeed, incomplete inhibition of a network of cooperating species is expected, not to promote, but to pull down the remaining species because of abrogation of positive feedback loops, as is motivated by recent ecological network studies (Coyte *et al.*, 2015). This would reduce, not increase, the overall tolerance of the multispecies biofilm (Feldman *et al.*, 2016). Our models only provide examples of multispecies biofilms in which specific species strongly suppress other species. Inhibition of the stronger competitors consequently reduces the competition that is experienced by the suppressed species and leads to an increased tolerance of the weaker competitors. However, the idea that antimicrobial tolerance in multispecies biofilms is connected to a reduction in competition should not be limited to this situation, as one can easily imagine that antimicrobial treatment can also reduce competition between equal competitors. For example, in the case of equally competing species that only produce

their toxins when the population density of the other species is sufficiently high (Cornforth and Foster, 2013), a reduction of the population size by antimicrobial treatment would interfere with toxin production, reduce competition and ultimately lead to increased antimicrobial tolerance compared to mono-culture.

Based on the commonly found prevalence of competitive interactions within multispecies biofilms, it is expected that reduction in competition might often be the cause of increased tolerance. However, little is known about this because previous work mainly focused on characterizing the antimicrobial tolerance of mono- and multispecies biofilms, without explicitly classifying the changes in inter-species interactions before and after treatment. In a number of studies, only the overall activity against the multispecies biofilm and the activity against the mono-cultures was measured, while information on individual species in co-culture is essential to understand the inter-species interactions (Adam *et al.*, 2002; Burmølle *et al.*, 2006; Baffone *et al.*, 2011; Simões *et al.*, 2010; Lopes *et al.*, 2012). Similarly, only determining the inhibition of each species in the co-culture without looking at the effects in mono-culture (Norwood and Gilmour, 2000; Hill *et al.*, 2010; DeLeon *et al.*, 2014; Sun *et al.*, 2008; Feldman *et al.*, 2016) or only focusing on specific species within the multispecies biofilm (Kumar and Peng, 2015; Jagmann *et al.*, 2015; Shakeri *et al.*, 2007) does not allow to study all changes in inter-species interactions. Nevertheless, a few studies have been conducted in which the tolerance of each species was examined individually, both under mono- and co-culture biofilm conditions (Harriott and Noverr, 2009; Van der Veen and Abee, 2011; Simões *et al.*, 2009; Whiteley *et al.*, 2001; Leriche *et al.*, 2003; Luppens *et al.*, 2008; Wang *et al.*, 2013; Schwering *et al.*, 2013; Elvers *et al.*, 2002). While the obtained data would allow to perform a detailed analysis of the changes in inter-species interactions as proposed in this paper, this analysis is generally missing and the representation of the data in most cases did not allow us to interpret the data a posteriori. Nevertheless, one study on tolerance of a 7-species biofilm provided sufficient data and is consistent with our mechanism of ‘increased tolerance due to reduced competition’ (Elvers *et al.*, 2002). Some of the bacterial species experienced a reduced growth due to competition in the untreated multispecies biofilm, while

antimicrobial treatment restored their growth in the multispecies biofilm to the level of the untreated monoculture biofilms. In contrast, but also consistent with our rationale, a reduction in antimicrobial tolerance under multispecies conditions has been explicitly associated with a reduction of (probably rare) cooperative inter-species interactions (Feldman *et al.*, 2016).

In our final model, we found that the presence of competitors can also directly enhance the inherent tolerance of other species by driving specific tolerance phenotypes. This could either be attributed to (i) protective effects of specific species on other species or to (ii) direct changes in tolerance phenotypes of specific species as a response to competitors. A previously described example of a protective effect occurs between competing *Pseudomonas aeruginosa* and *Staphylococcus aureus* species (Hoffman *et al.* 2006). Respiration of *S. aureus* was found to be inhibited by a competitive interaction involving the exoproduct 4-hydroxy-2-heptylquinoline-N-oxide of *P. aeruginosa*. As a consequence, aminoglycoside antibiotics were no longer taken up by *S. aureus* cells and their tolerance to these antibiotics increased. Additionally, the presence of *P. aeruginosa* on a long term increased the production of highly resistant small-colony variants of *S. aureus*, which further improved the antimicrobial tolerance of *S. aureus*. In addition, it is becoming increasingly clear that bacteria can also directly sense the presence of competitors and respond appropriately (i.e. ‘competition sensing’) (Cornforth and Foster, 2013). Recent studies indicate that these responses can include upregulated biofilm formation (Oliveira *et al.*, 2015), increased antibiotics or toxin production (Le Roux *et al.*, 2015; Abrudan *et al.*, 2015; Rosenberg *et al.*, 2016), altered secretion of specific secondary metabolites (Traxler *et al.*, 2013), but also increased antibiotic tolerance (Abrudan *et al.*, 2015; Roberfroid *et al.*, personal communication).

In conclusion, due to the their commonly observed increased antimicrobial tolerance, multispecies biofilms remain challenging to eradicate. Accordingly, we found multispecies biofilms to be a serious problem in breweries, as emphasized by the high microbial load of the isolated biofilm samples, both before and after CIP. An increased knowledge of the properties of these multispecies

biofilms may aid to improve their control. Our study demonstrates that competitive inter-species interactions dominate within multispecies biofilms and have a strong influence on the outcome of antimicrobial treatment. Specifically, we found that strongly suppressed species can bloom after inhibition of superior competitors by antimicrobial treatment, which results in increased tolerance. To avoid such unwanted effects of changing inter-species interactions, it would be useful to develop combination therapies that completely inhibit all species. Nevertheless, we also observed that the presence of competitors can increase the intrinsic tolerance of species by driving specific tolerance phenotypes. This means that antimicrobials that are completely effective against mono-culture biofilms are not necessarily effective against the same species in co-culture. Our study therefore underlines the need to further investigate and interfere with the mechanisms behind these specific tolerance phenotypes.

ACKNOWLEDGEMENTS

We would like to thank Stijn Robijns, Sandra Van Puyvelde and Bram Lories for their valuable comments and for their assistance in sampling of brewery biofilms. We thank K. Deflem and D. De Coster for experimental assistance. This work was supported by the KU Leuven Research Fund (STG/16/022), by the Institute for the Promotion of Innovation through Science and Technology in Flanders under grant IWT-SBO 120050 (NEMOA) and by FWO-Vlaanderen (W0.009.16N). IP is a research assistant of the IWT-Vlaanderen (SB/131721). HS acknowledges the receipt of a postdoctoral fellowship from FWO-Vlaanderen (PDO/11).

REFERENCES

- Abrudan MI, Smakman F, Grimbergen AJ, Westhoff S, Miller EL, van Wezel GP, *et al.* (2015). Socially mediated induction and suppression of antibiosis during bacterial coexistence. *Proc Natl Acad Sci U S A* **112**: 11054–11059.
- Adam B, Baillie GS, Douglas LJ. (2002). Mixed species biofilms of *Candida albicans* and *Staphylococcus epidermidis*. *J Med Microbiol* **51**: 344–349.
- Antoniani D, Bocci P, Maciag A, Raffaelli N, Landini P. (2010). Monitoring of diguanylate cyclase activity and of cyclic-di-GMP biosynthesis by whole-cell assays suitable for high-throughput screening of biofilm inhibitors. *Appl Microbiol Biotechnol* **85**: 1095–104.
- Baffone W, Sorgente G, Campana R, Patrone V, Sisti D, Falcioni T. (2011). Comparative effect of chlorhexidine and some mouthrinses on bacterial biofilm formation on titanium surface. *Curr Microbiol* **62**: 445–451.
- Barraud N, Hassett DJ, Hwang S-H, Rice SA, Kjelleberg S, Webb JS. (2006). Involvement of nitric oxide in biofilm dispersal of *Pseudomonas aeruginosa*. *J Bacteriol* **188**: 7344–53.
- Burmølle M, Hansen LH, Sørensen SJ. (2007). Establishment and early succession of a multispecies biofilm composed of soil bacteria. *Microb Ecol* **54**: 352–362.
- Burmølle M, Ren D, Bjarnsholt T, Sørensen SJ. (2014). Interactions in multispecies biofilms: do they actually matter? *Trends Microbiol* **22**: 84–91.
- Burmølle M, Webb JS, Rao D, Hansen LH, Sørensen SJ, Kjelleberg S. (2006). Enhanced biofilm formation and increased resistance to antimicrobial agents and bacterial invasion are caused by synergistic interactions in multispecies biofilms. *Appl Environ Microbiol* **72**: 3916–23.
- Cai H, Archambault M, Prescott JF. (2003). 16S Ribosomal RNA Sequence--Based Identification of Veterinary Clinical Bacteria. *J Vet Diagnostic Investig* **15**: 465–469.

Cardinale BJ, Wright JP, Cadotte MW, Carroll IT, Hector A, Srivastava DS, *et al.* (2007). Impacts of plant diversity on biomass production increase through time because of species complementarity. *Proc Natl Acad Sci U S A* **104**: 18123–8.

Ceri H, Olson ME, Stremick C, Read RR, Morck D, Buret A. (1999). The Calgary Biofilm Device: new technology for rapid determination of antibiotic susceptibilities of bacterial biofilms. *J Clin Microbiol* **37**: 1771–6.

Chorianopoulos NG, Giaouris ED, Skandamis PN, Haroutounian SA, Nychas GJE. (2008). Disinfectant test against monoculture and mixed-culture biofilms composed of technological, spoilage and pathogenic bacteria: bactericidal effect of essential oil and hydrosol of *Satureja thymbra* and comparison with standard acid-base sanitizers. *J Appl Microbiol* **104**: 1586–1596.

Cornforth DM, Foster KR. (2013). Competition sensing: the social side of bacterial stress responses. *Nat Rev Microbiol* **11**: 285–93.

Cotter PA, Stibitz S. (2007). c-di-GMP-mediated regulation of virulence and biofilm formation. *Curr Opin Microbiol* **10**: 17–23.

Coyte KZ, Schluter J, Foster KR. (2015). The ecology of the microbiome: Networks, competition, and stability. *Science (80-)* **350**: 663–666.

DeLeon S, Clinton A, Fowler H, Everett J, Horswill AR, Rumbaugh KP. (2014). Synergistic interactions of *Pseudomonas aeruginosa* and *Staphylococcus aureus* in an In vitro wound model. *Infect Immun* **82**: 4718–4728.

Elias S, Banin E. (2012). Multi-species biofilms: living with friendly neighbors. *FEMS Microbiol Rev* **36**: 990–1004.

Elvers KT, Leeming K, Lappin-Scott HM. (2002). Binary and mixed population biofilms: time-lapse image analysis and disinfection with biocides. *J Ind Microbiol Biotechnol* **29**: 331–338.

624 Feldman M, Ginsburg I, Al-Quntar A, Steinberg D. (2016). Thiazolidinedione-8 Alters Symbiotic
625 Relationship in *C. albicans*-*S. mutans* Dual Species Biofilm. *Front Microbiol* **7**: 140.

626 Fiegna F, Moreno-Letelier A, Bell T, Barraclough TG. (2015). Evolution of species interactions
627 determines microbial community productivity in new environments. *Isme J* **9**: 1235–1245.

628 Foster KR, Bell T. (2012). Competition, not cooperation, dominates interactions among culturable
629 microbial species. *Curr Biol* **22**: 1845–50.

630 Fox JW. (2005). Interpreting the ‘selection effect’ of biodiversity on ecosystem function. *Ecol Lett* **8**:
631 846–856.

632 Ghoul M, Mitri S. (2016). The Ecology and Evolution of Microbial Competition. *Trends Microbiol* **24**:
633 833–845.

634 Giaouris E, Chorianopoulos N, Doulgeraki A, Nychas GJ. (2013). Co-Culture with *Listeria*
635 monocytogenes within a Dual-Species Biofilm Community Strongly Increases Resistance of
636 *Pseudomonas putida* to Benzalkonium Chloride. *PLoS One* **8**: 1–14.

637 Gkana EN, Giaouris ED, Doulgeraki AI, Kathariou S, Nychas GJE. (2017). Biofilm formation by
638 *Salmonella* Typhimurium and *Staphylococcus aureus* on stainless steel under either mono- or dual-
639 species multi-strain conditions and resistance of sessile communities to sub-lethal chemical
640 disinfection. *Food Control* **73**: 838–846.

641 Hall-Stoodley L, Costerton JW, Stoodley P. (2004). Bacterial biofilms: from the natural environment
642 to infectious diseases. *Nat Rev Microbiol* **2**: 95–108.

643 Harcombe W. (2010). Novel cooperation experimentally evolved between species. *Evolution* **64**:
644 2166–72.

645 Harriott MM, Noverr MC. (2009). *Candida albicans* and *Staphylococcus aureus* form polymicrobial
646 biofilms: Effects on antimicrobial resistance. *Antimicrob Agents Chemother* **53**: 3914–3922.

647 Hengge R. (2009). Principles of c-di-GMP signalling in bacteria. *Nat Rev Microbiol* **7**: 263–273.

648 Hill KE, Malic S, McKee R, Rennison T, Harding KG, Williams DW, *et al.* (2010). An in vitro model of
649 chronic wound biofilms to test wound dressings and assess antimicrobial susceptibilities. *J*
650 *Antimicrob Chemother* **65**: 1195–1206.

651 Hoffman LR, Déziel E, D’Argenio D a, Lépine F, Emerson J, McNamara S, *et al.* (2006). Selection for
652 *Staphylococcus aureus* small-colony variants due to growth in the presence of *Pseudomonas*
653 *aeruginosa*. *Proc Natl Acad Sci U S A* **103**: 19890–19895.

654 Jagmann N, Henke SF, Philipp B. (2015). Cells of *Escherichia coli* are protected against severe
655 chemical stress by co-habiting cell aggregates formed by *Pseudomonas aeruginosa*. *Appl Microbiol*
656 *Biotechnol* **99**: 8285–8294.

657 Kart D, Tavernier S, Van Acker H, Nelis HJ, Coenye T. (2014). Activity of disinfectants against
658 multispecies biofilms formed by *Staphylococcus aureus* , *Candida albicans* and *Pseudomonas*
659 *aeruginosa*. *Biofouling* **30**: 377–383.

660 Kelsic ED, Zhao J, Vetsigian K, Kishony R. (2015). Counteraction of antibiotic production and
661 degradation stabilizes microbial communities. *Nature* **521**: 516–519.

662 Kim W, Racimo F, Schluter J, Levy SB, Foster KR. (2014). Importance of positioning for microbial
663 evolution. *Proc Natl Acad Sci U S A* **111**: E1639-47.

664 Kostaki M, Chorianopoulos N, Braxou E, Nychas GJ, Giaouris E. (2012). Differential biofilm formation
665 and chemical disinfection resistance of sessile cells of *Listeria monocytogenes* strains under
666 monospecies and dual-species (with *Salmonella enterica*) conditions. *Appl Environ Microbiol* **78**:
667 2586–2595.

668 Kumar A, Peng TY. (2015). Presence of *Pseudomonas aeruginosa* influences biofilm formation and
669 surface protein expression of *Staphylococcus aureus*. *Environ Microbiol* **17**: 4459–4468.

670 Lee KWK, Periasamy S, Mukherjee M, Xie C, Kjelleberg S, Rice S a. (2014). Biofilm development and
671 enhanced stress resistance of a model, mixed-species community biofilm. *ISME J* **8**: 894–907.

672 Leriche V, Briandet R, Carpentier B. (2003). Ecology of mixed biofilms subjected daily to a chlorinated
673 alkaline solution: spatial distribution of bacterial species suggests a protective effect of one species
674 to another. *Environ Microbiol* **5**: 64–71.

675 Lindsay D, Brözel VS, Mostert JF, Von Holy A. (2002). Differential efficacy of a chlorine dioxide-
676 containing sanitizer against single species and binary biofilms of a dairy-associated *Bacillus cereus*
677 and a *Pseudomonas fluorescens* isolate. *J Appl Microbiol* **92**: 352–361.

678 Lopes SP, Ceri H, Azevedo NF, Pereira MO. (2012). Antibiotic resistance of mixed biofilms in cystic
679 fibrosis: impact of emerging microorganisms on treatment of infection. *Int J Antimicrob Agents* **40**:
680 260–263.

681 Loreau M. (2000). Biodiversity and ecosystem functioning: recent theoretical advances. *Oikos* **91**: 3–
682 17.

683 Loreau M, Hector A. (2001). Partitioning selection and complementarity in biodiversity experiments.
684 *Nature* **412**: 72–6.

685 Loreau M, de Mazancourt C. (2013). Biodiversity and ecosystem stability: a synthesis of underlying
686 mechanisms. *Ecol Lett* **16**: 106–115.

687 Lories B, Parijs I, Foster KR, Steenackers HP. (2017) Meeting report on the ASM Conference on
688 Mechanisms of Interbacterial Cooperation and Competition. *J Bacteriol.* (in press)

689 Luppens SBI, Kara D, Bandounas L, Jonker MJ, Wittink FR a, Bruning O, *et al.* (2008). Effect of
690 *Veillonella parvula* on the antimicrobial resistance and gene expression of *Streptococcus mutans*
691 grown in a dual-species biofilm. *Oral Microbiol Immunol* **23**: 183–9.

692 Lynch AS, Abbanat D. (2010). New antibiotic agents and approaches to treat biofilm-associated

693 infections. *Expert Opin Ther Pat* **20**: 1373–87.

694 Madsen JS, Røder HL, Russel J, Sørensen H, Burmølle M, Sørensen SJ. (2016). Coexistence facilitates
695 interspecific biofilm formation in complex microbial communities. *Environ Microbiol* **18**: 2565–2574.

696 Maifreni M, Frigo F, Bartolomeoli I, Buiatti S, Picon S, Marino M. (2015). Bacterial biofilm as a
697 possible source of contamination in the microbrewery environment. *Food Control* **50**: 809–814.

698 Mamvura T, Iyuke S. (2011). Soil Films in the Beverage Industry: A Review. *J Inst Brew* **117**: 608–616.

699 Mitri S, Foster KR. (2013). The genotypic view of social interactions in microbial communities. *Annu*
700 *Rev Genet* **47**: 247–73.

701 Morin X, Fahse L, Scherer-Lorenzen M, Bugmann H. (2011). Tree species richness promotes
702 productivity in temperate forests through strong complementarity between species. *Ecol Lett* **14**:
703 1211–1219.

704 Mozina SS, Klancknik A, Raspor P. (2013). Mechanisms of microbial resistance in biofilms. In: *Biofilms*
705 *in Bioengineering*. pp 311–332.

706 Nithya C, Devi MG, Karutha Pandian S. (2011). A novel compound from the marine bacterium
707 *Bacillus pumilus* S6-15 inhibits biofilm formation in gram-positive and gram-negative species.
708 *Biofouling* **27**: 519–28.

709 Norwood DE, Gilmour A. (2000). The growth and resistance to sodium hypochlorite of *Listeria*
710 *monocytogenes* in a steady-state multispecies biofilm. *J Appl Microbiol* **88**: 512–520.

711 Oliveira NM, Martinez-Garcia E, Xavier J, Durham WM, Kolter R, Kim W, *et al.* (2015). Biofilm
712 formation as a response to ecological competition. *PLoS Biol* **13**: 1–23.

713 Opperman TJ, Kwasny SM, Williams JD, Khan AR, Peet NP, Moir DT, *et al.* (2009). Aryl rhodanines
714 specifically inhibit staphylococcal and enterococcal biofilm formation. *Antimicrob Agents Chemother*
715 **53**: 4357–67.

716 Pfeiffer T, Schuster S, Bonhoeffer S. (2001). Cooperation and Competition in the Evolution of ATP-
717 Producing Pathways. *Science (80-)* **292**: 504–507.

718 Polley HW, Wilsey BJ, Derner JD. (2003). Do species evenness and plant density influence the
719 magnitude of selection and complementarity effects in annual plant species mixtures? *Ecol Lett* **6**:
720 248–256.

721 Rakoff-Nahoum S, Foster KR, Comstock LE. (2016). The evolution of cooperation within the gut
722 microbiota. *Nature* **533**: 255–259.

723 Ren D, Madsen JS, Sørensen SJ, Burmølle M. (2015). High prevalence of biofilm synergy among
724 bacterial soil isolates in cocultures indicates bacterial interspecific cooperation. *ISME J* **9**: 81–89.

725 Riley MA, Gordon DM. (1999). The ecological role of bacteriocins in bacterial competition. *Trends*
726 *Microbiol* **7**: 129–133.

727 Rivett DW, Scheuerl T, Culbert CT, Mombrikotb SB, Johnstone E, Barraclough TG, *et al.* (2016).
728 Resource-dependent attenuation of species interactions during bacterial succession. *ISME J* **10**:
729 2259–2268.

730 Røder HL, Raghupathi PK, Herschend J, Brejnrod A, Knøchel S, Sørensen SJ, *et al.* (2015). Interspecies
731 interactions result in enhanced biofilm formation by co-cultures of bacteria isolated from a food
732 processing environment. *Food Microbiol* **51**: 18–24.

733 Røder HL, Sørensen SJ, Burmølle M. (2016). Studying Bacterial Multispecies Biofilms: Where to Start?
734 *Trends Microbiol* **24**: 503–513.

735 Romling U, Galperin MY, Gomelsky M. (2013). Cyclic di-GMP: the first 25 years of a universal
736 bacterial second messenger. *Microbiol Mol Biol Rev* **77**: 1–52.

737 Rosenberg G, Steinberg N, Oppenheimer-shaan Y, Olender T, Doron S, Ben-ari J. (2016). Not so
738 simple , not so subtle : the interspecies competition between *Bacillus simplex* and *Bacillus subtilis*

739 and its impact on the evolution of biofilms. *Biofilms and Microbiomes* **2**: 1–11.

740 Le Roux M, Kirkpatrick RL, Montauti EI, Tran BQ, Brook Peterson S, Harding BN, *et al.* (2015). Kin cell
741 lysis is a danger signal that activates antibacterial pathways of pseudomonas aeruginosa. *Elife* **2015**:
742 1–65.

743 Russell AB, Peterson SB, Mougous JD. (2014). Type VI secretion system effectors: poisons with a
744 purpose. *Nat Rev Microbiol* **12**: 137–148.

745 Scholz RL, Greenberg EP. (2015). Sociality in Escherichia coli: Enterochelin is a private good at low
746 cell density and can be shared at high cell density. *J Bacteriol* **197**: 2122–2128.

747 Schwering M, Song J, Louie M, Turner RJ, Ceri H. (2013). Multi-species biofilms defined from drinking
748 water microorganisms provide increased protection against chlorine disinfection. *Biofouling* **29**:
749 917–28.

750 Shakeri S, Kermanshahi RK, Moghaddam MM, Emtiazi G. (2007). Assessment of biofilm cell removal
751 and killing and biocide efficacy using the microtiter plate test. *Biofouling* **23**: 79–86.

752 Simões LC, Simões M, Vieira MJ. (2010). Influence of the diversity of bacterial isolates from drinking
753 water on resistance of biofilms to disinfection. *Appl Environ Microbiol* **76**: 6673–6679.

754 Simões M, Simões LC, Vieira MJ. (2009). Species association increases biofilm resistance to chemical
755 and mechanical treatments. *Water Res* **43**: 229–237.

756 Spehn EM, Hector A, Joshi J, Scherer-Lorenzen M, Schmid B, Bazeley-White E, *et al.* (2005).
757 Ecosystem effects of biodiversity manipulations in European grasslands. *Ecol Monogr* **75**: 37–63.

758 Stoodley P, Sauer K, Davies DG, Costerton JW. (2002). Biofilms as complex differentiated
759 communities. *Annu Rev Microbiol* **56**: 187–209.

760 Storgards E, Tapani K. (2006). Microbial attachment and biofilm formation in brewery bottling
761 plants. *Am Soc Brew Chem* **64**: 8–15.

762 Sun Y, Dowd SE, Smith E, Rhoads DD, Wolcott RD. (2008). In vitro multispecies Lubbock chronic
763 wound biofilm model. *Wound Repair Regen* **16**: 805–13.

764 Tan CH, Kelvin Lee KW, Burmølle M, Kjelleberg S, Rice SA, Lee KWK, *et al.* (2016). All Together Now:
765 Experimental Multispecies Biofilm Model Systems. *Environ Microbiol* **13**: 385–392.

766 Traxler MF, Watrous JD, Alexandrov T, Dorrestein PC, Kolter R. (2013). Interspecies interactions
767 stimulate diversification of the *Streptomyces coelicolor* secreted metabolome. *MBio* **4**: 1–12.

768 Turnbull LA, Levine JM, Loreau M, Hector A. (2013). Coexistence, niches and biodiversity effects on
769 ecosystem functioning. *Ecol Lett* **16**: 116–127.

770 Van der Veen S, Abee T. (2011). Mixed species biofilms of *Listeria monocytogenes* and *Lactobacillus*
771 *plantarum* show enhanced resistance to benzalkonium chloride and peracetic acid. *Int J Food*
772 *Microbiol* **144**: 421–431.

773 Wang R, Kalchayanand N, Schmidt JW, Haray DM. (2013). Mixed Biofilm Formation by Shiga Toxin-
774 Producing *Escherichia coli* and *Salmonella enterica* Serovar Typhimurium Enhanced Bacterial
775 Resistance to Sanitization due to Extracellular Polymeric Substances. *J Food Prot* **76**: 1513–1522.

776 West SA, Griffin AS, Gardner A. (2007). Social semantics: Altruism, cooperation, mutualism, strong
777 reciprocity and group selection. *J Evol Biol* **20**: 415–432.

778 White TJ, Bruns S, Lee S, Taylor J. (1990). Amplification and direct sequencing of fungal ribosomal
779 RNA genes for phylogenetics. *PCR Protoc A Guid to Methods Appl* 315–322.

780 Whiteley M, Ott JR, Weaver EA, McLean RJC. (2001). Effects of community composition and growth
781 rate on aquifer biofilm bacteria and their susceptibility to betadine disinfection. *Environ Microbiol* **3**:
782 43–52.

783 Yassin SA, German MJ, Rolland SL, Rickard AH, Jakubovics NS. (2016). Inhibition of multispecies
784 biofilms by a fluoride-releasing dental prosthesis copolymer. *J Dent* **48**: 62–70.