

### Oncolmmunology



U

ISSN: (Print) 2162-402X (Online) Journal homepage: http://www.tandfonline.com/loi/koni20

### A genetic IFN/STAT1/FAS axis determines CD4 T stem cell memory levels and apoptosis in healthy controls and Adult T-cell Leukemia patients

Ricardo Khouri, Gilvanéia Silva-Santos, Tim Dierckx, Soraya Maria Menezes, Daniele Decanine, Kristof Theys, Aline Clara Silva, Lourdes Farré, Achiléa Bittencourt, Massimo Mangino, Mario Roederer, Anne-Mieke Vandamme & Johan Van Weyenbergh

To cite this article: Ricardo Khouri, Gilvanéia Silva-Santos, Tim Dierckx, Soraya Maria Menezes, Daniele Decanine, Kristof Theys, Aline Clara Silva, Lourdes Farré, Achiléa Bittencourt, Massimo Mangino, Mario Roederer, Anne-Mieke Vandamme & Johan Van Weyenbergh (2018): A genetic IFN/STAT1/FAS axis determines CD4 T stem cell memory levels and apoptosis in healthy controls and Adult T-cell Leukemia patients, Oncolmmunology, DOI: 10.1080/2162402X.2018.1426423

To link to this article: https://doi.org/10.1080/2162402X.2018.1426423

9	© 2018 The Author(s). Published with license by Taylor & Francis Group, LLC© Ricardo Khourim, Gilvanéia Silva-Santos,	+	View supplementary material 🕼
	Tim Dierckx, Soraya Maria Menezes, Daniele Decanine, Kristof Theys, Aline Clara Silva, Lourdes Farré, Achiléa Bittencourt, Massimo Mangino, Mario Roederer, Anne-Mieke Vandamme and Johan Van Weyenbergh		
	Accepted author version posted online: 15 Jan 2018. Published online: 13 Feb 2018.		Submit your article to this journal 🖸
111	Article views: 25	Q	View related articles $\square$
	View Crossmark data 🗹		

#### **ORIGINAL RESEARCH**

OPEN ACCESS OPEN ACCESS

### A genetic IFN/STAT1/FAS axis determines CD4 T stem cell memory levels and apoptosis in healthy controls and Adult T-cell Leukemia patients

Ricardo Khouri <sup>(Da,b</sup>, Gilvanéia Silva-Santos<sup>b</sup>, Tim Dierckx<sup>a</sup>, Soraya Maria Menezes<sup>a</sup>, Daniele Decanine<sup>b</sup>, Kristof Theys<sup>a</sup>, Aline Clara Silva<sup>b</sup>, Lourdes Farré<sup>b</sup>, Achiléa Bittencourt<sup>c</sup>, Massimo Mangino<sup>d</sup>, Mario Roederer<sup>e</sup>, Anne-Mieke Vandamme <sup>(Da,f</sup>, and Johan Van Weyenbergh<sup>a</sup>

<sup>a</sup>KU Leuven – University of Leuven, Department of Microbiology and Immunology, Rega Institute for Medical Research, Clinical and Epidemiological Virology, Leuven, Belgium; <sup>b</sup>Instituto Gonçalo Moniz (IGM) - Fundação Oswaldo Cruz (FIOCRUZ), Salvador-Bahia, Brazil; <sup>c</sup>Department of Pathology, Complexo Hospitalar Universitário Prof Edgard Santos, Universidade Federal da Bahia (UFBA), Salvador-Bahia, Brazil; <sup>d</sup>Department of Twin Research & Genetic Epidemiology, King's College, London, UK; <sup>e</sup>Immunotechnology Section, Vaccine Research Center, NIAID, NIH, Bethesda-MD, USA; <sup>f</sup>Center for Global Health and Tropical Medicine, Unidade de Microbiologia, Instituto de Highne e Medicina Tropical, Universidade, Nova de Lisboa, Lisbon, Portugal

#### ABSTRACT

Adult T-cell leukemia (ATL) is an aggressive, chemotherapy-resistant  $CD4^+CD25^+$  leukemia caused by HTLV-1 infection, which usually develops in a minority of patients several decades after infection. IFN + AZT combination therapy has shown clinical benefit in ATL, although its mechanism of action remains unclear. We have previously shown that an IFN-responsive *FAS* promoter polymorphism in a STAT1 binding site (rs1800682) is associated to ATL susceptibility and survival. Recently, CD4 T stem cell memory ( $T_{SCM}$ ) Fas<sup>hi</sup> cells have been identified as the hierarchical cellular apex of ATL, but a possible link between FAS, apoptosis, proliferation and IFN response in ATL has not been studied.

In this study, we found significant ex vivo antiproliferative, antiviral and immunomodulatory effects of IFN- $\alpha$  treatment in short-term culture of primary mononuclear cells from ATL patients (n = 25). Bayesian Network analysis allowed us to integrate ex vivo IFN- $\alpha$  response with clinical, genetic and immunological data from ATL patients, thereby revealing a central role for *FAS* -670 polymorphism and apoptosis in the coordinated mechanism of action of IFN- $\alpha$ . *FAS* genotype-dependence of IFN-induced apoptosis was experimentally validated in an independent cohort of healthy controls (n = 20). The same *FAS* -670 polymorphism also determined CD4 T<sub>SCM</sub> levels in a genome-wide twin study (p = 7 × 10<sup>-11</sup>, n = 460), confirming a genetic link between apoptosis and T<sub>SCM</sub> levels. Transcriptomic analysis and cell type deconvolution confirmed the *FAS* genotype/T<sub>SCM</sub> link and IFN- $\alpha$ -induced downregulation of CD4 T<sub>SCM</sub>-specific genes in ATL patient cells.

In conclusion, ex vivo IFN- $\alpha$  treatment exerts a pleiotropic effect on primary ATL cells, with a genetic IFN/STAT1/Fas axis determining apoptosis vs. proliferation and underscoring the CD4 T<sub>SCM</sub> model of ATL leukemogenesis.

#### Introduction

Adult T-cell leukemia (ATL) is characterized by circulating CD4<sup>+</sup>CD25<sup>+</sup> T-cells<sup>1</sup> and is etiologically linked to infection with Human T-cell Leukemia Virus 1 (HTLV-1), the first isolated human pathogenic retrovirus.<sup>2</sup> The estimated lifetime risk is about 5% in infected individuals,<sup>3</sup> with a long incubation period after HTLV-1 infection, therefore infection early in life is fundamental in the development of ATL.<sup>4</sup> ATL is classified according to clinical and laboratory criteria as smoldering, chronic, lymphoma, or acute subtypes.<sup>5</sup> The median survival time for acute and lymphoma (aggressive) subtypes is less than one year, whereas patients with chronic and smoldering (indolent) subtypes survive longer.<sup>6</sup> In addition, an atypical aggressive, primary cutaneous tumoral form (PCT) of ATL with a shorter survival has been described.<sup>7</sup> The exceptional oncogenicity of HTLV-1, as compared to other viruses or infectious agents,<sup>8</sup> or even to its closest relative HTLV-2,<sup>9</sup>, is most likely due to its ability to deregulate several host cell signaling pathways. In sharp contrast to HIV, plasma viral load for HTLV-1 is mostly undetectable and proviral replication is driven by spontaneous lymphoproliferation (reviewed in<sup>3</sup>), which implies that ATL might be explained mainly by host factors such as immune response and host genetics.

Regarding host immune response, interleukin 2 (IL-2) promotes proliferation of IL-2 receptor-positive (IL- $2R^+/CD25^+$ ) T-

Supplemental data for this article can be accessed on the publisher's website.

#### ARTICLE HISTORY

Received 17 January 2017 Revised 04 January 2018 Accepted 07 January 2018

#### **KEYWORDS**

CD95; HTLV-1; genetics; interferon; leukemia; lymphoma; oncogenesis; retrovirus; signaling; stem cell; twins



**CONTACT** Johan Van Weyenbergh 🔊 j.vw@live.be, johan.vanweyenbergh@kuleuven.be 🖻 Clinical and Epidemiological Virology, Rega Institute for Medical Research, Department of Microbiology

<sup>&</sup>amp; Immunology, KU Leuven, Herestraat 49, box 1040 B-3000 Leuven, Belgium.

<sup>© 2018</sup> Ricardo Khourim, Gilvanéia Silva-Santos, Tim Dierckx, Soraya Maria Menezes, Daniele Decanine, Kristof Theys, Aline Clara Silva, Lourdes Farré, Achiléa Bittencourt, Massimo Mangino, Mario Roederer, Anne-Mieke Vandamme and Johan Van Weyenbergh. Published with license by Taylor & Francis Group, LLC

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (http://creativecommons.org/licenses/by-nc-nd/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

cells. Membrane-bound CD25 is present on all activated T-cells as well as malignant T-cells infected by HTLV-1. Serum levels of IL-2R/CD25 positively correlate with circulating ATL cells (CD4<sup>+</sup>CD25<sup>+</sup>) and disease severity.<sup>10</sup> In addition, systemic IL-6 levels correlate with aggressive clinical forms and short survival.<sup>11</sup>

Regarding host genetics, FAS (TNFRSF6/Apo-1/CD95) and TP53 polymorphisms have been associated to ATL disease progression.<sup>12,13</sup> In addition, FAS mutations have been demonstrated in ATL<sup>14,15</sup> as well as in other leukemias.<sup>16</sup> The FAS -670 A/G polymorphism is situated in a STAT1 binding site, for which the A allele has a higher binding affinity.<sup>17,18</sup> We have previously shown that the A allele corresponds to increased IFN-induced Fas mRNA, and is significantly overrepresented in ATL patients, as compared to healthy controls or HTLV-1-infected asymptomatic individuals.<sup>12</sup> In addition, ATL patients with the AA genotype were significantly more likely to develop aggressive (acute/lymphoma) clinical forms.<sup>12</sup> Interestingly, a specific subset of Fashi cells has recently been implicated in ATL leukemogenesis, as CD4 T stem cell memory (T<sub>SCM</sub>) cells were identified as its hierarchical cellular apex.<sup>19</sup>

A combination of interferon- $\alpha$  (IFN- $\alpha$ ) and zidovudine (AZT) has been found effective in pioneer ATL trials, and was confirmed by meta-analysis for non-lymphoma subtypes.<sup>20-22</sup> In spite of this clinical benefit of IFN- $\alpha$  + AZT combination therapy in ATL, its mechanisms of action remain elusive. IFN- $\alpha$  has anti-proliferative, pro-apoptotic, antiviral and immunomodulatory activity in many human cells, but limited in vitro, ex vivo or in vivo data are available in ATL.<sup>23</sup> HTLV-1 infected and/or ATL-derived cell lines are refractory to IFN-a-induced cell death in the absence or presence of  $AZT^{23-25}$  and to IFN- $\beta$ in vitro.<sup>26</sup> However, we have recently demonstrated a superior antiproliferative and pro-apoptotic activity of IFN- $\beta$  vs. IFN- $\alpha$ in primary ATL patient cells.<sup>27</sup> Moreover, a wealth of studies has demonstrated both pro- and anti-apoptotic activity of IFN- $\alpha/\beta$  in vitro and in vivo,<sup>28-33</sup> resulting in an "IFN apoptotic paradox" on the effect of type I IFN (IFN- $\alpha/\beta$ ) on programmed cell death. On the other hand, the clinical benefit of type I IFN has been convincingly demonstrated in several other leukemias, such as hairy-cell leukemia, chronic myeloid leukemia as well as myeloproliferative neoplasms, including essential thrombocytosis, polycythemia vera, and myelofibrosis.34-36

We therefore engaged in a systematic study of IFN- $\alpha$  response and its possible link to Fas levels, apoptosis, proliferation, viral protein expression and immune activation in ATL.

#### Methods

#### Patient recruitment

This study was approved by the Ethics Review Board of HUPES, according to the Declaration of Helsinki principles. All study participants signed informed consent. In this prospective study, consecutive patients were recruited between 2001 and 2005 at "Hospital Universitário Professor Edgar Santos" (HUPES, Salvador-Bahia), according to previously described inclusion and exclusion criteria.<sup>7</sup> Samples were obtained from 25 patients with clinically definite ATL

(Shimoyama criteria<sup>5</sup>), with serology, inverted PCR and/or flow cytometry as described.<sup>12,37,38</sup> All ATL patients were HTLV-1 seropositive and HIV/HTLV-2 seronegative. Standardized patient treatment was in agreement to a published international consensus.<sup>6</sup> Smoldering ATL forms were left untreated but under "watchful waiting", until possible disease progression, whereas acute/chronic leukemic patients received IFN- $\alpha$ +AZT combination therapy and lymphoma patients received chemotherapy. Samples were obtained before treatment and at least five-year follow-up was available for each patient. Clinical, demographic and flow cytometry data of ATL patients are summarized in Table I. Twenty healthy controls from the same endemic area (seronegative for HTLV-1/HTLV-2/HIV) were recruited in parallel.

#### **Ex vivo** *IFN-\alpha stimulation*

PBMCs from patients and healthy controls were purified from heparinized venous blood by Ficoll-Hypaque gradient (Sigma-Aldrich). Cells were plated in 24-well tissue culture plates (Costar, Corning Incorporated) at  $4 \times 10^6$  cells/ml in RPMI1640 medium with 2mM L-glutamine, gentamycin (50 µg/ml) and 10% heat-inactivated fetal calf serum (Gibco) and incubated at 37°C, 5% CO2, for 48h in the presence or absence of IFN- $\alpha$ 2A (1,000 U/ml, Blausiegel Ltda., SP-Brazil). Cells were immediately processed for bioassays (apoptosis, proliferation), flow cytometry or RNA extraction.

#### Flow cytometry

PBMC from Brazilian patients and controls were resuspended at 200,000 cells/50  $\mu$ l (1% BSA + 0.1% NaN<sub>3</sub> in PBS) and incubated for 30 min on ice with antibodies (CD3/CD4/CD25/ CD95) and corresponding isotype controls (BD Biosciences). Samples were acquired using FACSort (BD Biosciences) and analyzed using CellQuest software. CD4 T<sub>SCM</sub> phenotyping (CD3<sup>+</sup>CD4<sup>+</sup>CD8<sup>-</sup>CD45RA<sup>+</sup>CD57<sup>-</sup>CD95<sup>+</sup>CD27<sup>+</sup>CD127 <sup>+</sup>CD28<sup>+</sup>) and quantification was performed in the UK Twin cohort,<sup>39,40</sup> using the gating strategy outlined in Figure 1.

## FAS genotyping of Brazilian ATL patients and healthy controls

Genotyping for *FAS* -670 polymorphism by PCR-restriction fragment length polymorphism was performed as described.<sup>12</sup> Patient genotype distribution (Table I) is in agreement with our previous findings,<sup>12</sup> with the AA genotype being overrepresented in ATL patients (37%), as compared to healthy controls (10% AA, 30% GA, 60% GG, n = 20).

#### Combined GWAS and mass flow cytometry study of UK Twin cohort

This study was approved by the NIAID (NIH) IRB and London-Westminster NHS Research Ethics Committee; all participants provided informed consent. The discovery stage comprised 497 female participants from the UK Adult Twin Register, TwinsUK, with full genotyping data on 460 subjects.



Figure 1. Gating strategy for Tscm cells. CD4 T<sub>SCM</sub> phenotyping and quantification in the UK Twin cohort<sup>39</sup> was performed by consecutive gating on CD3<sup>+</sup>CD4<sup>+</sup>CD8<sup>-</sup>, naïve (CD45RA<sup>+</sup>) and CD57<sup>-</sup>CD95<sup>+</sup>CD27<sup>+</sup>CD127<sup>+</sup>CD28<sup>+</sup> lymphocytes.

The TwinsUK cohort is described in detail in.<sup>39</sup> GWAS and immunophenotyping are described in detail in.<sup>40</sup>

#### **Microarray analysis**

Total RNA from PBMCs was extracted according to manufacturer's protocol (RNeasy kit QIAgen, Venlo, Netherlands) from ATL patients (n = 8, with and without IFN- $\alpha$  treatment). Whole genome microarray was performed at the VIB Nucleomics Facility (Leuven, Belgium), using the Human Gene 1.0 ST Array with the WT PLUS reagent kit

Table 1. Clinical, der	nographic and	molecular	features o	f ATL	patients.
------------------------	---------------	-----------	------------	-------	-----------

Features	Proportion of ATL patients			
Clinical forms:				
Smoldering	48% (12/25)			
Chronic	8% (2/25)			
Acute	28% (7/25)			
Lymphoma	8% (2/25)			
PCT	8% (2/25)			
FAS -670 genotype				
GG	37% (7/19)			
GA	26% (5/19)			
AA	37% (7/19)			
Gender (Proportion of female)	56% (14/25)			
Age at diagnosis (Years)	48 (IQR = 39-61.5) (N = 25)			
CD4 <sup>+</sup> (%)	63.88%(IQR = 41.54-83.97) (N = 21)			
CD4 <sup>+</sup> CD25 <sup>+</sup> (%)	9.48%(IQR = 2.18-24.01) (N = 20)			
Fas <sup>hi</sup> (%)	65.80%(IQR = 34.58-83.59) (N = 22)			

(Affymetrix, Santa Clara, CA) according to manufacturer's instructions. Data were analyzed as previously described<sup>27</sup> and available at the National Center for Biotechnology Information Gene Expression Omnibus under accession number GSE85487. Expression data from GSE23321 was used to recapitulate the differentiation gradient demonstrated for CD8  $T_N$ ,  $T_{SCM}$ ,  $T_{CM}$  and  $T_{EM}$  cell populations<sup>41</sup> through Principal Component Analysis, using the published 278  $T_{SCM}$ -specific genes determined by Gattinoni et al.<sup>41</sup>

#### Measurement of cytokine production

Cytokine levels in 48h culture supernatants (kept frozen at  $-80^{\circ}$ C immediately after harvesting) were measured using Cytometric Bead Array Human Th1/Th2 Cytokine (BD Biosciences), following the manufacturer's instructions.

#### **Quantification of HTLV-1 p19 expression**

Viral p19 protein levels<sup>24,33</sup> in culture supernatants were measured using the HTLV-I/II p19 antigen ELISA (ZeptoMetrix), following the manufacturer's instructions.

#### **Proliferation assays**

PBMCs from ATL patients proliferate spontaneously upon culturing, while in healthy controls, proliferation was induced by



**Figure 2.** Pleiotropic *ex vivo* effect of IFN- $\alpha$  in PBMC of ATL patients. PBMC from ATL patients were treated *ex vivo* with IFN- $\alpha$  (1000 U/ml) and assessed for (A) proliferation (<sup>3</sup>H-thymidine incorporation), (B) apoptosis (% of annexin V+ cells), (C) virus expression (p19 pg/ml) and (D, E, F) immunomodulation (IL-2, IL-6 and FasL expression, respectively), as described in Methods. A scatter dot plot of within-patient paired observations connected by lines of control non-treated and *ex vivo* IFN- $\alpha$ -treated PBMC (Wilcoxon matched pairs test, \*p < 0.05, \*\*p < 0.01), and their IFN- $\alpha$  response (IFN- $\alpha$  treated/control value) are indicated in each graph. No IFN- $\alpha$  response is indicated by a dotted line.

anti-CD3 antibody stimulation. *Ex vivo* IFN-  $\alpha$  stimulation (120h) was as above, except for plating in 96-well U-bottom plates (200  $\mu$ l/well at 1 × 10<sup>6</sup> cells/ml). Lymphoproliferation was quantified by [<sup>3</sup>H]-thymidine incorporation after a 12–16h pulse (1 $\mu$ Ci/well), using gas phase scintillation (Direct Beta Counter Matrix9600, PerkinElmer Life Sciences). Results are expressed as the mean counts per minute in triplicate cultures.

#### Apoptosis assays

Apoptosis was measured by microscopic quantification of nuclear fragmentation in at least 100 cells (duplicate Hoechst33432 or hematoxylin/eosin staining), as well as flow cytometry (annexin V staining), as above.

#### Bayesian network learning

A Bayesian network (BN) is a probabilistic graphical model that describes statistical conditional dependencies between multiple variables. Although originally developed for large pharmacogenetic data sets, Bayesian network learning can discover robust interactions between variables in small datasets.<sup>42,43</sup> Dependencies are visualized in a directed acyclic graph and form the qualitative component of the BN. In this graph, each node corresponds to an attribute, and a direct arc between nodes

represents a direct influence. Mathematically, a Bayesian network provides a refractoring of the Joint Probability Distribution (JPD) of the data, using Bayes' rules. As a BN simplifies the JPD, it provides an effective model that summarizes statistical properties of the data. In this way, the best Bayesian network is searched that explains a maximum of the observed associations in the data using a minimum number of direct influences. Bayesian network learning was performed using Bcourse software adapted by Deforche et al.<sup>44</sup> Herein, we learned Bayesian networks from observation of continuous attributes discretized as quartiles (age, proportion of CD4<sup>+</sup>, CD4<sup>+</sup>CD25<sup>+</sup>, and Fas<sup>+</sup> cells). Gender was discretized as Male/ Female, clinical forms as smoldering, chronic, acute and lymphoma/PCT. FAS -670 genotypes were discretized as GG/GA/ AA. All IFN- $\alpha$  responses (proliferation/apoptosis/p19/IL-2/IL-6/FasL, Fig. 2) were discretized as positive, negative or neutral. The confidence interval of neutral effect  $(1\pm0.19)$  was determined based on 80% power to detect a difference between positive and negative effect for the variable available for the lowest number of patients (i.e. 12 patients for apoptosis). In this nonlinear model, the arcs (dependency) were scored based on the stability of the conditional dependency, assessed with a nonparametric bootstrap (100x replicates).45 All arcs with bootstrap over 60% were considered and depicted in the consensus network.

#### Statistical analysis

Parametric and non-parametric tests were used according to Kolmogorov-Smirnov test for normality: Pearson/Spearman correlation, t-test/Mann-Whitney/Wilcoxon tests, all two-tailed (GraphPad Prism 5.0 software), differences were considered significant at p<0.05. To maximize data mining output, outliers (Tukey's test) were included for all univariate and multivariate analyses (Fig. 2–6) and were excluded only for post-hoc testing of the models (Supplementary Fig. S1-S2).

#### Results

# Ex vivo antiproliferative, pro-apoptotic, antiviral and immunomodulatory effect of IFN- $\alpha$ in PBMC of ATL patients

Freshly isolated primary cells (PBMC) of ATL patients were treated *ex vivo* with or without IFN- $\alpha$  for 48h and possible proapoptotic, anti-proliferative, antiviral and/or immunomodulatory activity was quantified (Fig. 2). To take into account strong inter-patient variation, data were normalized and expressed as IFN- $\alpha$  response (IFN- $\alpha$  value/control value). IFN- $\alpha$  significantly decreased *ex vivo* lymphoproliferation by a median of 41% (p = 0.027) (Fig. 2A). In contrast, the pro-apoptotic effect of IFN- $\alpha$  (median 43% increase) did not reach statistical significance (p = 0.11), but a large inter-patient variability was observed (Fig. 2B). As an antiviral readout,<sup>24,33</sup> IFN- $\alpha$  significantly decreased viral p19 protein levels in cell culture supernatants (median 60% decrease, p = 0.0015) (Fig. 2C). The immunomodulatory effect of IFN- $\alpha$  was examined by measuring Th1/Th2 cytokines in supernatants. No significant



**Figure 3.** IFN-*α* response is dependent on clinical and molecular variables: A Bayesian Network approach. Annotated Bayesian Network including the following 13 clinical, molecular and cellular attributes: Clinical forms, Age, Gender, *FAS* -670 genotype, CD4<sup>+</sup> %, CD4<sup>+</sup>CD25<sup>+</sup> %, Fas<sup>hi</sup> % (see Table 1) and all six IFN-*α* response measures (proliferation, apoptosis, p19, IL-2, IL-6 and FasL, see Fig. 2). Variables that are strongly interdependent are shown by arcs, solid arcs when the association is positive, dashed arcs when negative. The stability of the dependency was assessed with a non-parametric bootstrap (100x replicates). All arcs with bootstrap over 60% are depicted in the network. Apoptosis triggered by IFN-*α* is strongly dependent on the presence of the *FAS* -670 A allele, linked to aggressive clinical forms of ATL (blue circles and arcs), and inversely linked to IFN-*α* induced reduction of IL-2 levels. Likewise, with regard to reduction of IL-6, IFN-*α* response is more pronounced when there are fewer Fas<sup>+</sup> cells and when ATL is less aggressive; while IFN-*α* induced FasL induction is more pronounced when there are fewer CD4<sup>+</sup>CD25<sup>+</sup> cells (comprising ATL cells).

differences were observed for TNF- $\alpha$ , IFN- $\gamma$ , IL-2, IL-4 and IL-10 production (p > 0.05 for all, data not shown). However, a median 74% decrease in IL-2 production was observed after IFN- $\alpha$  treatment in all 3 IL-2 producing patients (Fig. 2D). Furthermore, IFN- $\alpha$  treatment significantly reduced IL-6 levels (median 99% decrease, p = 0.0039) (Fig. 2E). Fas ligand (FasL) levels were not significantly increased after IFN- $\alpha$  treatment (median 24% increase, p = 0.055) (Fig. 2F). To investigate possible interdependencies between the different biological measures, correlation between all six measures of IFN- $\alpha$  response was assessed. In these pairwise comparisons, we found that IFN- $\alpha$ -induced apoptotic and FasL response were positively correlated (r = 0.75, p = 0.013). Similarly, IFN- $\alpha$ -induced antiproliferative and antiviral activities were also positively correlated (r = 0.77, p = 0.003) (Supplementary Fig. S1A-B). Together, these results argue for a pleiotropic effect of IFN- $\alpha$  in primary ATL cells. Therefore, we decided to model the interdependencies between molecular, cellular and clinical data using a data mining approach.

### IFN- $\alpha$ response is dependent on clinical and molecular variables: A Bayesian Network approach

Clinical, cellular and molecular attributes were included and discretized as detailed in Methods. Among all 13 attributes, Bayesian network learning discovered several robust interactions: 6 arcs had a bootstrap support over 60% (Fig. 3). The most connected variable was IFN- $\alpha$ -triggered apoptosis, directly linked to FAS -670 genotype, aggressive ATL, and IFN- $\alpha$ -triggered IL-2 reduction. This confirmed that IFN- $\alpha$  triggered apoptosis does play a role in the overall mechanism of action of IFN-  $\alpha$ , but that it is heavily dependent on other variables, explaining why it was not significant in univariate analysis (Fig. 2B). This Bayesian network suggests that IFN- $\alpha$ -triggered apoptosis is strongly dependent on FAS -670 genotype and especially so in aggressive clinical forms of ATL (blue lines in Fig. 3), confirming our earlier findings.<sup>12</sup> In addition, IFN- $\alpha$ -triggered IL-6 reduction depends on the proportion of Fas<sup>+</sup> cells, while IFN- $\alpha$ -induced FasL levels depend on the proportion of CD4<sup>+</sup>CD25<sup>+</sup> (comprising ATL) cells, consistent with the reported importance of IL-6 and Fas in ATL pathogenesis.<sup>11-15</sup>

#### **Pro-apoptotic effect of IFN-** $\alpha$ is FAS -670 genotypedependent in both ATL patients and healthy controls

To confirm the dependencies seen in the BN, we examined IFN- $\alpha$  response in function of *FAS* -670 genotype. Indeed, the presence of the *FAS* -670 A allele was highly associated with IFN- $\alpha$ -inducible apoptosis (p = 0.0069) (Fig. 4A) but not antiproliferative or antiviral IFN- $\alpha$  effect (data not shown). No significant difference in IFN- $\alpha$  response was observed between homozygous AA and heterozygous GA individuals, indicating that one functional allele A is sufficient to determine the proapoptotic phenotype.

Due to its unusually high penetrance (80% of phenotypic difference explained by genotype), *FAS* -670 genotype-dependent apoptosis upon IFN- $\alpha$  stimulation was tested in an additional cohort of twenty healthy donors from the same endemic area. Lymphocytes from healthy donors were mostly Fas<sup>lo</sup>



**Figure 4.** *FAS* -670 genotype-dependent effects of IFN- $\alpha$  induced *ex vivo* apoptosis in PBMCs from ATL patients and healthy donors. PBMCs from ATL patients and healthy donors were treated *ex vivo* with IFN- $\alpha$  (1000U/ml) (see methods). Apoptosis (A and B, % annexinV+ cells) and FasL levels in supernatants (C and D) were stratified according to *FAS* -670 genotype (GG vs. GA+AA). (A) IFN- $\alpha$ -induced apoptosis (IFN- $\alpha$  value/control value) in PBMCs of ATL patients depends on *FAS* -670 genotype (unpaired t test, \*\*p = 0.0069). (B) IFN- $\alpha$ - induced apoptosis in PBMCs of healthy donors depends on *FAS* -670 genotype (Wilcoxon signed rank test, \*p = 0.023). (C) IFN- $\alpha$  does not increase FasL levels in supernatants of PBMCs of ATL patients, independent of *FAS* -670 genotype. (D) IFN- $\alpha$  significantly increases FasL levels in supernatants of PBMCs of ATL patients, independent of *FAS* -670 genotype. (D) IFN- $\alpha$  value/control value) does not differ according to *FAS* -670 genotype.

(16.8±2.0% Fas<sup>+</sup>, n = 20). Therefore, Fas-expressing cells were gated separately for apoptosis quantification by annexinV staining. A strikingly similar genotype-phenotype effect was observed in healthy donors (Fig. 4B), where IFN- $\alpha$  again selectively induced ex vivo apoptosis in AA and GA donors (p = 0.023, n = 8), but not in GG donors (p = 0.23, n = 12), confirming high penetrance (90%) of the phenotype.

As shown in Figure 2, there is no significant IFN- $\alpha$ -induced increase in FasL levels in ATL, and in Figure 3C it is shown that this remains so, independent of FAS -670 genotype. In



**Figure 5.** *FAS* -670 genotype-dependent proliferation in PBMC from healthy donors and ATL patients. Spontaneous proliferation of PBMCs from healthy donors and ATL patients and anti-CD3 induced proliferation for healthy donors were stratified according to *FAS* -670 genotype (GG vs. GA + AA) (Mann-Whitney test, \*p < 0.05).

healthy donors, FasL levels can be induced by IFN- $\alpha$ , again independent of *FAS* -670 genotype (Fig. 4D). Of note, compared to healthy controls, FasL levels in ATL patients are constitutively high, at levels that seem not to be further inducible. Taken together, these data suggest a pivotal role of the Fas receptor, rather than its ligand in IFN- $\alpha$ -induced apoptosis in both ATL patients and in healthy donors.

#### FAS -670 genotype determines TCR-triggered proliferation in PBMC of healthy donors

To investigate the possible effect of *FAS* -670 genotype on lymphoproliferation, PBMC from genotyped healthy donors were treated with anti-CD3 antibody to induce TCR-triggered polyclonal proliferation. Since this is also known to result in activation-induced, Fas-mediated cell death,<sup>46–48</sup> we expected a decrease in proliferation for the "pro-apoptotic" GA/AA genotypes. However, the presence of *FAS* -670 GA/AA genotypes resulted in significantly higher anti-CD3-stimulated proliferation (p = 0.049), as compared to GG homozygotes (Fig. 5), suggesting Fas-mediated proliferation may be independent of Fas-mediated apoptosis. Interestingly, a 3.4 fold increase in spontaneous lymphoproliferation was also observed in ATL patients with a functional *FAS* -670 A allele, although not statistically significant (p = 0.11) (Fig. 5).

## CD4<sup>+</sup>CD25<sup>+</sup> levels correlate positively with IFN- $\alpha$ induced apoptosis

In order to appreciate whether a different T-cell population might be proliferating vs. undergoing apoptosis, CD4<sup>+</sup> and



**Figure 6.** *FAS* -670 genotype determines systemic CD4  $T_{SCM}$  levels in a large twin study. (A) *FAS* -670 genotype is significantly associated with circulating CD4  $T_{SCM}$  levels ( $p = 7 \times 10^{-11}$ , n = 460). Individuals with the functional (i.e. IFN-responsive) AA genotype exhibited a two-fold increase in circulating CD4  $T_{SCM}$  levels (5% of total PBMC), as compared to the non-functional GG genotype (2.5% of total PBMC). However, *FAS* -670 genotype was not associated to either activated CD4 + CD25 + cells (B), or total CD4 cells (C). Twin study and GWAS statistics using GenABEL software package as described.<sup>39,40</sup>

CD4<sup>+</sup>CD25<sup>+</sup> levels were correlated with IFN- $\alpha$ -proliferative/ apoptotic response in both healthy donors and ATL patients. A significant positive correlation between CD4<sup>+</sup>CD25<sup>+</sup> levels was found for IFN- $\alpha$ -induced apoptosis, but not proliferation, in both healthy donors (p = 0.044) and ATL patients (p = 0.0037) (Suppl. Fig. 2). Since we showed above that *FAS* genotype is significantly associated to *ex vivo* IFN- $\alpha$ -induced apoptosis and TCR-triggered proliferation, this suggests a differential effect on different T-cell types, which alerted us to the possible involvement of the recently described Fas<sup>hi</sup> memory stem cell subset in our *ex vivo* ATL model.

### FAS -670 genotype determines systemic CD4 T<sub>SCM</sub> levels in a large twin study

A recent report has identified CD4  $T_{SCM}$  cells as the hierarchical apex of ATL.<sup>19</sup> Due to the prospective design of our ATL cohort, patient cells were no longer available to test a possible association between *FAS* -670 genotype and CD4  $T_{SCM}$ . To investigate this possible association in healthy

donors, a large cohort would be required to achieve sufficient statistical power. Therefore, we tested this hypothesis in the unique twin cohort recently analyzed by simultaneous GWAS and mass cytometry.<sup>40</sup> As shown in Figure 6A, *FAS* -670 genotype was significantly associated with circulating CD4 T<sub>SCM</sub> levels ( $p = 7 \times 10^{-11}$ , n = 460). Individuals with the functional (i.e. IFN-responsive) AA genotype exhibited a two-fold increase in circulating CD4 T<sub>SCM</sub> levels, as compared to the non-functional GG. However, *FAS* -670 genotype was not associated to levels of either activated (CD4<sup>+</sup>CD25<sup>+</sup> cells, Fig. 6B) or total CD4 cells (Fig. 6C), highlighting its cell-type specificity.

### Transcriptomic analysis and cell type deconvolution confirms IFN/FAS/T<sub>SCM</sub> link

Since the T<sub>SCM</sub> cellular phenotype was unknown at the time of patient recruitment for this study, our flow cytometry analysis was limited to broader, overlapping phenotypes (CD4/CD8/ CD25/CD95). However, using archived RNA samples from a subset of 8 ATL patients with paired IFN-*a*-treated samples, we were able to perform transcriptome profiling by microarray and cell type deconvolution using CIBERSORT.<sup>49</sup> Thus, using validated cell type-specific gene sets, the relative size of total CD8 cells and CD4 subsets (naïve, memory resting, memory activated and Treg) was predicted in silico and correlated to patient survival, apoptosis and proliferation. As shown in Fig. 7A, unsupervised hierarchical clustering revealed the strong link between FAS genotype, IFN-induced apoptosis, CD4 naïve cells and CD4 resting memory cells. This is in agreement with Gattinoni et al., with T<sub>SCM</sub> displaying an intermediate transcriptome profile between naïve and central memory T cells.<sup>41</sup> Therefore, we used 278 T<sub>SCM</sub>-specific genes identified by Gattinoni et al.<sup>41</sup> (Suppl. Table I) to perform principal component analysis of purified naive T cells, memory stem T cells, central memory T cells and effector memory T cells (n = 3each, from<sup>41</sup>). and primary cells of three distinct ATL patients from this study cohort with GG, GA or AA genotypes (one each). As shown in Fig. 7B, the first principal component clearly recapitulates the described differentiation gradient of  $T_N \rightarrow T_{SCM} \rightarrow T_{CM} \rightarrow T_{EM}$ . Clustering (k-means) allowed to distinguish all four memory subsets, and to discriminate between functional (AA, GA) and non-functional (GG) FAS genotypes in ATL patients. Strikingly, only AA and GA patient samples clustered with T<sub>SCM</sub> cells, whereas the GG patient sample did not, replicating the intricate link between FAS -670 genotype and T<sub>SCM</sub> phenotype we identified in the large twin cohort (Fig. 6A). Clustering results were robust to the inclusion of our patient samples (predominantly CD4 subpopulations, by flow cytometry and CIBERSORT) with the published data used to define T<sub>SCM</sub>-specific genes in CD8 subpopulations (Supplementary Figure 3). suggesting the "stemness" of CD4 and CD8 cells might be conserved at the transcriptional level. To test the hypothesis of IFN-induced apoptosis of T<sub>SCM</sub> cells, we quantified the effect of IFN- $\alpha$  upon overall vs. T<sub>SCM</sub>-specific gene expression in the ATL transcriptome. When analyzing all patients together, irrespective of genotypes, IFN- $\alpha$  significantly decreased transcript levels of four out of six Naive/T<sub>SCM</sub> signature genes<sup>41</sup> (data not shown), including transcription



**Figure 7.** Transcriptomic analysis and cell type deconvolution confirms IFN/FAS/T<sub>SCM</sub> link in ATL primary cells. (A) Transcriptome profiling by microarray and cell type deconvolution (CIBERSORT) was used to quantify total CD8 cells and CD4 subsets (naïve, memory resting, memory activated and T<sub>reg</sub>). Data were correlated to patient survival, apoptosis and proliferation, followed by unsupervised hierarchical clustering. (B) For Principal Component (PC) analysis, we used microarray signal intensity data of 278 selected T<sub>SCM</sub> genes of purified naive (T<sub>N</sub> cells) (n = 3), stem cell memory (T<sub>SCM</sub> cells) (n = 3), central memory (T<sub>CM</sub> cells) (n = 3) and effector memory T cells (T<sub>EM</sub> cells) (n = 3) from the Gattinoni study,<sup>41</sup> and primary cells of three distinct ATL patients with *FAS* -670 GG, GA or AA genotypes (one each) from this study cohort. Clusters were defined by k-means algorithm. (C) Confirmation of antiviral, anti-inflammatory and antiproliferative effect of IFN- $\alpha$  in ATL patients (m = 8) at the transcriptional level, by upregulation of antiveral effector molecules (*BST2/TRIM5*) and downregulated as log fold-change (with a cut-off at -0.5) in ATL patients with FDR correction, *p* = 0.014 and *p* = 0.0011, respectively. \**p* < 0.05, \*\**p* < 0.01

regulators LEF1 (p = 0.005, n = 8), CERS6 (p = 0.036, n = 8) and TAF4B (p = 0.004, n = 8), whereas no significant effect was observed for T-cell effector or memory signature genes (e.g. Eomes, T-bet, PRDM1, data not shown). Furthermore, we were able to confirm the antiviral, anti-inflammatory and antiproliferative effects of IFN- $\alpha$  (Fig. 2) at the transcriptional level (Fig. 7C), as evidenced by pronounced upregulation of antiretroviral effectors (BST2/TRIM5) and downregulation of cytokines (IL2/IL6) and proliferation markers Ki67 and CD71 (encoded by MKI67/TFRC genes), in agreement with our previous results.<sup>27</sup> When considering FAS genotype, we found IFNinduced downregulation of T<sub>SCM</sub>-specific genes was dependent on the number of functional A alleles (0-1-2 for GG-GA-AA genotypes, ANOVA with post-test for linear trend, p = 0.001). As shown in Fig. 7D, IFN- $\alpha$  caused the strongest downregulation of  $T_{SCM}$  genes in AA genotype (>GA, p = 0.014; >GG, p = 0.0011), argueing in favour of Fas-dependent T<sub>SCM</sub> cell death. Of note, we also confirmed our previous observation<sup>12</sup> of a significant negative correlation between the number of functional FAS -670 A alleles and survival (r = -0.72, p = 0.030, n = 8), indicating this small subset is representative of the larger cohort.

### Proposed model for a coordinated mechanism of action of IFN- $\alpha$ in ATL

Endogenous IFN- $\alpha$  has been shown to be produced by plasmacytoid dendritic cells (pDC) upon stimulation with free virions,<sup>50</sup> which can be assumed to occur in vivo during ATL pathogenesis and/or disease progression.<sup>51</sup> Furthermore, exogenous IFN- $\alpha$  treatment exerts pleiotropic effects as demonstrated in this study. A simplified model (Fig. 8) illustrates how endogenous or exogenous IFN- $\alpha$  might exert its coordinated effects in vivo. Our data strongly suggest that IFN- $\alpha$ -induced apoptosis, intrinsically linked to the presence of the FAS -670 A allele, is driving the antiviral and antiproliferative activity of IFN- $\alpha$ . The FAS -670 A allele is also linked to increased CD4 T<sub>SCM</sub> cells (Fig. 6A), resulting in an ATL leukemic phenotype characterized as CD4<sup>+</sup>CD25<sup>+</sup>CD3<sup>lo</sup> and Fas<sup>hi</sup>.(<sup>52,53</sup> and this study) In addition, IFN- $\alpha$  results in a decrease of both IL-2 (growth factor for CD4<sup>+</sup>CD25<sup>+</sup> cells) and IL-6, high levels of which have been previously associated with decreased ATL survival.<sup>11</sup> Furthermore, a direct activity of IFN- $\alpha$  on CD4<sup>+</sup>CD25<sup>+</sup> cells can be assumed (Suppl. Fig. 2), which might be



**Figure 8.** Proposed model for coordinated mechanism of action of IFN- $\alpha$  in ATL. Endogenous (from plasmacytoid dendritic cells – pDC) or exogenous IFN- $\alpha$  (*ex vivo* or *in vivo* treatment) elicits pro-apoptotic, antiproliferative, antiviral and immuno-modulatory effects on CD4<sup>+</sup>CD25<sup>+</sup> cells, the major HTLV-1-infected reservoir, as well as on leukemic CD4<sup>+</sup>CD25<sup>+</sup>CD3<sup>lo</sup>Fas<sup>+</sup> cells. Red arrows indicate the link to *FAS* -670 polymorphism and T<sub>SCM</sub> cells (this study,<sup>12,19</sup>).

independent of *FAS* genotype (Fig. 6B). Apoptosis of  $CD4^+CD25^+$  cells, the main reservoir of infected cells,<sup>54</sup> likely results in decreased viral protein levels, as well as decreased  $CD4^+CD25^+$  proliferation. Due to the high significance of the *FAS*/CD4 T<sub>SCM</sub> connection, beyond the genome-wide association threshold (Fig. 6A), it is tempting to speculate that the model proposed here would be applicable and of clinical relevance in other leukemias, as well as other viral pathologies treated with IFN- $\alpha$ .

#### Discussion

Conflicting data exist on the cellular and molecular mechanisms of action of type I IFN (IFN- $\alpha$  and IFN- $\beta$ ) in several major human diseases<sup>23-36</sup> including ATL. This HTLV-1associated adult T-cell leukemia responds at varying degrees to IFN- $\alpha$  + AZT combination therapy.<sup>20-22</sup> In this study, we show that ex vivo IFN- $\alpha$  treatment exerts a pleiotropic and strongly heterogeneous response in ATL primary cells. IFN- $\alpha$  significantly decreased proliferation, viral protein levels and IL-6 levels for a majority of patients, while it induced apoptosis and reduced IL-2 levels for a subset of patients (Fig 2). These results are in contrast with our results in another HTLV-1-associated disease (HTLV-1associated myelopathy/Tropical spastic paraparesis), where we found no significant effect of ex vivo IFN- $\alpha$  upon proliferation, cell death or IL-6 production.<sup>25</sup> In addition, a recently published systems biology approach revealed an IFN-inducible gene signature in HAM/TSP patients, surprisingly paralleled by limited antiviral activity.<sup>55</sup> Therefore, we set out to model the interdependencies between ex vivo molecular and cellular IFN- $\alpha$ -response and the *in vivo* clinical and immunological data in ATL. In this study, we reveal a high-penetrance, IFN-dependent, pro-apoptotic phenotype for the functional A allele of the *FAS* -670 A/G polymorphism in both ATL patients and healthy controls. Strikingly, the association of this A allele with IFN-induced apoptosis underscores the recently established "CD4  $T_{SCM}$  apex" model for ATL,<sup>19</sup> since we also found a strong link between the A allele and higher *in vivo* circulating CD4  $T_{SCM}$  levels in a large twin study (Fig. 6A).

Since ATL is a relatively rare disease, even in highly endemic areas such as Salvador-Bahia, Brasil,<sup>56</sup> only 25 patients with clinically definite ATL could be recruited during a five-year period, followed by a five-year clinical follow-up, which precluded confirmation of our data in a second cohort of ATL patients. However, we were able to confirm the IFN-dependent link between CD4 T<sub>SCM</sub> cells, Fas genotype and apoptosis in a subset of ATL patients by transciptomic analysis (Fig. 7A-D). In addition, the dependence of the IFN- $\alpha$  pro-apoptotic effect on the presence of the FAS -670 A allele was validated in healthy donors from the same endemic area. The remarkably high penetrance (80% in ATL, 90% in normal donors) of the FAS polymorphism suggests a decisive effect of the receptor in Fas/FasL signaling, at least upon IFN- $\alpha$  stimulation. Although FasL polymorphisms have also been significantly associated with an in vitro pro-apoptotic phenotype in cervical cancer,<sup>57</sup> three lines of evidence support our receptor-mediated hypothesis for the genotype-phenotype association of IFNinduced apoptosis in this study. First, FASL -844 polymorphism was not associated to in vitro apoptosis in both ATL patients and healthy controls (Silva-Santos et al., unpublished). Second, IFN- $\alpha$  strongly increased FasL levels in 9 out of 13 ATL patients (Fig. 2F) and in all healthy controls tested (Fig. 4D, p < 0.001), resulting in excess ligand and hence indicating receptor signaling as the limiting step. Third, a recent GWAS twin study linking 80.000 immune phenotypes to SNPs identified polymorphisms in FAS, but not FASL, as significantly associated with circulating T-cell subset levels.<sup>40</sup> Following up on this study, we show that the FAS -670 polymorphism is significantly associated to circulating CD4 T<sub>SCM</sub>, but not CD4<sup>+</sup>CD25<sup>+</sup> or total CD4 levels (Fig. 6A-B-C).

Surprisingly, functional GA/AA genotypes displayed higher ex vivo proliferation upon TCR stimulation (Fig. 3D), which somehow contrasts with the pro-apoptotic genotype/phenotype, but is consistent with a proliferative T<sub>SCM</sub> phenotype in vitro.<sup>41</sup> Non-apoptotic Fas signaling has indeed been demonstrated to stimulate cellular proliferation in primary CD4 cells,<sup>58</sup> but also phenotypic differentiation of naïve CD8 T cells into memory subsets, through an Akt-driven mechanism.<sup>59</sup> Similar to these observations, we have recently identified a Fas<sup>hi</sup> lymphoproliferative phenotype in another HTLV-1-associated pathology, the neuroinflammatory disease HAM/TSP<sup>60</sup>. Likewise, it has been shown that most ATL cells display a Fashi phenotype,<sup>61</sup> whereas most T-cell leukemias display decreased Fas expression and/or mutations in Fas/FasL pro-apoptotic signaling.<sup>15</sup> Increased Fas expression might induce survival of (pre-)leukemic cells, through increased proliferation of CD4<sup>+</sup>Fas<sup>+</sup> T-cell clones upon antigenic stimulation and eventually escape apoptosis,<sup>62</sup> through increased c-FLIP or similar

pathways.<sup>63,64</sup> In fact, Fas was previously identified as a protooncogene, capable of inducing tumorogenesis in a different type of cancer.<sup>65</sup>

Our results indicate that this Fas-mediated dichotomy of apoptosis vs. proliferation, previously described in vitro in CD4<sup>+</sup> cells,<sup>58</sup> might be implicated in ATL pathogenesis, by tipping the balance between leukemic CD4<sup>+</sup>CD25<sup>+</sup>CD3<sup>lo</sup>(Fas<sup>hi</sup>) and anti-leukemic CD8<sup>+</sup> cells (Fig. 3 and Fig. 7A). Conversely, high levels of leukemic and/or Treg cells with a CD4<sup>+</sup>CD25<sup>+</sup> phenotype were positively correlated with IFN- $\alpha$ -induced apoptosis (Suppl. Fig. 2). Elimination of both cell types will most probably result in clinical benefit in ATL patients either directly or indirectly, since Tregs have been shown to inhibit the ex vivo CTL response in ATL patients,<sup>66</sup> and negatively correlate to patient survival in our analysis (Fig. 7A). Taken together, the functionality of the FAS -670 polymorphism resolves the IFN- $\alpha$  apoptotic paradox and reveals this polymorphism as a candidate pharmacogenetic marker for several other IFN- $\alpha$ -treated leukemias and cancers.<sup>67</sup> Thus, our model does not contradict the previous findings of absent cytotoxic or antiproliferative activity of IFNs<sup>23,26</sup> but highlights the role of FAS genotype in determining cell fate: proliferation vs. apoptosis.<sup>58–65,67</sup>

Moreover, patient-derived individual T-cell clones represent only a small percentage of the whole T-cell population, whereas the majority has been eliminated, either through "death by neglect", activation-induced cell death or direct CD8<sup>+</sup> T-cellmediated killing.<sup>46-48,61-68</sup> Therefore, our *ex vivo* results are not in contrast, but complimentary to previous *in vitro* results in patientderived T-cell clones.<sup>26,64,68</sup> In addition, the use of total PBMCs also allowed us to measure and integrate the impact of other cellular subtypes, such as DCs,<sup>50,51</sup> monocytes<sup>69</sup> and NK cells,<sup>70</sup> among others, upon apoptosis, proliferation and antiviral response. It is well described that pDC are the main source of IFN- $\alpha$  production *in vivo*.<sup>52</sup> Therefore, we believe our *ex vivo* model, in spite of its limitations, reflects the *in vivo* situation in ATL patients.

In conclusion, we show that a functional *FAS* polymorphism resolves the current "IFN apoptotic paradox", by determining *ex vivo* apoptosis in both ATL patients and healthy controls with high penetrance. The same polymorphism also determines lymphoproliferation and CD4  $T_{SCM}$  levels in healthy controls. In line with the CD4  $T_{SCM}$  ATL origin, our data help explain both *in vivo* IFN+AZT responsiveness and chemotherapy resistance in ATL patients. *FAS* -670 polymorphism thus represents a promising pharmacogenetic marker in the clinical follow-up of several major human diseases currently or previously treated by type I IFN, such as other leukemias, melanoma, HCV infection and multiple sclerosis.

#### Authorship and conflict of interest statements

RK and JVW designed research; RK, GSS, DD, SMM, ACS and JVW performed research; TD, KT, and AMV contributed vital new analytical tools; AB and LF provided patient samples; MM and MR provided genetic and flow cytometry data from the twin cohort; RK and JVW analyzed data and wrote the paper. None of the authors has any conflict of interest to declare.

#### Acknowledgments

This work was supported by: Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), "Science without Borders" Program,

PRONEX, FWO (Grant: ZKC1280-00-W10 and G0D6817N), Alban Program (European Union Program of High Level Scholarships for Latin America, scholarship no. E06D103200BR), VLAIO (TD scholarship) and KU Leuven (IRO and "Vaast Leysen Leerstoel voor Infectieziekten in Ontwikkelingslanden"). KT is funded as a Postdoctoral Researcher by the Research Foundation – Flanders (FWO).

#### Funding

Fonds Wetenschappelijk Onderzoek (G0D6817N) CNPq-FAPESB (PRO-NEX) VLAIO (PhD scholarship TD) Onderzoeksraad, KU Leuven (IRO) Onderzoeksraad, KU Leuven (Vaast Leysen Leerstoel)

#### ORCID

Ricardo Khouri ( http://orcid.org/0000-0001-5664-4436 Anne-Mieke Vandamme ( http://orcid.org/0000-0002-6594-2766

#### References

- Uchiyama T, Yodoi J, Sagawa K, Takatsuki K, Uchino H. Adult T-cell leukemia: clinical and hematologic features of 16 cases. Blood. 1977;50(3):481–92. PMID:301762.
- Poiesz BJ, Ruscetti FW, Gazdar AF, Bunn PA, Minna JD, Gallo RC. Detection and isolation of type C retrovirus particles from fresh and cultured lymphocytes of a patient with cutaneous T-cell lymphoma. Proc Natl Acad Sci United States Am. 1980;77(12):7415–9. doi: 10.1073/pnas.77.12.7415.
- Verdonck K, González E, Van Dooren S, Vandamme AM, Vanham G, Gotuzzo E. Human T-lymphotropic virus 1: recent knowledge about an ancient infection. Lancet Infect Dis. 2007;7(4):266–81.doi: 10.1016/ S1473-3099(07)70081-6 PMID:17376384.
- Cleghorn FR, Manns A, Falk R, Hartge P, Hanchard B, Jack N, Williams E, Jaffe E, White F, Bartholomew C, et al. Effect of human Tlymphotropic virus type I infection on non-Hodgkin's lymphoma incidence. J Natl Cancer Inst. 1995;87(13):1009–14. doi:10.1093/jnci/ 87.13.1009. PMID:7629870.
- Shimoyama M. Diagnostic criteria and classification of clinical subtypes of adult T-cell leukaemia-lymphoma. A report from the Lymphoma Study Group (1984-87). Br J Haematol. 1991;79(3):428–37. doi:10.1111/j.1365-2141.1991.tb08051.x. PMID:1751370.
- Tsukasaki K, Hermine O, Bazarbachi A, Ratner L, Ramos JC, Harrington W, Jr, O'Mahony D, Janik JE, Bittencourt AL, Taylor GP, et al. Definition, prognostic factors, treatment, and response criteria of adult T-cell leukemia-lymphoma: a proposal from an international consensus meeting. J Clin Oncol. 2009;27(3):453–9. doi:10.1200/ JCO.2008.18.2428. PMID:19064971.
- Bittencourt AL, da Graças Vieira M, Brites CR, Farre L, Barbosa HS. Adult T-cell leukemia/lymphoma in Bahia, Brazil: analysis of prognostic factors in a group of 70 patients. Am J Clin Pathol. 2007;128 (5):875–82. doi:10.1309/2YGD1P0QCVCWBLDX. PMID:17951212.
- Tagaya Y, Gallo RC. The Exceptional Oncogenicity of HTLV-1. Front Microbiol. 2017;8:1425. doi:10.3389/fmicb.2017.01425. PMID:28824561.
- Ciminale V, Rende F, Bertazzoni U, Romanelli MG. HTLV-1 and HTLV-2: highly similar viruses with distinct oncogenic properties. Front Microbiol. 2014;5:398. doi:10.3389/fmicb.2014.00398. PMID:25120538.
- Yasuda N, Lai PK, Ip SH, Kung PC, Hinuma Y, Matsuoka M, Hattori T, Takatsuki K, Purtilo D. Soluble interleukin 2 receptors in sera of Japanese patients with adult T cell leukemia mark activity of disease. Blood. 1988;71(4):1021–6. PMID:2895675.
- Yamamura M, Yamada Y, Momita S, Kamihira S, Tomonaga M. Circulating interleukin-6 levels are elevated in adult T-cell leukaemia/ lymphoma patients and correlate with adverse clinical features and survival. Br. J. Haematol. 1998;100(1):129–34. doi:10.1046/j.1365-2141.1998.00538.x. PMID:9450801.
- Farre L, Bittencourt AL, Silva-Santos G, Almeida A, Silva AC, Decanine D, Soares GM, Alcantara LC, Jr, Van Dooren S, Galvão-Castro B, et al. Fas 670 promoter polymorphism is associated to

susceptibility, clinical presentation, and survival in adult T cell leukemia. J Leukoc Biol. 2008;83(1):220–2. doi:10.1189/jlb.0407198. PMID:17962369.

- Takeuchi S, Matsushita M, Tsukasaki K, Takeuchi N, Tomonaga M, Komatsu N, Ikezoe T, Uehara Y, Koeffler HP. P53 codon 72 polymorphism is associated with disease progression in adult T-cell leukaemia/lymphoma. Br J Haematol. 2005;131(4):552–3. doi:10.1111/ j.1365-2141.2005.05798.x. PMID:16281948.
- Tamiya S, Etoh K, Suzushima H, Takatsuki K, Matsuoka M. Mutation of CD95 (Fas/Apo-1) gene in adult T-cell leukemia cells. Blood. 1998;91(10):3935–42. PMID:9573032.
- Kataoka K, Nagata Y, Kitanaka A, Shiraishi Y, Shimamura T, Yasunaga J, Totoki Y, Chiba K, Sato-Otsubo A, Nagae G, et al. Integrated molecular analysis of adult T cell leukemia/lymphoma. Nat Genet. 2015;47(11):1304–15. doi:10.1038/ng.3415. PMID:26437031.
- Beltinger C, Kurz E, Böhler T, Schrappe M, Ludwig WD, Debatin KM. CD95 (APO-1/Fas) mutations in childhood T-lineage acute lymphoblastic leukemia. Blood. 1998;91(10):3943-51. PMID:9573033.
- Huang QR, Morris D, Manolios N. Identification and characterization of polymorphisms in the promoter region of the human Apo-1/Fas (CD95) gene. Mol Immunol. 1997;34(8-9):577–82. doi:10.1016/ S0161-5890(97)00081-3. PMID:9393960.
- Kanemitsu S, Ihara K, Saifddin A, Otsuka T, Takeuchi T, Nagayama J, Kuwano M, Hara T. A functional polymorphism in fas (CD95/APO-1) gene promoter associated with systemic lupus erythematosus. J. Rheumatol. 2002;29(6):1183–8. PMID:12064832.
- Nagai Y, Kawahara M, Hishizawa M, Shimazu Y, Sugino N, Fujii S, Kadowaki N, Takaori-Kondo A. T memory stem cells are the hierarchical apex of adult T-cell leukemia. Blood. 2015;125(23):3527–35. doi:10.1182/blood-2014-10-607465. PMID:25847015.
- Gill PS, Harrington W, Kaplan MH, Ribeiro RC, Bennett JM, Liebman HA, Bernstein-Singer M, Espina BM, Cabral L, Allen S, et al. Treatment of adult T-cell leukemia-lymphoma with a combination of interferon alfa and zidovudine. N Engl J Med. 1995;332(26):1744–8. doi:10.1056/NEJM199506293322603. PMID:7760890.
- Hermine O, Bouscary D, Gessain A, Turlure P, Leblond V, Franck N, Buzyn-Veil A, Rio B, Macintyre E, Dreyfus F, et al. Brief report: treatment of adult T-cell leukemia-lymphoma with zidovudine and interferon alfa. N Engl J Med. 1995;332(26):1749–51. doi:10.1056/ NEJM199506293322604. PMID:7760891.
- Bazarbachi A, Plumelle Y, Carlos Ramos J, Tortevoye P, Otrock Z, Taylor G, Gessain A, Harrington W, Panelatti G, Hermine O. Metaanalysis on the use of zidovudine and interferon-alfa in adult T-cell leukemia/lymphoma showing improved survival in the leukemic subtypes. J Clin Oncol. 2010;28(27):4177–83. doi:10.1200/ JCO.2010.28.0669. PMID:20585095.
- Bazarbachi A, Nasr R, El-Sabban ME, Mahé A, Mahieux R, Gessain A, Darwiche N, Dbaibo G, Kersual J, Zermati Y, et al. Evidence against a direct cytotoxic effect of alpha interferon and zidovudine in HTLV-I associated adult T cell leukemia/lymphoma. Leukemia. 2000;14 (4):716–21. doi:10.1038/sj.leu.2401742. PMID:10764160.
- Moens B, Pannecouque C, López G, Talledo M, Gotuzzo E, Khouri R, Bittencourt A, Farré L, Galvão-Castro B, Vandamme AM, et al. Simultaneous RNA quantification of human and retroviral genomes reveals intact interferon signaling in HTLV-1-infected CD4+ T cell lines. Virol J. 2012;9:171. doi:10.1186/1743-422X-9-171. PMID:22917064.
- 25. Moens B, Decanine D, Menezes SM, Khouri R, Silva-Santos G, Lopez G, Alvarez C, Talledo M, Gotuzzo E, de Almeida Kruschewsky R, et al. Ascorbic acid has superior ex vivo antiproliferative, cell death-inducing and immunomodulatory effects over IFN-α in HTLV-1-associated myelopathy. PLoS Negl Trop Dis. 2012;6(7):e1729. doi:10.1371/journal.pntd.0001729. PMID:22848768.
- Smith D, Buckle GJ, Hafler DA, Frank DA, Höllsberg P. HTLV-Iinfected T cells evade the antiproliferative action of IFN-beta. Virology. 1999;257(2):314–21. doi:10.1006/viro.1999.9679. PMID:10329542.
- 27. Dierckx T, Khouri R, Menezes SM, Decanine D, Farre L, Bittencourt A, Vandamme AM, Van Weyenbergh J. IFN- $\beta$  induces greater antiproliferative and proapoptotic effects and increased p53 signaling compared with IFN- $\alpha$  in PBMCs of Adult T-cell Leukemia/

Lymphoma patients. Blood Cancer J. 2017;7(1):e519.25. doi:10.1038/ bcj.2016.126.

- Kaser A, Nagata S, Tilg H. Interferon alpha augments activationinduced T cell death by upregulation of Fas (CD95/APO-1) and Fas ligand expression. Cytokine. 1999;11(10):736–43. doi:10.1006/ cyto.1998.0484. PMID:10525311.
- Van Weyenbergh J, Wietzerbin J, Rouillard D, Barral-Netto M, Liblau R. Treatment of multiple sclerosis patients with interferon-beta primes monocyte-derived macrophages for apoptotic cell death. J Leukoc Biol. 2001;70(5):745–8. PMID:11698494.
- Gniadek P, Aktas O, Wandinger K-P, Bellmann-Strobl J, Wengert O, Weber A, von Wussow P, Obert HJ, Zipp F. Systemic IFN-beta treatment induces apoptosis of peripheral immune cells in MS patients. J Neuroimmunol. 2003;137(1-2):187–96. doi:10.1016/S0165-5728(03) 00074-2. PMID:12667663.
- Zipp F, Beyer M, Gelderblom H, Wernet D, Zschenderlein R, Weller M. No induction of apoptosis by IFN-beta in human antigen-specific T cells. Neurology. 2000;54(2):485–7. doi:10.1212/WNL.54.2.485. PMID:10668719.
- 32. Dondi E, Roué G, Yuste VJ, Susin SA, Pellegrini S. A dual role of IFNalpha in the balance between proliferation and death of human CD4+ T lymphocytes during primary response. J Immunol. 2004;173 (6):3740–7. doi:10.4049/jimmunol.173.6.3740. PMID:15356120.
- Feng X, Vander HN, Ratner L. Alpha interferon inhibits human T-cell leukemia virus type 1 assembly by preventing Gag interaction with rafts. J Virol. 2003;77(24):13389–95. doi:10.1128/JVI.77.24.13389-13395.2003. PMID:14645593.
- Kiladjian J-J, Giraudier S, Cassinat B. Interferon-alpha for the therapy of myeloproliferative neoplasms: targeting the malignant clone. Leukemia. 2016;30(4):776–81. doi:10.1038/leu.2015.326. PMID:26601783.
- 35. Mullally A, Bruedigam C, Poveromo L, Heidel FH, Purdon A, Vu T, Austin R, Heckl D, Breyfogle LJ, Kuhn CP, et al. Depletion of Jak2V617F myeloproliferative neoplasm-propagating stem cells by interferon- $\alpha$  in a murine model of polycythemia vera. Blood. 2013;121(18):3692–702. doi:10.1182/blood-2012-05-432989. PMID:23487027.
- Talpaz M, Mercer J, Hehlmann R. The interferon-alpha revival in CML. Ann. Hematol. 2015;94 Suppl 2:S195–207. doi:10.1007/s00277-015-2326-y. PMID:25814086.
- Bittencourt AL, Barbosa HS, Requião C, da Silva AC, Vandamme AM, Van Weyenbergh J, Farré L. Adult T-cell leukemia/lymphoma with a mixed CD4+ and CD8+ phenotype and indolent course. J Clin Oncol. 2007;25(17):2480–2. doi:10.1200/JCO.2007.11.3043. PMID:17557960.
- Farre L, de Oliveira M, de FP, Primo J, Vandamme AM, Van Weyenbergh J, Bittencourt AL. Early sequential development of infective dermatitis, human T cell lymphotropic virus type 1-associated myelopathy, and adult T cell leukemia/lymphoma. Clin Infect Dis. 2008;46(3):440–2. doi:10.1086/524695. PMID:18173359.
- 39. Moayyeri A, Hammond CJ, Hart DJ, Spector TD. Effects of age on genetic influence on bone loss over 17 years in women: the Healthy Ageing Twin Study (HATS). J Bone Miner Res. 2012;27(10):2170–8. doi:10.1002/jbmr.1659. PMID:22589082.
- 40. Roederer M, Quaye L, Mangino M, Beddall MH, Mahnke Y, Chattopadhyay P, Tosi I, Napolitano L, Terranova Barberio M, Menni C, et al. The genetic architecture of the human immune system: A bioresource for autoimmunity and disease pathogenesis. Cell. 2015;161(2):387–403. doi:10.1016/j.cell.2015.02.046. PMID:25772697.
- Gattinoni L, Lugli E, Ji Y, Pos Z, Paulos CM, Quigley MF, Almeida JR, Gostick E, Yu Z, Carpenito C, et al. A human memory T cell subset with stem cell-like properties. Nat Med. 2011;17(10):1290–7. doi:10.1038/nm.2446. PMID:21926977.
- 42. Kouri V, Khouri R, Alemán Y, Abrahantes Y, Vercauteren J, Pineda-Peña AC, Theys K, Megens S, Moutschen M, Pfeifer N, et al. CRF19\_cpx is an Evolutionary fit HIV-1 Variant Strongly Associated With Rapid Progression to AIDS in Cuba. EBioMedicine. 2015;2 (3):244–54. doi:10.1016/j.ebiom.2015.01.015. PMID:26137563.
- Snoek J, Casado C, Colombo S, et al. A Bayesian network approach to study host and viral genetic correlates of HIV-1 disease progression. Retrovirology. 2011;8(Suppl 2):P70. doi:10.1186/1742-4690-8-S2-P70.
- 44. Deforche K, Silander T, Camacho R, Grossman Z, Soares MA, Van Laethem K, Kantor R, Moreau Y, Vandamme AM. Analysis of HIV-1 pol sequences using Bayesian Networks: implications for

drug resistance. Bioinformatics. 2006;22(24):2975–9. doi:10.1093/ bioinformatics/btl508. PMID:17021157.

- 45. Friedman N, Goldszmidt M, Wyner A. Data Analysis with Bayesian Networks: A Bootstrap Approach. CoRR. 2013;abs-1301-6695. http://dblp.org/rec/bib/journals/corr/abs-1301-6695
- Dhein J, Walczak H, Bäumler C, Debatin KM, Krammer PH. Autocrine T-cell suicide mediated by APO-1/(Fas/CD95). Nature. 1995;373(6513):438–41. doi:10.1038/373438a0. PMID:7530335.
- 47. Brunner T, Mogil RJ, LaFace D, Yoo NJ, Mahboubi A, Echeverri F, Martin SJ, Force WR, Lynch DH, Ware CF, et al. Cell-autonomous Fas (CD95)/Fas-ligand interaction mediates activation-induced apoptosis in T-cell hybridomas. Nature. 1995;373(6513):441–4. doi:10.1038/373441a0. PMID:7530336.
- Ju ST, Panka DJ, Cui H, Ettinger R, el-Khatib M, Sherr DH, Stanger BZ, Marshak-Rothstein A. Fas(CD95)/FasL interactions required for programmed cell death after T-cell activation. Nature. 1995;373 (6513):444–8. doi:10.1038/373444a0. PMID:7530337.
- Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, Hoang CD, Diehn M, Alizadeh AA. Robust enumeration of cell subsets from tissue expression profiles. Nat Methods. 2015;12(5):453–7. doi:10.1038/nmeth.3337. PMID:25822800.
- Siegal FP, Kadowaki N, Shodell M, Fitzgerald-Bocarsly PA, Shah K, Ho S, Antonenko S, Liu YJ. The nature of the principal type 1 interferon-producing cells in human blood. Science. 1999;284(5421):1835– 7. doi:10.1126/science.284.5421.1835. PMID:10364556.
- Jones KS, Petrow-Sadowski C, Huang YK, Bertolette DC, Ruscetti FW. Cell-free HTLV-1 infects dendritic cells leading to transmission and transformation of CD4(+) T cells. Nat Med. 2008;14(4):429–36. doi:10.1038/nm1745. PMID:18376405.
- 52. Yokote T, Akioka T, Oka S, Hara S, Kobayashi K, Nakajima H, Yamano T, Ikemoto T, Shimizu A, Tsuji M, et al. Flow cytometric immunophenotyping of adult T-cell leukemia/lymphoma using CD3 gating. Am J Clin Pathol. 2005;124(2):199–204. doi:10.1309/ KEN4MXM5Y9A1GEMP. PMID:16040289.
- 53. Tian Y, Kobayashi S, Ohno N, Isobe M, Tsuda M, Zaike Y, Watanabe N, Tani K, Tojo A, Uchimaru K. Leukemic T cells are specifically enriched in a unique CD3(dim) CD7(low) subpopulation of CD4(+) T cells in acute-type adult T-cell leukemia. Cancer Sci. 2011;102 (3):569–77. doi:10.1111/j.1349-7006.2010.01833.x. PMID:21205081.
- 54. Yamano Y, Cohen CJ, Takenouchi N, Yao K, Tomaru U, Li HC, Reiter Y, Jacobson S. Increased expression of human T lymphocyte virus type I (HTLV-I) Tax11-19 peptide-human histocompatibility leukocyte antigen A\*201 complexes on CD4+ CD25+ T Cells detected by peptide-specific, major histocompatibility complex-restricted antibodies in patients with HTLV-I-associated neurologic disease. J Exp Med. 2004;199(10):1367–77. doi:10.1084/ jem.20032042. PMID:15136590.
- Tattermusch S, Skinner JA, Chaussabel D, Banchereau J, Berry MP, McNab FW, O'Garra A, Taylor GP, Bangham CR. Systems biology approaches reveal a specific interferon-inducible signature in HTLV-1 associated myelopathy. PLoS Pathog. 2012;8(1):e1002480. doi:10.1371/journal.ppat.1002480. PMID:22291590.
- Dourado I, Alcantara LCJ, Barreto ML, da Gloria Teixeira M, Galvão-Castro B. HTLV-I in the general population of Salvador, Brazil: a city with African ethnic and sociodemographic characteristics. J Acquir Immune Defic Syndr. 2003;34(5):527–31. doi:10.1097/00126334-200312150-00013. PMID:14657765.
- 57. Sun T, Zhou Y, Li H, Han X, Shi Y, Wang L, Miao X, Tan W, Zhao D, Zhang X, et al. FASL -844C polymorphism is associated with increased

activation-induced T cell death and risk of cervical cancer. J. Exp. Med. 2005;202(7):967–74. doi:10.1084/jem.20050707. PMID:16186185.

- Paulsen M, Valentin S, Mathew B, Adam-Klages S, Bertsch U, Lavrik I, Krammer PH, Kabelitz D, Janssen O. Modulation of CD4+ T-cell activation by CD95 co-stimulation. Cell Death Differ. 2011;18(4):619– 31. doi:10.1038/cdd.2010.134. PMID:21052094.
- Klebanoff CA, Scott CD, Leonardi AJ, Yamamoto TN, Cruz AC, Ouyang C, Ramaswamy M, Roychoudhuri R, Ji Y, Eil RL, et al. Memory T cell-driven differentiation of naive cells impairs adoptive immunotherapy. J Clin Invest. 2016;126(1):318–34. doi:10.1172/JCI81217. PMID:26657860.
- Menezes SM, Leal FE, Dierckx T, Khouri R, Decanine D, Silva-Santos G, Schnitman SV, Kruschewsky R, López G, Alvarez C, et al. A Fas(hi) Lymphoproliferative Phenotype Reveals Non-Apoptotic Fas Signaling in HTLV-1-Associated Neuroinflammation. Front Immunol. 2017;8:97. doi:10.3389/fimmu.2017.00097. PMID:28261198.
- Debatin KM, Goldman CK, Waldmann TA, Krammer PH. APO-1induced apoptosis of leukemia cells from patients with adult T-cell leukemia. Blood. 1993;81(11):2972–7. PMID:7684622.
- Strauss G, Knape I, Melzner I, Debatin K-M. Constitutive caspase activation and impaired death-inducing signaling complex formation in CD95resistant, long-term activated, antigen-specific T cells. J. Immunol. 2003;171(3):1172–82. doi:10.4049/jimmunol.171.3.1172. PMID:12874203.
- Krueger A, Fas SC, Giaisi M, Bleumink M, Merling A, Stumpf C, Baumann S, Holtkotte D, Bosch V, Krammer PH, et al. HTLV-1 Tax protects against CD95-mediated apoptosis by induction of the cellular FLICE-inhibitory protein (c-FLIP). Blood. 2006;107(10):3933–9. doi:10.1182/blood-2005-06-2567. PMID:16403915.
- 64. Zane L, Sibon D, Legras C, Lachuer J, Wierinckx A, Mehlen P, Delfau-Larue MH, Gessain A, Gout O, Pinatel C, et al. Clonal expansion of HTLV-1 positive CD8+ cells relies on cIAP-2 but not on c-FLIP expression. Virology. 2010;407(2):341–51. doi:10.1016/j.virol.2010.07.023. PMID:20863547.
- Chen L, Park S-M, Tumanov AV, Hau A, Sawada K, Feig C, Turner JR, Fu YX, Romero IL, Lengyel E, et al. CD95 promotes tumour growth. Nature. 2010;465(7297):492–6. doi:10.1038/nature09075. PMID:20505730.
- Toulza F, Heaps A, Tanaka Y, Taylor GP, Bangham CRM. High frequency of CD4+FoxP3+ cells in HTLV-1 infection: inverse correlation with HTLV-1-specific CTL response. Blood. 2008;111(10):5047–53. doi:10.1182/blood-2007-10-118539. PMID:18094326.
- 67. Sibley K, Rollinson S, Allan JM, Smith AG, Law GR, Roddam PL, Skibola CF, Smith MT, Morgan GJ. Functional FAS promoter polymorphisms are associated with increased risk of acute myeloid leukemia. Cancer Res. 2003;63(15):4327–30. PMID:12907599.
- Sibon D, Gabet A-S, Zandecki M, Pinatel C, Thête J, Delfau-Larue MH, Rabaaoui S, Gessain A, Gout O, Jacobson S, et al. HTLV-1 propels untransformed CD4 lymphocytes into the cell cycle while protecting CD8 cells from death. J Clin Invest. 2006;116(4):974–83. doi:10.1172/JCI27198. PMID:16585963.
- Enose-Akahata Y, Oh U, Grant C, Jacobson S. Retrovirally induced CTL degranulation mediated by IL-15 expression and infection of mononuclear phagocytes in patients with HTLV-I-associated neurologic disease. Blood. 2008;112(6):2400–10. doi:10.1182/blood-2008-02-138529. PMID:18509087.
- Norris PJ, Hirschkorn DF, DeVita DA, Lee TH, Murphy EL. Human T cell leukemia virus type 1 infection drives spontaneous proliferation of natural killer cells. Virulence. 2010;1(1):19–28. doi:10.4161/ viru.1.1.9868. PMID:20640055.