

Probabilistic solutions for a class of deterministic optimal allocation problems

Ka Chun Cheung* Jan Dhaene† Yian Rong‡
Sheung Chi Phillip Yam§

Abstract

We revisit the general problem of minimizing a separable convex function with both a budget constraint and a set of box constraints. This optimization problem arises naturally in many resource allocation problems in engineering, economics, finance and insurance. Existing literature tackles this problem by using the traditional Kuhn-Tucker theory, which leads to either iterative schemes or yields explicit solutions only under some special classes of convex functions owe to the presence of box constraints. This paper presents a novel approach of solving this constrained minimization problem by using the theory of comonotonicity. The key step is to apply an integral representation result to express each convex function as the stop-loss transform of some suitable random variable. By using this approach, we can derive and characterize not only the explicit solution, but also obtain its geometric meaning and some other qualitative properties.

Keywords: Optimal allocation, Constrained optimization, Comonotonicity, Stop-loss transform

MSC: 90B99, 90C30

1 Introduction and problem formulation

Let (X_1, \dots, X_n) be the portfolio of risks (i.e., random variables representing losses) we are facing. A provision of d dollars is available to be allocated among these n risks. We use the function $f_i(d_i)$ to model the level of riskiness of the risk X_i if d_i dollars is allocated to X_i . When more capital is allocated to risk i , the position is considered to be safer, and hence the corresponding risk level is less. This means that f_i should be a decreasing function. It is also natural to assume that the decrement is diminishing per unit of growth. Accordingly, f_i is both decreasing and convex. As a typical example, we may take $f_i(d_i) := \rho((X_i - d_i)_+)$, where ρ is some convex and increasing functional. The amount $\rho((X_i - d_i)_+)$ could be

*The University of Hong Kong, Hong Kong. Email: kccg@hku.hk

†KU Leuven, Leuven, Belgium. Email: Jan.Dhaene@econ.kuleuven.be

‡The University of Hong Kong, Hong Kong. Email: yrong@hku.hk

§The Chinese University of Hong Kong. Email: scpyam@sta.cuhk.edu.hk

interpreted for example as the provision required by the residual risk $(X_i - d_i)_+$ after d_i dollars has been allocated to the potential loss X_i .

Alternatively, instead of treating f_i as a measurement of risk, we can think of it as a penalty function in the sense that allocated capital is expected to be as close as possible to the loss being allocated to, and that f_i penalizes the deviation. This point of view was adopted by Zaks et al. (2006) and Frostig et al. (2007) in determining the fair price of a heterogeneous portfolio. Common examples of penalty functions include quadratic deviations and absolute deviations. With this interpretation, it is natural to assume that f_i is convex, but not necessarily decreasing.

The considerations above lead us to study the minimization of the total required provision or total penalty by determining the optimal amount d_i^* needed for each risk:

$$\min_{d_1 + \dots + d_n = d} \sum_{i=1}^n f_i(d_i).$$

In the case of provision allocations and many other, it is common to require, apart from $d_1 + \dots + d_n = d$, that each allocation d_i is positive. This consideration leads to formulate the following general problem:

$$\min_{(d_1, \dots, d_n) \in \mathcal{A}(d)} \sum_{i=1}^n f_i(d_i), \quad (1)$$

where the set of admissible allocations equals

$$\mathcal{A}(d) = \{(d_1, \dots, d_n) \in \mathbb{R}^n \mid d_1 + \dots + d_n = d, d_i \in [l_i, u_i], i = 1, \dots, n\},$$

in where l_i, u_i are given fixed constants with $l_i < u_i$. In other words, $\mathcal{A}(d)$ is the intersection of the hyperplane $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_1 + \dots + x_n = d\}$ and the n -dimensional box $\prod_{i=1}^n [l_i, u_i]$. The constraints $l_i \leq d_i \leq u_i$ are commonly referred to as *box constraints*. To ensure that $\mathcal{A}(d)$ is non-empty and is not a singleton, we assume that

$$l_1 + \dots + l_n < d < u_1 + \dots + u_n.$$

Throughout this paper, we assume that each f_i is convex and continuous on $[l_i, u_i]$, but is not necessarily decreasing. For simplicity, we also assume that $(f_i)'_+(l_i)$ and $(f_i)'_-(u_i)$ are finite for all i , where f'_+ and f'_- denote the right-hand and left-hand derivative of any convex function f . Since $\mathcal{A}(d)$ is non-empty and compact and the objective function is continuous, a solution always exists.

The linear constraint $d_1 + \dots + d_n = d$ can easily be extended to a more general linear constraint of the form

$$c_1 d_1 + \dots + c_n d_n = d \quad (2)$$

where c_1, \dots, c_n are some fixed constants. We may assume that all of them are non-zero: if $c_i = 0$ for some i , we can minimize $f_i(d_i)$ over $d_i \in [l_i, u_i]$ separately as a one-dimensional problem. With this new constraint, we have the following more general minimization problem:

$$\min_{c_1 d_1 + \dots + c_n d_n = d, l_i \leq d_i \leq u_i} \sum_{i=1}^n f_i(d_i). \quad (3)$$

Assume that the admissible set is non-empty. Let $\tilde{d}_i := c_i d_i$ and $\tilde{f}_i(x) := f_i(x/c_i)$ for all i , then Problem (3) becomes

$$\min_{\tilde{d}_1 + \dots + \tilde{d}_n = d, \tilde{d}_i \in [\tilde{l}_i, \tilde{u}_i]} \sum_{i=1}^n \tilde{f}_i(\tilde{d}_i),$$

where $[\tilde{l}_i, \tilde{u}_i] = [cl_i, cu_i]$ if $c_i > 0$, and $[\tilde{l}_i, \tilde{u}_i] = [cu_i, cl_i]$ if $c_i < 0$. With this transformation, Problem (3) can be treated as a special case of Problem (1).

Capital allocation rules in the form of (1) or (3) are fairly general. It covers the various optimization-based models proposed and studied in Dhaene et al. (2012) and Zaks (2013), and also the insurance pricing models studied in Zaks et al. (2006) and Frostig et al. (2007). These problems also arise naturally in many optimization models in economics, operation management, finance, marketing, etc. We refer to Bitran and Hax (1981), Luss and Gupta (1975), Stefanov (2005) and the references therein, for various applications and discussions. Existing solution methods include standard convex programming (Kuhn-Tucker theory), dynamic programming (for instance, Wilkinson and Gupta (1969)) and the iterative method (e.g. Luss and Gupta (1975), Stefanov (2005)). The classical Kuhn-Tucker theory lies at the heart of all these different approaches. Similar allocation problems for future not-yet-realized risk or payoff, rather than for the current deterministic capital in the current context, has also gained considerable attention in recent years, we refer to Rüschemdorf (2013) for a comprehensive discussion.

If the box constraints $d_i \in [l_i, u_i]$ are removed, and each f_i is defined on the whole real line, Problem (1) is just the classical “infimum-convolution” of the convex functions f_1, \dots, f_n in convex analysis, which is well-studied in the literature (see, for instance, Rockafellar (1970)). The introduction of the box constraints $d_i \in [l_i, u_i]$, which consist of a total of $2n$ one-sided inequality constraints, makes the problem more difficult and possibly non-tractable analytically.

This paper presents an alternative method to solve Problem (1). Instead of using the traditional Lagrangian technique, we first express each convex function f_i as the stop-loss transform of some random variable, so that the objective function becomes a sum of stop-loss transform. The box constraints can be effectively captured and removed by carefully choosing the random variables. Problem (1) then becomes the minimization of a sum of stop-loss transform subject to a homogeneous linear constraint. We demonstrate how the theory of comonotonicity can be used to solve such minimization problem effectively, with the solution set being completely characterized and explicitly expressed in an intuitive geometric way. While the transformation of convex functions into stop-loss transforms is not new and has been applied in different areas, its application in optimization has not been explored in the literature except in the recent paper Cheung et al. (2014). Further advantages of this approach are that we can easily obtain various qualitative properties of solutions of Problem (1), such as their uniqueness and conditions where the box constraints are binding; furthermore, some well-known results in convex analysis associated with infimum-convolution can be derived easily.

This paper is organized as follows. Section 2 reviews basic properties of the notion of comonotonicity. Special focus is put on the geometry of the support of a comonotonic random vector; in particular, we study how the support of a comonotonic random vector intersects with a given hyperplane. Section 3 studies a special case of Problem (1), in which each f_i is a stop-

loss transform. We indicate how Problem (1) can be solved using simple geometric arguments and the results in Section 2. In Section 4, we explain how one can express a given convex function as the stop-loss transform of some random variable. Section 5 combines the ideas in Sections 3 and 4 together to present a novel way to solve Problem (1). We not only provide a complete and explicit characterization of the solutions to Problem (1), but also derive some qualitative properties of the solutions. In Section 6, we give examples to demonstrate how the theory can be applied. Section 7 indicates several variants of Problem (1) to illustrate the flexibility of our model. Finally, Section 8 discusses how the techniques can be used when the box constraints are removed, in which case Problem (1) is related to the notion of infimum-convolution in convex analysis.

2 Supports of comonotonic random vectors

This section reviews some properties of comonotonicity that are pertinent to our later analysis. The focus here is on the behavior of the support of a comonotonic random vector, and how it intersects with a hyperplane with normal $\mathbf{1}$. In fact, it will be shown in Section 5 that the solution of Problem (1) can always be expressed as an intersection of such kind.

Following Kaas et al. (2000), a set $A \subset \mathbb{R}^n$ is said to be comonotonic if any two points in A can be ordered componentwise, that is, for any (x_1, \dots, x_n) and (y_1, \dots, y_n) in A , $(x_i - y_i)(x_j - y_j) \geq 0$ for any $i, j \in \{1, \dots, n\}$. A random vector (X_1, \dots, X_n) is said to be *comonotonic* if there is some comonotonic set $A \subset \mathbb{R}^n$ so that $\mathbb{P}((X_1, \dots, X_n) \in A) = 1$. Comonotonicity of (X_1, \dots, X_n) is equivalent to $(X_1, \dots, X_n) \stackrel{d}{=} (F_{X_1}^{-1}(U), \dots, F_{X_n}^{-1}(U))$ for any uniform $(0, 1)$ random variable U . In this paper, for any given random vector $\mathbf{X} = (X_1, \dots, X_n)$, we define its comonotonic modification $\mathbf{X}^c = (X_1^c, \dots, X_n^c)$ as $\mathbf{X}^c := (F_{X_1}^{-1}(U), \dots, F_{X_n}^{-1}(U))$, where U is an arbitrary uniform $(0, 1)$ random variable. By construction, a comonotonic modification is always comonotonic and has the same marginal distributions as the original random vector. For a comprehensive overview of the theory on comonotonicity, we refer to Dhaene et al. (2002).

Throughout this paper, F_X^{-1} denotes the left-continuous inverse of the distribution function F_X of any random variable X :

$$F_X^{-1}(p) := \inf\{x \in \mathbb{R} \mid F_X(x) \geq p\}, \quad 0 \leq p \leq 1.$$

Similarly, the right-continuous inverse distribution function is defined as

$$F_X^{-1+}(p) := \sup\{x \in \mathbb{R} \mid F_X(x) \leq p\}, \quad 0 \leq p \leq 1.$$

With the convention that $\inf \emptyset = +\infty$ and $\sup \emptyset = -\infty$, $F_X^{-1+}(0)$ and $F_X^{-1}(1)$ are the essential infimum and essential supremum of X respectively. By definition, $F_X^{-1}(0) = -\infty$ and $F_X^{-1+}(1) = \infty$ regardless of the actual distribution of X . For our later purpose, we also need the notion of α -mixed inverse distribution function. Following Kaas et al. (2000), it is defined as

$$F_X^{-1(\alpha)}(p) := \alpha F_X^{-1}(p) + (1 - \alpha) F_X^{-1+}(p), \quad 0 \leq p \leq 1, 0 \leq \alpha \leq 1.$$

For consistency, we also adopt the convention $0 \cdot \pm\infty = 0$ so that $F_X^{-1(0)}(p) = F_X^{-1+}(p)$ and $F_X^{-1(1)}(p) = F_X^{-1}(p)$ for any $0 \leq p \leq 1$.

In the remainder of this section, $\mathbf{X}^c = (X_1^c, \dots, X_n^c)$ denotes a fixed comonotonic random vector with marginal distribution functions F_1, \dots, F_n , and S^c is the comonotonic sum $X_1^c + \dots + X_n^c$. A fundamental property of comonotonicity (see for instance Denneberg (1994) or Dhaene et al. (2002)) is that the inverse distribution function of S^c can be computed explicitly as follows:

$$F_{S^c}^{-1(\alpha)}(p) = F_1^{-1(\alpha)}(p) + \dots + F_n^{-1(\alpha)}(p), \quad 0 \leq p \leq 1, 0 \leq \alpha \leq 1. \quad (4)$$

By definition, it is clear that

$$\text{msupp}(\mathbf{X}^c) := \{(F_1^{-1}(u), \dots, F_n^{-1}(u)) \mid 0 < u < 1\}$$

is a comonotonic set in \mathbb{R}^n and is a support¹ of \mathbf{X}^c . We say that $\mathbf{s} \in \mathbb{R}^n$ is a *comonotonic support point* of \mathbf{X}^c , if $\{\mathbf{s}\} \cup \text{msupp}(\mathbf{X}^c)$ is again comonotonic. In other words, adding a comonotonic support point to $\text{msupp}(\mathbf{X}^c)$ will not destroy its comonotonicity. The collection of all comonotonic support points of \mathbf{X}^c will be denoted as $\text{csupp}(\mathbf{X}^c)$:

$$\text{csupp}(\mathbf{X}^c) := \{\mathbf{s} \in \mathbb{R}^n \mid \{\mathbf{s}\} \cup \text{msupp}(\mathbf{X}^c) \text{ is comonotonic}\}.$$

Obviously, $\text{msupp}(\mathbf{X}^c) \subset \text{csupp}(\mathbf{X}^c)$, and $\text{msupp}(\mathbf{X}^c) \cup \{\mathbf{s}\}$ is also a comonotonic support of \mathbf{X}^c for every $\mathbf{s} \in \text{csupp}(\mathbf{X}^c)$. However, $\text{csupp}(\mathbf{X}^c)$ is not necessarily comonotonic.

For any $d \in \mathbb{R}$, we denote by $\ell(d)$ the hyperplane

$$\{\mathbf{d} = (d_1, \dots, d_n) \in \mathbb{R}^n \mid d_1 + \dots + d_n = d\}.$$

As indicated earlier, we are interested in finding the intersection $\ell(d) \cap \text{csupp}(\mathbf{X}^c)$, which will be denoted as $i(d, \mathbf{X}^c)$:

$$i(d, \mathbf{X}^c) := \{\mathbf{s} \in \mathbb{R}^n \mid \{\mathbf{s}\} \cup \text{msupp}(\mathbf{X}^c) \text{ is comonotonic and } s_1 + \dots + s_n = d\}.$$




Figure 1: The set $i(d, \mathbf{X}^c)$ is the portion of the line $d_1 + d_2 = d$ that lies inside $\text{csupp}(\mathbf{X}^c)$.

¹By a support of a random variable or a random vector Y , we mean any Borel measurable set A such that $\mathbb{P}(Y \in A) = 1$.

First, we observe that if $d < F_{S^c}^{-1+}(0)$ or $d > F_{S^c}^{-1}(1)$, the cardinality of $i(d, \mathbf{X}^c)$ is infinity. For if $d < F_{S^c}^{-1+}(0) = F_1^{-1+}(0) + \dots + F_n^{-1+}(0)$, there are infinitely many possible ways to decompose d into a sum $d_1 + \dots + d_n$ such that $d_i \leq F_i^{-1+}(0)$ for each i . Such a decomposition belongs to $i(d, \mathbf{X}^c)$ by definition. The case for $d > F_{S^c}^{-1}(1)$ is similar.

The following result characterizes the set $i(d, \mathbf{X}^c)$ for the more interesting and relevant case where $F_{S^c}^{-1+}(0) \leq d \leq F_{S^c}^{-1}(1)$. Its geometric meaning is clear and is described in Figure 1: $i(d, \mathbf{X}^c)$ is simply the portion of the hyperplane $\ell(d)$ that lies inside the box of comonotonic support points.

Proposition 2.1. *Suppose that d is a real number such that $F_{S^c}^{-1+}(0) \leq d \leq F_{S^c}^{-1}(1)$. Then*

$$i(d, \mathbf{X}^c) = \{(d_1, \dots, d_n) \in \ell(d) \mid F_i^{-1}(F_{S^c}(d)) \leq d_i \leq F_i^{-1+}(F_{S^c}(d)) \text{ for all } i\}. \quad (5)$$

Proof: We first assume that $\sum_{i=1}^n F_i^{-1+}(0) < d < \sum_{i=1}^n F_i^{-1}(1)$. In this case, $0 < F_{S^c}(d) < 1$. Suppose $\mathbf{d} = (d_1, \dots, d_n)$ belongs to the set on the right hand side of (5). Then for any $i = 1, \dots, n$, we have $d_i \geq F_i^{-1}(p)$ when $p \in (0, F_S(d)]$ and $d_i \leq F_i^{-1}(p)$ when $p \in (F_{S^c}(d), 1)$, so $\mathbf{d} \cup \text{msupp}(\mathbf{X}^c)$ is comonotonic. Therefore, \mathbf{d} is a comonotonic support point of \mathbf{X}^c and hence it lies in $i(d, \mathbf{X}^c)$.

Now we suppose that $\mathbf{d} = (d_1, \dots, d_n) \in i(d, \mathbf{X}^c)$. Notice that

$$\sum_{i=1}^n F_i^{-1}(F_{S^c}(d)) = F_{S^c}^{-1}(F_{S^c}(d)) \leq d \leq F_{S^c}^{-1+}(F_{S^c}(d)) = \sum_{i=1}^n F_i^{-1+}(F_{S^c}(d)). \quad (6)$$

Since $\mathbf{d} \in \text{csupp}(\mathbf{X}^c)$, either $F_i^{-1}(F_{S^c}(d)) \leq d_i$ for all i or $F_i^{-1}(F_{S^c}(d)) > d_i$ for all i . The second possibility is ruled out by the second inequality in (6) and the condition that $\mathbf{d} \in \ell(d)$, unless $d_i = F_i^{-1}(F_{S^c}(d))$ for all i . Therefore, $F_i^{-1}(F_{S^c}(d)) \leq d_i$ for all i . By the same argument, $d_i \leq F_i^{-1+}(F_{S^c}(d))$ for all i . This proves the reverse inclusion.

If $d = F_{S^c}^{-1+}(0) = \sum_{i=1}^n F_i^{-1+}(0) \in \mathbb{R}$, the only way to decompose d into a sum $d = d_1 + \dots + d_n$ in such a way that $\mathbf{d} \cup \text{msupp}(\mathbf{X}^c)$ is comonotonic is given by $d = \sum_{i=1}^n F_i^{-1+}(0)$. In this case, $i(d, \mathbf{X}^c) = \{(F_1^{-1+}(0), \dots, F_n^{-1+}(0))\}$. If $F_{S^c}(d) = 0$, the right hand side of (5) becomes

$$\{(d_1, \dots, d_n) \in \mathbb{R}^n \mid d_1 + \dots + d_n = d, -\infty < d_i \leq F_i^{-1+}(0) \text{ for all } i\},$$

which contains $(F_1^{-1+}(0), \dots, F_n^{-1+}(0))$ only; if $F_{S^c}(d) > 0$, the right hand side of (5) becomes

$$\{(d_1, \dots, d_n) \in \mathbb{R}^n \mid d_1 + \dots + d_n = d, F_i^{-1+}(0) \leq d_i \leq F_i^{-1+}(F_{S^c}(d)) \text{ for all } i\},$$

again, this set contains $(F_1^{-1+}(0), \dots, F_n^{-1+}(0))$ only.

The proof for the case where $d = F_{S^c}^{-1}(1) = \sum_{i=1}^n F_i^{-1}(1) \in \mathbb{R}$ is similar and so it is omitted. \square

To study the cardinality $i(d, \mathbf{X}^c)$, we introduce the following subset of \mathbb{R} :

$$s(\mathbf{X}^c) := \left\{ d \in \mathbb{R} \mid d = \sum_{i=1}^n F_i^{-1}(p), \quad p \in (0, 1], \text{ or } d = \sum_{i=1}^n F_i^{-1+}(p), \quad p \in [0, 1) \right\}.$$

Corollary 2.2. *Suppose that d is a real number such that $F_{S^c}^{-1+}(0) \leq d \leq F_{S^c}^{-1}(1)$. If $d \in s(\mathbf{X}^c)$, then $\text{card}(i(d, \mathbf{X}^c)) = 1$; otherwise, if $d \notin s(\mathbf{X}^c)$,*

- (a) $\text{card}(i(d, \mathbf{X}^c)) = \infty$ if there are more than one of F_i^{-1} , $i = 1, \dots, n$, jump at $F_{S^c}(d)$;
(b) $\text{card}(i(d, \mathbf{X}^c)) = 1$ if exactly one of F_i^{-1} , $i = 1, \dots, n$, jumps at $F_{S^c}(d)$.

Proof: For the first assertion, consider $d \in s(\mathbf{X}^c)$, and suppose that $d = \sum_{i=1}^n F_i^{-1}(p)$ for some $p \in (0, 1]$ (the argument for the case where $d = \sum_{i=1}^n F_i^{-1+}(p)$ for some $p \in [0, 1)$ is similar). Obviously, $(F_1^{-1}(p), \dots, F_n^{-1}(p)) \in i(d, \mathbf{X}^c)$. If $\mathbf{s} = (s_1, \dots, s_n)$ is a different point in $i(d, \mathbf{X}^c)$, then by the definition of comonotonicity, either $F_i^{-1}(p) \leq s_i$ for all i , or $F_i^{-1}(p) \geq s_i$ for all i , with the inequality being strict for at least one i in both possibilities. At the same time, it is required that $\sum_{i=1}^n F_i^{-1}(p) = \sum_{i=1}^n s_i = d$. Clearly, such a point \mathbf{s} does not exist, so $i(d, \mathbf{X}^c)$ is a singleton.

For the second assertion, note that $d \notin s(\mathbf{X}^c)$ implies that $F_{S^c}^{-1+}(0) < d < F_{S^c}^{-1}(1)$, or equivalently $0 < F_{S^c}(d) < 1$. In this case, at least one of the F_i^{-1} , $i = 1, \dots, n$, jumps at $F_{S^c}(d)$. If not, $F_i^{-1}(F_{S^c}(d)) = F_i^{-1+}(F_{S^c}(d))$ for all i , summing over i yields $F_{S^c}^{-1}(F_{S^c}(d)) = F_{S^c}^{-1+}(F_{S^c}(d))$ and hence $d = F_{S^c}^{-1}(F_{S^c}(d)) = F_1^{-1}(F_{S^c}(d)) + \dots + F_n^{-1}(F_{S^c}(d))$, which contradicts the assumption that $d \notin s(\mathbf{X}^c)$. Now the result follows from (5). \square

Corollary 2.3. *Suppose that $F_{S^c}^{-1+}(0) < d < F_{S^c}^{-1}(1)$. If we define $d_i^* = F_i^{-1(\alpha)}(F_{S^c}(d))$ for $i = 1, \dots, n$, where $\alpha \in [0, 1]$ is a solution of $F_{S^c}^{-1(\alpha)}(F_{S^c}(d)) = d$, then $(d_1^*, \dots, d_n^*) \in i(d, \mathbf{X}^c)$.*

Proof: This corollary follows from (5) and the fact that $F_i^{-1}(F_{S^c}(d)) \leq d_i^* \leq F_i^{-1+}(F_{S^c}(d))$ for all i . \square

3 A canonical optimal capital allocation problem

Before solving Problem (1), we first consider the well known canonical special case where $f_i(d_i)$ takes the form of $\mathbb{E}[(X_i - d_i)_+]$ for a given integrable random variable X_i :

$$\min_{d_1 + \dots + d_n = d} \sum_{i=1}^n \mathbb{E}[(X_i - d_i)_+]. \quad (7)$$

Notice that $d_1 + \dots + d_n = d$ is the only constraint in Problem (7). We do not impose any box constraint to restrict the individual allocations. In the next two sections, we will demonstrate that Problem (1) with box constraints can always be transformed into the form of Problem (7) without box constraints. Therefore, it is instructive to review this problem and to understand how comonotonicity comes to play.

Later in this section, we consider a much more general version of this canonical allocation problem in which the expectation will be replaced by a general risk measure.

Using the notion of comonotonicity, Kaas et al. (2002) used a simple geometric argument to prove that for $\sum_{i=1}^n F_{X_i}^{-1+}(0) < d < \sum_{i=1}^n F_{X_i}^{-1}(1)$, a solution to Problem (7) is given by

$$d_i^* = F_{X_i}^{-1(\alpha)}(F_{S^c}(d)), \quad i = 1, \dots, n, \quad (8)$$

in which $\alpha \in [0, 1]$ is a solution of the equation $F_{S^c}^{-1(\alpha)}(F_{S^c}(d)) = d$, and $S^c := X_1^c + \dots + X_n^c$ where $\mathbf{X}^c = (X_1^c, \dots, X_n^c)$ is a comonotonic modification of (X_1, \dots, X_n) . It is easy to see that

this solution is indeed a comonotonic support point of \mathbf{X}^c , and hence it belongs to $i(d, \mathbf{X}^c)$. The following theorem gives a full characterization of the solution set of Problem (7) for any $d \in \mathbb{R}$ in terms of $i(d, \mathbf{X}^c)$, which covers (8), the result from Kaas et al. (2002), as a special case.

Theorem 3.1. *For any $d \in \mathbb{R}$, the solution set of Problem (7) is $i(d, \mathbf{X}^c)$.*

Proof: Let U be uniform(0, 1), $\mathbf{X}^c := (F_{X_1}^{-1}(U), \dots, F_{X_n}^{-1}(U))$, and $S^c := F_{X_1}^{-1}(U) + \dots + F_{X_n}^{-1}(U)$. First, we may replace each X_i in Problem (7) by $F_{X_i}^{-1}(U)$. For any $d_1 + \dots + d_n = d$, we have

$$\sum_{i=1}^n \mathbb{E}[(F_{X_i}^{-1}(U) - d_i)_+] \geq \mathbb{E} \left[\left(\sum_{i=1}^n F_{X_i}^{-1}(U) - \sum_{i=1}^n d_i \right)_+ \right] = \mathbb{E}[(S^c - d)_+],$$

so $\mathbb{E}[(S^c - d)_+]$ is a lower bound of the objective function in (7).

For any $\mathbf{d} = (d_1, \dots, d_n) \in i(d, \mathbf{X}^c)$, $\{\mathbf{d}\} \cup \text{msupp}(\mathbf{X}^c)$ is comonotonic by definition, so either $(F_{X_i}^{-1}(U) - d_i)_+ = F_{X_i}^{-1}(U) - d_i$ for all i simultaneously, or $(F_{X_i}^{-1}(U) - d_i)_+ = 0$ for all i simultaneously. In either case, $\sum (F_{X_i}^{-1}(U) - d_i)_+ = (S^c - d)_+$, so \mathbf{d} is a solution of Problem (7).

Finally, suppose $\mathbf{d} \in \ell(d)$ but $\mathbf{d} \notin \text{csupp}(\mathbf{X}^c)$. Then there exist some $i, j \in \{1, \dots, n\}$ and $u \in (0, 1)$ such that $(F_{X_i}^{-1}(u) - d_i)(F_{X_j}^{-1}(u) - d_j) < 0$. By the left continuity of F^{-1} , this strict inequality continues to hold on $[u - \varepsilon, u]$ for some $\varepsilon > 0$. Therefore, $\sum (F_{X_i}^{-1}(U) - d_i)_+ > (S^c - d)_+$ with a strictly positive probability, and so \mathbf{d} is not optimal. \square

We remark that similar results can be found in Chen et al. (2015), in which the authors discuss the issue of (non)-uniqueness of the solution of Problem (7). Without recourse to the geometric notion of $i(d, \mathbf{X}^c)$, they show directly that the solution set is given by $i(d, \mathbf{X}^c)$.

The following result is a direct consequence of Theorem 3.1, Corollary 2.2 and Corollary 2.3.

Corollary 3.2. *If $d \in s(\mathbf{X}^c)$, then Problem (7) admits a unique solution, and if $F_{S^c}^{-1+}(0) < d < F_{S^c}^{-1}(1)$, then any solution (d_1^*, \dots, d_n^*) of Problem (7) satisfies $F_{X_i}^{-1+}(0) \leq d_i^* \leq F_{X_i}^{-1}(1)$.*

Upon examining first part of the proof of Theorem 3.1 carefully, one can see that $i(d, \mathbf{X}^c)$ is indeed not only the solution set of Problem (7), but is also contained in the solution set of the problem

$$\min_{d_1 + \dots + d_n = d} \sum_{i=1}^n (X_i^c(\omega) - d_i)_+,$$

for all $\omega \in \Omega$, where $\mathbf{X}^c = (F_{X_1}^{-1}(U), \dots, F_{X_n}^{-1}(U))$ and U is a uniform(0, 1) random variable.

This observation leads to the following generalization of Problem (7) in terms of a risk measure.

Proposition 3.3. *Let ρ be a risk measure which is (i) law invariant, (ii) increasing in the sense that $Z_1 \leq_{st} Z_2$ (that is, $\mathbb{P}(Z_1 \leq t) \geq \mathbb{P}(Z_2 \leq t)$ for all t) implies that $\rho(Z_1) \leq \rho(Z_2)$, and (iii) comonotonic additive. For any $d \in \mathbb{R}$, the solution set of the problem*

$$\min_{d_1 + \dots + d_n = d} \sum_{i=1}^n \rho((X_i - d_i)_+) \tag{9}$$

contains $i(d, \mathbf{X}^c)$.

Proof: By law-invariance and comonotonic additivity, the objection function in Problem (9) equals

$$\rho \left(\sum_{i=1}^n (F_{X_i}^{-1}(U) - d_i)_+ \right),$$

where U is a uniform $(0, 1)$ random variable. Since $\sum_{i=1}^n (F_{X_i}^{-1}(U(\omega)) - d_i)_+$ is minimized under the constraint $\sum_i d_i = d$ on the set $i(d, \mathbf{X}^c)$ for all $\omega \in \Omega$, so in view of the monotonicity of ρ , we conclude that the solution set of Problem (9) contains $i(d, \mathbf{X}^c)$. \square

4 Representing convex functions as stop-loss transforms

In this section, we show that Problem (1) can be transformed into Problem (7) for some suitable random variables X_1, \dots, X_n , and hence the solution set of Problem (1) equals $i(d, \mathbf{X}^c)$ by Theorem 3.1.

Recall that each f_i in Problem (1) is assumed to be convex and continuous on $[l_i, u_i]$, with finite right-hand derivative at l_i and finite left-hand derivative at u_i .

Proposition 4.1. *Let $g : [l, u] \rightarrow \mathbb{R}$ be a continuous and convex function with*

$$-1 \leq g'_+(l) \leq g'_-(u) \leq 1. \quad (10)$$

Then there exists a random variable X with $\mathbb{P}(l \leq X \leq u) = 1$ such that

$$g(x) = g(l) + x + l + 2\mathbb{E}[(X - x)_+] - 2\mathbb{E}(X) \text{ for any } x \in [l, u].$$

Moreover, the distribution function of X is given by

$$F_X(x) = \begin{cases} 0 & \text{if } x < l, \\ (1 + g'_+(x))/2 & \text{if } l \leq x < u, \\ 1 & \text{if } x \geq u. \end{cases} \quad (11)$$

Proof: By convexity, $g'_+(x)$ is right-continuous and increasing on $[l, u)$. Condition (10) ensures that the expression in (11) is a genuine distribution function. We denote by X an arbitrary random variable with such a distribution function. It is clear that $\mathbb{P}(l \leq X \leq u) = 1$, and there is a possible jump at l and at u . For any $x \in [l, u]$, we have

$$\begin{aligned} & x + 2\mathbb{E}[(X - x)_+] - 2\mathbb{E}(X) \\ &= \mathbb{E}|X - x| - \mathbb{E}(X) \\ &= \int_0^{x-l} F_{X-l}(t) dt + \int_{x-l}^{u-l} (1 - F_{X-l}(t)) dt - \int_0^{u-l} (1 - F_{X-l}(t)) dt - l \\ &= g(x) - g(l) - l. \end{aligned}$$

Rearranging this equation yields the desired result. \square

By the same argument, we have the following variant of Proposition 4.1, in which the upper end point u of the domain of g is infinity:

Proposition 4.2. *Let $g : [l, \infty) \rightarrow \mathbb{R}$ be a decreasing convex function with*

$$-1 \leq g'_+(l) \leq \lim_{x \rightarrow \infty} g'_+(x) = 0.$$

Let $L := \lim_{x \rightarrow \infty} g(x) \in \mathbb{R}$. Then there exists a random variable with $\mathbb{P}(X \geq l) = 1$ such that

$$g(x) = L + \mathbb{E}[(X - x)_+] \text{ for any } x \geq l.$$

Moreover, the distribution function of X is given by

$$F_X(x) = \begin{cases} 0 & \text{if } x < l, \\ 1 + g'_+(x) & \text{if } x \geq l. \end{cases}$$

A common requirement in these two propositions is that the convex function concerned has bounded derivative on the relevant domain. Such requirement enables us to rescale linearly the right-hand derivative into a distribution function of a random variable. Our methodology remains valid even if the derivative is not bounded. In that case, instead of rescaling the right-hand derivative into a distribution function of a random variable, we can directly treat the right-hand derivative as the distribution function of a Radon measure on \mathbb{R} , see page 16 of Föllmer and Schied (2004) and page 545 of Revuz and Yor (1999). Comonotonicity of real-valued measurable maps on a general measurable space can be defined in exactly the same way as it is defined for random variables on a probability space, Property (4) on the additivity of the inverse distribution functions of comonotonic sums is also valid. We choose not to pursue such generality in order to put the focus on the ideas and techniques rather than on technicalities. Interested readers can easily work out the details for the general case.

5 Solution set of minimization Problem (1)

Now we return to our optimal capital allocation Problem (1):

$$\min_{\mathbf{d} \in \mathcal{A}(d)} \sum_{i=1}^n f_i(d_i),$$

where the functions f_i are convex and continuous on $[l_i, u_i]$ with $-\infty < (f_i)'_+(l_i) \leq (f_i)'_-(u_i) < \infty$, and

$$\mathcal{A}(d) = \{(d_1, \dots, d_n) \in \mathbb{R}^n \mid d_1 + \dots + d_n = d, d_i \in [l_i, u_i], i = 1, \dots, n\}$$

is the set of admissible allocations with $l_1 + \dots + l_n < d < u_1 + \dots + u_n$. Since condition (10) of Proposition 4.1 may not be satisfied by f_i , a simple rescaling is needed. To this end, take ν to be any number that is strictly larger than ν^* , which is defined by

$$\nu^* := \max_{1 \leq i \leq n} (|(f_i)'_+(l_i)| \vee |(f_i)'_-(u_i)|) \in \mathbb{R}, \quad (12)$$

and define the functions

$$\tilde{f}_i(x) := f_i(x)/\nu \quad \text{for } x \in [l_i, u_i] \text{ and } i = 1, \dots, n. \quad (13)$$

The functions \tilde{f}_i satisfy all conditions of Proposition 4.1, and hence there exist random variables X_1, \dots, X_n such that for $i = 1, \dots, n$,

$$\tilde{f}_i(x) = \tilde{f}_i(l_i) + x + l_i + 2\mathbb{E}[(X_i - x)_+] - 2\mathbb{E}(X_i), \quad x \in [l_i, u_i],$$

where the distribution function of X_i is given by

$$F_{X_i}(x) = \begin{cases} 0 & \text{if } x < l_i, \\ (1 + (\tilde{f}_i)'_+(x)/\nu)/2 & \text{if } l_i \leq x < u_i, \\ 1 & \text{if } x \geq u_i. \end{cases} \quad (14)$$

Moreover, as ν is chosen to be larger than ν^* , each X_i has a point mass at both of its essential infimum $F_{X_i}^{-1+}(0) = l_i$ and essential supremum $F_{X_i}^{-1}(1) = u_i$. With the above transformation, we find

$$\sum_{i=1}^n f_i(d_i) = C + 2\nu \sum_{i=1}^n \mathbb{E}[(X_i - d_i)_+]$$

for any $(d_1, \dots, d_n) \in \mathcal{A}(d)$, where C is some constant which is independent of (d_1, \dots, d_n) . Therefore, Problem (1) is equivalent to the following problem:

$$\min_{\mathbf{d} \in \mathcal{A}(d)} \sum_{i=1}^n \mathbb{E}[(X_i - d_i)_+], \quad (15)$$

in the sense that the two problems have the same solution sets.

One immediately notices the similarity between Problem (7) and Problem (15). The only difference between them is that Problem (15) requires that $d_i \in [l_i, u_i]$ for all i while Problem (7) does not. However, from Corollary 3.2 and (14), we know that any solution (d_1^*, \dots, d_n^*) of Problem (7) satisfies $d_i^* \in [F_{X_i}^{-1+}(0), F_{X_i}^{-1}(1)] = [l_i, u_i]$, so the box constraints on the individual allocations is automatically fulfilled. Combining this observation with Theorem 3.1 leads to following result:

Theorem 5.1. *The solution set of Problem (1) equals $i(d, \mathbf{X}^c)$, where the marginal distribution of X_i is given by (14). Moreover, if $d \in s(\mathbf{X}^c)$, the Problem (1) admits a unique solution.*

In the remainder of this section, S^c denotes the comonotonic sum $X_1^c + \dots + X_n^c$, where (X_1^c, \dots, X_n^c) is a comonotonic modification of (X_1, \dots, X_n) with marginal distributions given by (14).

Corollary 5.2. *The solution set to Problem (1) is given by*

$$\{(d_1, \dots, d_n) \mid d_1 + \dots + d_n = d, F_{X_i}^{-1}(F_{S^c}(d)) \leq d_i \leq F_{X_i}^{-1+}(F_{S^c}(d)) \text{ for all } i\}.$$

Moreover, $0 < F_{S^c}(d) < 1$.

Proof: Since ν is chosen to be strictly larger ν^* defined in (13), $F_{X_i}^{-1+}(0) = l_i$ and $F_{X_i}^{-1}(1) = u_i$ for each i and so $F_{S^c}^{-1+}(0) = \sum_i l_i < d < \sum_i u_i = F_{S^c}^{-1}(1) = nd$. This implies that $0 < F_{S^c}(d) < 1$. Now the result follows from Theorem 5.1 and Proposition 2.1. \square

Corollary 5.3. *Problem (1) has a unique solution given by $(F_{X_1}^{-1}(F_{S^c}(d)), \dots, F_{X_n}^{-1}(F_{S^c}(d)))$ if each f_i is strictly convex.*

Although this corollary is a standard result in the theory of convex minimization, here we will give a new and simple proof by using the theory of comonotonicity.

Proof: When each f_i is strictly convex, each F_{X_i} defined in (14) is strictly increasing on $[l_i, u_i]$, and hence $F_{X_i}^{-1}$ does not contain any discontinuity. In particular, this implies that $F_{X_i}^{-1}(F_{S^c}(d)) = F_{X_i}^{-1+}(F_{S^c}(d))$ and so by Corollary 5.2, Problem (1) has a unique solution given by $(F_{X_1}^{-1}(F_S(d)), \dots, F_{X_n}^{-1}(F_S(d)))$. \square

The next result can be found in Bitran and Hax (1981). Instead of proving it using Kuhn-Tucker theory, we demonstrate that it is a direct consequence of Corollary 5.2.

Corollary 5.4. *Suppose that f_i is strictly increasing on $[l_i, u_i]$ for $i \in J_I \subset \{1, \dots, n\}$ and is strictly decreasing on $[l_i, u_i]$ for $i \in J_D \subset \{1, \dots, n\}$. If (d_1^*, \dots, d_n^*) is a solution of Problem (1), then either $d_i^* = l_i$ for all $i \in J_I$, or $d_i^* = u_i$ for all $i \in J_D$, or both.*

Proof: If f_i is strictly increasing on $[l_i, u_i]$, the corresponding F_{X_i} in (14) jumps at l_i from 0 to $F_{X_i}(l_i) > 1/2$, and hence $F_{X_i}^{-1}(p) = F_{X_i}^{-1+}(p) = l_i$ on an interval containing $K_I \supset (0, 1/2]$. Similarly, if f_i is strictly decreasing on $[l_i, u_i]$, the corresponding F_{X_i} in (14) jumps at u_i from $F_{X_i}(u_i-) < 1/2$ to 1, and hence $F_{X_i}^{-1}(p) = F_{X_i}^{-1+}(p) = u_i$ on an interval containing $K_D \supset [1/2, 1)$. From Corollary 5.2, if $F_{S^c}(d) \in K_I$, then $d_i^* = l_i$ for all $i \in J_I$; if $F_{S^c}(d) \in K_D$, then $d_i^* = u_i$ for all $i \in J_D$. As $K_I \cup K_D = (0, 1)$, the result follows. \square

6 Examples

In this section, we provide three examples to illustrate how the theory developed above can be used to solve practically relevant problems.

Example 1 Let $X_i, i = 1, \dots, n$ be positive and integrable risks. We consider the following optimal capital allocation problem, which is a special case of Problem (9) studied in Section 3:

$$\min_{d_1 + \dots + d_n = d, d_i \geq 0} \sum_{i=1}^n \text{TVaR}_\alpha((X_i - d_i)_+), \quad d > 0. \quad (16)$$

Here, TVaR_α denotes the Tail Value-at-Risk at probability level $\alpha \in (0, 1)$ which is defined by

$$\text{TVaR}_\alpha(X) := \frac{1}{1 - \alpha} \int_\alpha^1 F_X^{-1}(p) dp$$

for any integrable random variable X . Since Tail Value-at-Risk is law-invariant, increasing, and comonotonic additive, it follows from Proposition 3.3 that the solution set of Problem (16) contains $i(d, \mathbf{X}^c)$. Our objective here is to derive this result by using the methodology developed in Sections 4 and 5.

To this end, we first define $f_i(d_i) := \text{TVaR}_\alpha((X_i - d_i)_+)$ for $i = 1, \dots, n$. Each $f_i : [0, \infty) \rightarrow$

\mathbb{R} is a decreasing and convex function, and it is straightforward to obtain

$$f'_i(d_i) = \begin{cases} -1 & \text{if } d_i < F_{X_i}^{-1}(\alpha), \\ -\frac{1-F_{R_i}(d_i)}{1-\alpha} & \text{if } d_i \geq F_{X_i}^{-1}(\alpha). \end{cases}$$

So it follows from Proposition 4.2 that we can express f_i as

$$f_i(d_i) = \mathbb{E}[(Y_i - d_i)_+], \quad d_i \geq 0,$$

where the distribution function of Y_i is given by

$$F_{Y_i}(x) = \begin{cases} 0 & \text{if } x < F_{X_i}^{-1}(\alpha), \\ \frac{F_{X_i}(x) - \alpha}{1 - \alpha} & \text{if } x \geq F_{X_i}^{-1}(\alpha). \end{cases} \quad (17)$$

Using this transformation, Problem (16) becomes

$$\min_{d_1 + \dots + d_n = d, d_i \geq 0} \sum_{i=1}^n \mathbb{E}[(Y_i - d_i)_+],$$

and hence Corollary 5.2 implies that the solution set of Problem (16) is given by $i(d, \mathbf{Y}^c)$:

$$\{(d_1, \dots, d_n) \mid d_1 + \dots + d_n = d, F_{Y_i}^{-1}(F_{S_Y^c}(d)) \leq d_i \leq F_{Y_i}^{-1+}(F_{S_Y^c}(d)) \text{ for all } i\}, \quad (18)$$

in which $S_Y^c := Y_1^c + \dots + Y_n^c$ and (Y_1^c, \dots, Y_n^c) is a comonotonic copy of (Y_1, \dots, Y_n) . From relationship (17), which is piecewise linear, it is easy to express $F_{Y_i}^{-1}$ in terms of $F_{X_i}^{-1}$, and $F_{S_Y^c}$ in terms of $F_{S_X^c}$ where S_X^c denotes the corresponding sum of a comonotonic copy of (X_1, \dots, X_n) . We then can easily show that $i(d, \mathbf{Y}^c)$ is identical to $i(d, \mathbf{X}^c)$.

Example 2 Consider the following optimal capital allocation problem:

$$\min_{d_1 + \dots + d_n = d, d_i \geq l_i} \sum_{i=1}^n s_i \exp(-m_i d_i), \quad (19)$$

where s_i, m_i are some strictly positive constants. Notice that $d_i \mapsto -s_i \exp(-m_i d_i)$ is a standard utility function with *constant absolute risk aversion*. It is assumed that

$$d > L := l_1 + \dots + l_n \quad (20)$$

in order to avoid that the problem is trivial or ill-posed. To simplify the notation, we define

$$\theta_i := \frac{\ln m_i s_i}{m_i}, \quad \Theta_i := \sum_{j=1}^i \theta_j, \quad A_i := 1 - m_i s_i \exp(-m_i l_i), \quad M_i := \sum_{j=1}^i \frac{1}{m_j}, \quad L_i := \sum_{j=i+1}^n l_j,$$

for $i = 1, \dots, n$. Note that $L_n := 0$ by convention. Without loss of generality, we assume that $A_1 \leq \dots \leq A_n$.

Proposition 6.1. *For any given $d > L$, define*

$$i^* := \inf \left\{ i \in \{1, \dots, n-1\} \mid d \leq L + \sum_{j=1}^i M_j (\ln(1 - A_j) - \ln(1 - A_{j+1})) \right\} \wedge n, \quad (21)$$

with the convention that $\inf \emptyset = \infty$. Then the solution to Problem (19) is given by

$$d_i^* = \begin{cases} l_i, & i = i^* + 1, \dots, n, \\ \theta_i - \frac{\Theta_{i^*} + L_{i^*} - d}{m_i M_{i^*}}, & i = 1, \dots, i^*. \end{cases}$$

Proof: We first notice that every $f_i(d_i) := s_i \exp(-m_i d_i)$ is strictly decreasing and strictly convex. Without loss of generality, we may assume that s_1, \dots, s_n have been rescaled properly such that $-1 \leq (f_i)'_+(l_i)$ for all i . Since $\lim_{x \rightarrow \infty} f_i(x)$ exists in \mathbb{R} and $\lim_{x \rightarrow \infty} (f_i)'_+(x) = 0$ for all i , it follows from Proposition 4.2 and Corollary 5.3 that Problem (19) has a unique solution (d_1^*, \dots, d_n^*) given by $d_i^* = F_{X_i}^{-1}(F_{S^c}(d))$ where $S^c := F_{X_1}^{-1}(U) + \dots + F_{X_n}^{-1}(U)$ for any uniform(0, 1) random variable U and

$$F_{X_i}(x) = \begin{cases} 0, & x < l_i, \\ 1 - m_i s_i \exp(-m_i x), & l_i \leq x. \end{cases}$$

It remains to compute $F_{S^c}(d)$ and $F_{X_i}^{-1}$.

The inverse of this distribution function is given by

$$F_{X_i}^{-1}(p) = \begin{cases} l_i, & 0 < p \leq A_i \\ \frac{1}{m_i} \ln \frac{m_i s_i}{1-p}, & A_i \leq p < 1, \end{cases} = \left(\frac{1}{m_i} \ln \frac{m_i s_i}{1-p} \right) \vee l_i.$$

Also, for any $p \in (0, 1)$, we find that

$$\begin{aligned} F_{S^c}^{-1}(p) &= F_{X_1}^{-1}(p) + \dots + F_{X_n}^{-1}(p) \\ &= \begin{cases} l_1 + \dots + l_n, & 0 < p \leq A_1, \\ \frac{1}{m_1} \ln \frac{m_1 s_1}{1-p} + l_2 + \dots + l_n, & A_1 \leq p \leq A_2, \\ \frac{1}{m_1} \ln \frac{m_1 s_1}{1-p} + \frac{1}{m_2} \ln \frac{m_2 s_2}{1-p} + l_3 + \dots + l_n, & A_2 \leq p \leq A_3, \\ \vdots \\ \frac{1}{m_1} \ln \frac{m_1 s_1}{1-p} + \dots + \frac{1}{m_n} \ln \frac{m_n s_n}{1-p}, & A_n \leq p < A_{n+1} := 1. \end{cases} \end{aligned} \quad (22)$$

Notice that $F_{S^c}^{-1}$ is continuous and is piecewise linear on $(0, A_1], [A_1, A_2], \dots, [A_n, A_{n+1})$. Simple algebraic manipulation shows that

$$F_{S^c}^{-1}(A_{i+1}) = L + \sum_{j=1}^i M_j (\ln(1 - A_j) - \ln(1 - A_{j+1})), \quad i = 1, \dots, n-1,$$

which is the expression in (21). The definition of i^* in (21) enables us to locate the exact “layer” that $F_{S^c}(d)$ belongs to, so that $A_{i^*} < F_{S^c}(d) \leq A_{i^*+1}$. By solving the equation $F_{S^c}^{-1}(F_{S^c}(d)) = d$ for $F_{S^c}(d)$ using this particular layer in (22), we obtain

$$F_{S^c}(d) = 1 - \exp\left(\frac{\Theta_{i^*} + L_{i^*} - d}{M_{i^*}}\right), \quad d > L. \quad (23)$$

Therefore, the solution to Problem (19) is given by

$$d_i^* = F_{X_i}^{-1}(F_{S^c}(d)) = \left(\frac{1}{m_i} \ln \frac{m_i s_i}{1 - F_{S^c}(d)} \right) \vee l_i, \quad i = 1, \dots, n.$$

Simplifying this expression yields the desired result. \square

For instance, if

$$L < d \leq \frac{1}{m_1} \ln \frac{m_1 s_1}{1 - A_2} + l_2 + \dots + l_n,$$

then $i^* = 1$ and $A_1 < F_{Sc}(d) \leq A_2$. Therefore, the solution is given by

$$d_i^* = \begin{cases} \theta_1 - \frac{\Theta_1 + L_1 - d}{m_1 M_1} = d - (l_2 + \dots + l_n), & i = 1, \\ l_i, & i = 2, \dots, n. \end{cases}$$

As another illustration, suppose that d is sufficiently large such that

$$d > \frac{1}{m_1} \ln \frac{m_1 s_1}{1 - A_n} + \dots + \frac{1}{m_n} \ln \frac{m_n s_n}{1 - A_n} = L + \sum_{j=1}^{n-1} M_j (\ln(1 - A_j) - \ln(1 - A_{j+1})),$$

then $i^* = n$ and $A_n < F_{Sc}(d) < 1$. Applying Proposition 6.1 yields that

$$d_i^* = \theta_i - \frac{\Theta_n - d}{m_i M_n} > l_i, \quad i = 1, \dots, n.$$

Example 3 In Dhaene et al. (2012), the following optimal capital allocation was considered:

$$\min_{d_1 + \dots + d_n = d} \sum_{i=1}^n \mathbb{E} \left[\frac{\zeta_i (Y_i - d_i)^2}{\nu_i} \right],$$

where ζ_1, \dots, ζ_n are positive random variables with mean 1, ν_1, \dots, ν_n are given strictly positive numbers summing to 1, and Y_1, \dots, Y_n are some square integrable random variables. We refer to Dhaene et al. (2012) for a detailed interpretation of this model. In that paper, it is shown that the optimal allocations are given by

$$d_i^* = \nu_i \left(d - \sum_{j=1}^n \mathbb{E}[\zeta_j Y_j] \right) + \mathbb{E}[\zeta_i Y_i], \quad i = 1, \dots, n.$$

Here, we want to add the box constraints $d_i \in [0, d]$ for all i to the minimization problem above. More precisely, we would like to apply the theory developed in the previous sections to solve the following problem:

$$\min_{d_1 + \dots + d_n = d, 0 \leq d_i \leq d} \sum_{i=1}^n \mathbb{E} \left[\frac{\zeta_i (Y_i - d_i)^2}{\nu_i} \right]. \quad (24)$$

To simplify our notation, we define $c_i := \mathbb{E}[\zeta_i Y_i]$ for $i = 1, \dots, n$ and assume without loss of generality that

$$\frac{c_1}{\nu_1} \geq \dots \geq \frac{c_n}{\nu_n}. \quad (25)$$

Proposition 6.2. *For any given $d > 0$, define*

$$i^* := \inf \left\{ i \in \{1, \dots, n-1\} \mid d \leq \sum_{j=1}^i \nu_j \left(\frac{c_j}{\nu_j} - \frac{c_{i+1}}{\nu_{i+1}} \right) \right\} \wedge n, \quad (26)$$

with the convention that $\inf \emptyset = \infty$. Then the solution to Problem (24) is given by

$$d_i^* = \begin{cases} \frac{\nu_i}{\sum_{j=1}^{i^*} \nu_j} \left(d - \sum_{j=1}^{i^*} c_j \right) + c_i, & i = 1, \dots, i^*, \\ 0, & i = i^* + 1, \dots, n. \end{cases}$$

Proof: We first let

$$f_i(x) := \mathbb{E} \left[\frac{\zeta_i(Y_i - x)^2}{\nu_i} \right], \quad x \in \mathbb{R}, i = 1, \dots, n.$$

Notice that the box constraints $d_i \in [0, d]$ for all i can be replaced by $d_i \in [0, u_i]$ for all i as long as each u_i is larger than d . In particular, we choose u_1, \dots, u_n greater than d such that

$$(f_1)'_+(u_1) = \dots = (f_n)'_+(u_n) \geq \max_{1 \leq i \leq n} (f_i)'_+(0),$$

which is equivalent to

$$\frac{2(u_1 - \mathbb{E}[\zeta_1 Y_1])}{\nu_1} = \dots = \frac{2(u_n - \mathbb{E}[\zeta_n Y_n])}{\nu_n} \geq \max_{1 \leq i \leq n} \left| \frac{-2\mathbb{E}[\zeta_i Y_i]}{\nu_i} \right|.$$

Moreover, we take ν to be the common value on the left-hand side of the inequality above. By Proposition 4.1, Problem (24) is equivalent to

$$\min_{d_1 + \dots + d_n = d, 0 \leq d_i \leq u_i} \sum_{i=1}^n \mathbb{E}[(X_i - d_i)_+],$$

where the distribution function of X_i is given by

$$F_{X_i}(x) = \begin{cases} 0 & \text{if } x < 0, \\ (1 + f'_+(x)/\nu)/2 & \text{if } 0 \leq x < u_i, \\ 1 & \text{if } x \geq u_i. \end{cases}$$

Our choice of u_1, \dots, u_n guarantees that none of the distribution function F_{X_i} has a point mass at u_i . Inverting the distribution function above yields that

$$F_{X_i}^{-1}(p) = \begin{cases} 0, & 0 < p \leq A_i \\ \nu\nu_i(p - 1/2) + c_i, & A_i \leq p < 1, \end{cases} = (\nu\nu_i(p - 1/2) + c_i)_+$$

where $A_i := 1/2 - \frac{c_i}{\nu\nu_i}$ for $i = 1, \dots, n$. From assumption (25), $A_1 \leq \dots \leq A_n$. Let S^c be the comonotonic sum $F_{X_1}^{-1}(U) + \dots + F_{X_n}^{-1}(U)$, where U is any uniform(0, 1) random variable. Then for $p \in (0, 1)$,

$$\begin{aligned} F_{S^c}^{-1}(p) &= F_{X_1}^{-1}(p) + \dots + F_{X_n}^{-1}(p) \\ &= \begin{cases} 0, & 0 < p \leq A_1, \\ \nu\nu_1(p - 1/2) + c_1, & A_1 \leq p \leq A_2, \\ \nu\nu_1(p - 1/2) + c_1 + \nu\nu_2(p - 1/2) + c_2, & A_2 \leq p \leq A_3, \\ \vdots & \\ \nu\nu_1(p - 1/2) + c_1 + \dots + \nu\nu_n(p - 1/2) + c_n, & A_n \leq p < 1. \end{cases} \end{aligned}$$

By a similar argument as in the proof of Proposition 6.1, we obtain

$$F_{S^c}(d) = \frac{1}{2} + \frac{d - \sum_{j=1}^{i^*} c_j}{\nu \sum_{i=1}^{i^*} \nu_j}. \quad (27)$$

From Corollary 5.3, the solution to Problem (24) is given by

$$d_i^* = F_{X_i}^{-1}(F_{S^c}(d)) = (\nu\nu_i(F_{S^c}(d) - 1/2) + c_i)_+, \quad i = 1, \dots, n.$$

Putting the expression of $F_{S^c}(d)$ in (27) in this formula yields the desired result. \square

As an illustration, consider the case where

$$0 < d \leq \nu_1 \left(\frac{c_1}{\nu_1} - \frac{c_2}{\nu_2} \right).$$

Then $i^* = 1$ and $d_1^* = d$ and $d_2^* = \dots = d_n^* = 0$. As another illustration, if

$$\sum_{j=1}^{n-1} \nu_j \left(\frac{c_j}{\nu_j} - \frac{c_n}{\nu_n} \right) < d,$$

then $i^* = n$ and

$$d_i^* = \nu_i \left(d - \sum_{j=1}^n c_j \right) + c_i, \quad i = 1, \dots, n,$$

which is the solution obtained in Dhaene et al. (2012) without any box constraints.

7 Some variants

7.1 Minimization of a weighted sum of stop-loss premiums

Consider the following variant of minimization Problem (7):

$$\min_{d_1 + \dots + d_n = d, l_i \leq d_i \leq u_i} \sum_{i=1}^n \nu_i \mathbb{E}[(Y_i - d_i)_+], \quad (28)$$

where ν_1, \dots, ν_n are some strictly positive constants that are not all equal, and Y_1, \dots, Y_n are some integrable random variables with possibly unbounded support.

We first remark that the support of each Y_i in Problem (28) can be assumed to be contained in $[l_i, u_i]$ without loss of generality. For if $d_i \in [l_i, u_i]$,

$$\mathbb{E}[(Y_i - d_i)_+] = \mathbb{E}[(Y_i \vee l_i - d_i)_+] = \mathbb{E}[((Y_i \vee l_i) \wedge u_i - d_i)_+] + \mathbb{E}[(Y_i - u_i)_+],$$

and thus Y_i can be replaced by $(Y_i \vee l_i) \wedge u_i$ in Problem (28) without changing the solution set. For the remainder of this section, we assume that the support of each Y_i is contained in $[l_i, u_i]$.

To solve Problem (28), one may simply treat it as a special case of Problem (1) by writing

$$f_i(d_i) := \nu_i \mathbb{E}[(Y_i - d_i)_+], \quad d_i \in [l_i, u_i],$$

and proceed as in Section 4 to express $f_i(d_i)$ as an affine function of $\mathbb{E}[(X_i - d_i)_+]$ for some suitable X_i so that the leading coefficients are equalized.

In what follows, we present a simple trick to accomplish this transformation by using suitable Bernoulli variables to “absorb” the coefficients ν_i . To explain this approach, we first assume, without loss of generality, that each ν_i is strictly less than 1. If not, we may simply replace ν_i by $\nu_i / \sum \nu_i$. Let Z_1, \dots, Z_n be Bernoulli variables which are independent of Y_1, \dots, Y_n so that

$$Z_i = \begin{cases} 1 & \text{with probability } \nu_i, \\ 0 & \text{with probability } 1 - \nu_i, \end{cases}$$

and $X_i := Z_i(Y_i - l_i) + l_i$. Then for any $l_i \leq d_i \leq u_i$,

$$\begin{aligned} \nu_i \mathbb{E}[(Y_i - d_i)_+] &= \mathbb{P}(Z_i = 1) \mathbb{E}[((Y_i - l_i) - (d_i - l_i))_+] \\ &= \mathbb{E}[(Z_i(Y_i - l_i) - (d_i - l_i))_+] \\ &= \mathbb{E}[(X_i - d_i)_+] \end{aligned}$$

by the assumed independence between Z_i and X_i .

7.2 Nonlinear constraints

In Problem (1), the linear constraint $d_1 + \dots + d_n = d$ can be replaced by a non-linear constraint of the form

$$h_1(d_1) + \dots + h_n(d_n) = d,$$

where each h_i is a 1-1 function so that the inverse h_i^{-1} is well-defined (on a suitable domain). Let $\tilde{d}_i := h_i(d_i)$ and $\tilde{f}_i(x) = f_i(h_i^{-1}(x))$ for all i . If each h_i^{-1} is convex and f_i is increasing and convex, or h_i^{-1} is concave and f_i is decreasing and convex, then \tilde{f}_i is convex as well, and hence the corresponding minimization problem is reduced to the form of Problem (1) again.

8 Connection with infimum-convolution

If the box constraints $d_i \in [l_i, u_i]$ for all i in Problem (1) are removed (but $d_1 + \dots + d_n = d$ is kept), and the domain of each real-valued convex function f_i is \mathbb{R} instead of $[l_i, u_i]$, then

$$(\wedge_{i=1}^n f_i)(d) := \min_{d_1 + \dots + d_n = d} \sum_{i=1}^n f_i(d_i) \quad (29)$$

is called the *infimum-convolution* of f_1, \dots, f_n in convex analysis (cf. Rockafellar (1970)). Notice that the minimization in (29) may not have a solution in general.

Proposition 8.1. *Consider some (d_1^*, \dots, d_n^*) with $d_1^* + \dots + d_n^* = d$. Then*

$$(\wedge_{i=1}^n f_i)(d) = \sum_{i=1}^n f_i(d_i^*)$$

if and only if

$$\partial f_1(d_1^*) \cap \dots \cap \partial f_n(d_n^*) \neq \emptyset. \quad (30)$$

Here, $\partial f_i(d_i) := [(f_i)'_-(d_i), (f_i)'_+(d_i)]$ is the subdifferential of f_i at d_i . The objective here is to prove this result using the perspective of comonotonicity and the theory we developed in previous sections.

Proof: We first prove the “if” part. Fix some (d_1^*, \dots, d_n^*) with $d_1^* + \dots + d_n^* = d$ and some u^* such that

$$u^* \in \partial f_1(d_1^*) \cap \dots \cap \partial f_n(d_n^*).$$

Choose any $u_1, \dots, u_n, l_1, \dots, l_n$ such that $l_i < d_i^* < u_i$ for all i , and let $\mathbf{X}^c = (X_1^c, \dots, X_n^c)$ be a comonotonic random vector with marginal distributions given by (14) for some large enough ν . By (14), $(f_i)'_-(d_i^*) = \nu(2F_{X_i}(d_i^* -) - 1)$ and $(f_i)'_+(d_i^*) = \nu(2F_{X_i}(d_i^*) - 1)$, and so

$$F_{X_i}(d_i^* -) \leq u^{**} \leq F_{X_i}(d_i^*) \quad \text{for all } i,$$

where $u^{**} := (u^*/\nu + 1)/2$. This can be rewritten as

$$F_{X_i}^{-1}(u^{**}) \leq d_i \leq F_{X_i}^{-1+}(u^{**}) \quad \text{for all } i.$$

In particular, this implies that $(d_1^*, \dots, d_n^*) \in i(d, \mathbf{X}^c)$. By Theorem 5.1, (d_1^*, \dots, d_n^*) solves the problem

$$\min_{d_1 + \dots + d_n = d, l_i \leq d_i \leq u_i} \sum_{i=1}^n f_i(d_i).$$

However, as the l_i 's and the u_i 's can be chosen arbitrarily small and large respectively, we conclude that

$$\min_{d_1 + \dots + d_n = d} \sum_{i=1}^n f_i(d_i) = \sum_{i=1}^n f_i(d_i^*).$$

This proves the “if” part.

For the “only if” part, suppose that $d_1^* + \dots + d_n^* = d$ and $(\bigwedge_{i=1}^n f_i)(d) = \sum_{i=1}^n f_i(d_i^*)$. Choose any $u_1, \dots, u_n, l_1, \dots, l_n$ such that $l_i < d_i^* < u_i$ for all i . Then (d_1^*, \dots, d_n^*) solves the problem

$$\min_{d_1 + \dots + d_n = d, l_i \leq d_i \leq u_i} \sum_{i=1}^n f_i(d_i).$$

It then follows from Theorem 5.1 and Proposition 2.1 that $F_{X_i}^{-1}(F_{S^c}(d)) \leq d_i^* \leq F_{X_i}^{-1+}(F_{S^c}(d))$ for all i , where (X_1^c, \dots, X_n^c) is a comonotonic random vector with marginal distributions given by (14) for some large enough ν , and $S^c := X_1^c + \dots + X_n^c$. From the proof of the “if” part, we have

$$\nu(2F_{S^c}(d) - 1) \in \partial f_1(d_1^*) \cap \dots \cap \partial f_n(d_n^*).$$

This shows that the right hand intersection is non-empty. \square

Proposition 8.1 is well-known. It gives necessary and sufficient conditions for (d_1^*, \dots, d_n^*) to be the solution of minimization problem (29).

9 Acknowledgments

Ka Chun Cheung acknowledges the financial support of the Research Grants Council of HKSAR (GRF Project 17324516). Jan Dhaene acknowledges the financial support of Onderzoeksfonds KU Leuven (GOA/13/002). Phillip Yam acknowledges the financial support from

The Hong Kong RGC GRF 14301015 with the project title: Advance in Mean Field Theory and The Hong Kong RGC GRF 14300717 with the project title: New kinds of Forward-backward Stochastic Systems with Applications. Phillip Yam also acknowledges the financial support from Department of Statistics of Columbia University in the City of New York during the period he was a visiting faculty member.

References

- [1] Bitran, G.R., Hax, A.C., (1981). Disaggregation and resource allocation using convex knapsack problems with bounded variables. *Management Science*, 27, 431-441.
- [2] Chen, X., Deelstra, G., Dhaene, J., Linders, D., Vanmaele, M., (2015). On an optimization problem related to static super-replicating strategies. *Journal of Computational and Applied Mathematics*, 278, 213-230
- [3] Cheung, K.C., Rong, Y., Yam, S.C., (2014). Borch's Theorem from the perspective of comonotonicity. *Insurance: Mathematics and Economics*, 54, 144-151.
- [4] Denneberg, D., (1994). *Non-additive Measure and Integral*. Kluwer Academic Publishers, Boston.
- [5] Dhaene, J., Denuit, M., Goovaerts, M.J., Kaas, R., Vyncke, D., (2002). The concept of comonotonicity in actuarial science and finance: theory. *Insurance: Mathematics and Economics*, 31, 3-33.
- [6] Dhaene, J., Tsanakas, A., Valdez, E., Vanduffel, S., (2012). Optimal capital allocation principles. *Journal of Risk and Insurance*, 79, 1-28.
- [7] Föllmer, H, Schied, A., (2004). *Stochastic Finance: An Introduction in Discrete Time*, 2nd Edition. de Gruyter, Berlin.
- [8] Frostig, E., Zaks, Y, Levikson, B., (2007). Optimal pricing for a heterogeneous portfolio for a given risk factor and convex distance measure. *Insurance: Mathematics and Economics*, 40, 459-467.
- [9] Kaas, R., Dhaene, J., Goovaerts, M., (2000). Upper and lower bounds for sums of random variables. *Insurance: Mathematics and Economics*, 27, 151-168.
- [10] Kaas, R., Dhaene, J., Vyncke, D., Goovaerts, M., Denuit, M., (2002). A simple geometric proof that comonotonic risks have the convex-largest sum. *ASTIN Bulletin*, 32, 71-80.
- [11] Luss, H., Gupta, S.K., (1975). Allocation of Effort Resources Among Competing Activities. *Operations Research*, 23, 360-366.
to the length of critical paths. *Journal of Applied Probability*, 16, 671-677.
- [12] Revuz, D., Yor, D., (1999). *Continuous Martingales and Brownian Motion*, 3rd Edition. Springer, Berlin.
- [13] Rockafellar, R., (1970). *Convex Analysis*. Princeton University Press, Princeton, N.J.

- [14] Rüschemdorf, L., (2013). *Mathematical Risk Analysis: Dependence, Risk Bounds, Optimal Allocations and Portfolios*. Springer, Heidelberg.
- [15] Stefanov, S., (2005). Convex separable minimization problems with a linear constraint and bounded variables. *International Journal of Mathematics and Mathematical Sciences*, 9, 1339-1363.
- [16] Wilkinson, C., Gupta, S.K., (1969). Allocating Promotional Effort to Competing Activities: A Dynamic Programming Approach. *IFORS Conference, Venice*, 419-432.
- [17] Zaks, Y, (2013). Optimal capital allocation - a generalization of the optimization problem. *Insurance Markets and Companies: Analyses and Actuarial Computations International Research Journal*, 4, 29-34.
- [18] Zaks, Y, Frostig, E., Levikson, B., (2006). Optimal pricing of a heterogeneous portfolio for a given risk level. *ASTIN Bulletin*, 36, 161-185.