

Maternal copy number variations in the *DMD* gene as secondary findings in non-invasive prenatal testing

Nathalie Brison^{1,2,#}, PhD, Jazz Storms^{2,3,#}, Darine Villela², PhD, Kristl G. Claey^{4,5}, MD PhD, Luc Dehaspe^{1,2}, PhD, Thomy de Ravel^{1,2}, MD PhD, Liesbeth De Waele^{6,7}, MD PhD, Nathalie Goemans⁶, MD PhD, Eric Legius^{1,2}, MD PhD, Hilde Peeters^{1,2}, MD PhD, Hilde Van Esch^{1,2}, MD PhD, Valerie Race^{1,2}, PhD, Joris Robert Vermeesch^{1,2}, PhD, Koenraad Devriendt^{1,2}, MD PhD, Kris Van Den Bogaert^{1,2}, PhD

¹Center for Human Genetics, University Hospitals Leuven, Leuven, Belgium

²Department of Human Genetics, KU Leuven, Leuven, Belgium

³Faculty of Medicine, KU Leuven, Leuven, Belgium

⁴Department of Neurology, University Hospitals Leuven, Leuven, Belgium

⁵Laboratory for Muscle Diseases and Neuropathies, Department of Neurosciences, KU Leuven, Leuven, Belgium

⁶Department of Pediatric Neurology, University Hospitals Leuven, Leuven, Belgium

⁷KU Leuven Department of Development and Regeneration, Campus Kulak Kortrijk, Kortrijk, Belgium

#shared first co-authors

Running title: Maternal CNV's in the *DMD* gene detected by NIPS

Word count : 3813

Table count : 2

Figure count : 3

Corresponding author : Kris Van Den Bogaert, Center for Human Genetics, UZ Leuven, Herestraat 49, 3000 Leuven, Belgium, [+32 16 340399](tel:+3216340399), kris.vandenbogaert@uzleuven.be

ABSTRACT

Purpose: Non-invasive prenatal testing (NIPS) using genome sequencing also reveals maternal copy number variations (CNV's). Those CNV's can be clinically actionable or might be harmful for the fetus if inherited. CNV's in the *DMD* gene potentially causing dystrophinopathies are amongst the most commonly observed maternal CNV's. We present our experience with maternal *DMD*-gene CNV's detected by NIPS.

Methods: We analyzed the data of maternal CNV's detected in the *DMD* gene revealed by NIPS.

Results: Out of 26.123 NIPS analyses, 16 maternal CNV's in the *DMD* gene were detected (1/1.632 pregnant women). Variant classification regarding pathogenicity and phenotypic severity was based on public databases, segregation analysis in the family and prediction of the effect on the reading frame. Ten CNV's were classified as pathogenic, four as benign, whereas two remained unclassified.

Conclusion: NIPS leverages CNV screening in the general population of pregnant women. We implemented a strategy for the interpretation and the returning of maternal CNV's in the *DMD* gene detected by NIPS.

Keywords: maternal CNV, *DMD* gene, secondary findings, NIPS

INTRODUCTION

Noninvasive prenatal testing (NIPS) based on the analysis of cell-free DNA (cfDNA) in the maternal serum has enabled the detection of fetal aneuploidies, typically trisomy 21, 13 and 18, with a high sensitivity and specificity^{1,2}. Because of its high accuracy, NIPS was rapidly implemented as part of routine prenatal care. Notably, whole-genome massively parallel sequencing allows not only the analysis of the viable trisomies, but also fetal aneuploidies of other chromosomes, and has the potential to detect smaller fetal chromosomal imbalances as well³⁻⁵. Moreover, since on average 90% of the cfDNA in the maternal circulation originates from maternal cells, a genome-wide cfDNA analysis equally results in a high resolution screen for maternal copy number variations (CNV's)⁶. Hence, NIPS has the potential to detect relevant maternal CNVs that can be clinically actionable or could be potentially harmful for the fetus if inherited⁶⁻⁸.

Duchenne Muscular Dystrophy (DMD) is an X-linked condition that presents as a progressive muscle disorder in young boys, affecting both skeletal and cardiac muscle⁹. With an incidence at birth of 1/3.800 to 1/6.300, it is one of the most common severe, untreatable neuromuscular disorders¹⁰. It is caused by a loss-of-function variant in the *DMD* gene (Xp21.2-p21.1), and represents the severe end of the spectrum of diseases collectively called dystrophinopathies. Becker Muscular Dystrophy (BMD) on the contrary is a milder dystrophinopathy, with onset later in childhood and with an incidence of 1 in 20.000 to 30.000 males. A milder phenotype is DMD-associated cardiomyopathy, also called X-linked dilated cardiomyopathy, with onset usually in young adults¹¹. Asymptomatic individuals with isolated elevated serum creatine kinase (CK) have also been reported. Most carrier females are asymptomatic, but there is an increased risk for muscle damage and dilated cardiomyopathy¹². Of interest, the *DMD* gene is one of the largest genes in the human genome, with 79 exons, and most of its variants correspond to deletions (60-70%) and duplications (5-10%), the remaining being single nucleotide variants. Given that a large proportion of variants in the *DMD* gene are inherited CNV's, genome-wide NIPS is expected to detect carrier mothers for DMD⁹.

When a CNV in the *DMD* gene is detected, sometimes it is challenging to predict the phenotype for male carriers. The most reliable phenotypic predictions are based on family history. However, in population-based genome-wide analysis, family histories of dystrophinopathy are lacking. To interpret the potential pathogenicity, different databases with genotype-phenotype correlations of *DMD*-gene variants can be consulted. The best known are the Leiden Open Variation Database (LOVD) and the UMD-DMD France Database^{13,14}. The effect of novel

hitherto unreported CNV's may be predicted using the frameshift-rule, which permits to distinguish DMD from BMD with a 91-92% accuracy⁹. In general, frameshift variants cause DMD whereas in-frame variants cause BMD¹³. However, exceptions exist, e.g. 2% of out-of-frame deletions and duplications will actually result in BMD and 7% of all DMD cases is caused by an in-frame variant. This indicates that other factors including size and position of the CNV may also play a role. The interpretation is further complicated by the variable expression observed for certain CNV's with apparently the same breakpoints¹³. Whether or not to report a maternal CNV detected by NIPS depends on reliable prediction of the phenotypic consequences of that specific CNV.

The Belgian Advisory Committee on Bioethics (9 May 2016) stated that secondary findings of clinical relevance detected during NIPS should be reported: "*Where this information may lead to a preventive or therapeutic intervention, it is important to also share this information with the patient in the context of clinical genetics. The failure to do so may be construed as serious negligence*"¹⁵. This prompted the Belgian Society for Human Genetics to issue guidelines on how to deal with clinically significant secondary findings, both maternal and fetal, detected by

NIPS¹⁶. According to these guidelines, carriership for X-linked recessive disorders will be communicated, irrespective of the sex of the fetus. Since July 1st 2017, NIPS is reimbursed by the National Health System for all pregnant women in Belgium. In this study, we report a systematic review of the cases where NIPS revealed a maternal CNV involving the *DMD* gene from a single genetic center, during a one year period following the universal reimbursement in Belgium. Since current expertise on this topic is limited, and guidelines are either lacking or based on expert opinion, here we propose a strategy to interpret and return variants in the *DMD* gene.

MATERIALS AND METHODS

Sample collection

This study was approved by the Ethical Committee at UZ/KU Leuven (MP004717). Written informed consent for returning secondary findings was obtained from all the participants by the referring physician. The study population consists of all pregnant women in Belgium undergoing a NIPS at the Center for Human Genetics, UZ Leuven, Leuven, Belgium in a one year period from July 1st 2017 to June 30th 2018.

Genetic technology

NIPS and CNV detection were carried out as described^{3,6,17}. Briefly, low-pass genome sequencing generated ~ 10 million single-end reads of 36 bp per sample, corresponding to a 0.1x coverage. These reads were mapped to human genome build 19 (hg19), and PCR duplicates were removed. The resulting genomic coordinates of read starts served as input to CNV caller SeqCBS²¹, in combination with a sex-matched control obtained by pooling read start coordinates from 50 male (46,XY) or 50 female (46,XX) “normal” pregnancies. To confirm the presence of the respective maternal CNV’s by chromosomal microarray analysis, DNA was extracted from maternal white blood cells, obtained from the stored buffy coats of the NIPS blood sample. Array-CGH was performed using the 8 × 60 K CytoSure ISCA v3 microarray (Oxford Gene Technology Oxford, UK) as described¹⁸. Array-CGH genomic coordinates are based on human genome build 19 (hg19).

Variant classification

The data were retrieved from patient and laboratory files. Phenotype predictions were based on the following principles. First, we assessed whether the CNV’s had already been described in patients with a dystrophinopathy, using two publicly available databases: (i) The UMD-DMD France database describes both published and unpublished genetic and clinical information about variants in the *DMD* gene. Molecular data were derived from 14 diagnostic laboratories in France. The database only includes CNV’s with known exon boundaries; (ii) The Leiden Open Variation Database (LOVD) contains genetic and in many cases also phenotypic data from over 16.000 patients with a *DMD* gene variant. Second, we determined whether the CNV was predicted to be in-frame or out-of-frame, using the online tool in the LOVD. Third, for the reported variants, segregation analysis within the family was proposed during the counselling, using as many available and potentially informative individuals as possible. Clinical information included a detailed history and, when available, serum creatine kinase values.

RESULTS

Maternal CNV’s in the *DMD* gene

Sixteen maternal CNV’s were detected in the *DMD* gene out of 26.123 NIPS analyses performed. This corresponds to an incidence of 1/1.632 pregnant women. The data are summarized in **Table 1**.

The genomic representation of the region containing the *DMD* gene, displaying the 16 maternal CNV's detected by NIPS analysis is illustrated in **Figure 1**. At 0.1x coverage, the performance of SeqCBS CNV calling can be expected to vary per CNV size and per region. Therefore, confirmation with an independent technique is necessary, especially for smaller CNVs and if the exonic content of a CNV determines the clinical outcome, as is the case for DMD/BMD. The size and position of all CNV's were validated by chromosome microarray analysis, using DNA extracted from maternal white blood cells (**Table 2**). The array platform includes probes for up to 502 highly-targeted developmental delay genes enabling detection of single exon aberrations. Higher probe density across the exons of the *DMD* gene allows improved detection of exonic CNV's. All 16 maternal CNV's detected by NIPS were confirmed by chromosomal microarray analysis. In four cases, the array-CGH further refined the CNV size and exonic content (family 4, 6, 8, and 12) (**Table 2**). **Figure 2** shows an example (family 4) of array-CGH refinement of one *DMD* CNV detected by NIPS.

Variant classification

Of the 16 families, 3 carried the same recurrent in-frame duplication of exons 10-27 (family 10, 11 and 12). Thus, in total, 14 different CNV's were observed. Nine of these 14 CNV's (family 1-9) were present in the databases with sufficient phenotypic data, enabling correct interpretation. In addition, family segregation data were available in 5 of these 9 families (family 1-5). In three families (family 1-3), the CNV had previously been diagnosed. In family 1, the mother has a son affected with DMD and was a known carrier of a pathogenic *DMD* deletion (exon 51). The pregnancy was established after preimplantation genetic diagnosis (PGD). In family 2, the CNV was detected as a secondary finding by microarray-CGH in her nephew with unexplained intellectual disability. This information was not known to us when the NIPS was performed. In family 3, segregation confirmed the presence of a mild BMD phenotype described in the literature. In two families (family 4 and 5) segregation was not informative; in family 4 the variant was maternally inherited, and in family 5 the variant is *de novo* so no clinical information was gained. In the remaining 4 families (6-9) no segregation analysis could be performed.

The five remaining CNV's (family 10-16) had not been reported before and further steps were needed to classify them. Segregation analysis enabled classification of 2. Three apparently unrelated women (family 10-12) carried a recurrent in-frame duplication of exons 10-27. The

presence of this CNV in some clinically unaffected adult male relatives allowed classification of the CNV as likely benign. Therefore, this variant is considered as a rare local benign CNV. Also in family 13, paternal inheritance allowed classification of this CNV as likely benign. One novel out-of-frame duplication (family 14, exons 51-62), for which family segregation was not informative, was classified as likely pathogenic for DMD. The variants detected in family 15 and family 16 remained unclassified.

Based on the experience we present here, we propose a strategy for the interpretation and reporting of CNV's in the *DMD* gene (**Figure 3**).

DISCUSSION

Here, we report our experience with NIPS and the incidental detection of maternal CNV's in the *DMD* gene in a one year period. Since NIPS is reimbursed for all pregnant women in the time period studied, this population is likely to be representative of the Belgian population of reproductive women, with a standard risk of having a child with a dystrophinopathy. In a one year period, we performed 26.123 NIPS tests and we observed 16 CNV's in the *DMD* gene. CNV's in DMD are thus present in 1/1.632 women.

Variant classification is challenging, especially in a setting where the *DMD* CNV is not ascertained through an affected family member. Nguyen *et al.* reported chromosomal microarray data of five women referred for diagnosis of intellectual disability with a CNV in the *DMD* gene¹⁹. In two, the CNV was paternally inherited and classified as benign, but in the other three, the CNV was not detected in any male family member. This led to challenges regarding disclosure of genetic risk to the family. Of the 16 families investigated in this study, 3 carried a recurrent in-frame duplication of exons 10-27. Thus, 14 different CNV's were observed. The most reliable predictor of the phenotype relies on previously known genotype-phenotype correlations, either from affected family members or from the databases. Of interest, only 9 of the 14 CNV's have been reported in public databases. Of these nine, four were associated with likely or known DMD, two with likely BMD and three with a variable phenotype. Segregation analysis in the family can contribute to variant classification. In two families the mother was a carrier of a previously detected familial *DMD* CNV: (i) a pathogenic out-of-frame DMD deletion (exon 51), resulting in the expected Duchenne muscular dystrophy phenotype in her son, and (ii) an in-frame duplication without associated phenotypic manifestations in this family. Segregation could be carried out in 3 families: in 2 families it was

not informative and in one it confirmed the mild BMD phenotype described in the literature. Of the five novel variants, segregation analysis could be performed for three. Based on this analysis two CNV's were classified as likely benign while one family was not informative. The reading frame rule predicts that out-of-frame CNV's have a more than 90% risk of being associated with DMD. The out-of-frame duplication of exons 51-62 was therefore reclassified from variant of unknown significance (VUS) to likely pathogenic. A major challenge in variant classification is the observation that for 3 of the detected CNV's, the databases report a variable phenotype. Variability in expression is well known in *DMD* gene variants. Several explanations have been put forward. One possibility is that the positions of the CNV's are not exact, due to an imperfect resolution of the technologies used^{13,20}. In a diagnostic setting, the phenotype is usually clinically evident and relies less on the underlying genetic defect. In contrast, imperfect breakpoint mapping may hamper the prediction of the phenotype in a family without affected individuals.

After exclusion of the two previously known carrier women, of the 16 pregnancies, 8 carried a CNV associated with a dystrophinopathy, conferring a 50% risk of an affected male fetus in 6 of them. From the incomplete follow-up data, we know that for 2 male fetuses at risk for DMD, an invasive test was performed, with one normal result and one affected result. However, even in pregnancies with a female fetus, this information can be important for the mother herself, given the increased risk of manifestations for future pregnancies and for relatives. In accordance with the guidelines from the Belgian Society for Human Genetics on secondary findings detected by NIPS, these results were reported.

One of the main challenges in managing secondary findings during prenatal diagnosis is maintaining the balance between providing actionable genetic information and not returning information which harms the patient²⁰. The actionability of returning CNV's with known or high risk of being associated with DMD is evident. However, this is less evident for BMD-associated variants. This is further complicated by the fact that the severity of BMD can be highly variable. Therefore, segregation analysis can aid in variant classification, as it did in some of the families in this study. However, this necessitates reporting and discussing the results, which may cause unnecessary anxiety. There are several arguments not to report unclassified variants, unless they are associated with a high risk of a severe phenotype, i.e. when they cause a frame-shift. First, novel CNV's, absent in the very extensive disease databases, are more likely to represent benign CNV's without phenotypic consequences. Second, segregation analysis is unlikely to provide evidence for severe phenotypic consequences, but rather for a

likely benign effect when observed in unaffected male adults. Classification as DMD- or BMD-associated can only be achieved in the case of a positive family history for the disease. A preconception family history should have revealed this. Third, after reporting the variants, genetic counselling was offered to all couples without any delay, together with physicians from the neuromuscular team when indicated. In one of the pregnancies where a novel CNV was found, the couple decided to terminate the pregnancy after their healthy 3 year old son was found with mildly elevated CK's. They opted not to wait for the results of the healthy maternal father, who was subsequently also found to be a carrier.

In summary, NIPS results can return relevant and clinically actionable variants to the mother. We show that NIPS leverages a screening in the general population of pregnant women, which enables CNV detection causing dystrophinopathies. Since no studies exist on CNV's in the *DMD* gene in women following NIPS, and current expertise on this topic is limited, we presented a strategy to interpret and return variants in the *DMD* gene.

Conflict of interest statement: No conflicts of interest

REFERENCES

1. Bianchi DW, Chiu RWK. Sequencing of Circulating Cell-free DNA during Pregnancy. *N Engl J Med*. 2018;379(5):464-473.
2. Diana W. Bianchi, M.D., R. Lamar Parker, M.D., Jeffrey Wentworth, M.D., Rajeevi Madankumar, M.D., Craig Saffer, M.D., Anita F. Das, Ph.D., Joseph A. Craig, M.D., Darya I. Chudova, Ph.D., Patricia L. Devers, M.S., C.G.C., Keith W. Jones, Ph.D., Kelly Oliver, B.S., Richard P. Rava, Ph.D., for the CARE Study Group. DNA sequencing versus Standard Prenatal Aneuploidy Screening.
3. Brison N, Neofytou M, Dehaspe L, Bayindir B, Van Den Bogaert K, Dardour L, Peeters H, Van Esch H, Van Buggenhout G, Vogels A, de Ravel T, Legius E, Devriendt K, Vermeesch JR. Predicting fetoplacental chromosomal mosaicism during non-invasive prenatal testing. *Prenat Diagn*. 2018 Mar;38(4):258-266.
4. Lefkowitz RB, Tynan JA, Liu T, Wu Y, Mazloom AR, Almasri E, Hogg G, Angkachatchai V, Zhao C, Grosu DS, McLennan G, Ehrich M. Clinical validation of a noninvasive prenatal test for genome-wide detection of fetal copy number variants. *Am J Obstet Gynecol*. 2016 Aug;215(2):227.e1-227.e16. doi: 10.1016/j.ajog.2016.02.030. Epub 2016 Feb 17.
5. Li R, Wan J, Zhang Y, Fu F, Ou Y, Jing X, Li J, Li D, Liao C. Detection of fetal copy number variants by non-invasive prenatal testing for common aneuploidies. *Ultrasound Obstet Gynecol*. 2016 Jan;47(1):53-7.
6. Brison N, Van Den Bogaert K, Dehaspe L, et al. Accuracy and clinical value of maternal incidental findings during noninvasive prenatal testing for fetal aneuploidies. *Genet Med*. 2017;19(3):306-313.
7. Amant F, Verheecke M, Wlodarska I, Dehaspe L, Brady P, Brison N, Van Den Bogaert K, Dierickx D, Vandecaveye V, Tousseyn T, Moerman P, Vanderstichele A, Vergote I, Neven P, Berteloot P, Putseys K, Danneels L, Vandenberghe P, Legius E, Vermeesch JR. Presymptomatic Identification of Cancers in Pregnant Women During Noninvasive Prenatal Testing. *JAMA Oncol*. 2015 Sep;1(6):814-9.
8. Bianchi DW, Chudova D, Sehnert AJ, Bhatt S, Murray K, Prosen TL, Garber JE, Wilkins-Haug L, Vora NL, Warsof S, Goldberg J, Ziainia T, Halks-Miller M. Noninvasive Prenatal Testing and Incidental Detection of Occult Maternal Malignancies. *JAMA*. 2015 Jul 14;314(2):162-9.

9. Darras BT, Urion DK, Ghosh PS. Dystrophinopathies. 2000 Sep 5 [updated 2018]. In: Adam MP, Ardinger HH, Pagon RA, et al. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2018. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1119/>
10. Landfeldt E, Lindgren P, Bell C, et al. The burden of Duchenne muscular dystrophy: an international, cross-sectional study. *Neurology* 2014;83(6):529-36.
11. Ferlini, Sewry C, Melis MA, Mateddu A, Muntoni F.- X-linked dilated cardiomyopathy and the dystrophin gene, *Neuromuscul Disord*. 1999; 9(5):339-346.
12. Ishizaki M, Kobayashi M, Adachi K, et al. Female dystrophinopathy: Review of current literature. *Neuromuscul Disord*. 2018;28(7):572-581.
13. Aartsma-Rus A, Van Deutekom JC, Fokkema IF, et al. Entries in the Leiden Duchenne muscular dystrophy mutation database: an overview of mutation types and paradoxical cases that confirm the reading-frame rule. *Muscle & Nerve*. 2006;34(2):135–144.
14. Tuffery-Giraud S, Bérourd C, Leturcq F, et al. Genotype-Phenotype Analysis in 2405 patients with a dystrophinopathy using the UMD-DMD Database. A model of Nationwide Knowledgebase. *Human Mutation*. 2009;30(6):934-945.
15. Belgian Advisory Committee on Bioethics. Opinion no. 66 - non-invasive prenatal testing (NIPT) .
https://www.health.belgium.be/sites/default/files/uploads/fields/fpshealth_theme_file/opinion_66_web_1.pdf
16. BeSHG. Belgian guidelines for managing incidental findings detected by NIPT. 2017.
<http://www.beshg.be/index.php?page=guidelines>
17. Bayindir B, Dehaspe L, Brison N, et al. Noninvasive prenatal testing using a novel analysis pipeline to screen for all autosomal fetal aneuploidies improves pregnancy management. *Eur J Hum Genet*. 2015;23(10):1286-1293.
18. Brady P, Brison N, Van Den Bogaert K, et al. Clinical implementation of NIPT - technical and biological challenges. *Clin Genet*. 2016;89(5):523-530.
19. Nguyen K, Putoux A, Busa T, et al. Incidental findings on array comparative genomic hybridization: detection of carrier females of dystrophinopathy without any family history. *Clin Genet*. 2015;87(5):488-491.
20. Aartsma-Rus A, Ginjaar IB, Bushby K. The importance of genetic diagnosis for Duchenne muscular dystrophy. *J Med Genet*. 2016;53(3):145-151.

21. Shen JJ, Zhang NR. Change-point model on nonhomogeneous Poisson processes with application in copy number profiling by next-generation DNA sequencing. *The Annals of Applied Statistics*. 2012;6(2):476-496.

FIGURE LEGENDS

Figure 1. Genomic representation of the Xp21.2-p21.1 region encompassing the *DMD* gene. UCSC genomic representation of the X chromosome encompassing the *DMD* gene at Xp21.2-p21.1 displaying the 16 maternal CNV's detected by NIPS. The duplications and deletions are represented as blue and red bars, respectively. CNV's, copy number variations; NIPS, non-invasive prenatal testing.

Figure 2. Example of one *DMD* CNV detected by NIPS. (A) Ideogram of chromosome X highlighting the Xp21.2-p21.1 region encompassing the *DMD* gene. (B) NIPS CNV result, represented as red bar, with the corresponding exons tracks included in the *DMD* CNV call. (C) Array-CGH result showing the refinement of *DMD* CNV size. Image of array-CGH data was extracted from CytoSure™ Interpret software. CNV, copy number variations; NIPS, non-invasive prenatal testing.

Figure 3. Flow-chart for managing maternal CNV's in the *DMD* gene detected by NIPS. Proposed strategy for the interpretation and reporting of CNV's in the *DMD* gene. CNV's, copy number variations; NIPS, non-invasive prenatal testing.

Table 1. Overview of the maternal CNV's detected by NIPS and classification steps.

ID	Type and Position		Evaluation by means of databases			Segregation	Reading frame	Final classification	Fetal sex
	Del/Dup	Exon	Leiden OVD	France UMD-DMD	Database Classification				
1	Del	51	DMD	DMD	DMD	son DMD	out-of-frame	DMD	M
2	Dup	45-53	78 BMD, 16DMD	48 BMD	likely BMD	nephew age 12 clinically NL CK 1454U/L father age 69 carrier NL	in-frame	mild BMD or likely benign	F
3	Del	10-16	1 BMD	1 BMD	likely BMD	father carrier mild BMD	in-frame	mild BMD	M
4	Del	14-21	1 DMD	absent	likely DMD	NI (maternal)	out-of-frame	likely DMD	M
5	Del	45-51	21 BMD, 12 DMD	6 BMD, 1 DMD	variable	NI (de novo)	in-frame	variable	M
6	Del	49	1 BMD, 7 DMD	absent	likely DMD	parents unavailable	in-frame	likely DMD	M
7	Del	50-55	2 DMD	absent	likely DMD	ND	in-frame	likely DMD	M
8	Del	48-49	47 BMD, 11 DMD	1 DCM, 30 BMD, 2 NL	variable	ND	in-frame	variable	M
9	Del	52-53	3 DMD, 1 BMD, 1 NL	absent	variable	ND	in-frame	variable	F
10	Dup	10-27	3 cases (no phenotype)	absent	VUS	father carrier NL	in-frame	likely benign	F
11	Dup	10-27	3 cases (no phenotype)	absent	VUS	NI (maternal)	in-frame	likely benign	F
12	Dup	10-27	3 cases (no phenotype)	absent	VUS	ND	in-frame	likely benign	M
13	Del	54-55	1 case (no phenotype)	absent	VUS	son age 3 clinically NL CK 881U/L father carrier NL	in-frame	likely benign	M
14	Dup	51-62	1 case (no phenotype)	absent	VUS	NI (maternal, son NL non carrier)	out-of-frame	likely DMD	F
15	Del	51-64	absent	absent	VUS	ND	in-frame	VUS	M
16	Del	56-59	absent	absent	VUS	ND	in-frame	VUS	M

Abbreviations : CNV's, copy number variations; ID, family identity; Del, deletion; Dup, duplication; DMD, Duchenne muscular dystrophy; BMD, Becker muscular dystrophy; DCM, dilated cardiomyopathy; VUS, variant of unknown significance; NI, not informative; ND, not determined; pheno, phenotype; NL, normal; M, male; F, female.

Table 2. Size and position of the CNV's: validation of chromosomal positions predicted by NIPS using chromosomal microarray analysis.

ID	Del/Dup	NIPS					Array-CGH Mother				
		Start	Stop	Size (kb)	Exon	Effect	Start	Stop	Size (kb)	Exon	Effect
1	Del	31,786,996	31,835,697	48,701	51	out-of-frame	31,780,126	31,821,260	41,134	51	out-of-frame
2	Dup	31,683,494	32,202,063	518,569	45-53	in-frame	31,605,940	32,118,239	512,299	45-53	in-frame
3	Del	32,580,346	32,681,449	101,103	10-16	in-frame	32,582,584	32,676,287	93,703	10-16	in-frame
4	Del	32,489,765	32,598,764	108,999	14-22	out-of-frame	32,500,113	32,592,155	92,042	14-21	out-of-frame
5	Del	31,750,628	32,008,715	258,087	45-51	in-frame	31,754,837	32,006,298	251,461	45-51	in-frame
6	Del	31,856,163	31,889,973	33,810	/	/	31,854,529	31,877,889	23,36	49	in-frame
7	Del	31,636,676	31,844,031	207,355	50-55	in-frame	31,642,684	31,838,230	195,546	50-55	in-frame
8	Del	31,855,215	31,935,595	80,380	48	in-frame	31,854,854	31,935,133	80,279	48-49	in-frame
9	Del	31,694,976	31,786,958	91,982	52-53	in-frame	31,696,408	31,776,368	79,96	52-53	in-frame
10	Dup	32,465,085	32,673,646	208,561	10-27	in-frame	32,465,982	32,664,559	198,577	10-27	in-frame
11	Dup	32,462,001	32,671,622	209,621	10-27	in-frame	32,465,982	32,664,559	198,577	10-27	in-frame
12	Dup	32,462,635	32,654,437	191,802	12-27	in-frame	32,465,982	32,664,559	198,577	10-27	in-frame
13	Del	31,602,998	31,695,378	92,380	54-55	in-frame	31,610,838	31,694,483	83,645	54-55	in-frame
14	Dup	31,323,743	31,815,934	492,191	51-62	out-of-frame	31,308,362	31,805,802	497,44	51-62	out-of-frame
15	Del	31,235,477	31,798,048	562,571	51-64	in-frame	31,240,957	31,792,296	551,339	51-64	in-frame
16	Del	31,480,299	31,639,279	158,980	56-59	in-frame	31,482,871	31,635,046	152,175	56-59	in-frame

Abbreviations: CNV's, copy number variations; NIPS, non-invasive prenatal testing; ID, family identity; Del, deletion; Dup, duplication

FIGURE 1




FIGURE 2




FIGURE 3

