

Parathyroid hormone and phosphate homeostasis in patients with Bartter and Gitelman syndrome: an international cross-sectional study

Maartje F.A. Verploegen¹, Rosa Vargas-Poussou², Stephen B. Walsh³, Harika Alpay⁴, Atefeh Amouzegar⁵, Gema Ariceta⁶, Bahriye Atmis⁷, Justine Bacchetta⁸, Peter Bány⁹, Stéphanie Baron¹⁰, Umut Selda Bayrakci¹¹, Hendrica Belge¹², Martine Besouw¹³, Anne Blanchard^{14,15}, Arend Bökenkamp¹⁶, Olivia Boyer¹⁷, Kathrin Burgmaier¹⁸, Lorenzo A. Calò¹⁹, Stéphane Decramer²⁰, Olivier Devuyst²¹, Maria van Dyck²², Pietro Manuel Ferraro^{23,24}, Marc Fila²⁵, Telma Francisco²⁶, Gian Marco Ghiggeri²⁷, Leire Gondra²⁸, Stefano Guarino²⁹, Nakysa Hooman³⁰, Ewout J. Hoorn³¹, Pascal Houillier^{15,32}, Konstantinos Kamperis³³, Jameela A. Kari³⁴, Martin Konrad³⁵, Elena Levtchenko³⁶, Laura Lucchetti³⁷, Francesca Lugani²⁷, Pierluigi Marzuillo²⁹, Barian Mohidin³, Thomas J. Neuhaus³⁸, Abdaldae Osman³⁹, Svetlana Papizh⁴⁰, Manel Perelló⁴¹, Maarten B Rookmaaker⁴², Valerie Said Conti⁴³, Fernando Santos⁴⁴, Ghalia Sawaf⁴⁵, Erkin Serdaroglu⁴⁶, Maria Szczepanska⁴⁷, Francesca Taroni⁴⁸, Rezan Topaloglu⁴⁹, Francesco Trepiccione⁵⁰, Enrico Vidal⁵¹, Elizabeth R. Wan³, Lutz Weber¹⁸, Zeynep Yuruk Yildirim⁵², Selçuk Yüksel⁵³, Galia Zlatanova⁵⁴, Detlef Bockenhauer^{3,39}, Francesco Emma³⁷, Tom Nijenhuis¹

1. Department of Nephrology, Radboud University Medical Center, Nijmegen, The Netherlands
2. Department of Genetics, Centre de Références MARHEA, Hôpital Européen Georges Pompidou Assistance Publique Hôpitaux de Paris, Paris, France
3. Department of Renal Medicine, University College London, London, UK
4. Division of Pediatric Nephrology, Faculty of Medicine, Marmara University, Istanbul, Turkey
5. Division of Nephrology, Department of Medicine, Firoozgar Clinical Research Development Center, Iran University of Medical Sciences, Tehran, Iran
6. Paediatric Nephrology Department, University Hospital Vall d'Hebron, Barcelona, Spain
7. Department of Pediatric Nephrology, Cukurova University Faculty of Medicine, Adana, Turkey
8. Department of Pediatric Nephrology, Rheumatology and Dermatology, Reference Center for Rare Renal Diseases, Reference Center for Rare Diseases of Calcium and Phosphate Metabolism. University Children's Hospital, Lyon, France
9. Division of Renal Medicine, Department of Clinical Science, Intervention and Technology, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden
10. Department of Physiology, Hôpital Européen Georges Pompidou Assistance Publique Hôpitaux de Paris, Paris, France

11. Department of Pediatric Nephrology, Ankara City Hospital, Üniversiteler Mahallesi Bilkent Caddesi, Çankaya/Ankara, Turkey
12. Center for Human Genetics, Institute of Pathology and Genetics, Gosselies, Belgium.
13. Department of Pediatric Nephrology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands
14. Clinical Research Center 1418, Centre de Références MARHEA, Hôpital Européen Georges Pompidou Assistance Publique Hôpitaux de Paris, Paris, France
15. Centre de Recherche des Cordeliers, INSERM, Sorbonne Université, Université de Paris, Paris, France
16. Department of Pediatric Nephrology, Emma Children's Hospital, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands
17. Department of Pediatric Nephrology, Necker Hospital, APHP, MARHEA, Imagine Institute, Paris University, Paris, France
18. Faculty of Medicine, University Hospital of Cologne, Children's and Adolescents' Hospital, Pediatric Nephrology, University of Cologne, Cologne, Germany
19. Department of Medicine (DIMED), Nephrology, Dialysis, Transplantation, University of Padova, Padova, Italy
20. Department of Pediatric Nephrology. Centre de Références SORARE, Toulouse University Hospital, Toulouse, France
21. Division of Nephrology, UCLouvain Medical School, Brussels, Belgium; Institute of Physiology, Mechanism of Inherited Kidney Disorders Group, University of Zurich, Zurich, Switzerland
22. Department of Pediatric Nephrology, University Hospital Leuven, Leuven, Belgium
23. U.O.S. Terapia Conservativa della Malattia Renale Cronica, U.O.C. Nefrologia, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy
24. Dipartimento Universitario di Medicina e Chirurgia Traslazionale, Università Cattolica del Sacro Cuore, Rome, Italy
25. Pediatric Nephrology, CHU Arnaud de Villeneuve, Montpellier University Hospital, Montpellier, France
26. Department of Pediatric Nephrology, Centro Hospitalar Universitário de Lisboa Central, Lisbon, Portugal
27. Division of Nephrology, Dialysis and Transplantation, Istituto Giannina Gaslini, IRCCS, Genoa, Italy.
28. Pediatric Nephrology Department, Cruces University Hospital, Barakaldo, Spain; Biocruces Bizkaia Health Research Institute, Barakaldo, Spain; Pediatric Department, University of the Basque Country UPV/EHU, Leioa, Spain


29. Department of Woman, Child and of General and Specialized Surgery, Università degli Studi della Campania "Luigi Vanvitelli", Naples, Italy
30. Ali-Asghar Clinical Research Development Center, Iran University of Medical Sciences, Tehran, Iran
31. Department of Internal Medicine, Division of Nephrology and Transplantation, Erasmus Medical Center, University Medical Center Rotterdam, Rotterdam, The Netherlands
32. Department of Physiology, Centre de Références MARHEA, Hôpital Européen Georges Pompidou Assistance Publique Hôpitaux de Paris, Paris, France
33. Department of Paediatrics and Adolescent Medicine, Aarhus University Hospital, Aarhus, Denmark
34. Pediatric Nephrology Center of Excellence and Pediatric Department, Faculty of Medicine, King Abdulaziz University, Jeddah, Kingdom of Saudi Arabia
35. Department of General Pediatrics, Pediatric Nephrology, University Hospital Münster, Germany
36. Department of Pediatric Nephrology & Department of Development and Regeneration, University Hospital Leuven, KU Leuven, Leuven, Belgium
37. Department of Pediatric Subspecialties, Division of Nephrology, Bambino Gesù Children's Hospital – IRCCS, Rome, Italy
38. Department of Pediatrics, Children's Hospital of Lucerne, Cantonal Hospital of Lucerne, Lucerne, Switzerland
39. Paediatric Nephrology Unit, Great Ormond Street Hospital, London, UK
40. Department of Hereditary and Acquired Kidney Diseases, Research and Clinical Institute for Pediatrics, Pirogov Russian National Research Medical University, Moscow, Russia
41. Nephrology Department, Hospital Universitari Vall d'Hebron, Universitat Autònoma de Barcelona, Barcelona, Spain
42. Department of Nephrology and Hypertension, University Medical Centre Utrecht, Utrecht, The Netherlands
43. Department of Pediatrics, Mater Dei Hospital Malta, Msida, Malta
44. Department of Pediatrics, Asturias Central University Hospital, University of Oviedo, Oviedo, Spain
45. Department of Pediatric Nephrology, Damascus Hospital, Damascus, Syria
46. Department of Pediatric Nephrology, Dr. Behçet Uz Children's Hospital, Izmir, Turkey
47. Department of Pediatrics, Faculty of Medical Sciences in Zabrze, SUM in Katowice, Poland
48. Pediatric Nephrology, Dialysis and Transplant Unit, Fondazione IRCCS Ca' Granda-Ospedale Maggiore Policlinico, Milan, Italy
49. Department of Pediatric Nephrology, Hacettepe University School of Medicine, Ankara, Turkey

50. Department of Translational Medical Sciences, University of Campania "L. Vanvitelli", Naples, Italy
51. Division of Pediatrics, Department of Medicine, University of Udine, Udine, Italy
52. Department of Pediatrics, Division of Pediatric Nephrology, Istanbul University Istanbul Faculty of Medicine, Istanbul, Turkey
53. Department of Pediatric Nephrology, Pamukkale University School of Medicine, Denizli, Turkey
54. University Children's Hospital Medical University, Sofia, Bulgaria

Correspondence to: Francesco Emma; E-mail: francesco.emma@opbg.net and Tom Nijenhuis; E-mail: Tom.Nijenhuis@radboudumc.nl

ORIGINAL UNEDITED MANUSCRIPT

GRAPHICAL ABSTRACT



ABSTRACT

Background. Small cohort studies have reported high parathyroid hormone (PTH) levels in patients with Bartter syndrome and lower serum phosphate levels have anecdotally been reported in patients with Gitelman syndrome. In this cross-sectional study, we assessed PTH and phosphate homeostasis in a large cohort of patients with salt-losing tubulopathies.

Methods. Clinical and laboratory data of 589 patients with Bartter and Gitelman syndrome were provided by members of the European Rare Kidney Diseases Reference Network (ERKNet) and the European Society for Pediatric Nephrology (ESPN).

Results. 285 patients with Bartter syndrome and 304 patients with Gitelman syndrome were included for analysis. Patients with Bartter syndrome type I & II had the highest median PTH level (7.5 pmol/l) and 56% had hyperparathyroidism (PTH >7.0 pmol/l). Serum calcium was slightly lower in Bartter syndrome type I & II patients with hyperparathyroidism (2.42 vs. 2.49 mmol/l; $p = 0.038$) compared to those with normal PTH levels and correlated inversely with PTH ($r_s -0.253$; $p = 0.009$). Serum phosphate and urinary phosphate excretion did not correlate with PTH. Overall, 22% of patients had low serum phosphate levels (phosphate – standard deviation score < -2), with the

highest prevalence in patients with Bartter syndrome type III (32%). Serum phosphate correlated with TmP/GFR (r_s 0.699; $p < 0.001$), suggesting renal phosphate wasting.

Conclusions. Hyperparathyroidism is frequent in patients with Bartter syndrome type I & II. Low serum phosphate is observed in a significant number of patients with Bartter and Gitelman syndrome and appears associated with renal phosphate wasting.

Keywords: Bartter syndrome, Gitelman syndrome, parathyroid hormone, phosphate, salt losing tubulopathies

KEY LEARNING POINTS

What is already known about this subject?

- High parathyroid hormone (PTH) levels have been observed in patients with Bartter syndrome in small cohort studies.
- In patients with Gitelman syndrome, low serum phosphate levels have anecdotally been reported.
- Data from large cohort studies on PTH and phosphate homeostasis in Bartter and Gitelman syndrome are lacking.

What this study adds?

- Up till now, this is the largest cohort of patients with salt-losing tubulopathies reported, comprising 285 patients with Bartter syndrome and 304 patients with Gitelman syndrome.
- Elevated PTH is frequently observed in patients with Bartter syndrome, especially in types I and II.
- In a significant number of patients with Bartter and Gitelman syndrome low serum phosphate was reported and is most likely related to a primarily PTH-independent renal phosphate leak.

What impact this may have on practice or on policy?

- This study provides unique insights into the prevalence of aberrant PTH and phosphate homeostasis in patients with Bartter and Gitelman syndrome.
- Further studies are necessary to understand the pathophysiology of these abnormalities and to demonstrate whether PTH and/or phosphate should be a treatment target in these salt-losing tubulopathies.

INTRODUCTION

Bartter and Gitelman syndrome are genetically distinct hereditary salt-losing tubulopathies that are characterized by hypokalemic metabolic alkalosis secondary to impaired reabsorption of sodium chloride (NaCl) in different segments of the renal tubule^{1,2}. Bartter syndrome is further classified into different subtypes. Bartter syndrome types I and II are caused by biallelic mutations in the *SLC12A1* gene encoding the NKCC2 sodium-potassium-chloride cotransporter and by biallelic mutations in the *KCNJ1* gene encoding the ROMK potassium channel, respectively, both of which are expressed in the thick ascending limb of the loop of Henle (TAL). Bartter syndrome type III results from biallelic mutations in the *CLCNKB* gene, which encodes the ClC-Kb chloride channel. Bartter syndrome type IV is caused by biallelic mutations in the *BSND* gene that encodes Barttin, or by molecular abnormalities affecting the two adjacent *CLCNKA* (encoding the chloride channel ClC-Ka) and *CLCNKB* genes. Barttin and ClC-Kb are expressed in the TAL and distal convoluted tubule (DCT); ClC-Ka is only expressed in the TAL. Finally, a transient X-linked form of Bartter syndrome (type V Bartter syndrome) has been described in male patients harboring mutations in the *MAGED2* gene, which alters the expression of the NKCC2 and NCC, the sodium chloride cotransporter. Gitelman syndrome is caused by biallelic mutations in the *SLC12A3* gene, encoding the NCC, of which the expression is restricted to the DCT.

While Bartter and Gitelman syndromes share similarities, including hypokalemia, metabolic alkalosis, hyperreninemic hyperaldosteronism, and low/normal blood pressure, differences in the tubular segments where NaCl reabsorption is compromised produce different clinical pictures². Patients with Gitelman syndrome usually present in late childhood or early adulthood. Typically, patients with Gitelman syndrome have hypomagnesemia and hypocalciuria. Conversely, type I and II Bartter syndrome patients usually present antenatally with polyhydramnios and postnatally with severe polyuria, hypercalciuria, and early onset nephrocalcinosis. Patients with type III Bartter syndrome have variable phenotypes that, in some cases, are difficult to distinguish from Gitelman syndrome. The large phenotypic spectrum of Bartter III patients is most likely explained by variable expression of the pathogenic gene product in the TAL and DCT, and by the possibility of partial compensation by other chloride channels, in particular ClC-Ka in the TAL.

In recent years, several investigators have observed abnormal levels of parathyroid hormone (PTH) and serum phosphate in patients with Bartter and Gitelman syndrome³⁻¹³. Patients with Bartter syndrome tend to have a high PTH³⁻⁹. However, the prevalence and pathophysiology of hyperparathyroidism in these Bartter patients are unclear. Furthermore, data have been in part contradictory since a tendency towards both hyper- and hypophosphatemia has been described in

Bartter syndrome^{3,5}. In patients with Gitelman syndrome, high PTH levels have only been described in patients that had parathyroid adenoma^{14,15}, while hypophosphatemia has been reported in several small studies¹⁰⁻¹³ and has been attributed to renal phosphate wasting^{12,13}.

To verify whether these observations from case reports and small series are prevalent in Bartter and Gitelman syndromes or just incidental findings, we analyzed PTH and phosphate levels in a very large cross-sectional cohort of patients with Bartter and Gitelman syndrome collected across Europe, including both pediatric and adult subjects.

MATERIALS AND METHODS

On behalf of the ERKNet Tubulopathies Working Group and of the ESPN Working Group on Inherited Renal Disorders, members of the European Rare Kidney Diseases Reference Network (ERKNet) and the European Society for Pediatric Nephrology (ESPN) were invited to provide retrospective clinical and laboratory data on patients with Bartter and Gitelman syndrome. Patients had either signed an informed consent form, allowing an anonymized transfer of clinical data within the ERKNet network, or consent was obtained according to local regulations and under the responsibility of local investigators. The data collection form included 38 items, of which some were mandatory, in particular those related to kidney function, PTH levels, and calcium-phosphate metabolism (see Supplementary data, Table S1). To minimize biases in estimating the prevalence of signs and symptoms, data collection was cross-sectional and centers were asked to provide only information corresponding to the last available patient visit. For example, if a given patient had transiently developed high PTH in the past, this information would not be captured to avoid overestimating the prevalence of hyperparathyroidism.

Laboratory data were converted to SI units. Outlier data were identified and verified with the referring clinician. Patients with incomplete data on creatinine or PTH were excluded from the analysis. To preclude measuring the effects of low GFR on phosphate and PTH levels, patients with decreased kidney function were excluded. Since estimated GFR (eGFR) data were not available, patients were excluded based on serum creatinine. To this end, we excluded all patients aged > 15 years with serum creatinine > 100 $\mu\text{mol/L}$ and patients aged ≤ 15 years with serum creatinine levels above an arbitrary value calculated with the following formula: $2.7 \times \text{age (years)} + 60$. Using these conservative cut-off values, we estimated by preliminary testing that most patients with $\text{eGFR} < 65 \text{ ml/min/1.73m}^2$ have been excluded. Overall, we excluded 76 patients with elevated creatinine (Figure 1). A data completeness list is provided in Supplementary data, Table S2.

Serum phosphate, alkaline phosphatase, and TmP/GFR are strongly influenced by age in the pediatric population. To be able to compare these variables in all patients, standard deviation scores (SDS) were calculated based on age-related reference values¹⁶⁻¹⁸. We used the following formula to calculate SDS for phosphate, alkaline phosphatase and TmP/GFR:

$$\text{SDS} = (\text{level} - \text{mean level for age}) / \text{SD for age}.$$

Each formula was based on published age-related reference values for phosphate, alkaline phosphatase and TmP/GFR. (see Supplementary Methods and Supplementary Data, Table S3 and S4, Figure S1 – S3).

Patients were stratified into 4 groups, based on the underlying genetic cause. These include patients with Bartter syndrome type I or II (n=107), Bartter syndrome type III (n=88), Bartter syndrome “not otherwise specified” (NOS) including patients with Bartter syndrome for which genetic data were not available (n=90), and Gitelman syndrome (n=304). Patients with Bartter syndrome type IV (n = 9), type V (n = 4), or Gitelman syndrome NOS (n= 32) were excluded from the final analysis because these tests were severely underpowered (Figure 1).

Data were analyzed using IBM SPSS Statistics Version 25. Categorical data are described as number and percent. Continuous data are presented as median with interquartile range (IQR). Data were analyzed using Chi-squared test, Fisher’s exact test, Mann-Whitney U test, and Kruskal Wallis test as appropriate. Spearman’s rank correlation was used for the analysis of the correlation between variables. Univariable and multivariable logistic regression were used to identify variables associated with elevated PTH; variables that reached a p-value < 0.10 at the univariable level were used in the multivariable model.

RESULTS

Overall characteristics

After excluding subjects with incomplete data, with serum creatinine above the threshold, or underpowered subgroups of patients, 589 patients from the original 770 patients collected were included for analysis (Figure 1).

Patient characteristics are in agreement with the known phenotypic characteristics of the diseases and are detailed in Table 1. The median age was 16.6 years [IQR 8.2 – 33.5]. Patients with Bartter syndrome were significantly younger than patients with Gitelman syndrome (median age 9.5 years

[3.5 – 16.3] versus 28.8 years [16.4 – 43.9] respectively; $p < 0.001$). Sex distribution was not significantly different in the four subgroups ($p = 0.113$). Overall, 52% of the patients were female. Nephrocalcinosis was reported more frequently in patients with Bartter syndrome type I & II compared to type III, with a prevalence of 89% and 18% respectively, and very rarely in patients with Gitelman syndrome, as expected. On average, patients with Bartter syndrome had higher urinary calcium/creatinine ratios than Gitelman patients ($p < 0.001$) and patients with Gitelman syndrome had lower serum magnesium level than patients with Bartter syndrome ($p < 0.001$). Indomethacin or other nonsteroidal anti-inflammatory drugs (NSAIDs) were prescribed more often to patients with Bartter syndrome than Gitelman syndrome ($p < 0.001$).

Elevated PTH levels were observed in a significant proportion of patients. Hyperparathyroidism (iPTH > 7.0 pmol/l) was present in 23% of patients (Supplementary data, Table S5). Patients with Bartter syndrome type I & II had the highest median iPTH levels (7.5 pmol/l [4.8 – 11.2]) (Table 1; Figure 2) and the prevalence of hyperparathyroidism was 56% in this group. Conversely, Gitelman patients had the lowest prevalence of hyperparathyroidism (7%) and even showed a 20% prevalence of hypoparathyroidism (iPTH < 2.0 pmol/l).

All disease subgroups had a negative median phosphate-SDS, demonstrating that median serum phosphate was lower in all subgroups than median age-specific serum phosphate levels (Table 1; Figure 2). Patients with Bartter syndrome type III had the lowest median phosphate-SDS of -1.34 [-2.31 – -0.15], which was significantly lower compared to phosphate-SDS in Bartter syndrome type I & II and Gitelman syndrome ($p = 0.032$ and $p = 0.011$ respectively; Figure 2). No differences were observed in tubular reabsorption of phosphate (TRP) ($p = 0.254$) between different disease subgroups. TmP/GFR-SDS was lowest in patients with Gitelman syndrome (-0.98 [-1.47 – -0.46]; Figure 2). This was significantly lower compared to Bartter syndrome type I & II ($p = 0.006$).

Median and IQR of serum calcium were within the normal range for all subgroups (Table 1; Figure 2). Serum calcium was significantly higher in patients with Bartter syndrome compared to Gitelman syndrome ($p < 0.001$). No significant differences in serum calcium were observed between Bartter syndrome subtypes.

Although data were available only in part of the entire cohort (Supplementary data, Table S2), 25-hydroxy vitamin D (25OH vitamin D) levels were lower in patients with Bartter syndrome compared to Gitelman syndrome ($p = 0.001$), and patients with Bartter syndrome used vitamin D supplements more often ($p < 0.001$; Table 1). No difference in serum calcium levels was found between patients who used vitamin D supplements and non-supplemented patients (data not shown).

Hyperparathyroidism and hypophosphatemia

A. Analysis of the entire cohort

The characteristics and laboratory data of patients according to the absence or presence of hyperparathyroidism are shown in Supplementary data, Table S6. Overall, 84% of patients with elevated iPTH had Bartter syndrome. Notably, 45% of the patients with hyperparathyroidism were patients with Bartter syndrome type I & II, while this disease subtype comprises only 18% of patients in the entire cohort. Several parameters, among which urinary calcium/creatinine ratio and serum magnesium, were significantly different when comparing patients with and without hyperparathyroidism (Supplementary data, Table S6) and iPTH was correlated with several parameters (Supplementary data, Table S7). However, these analyses appear markedly influenced by the unequal prevalence of hyperparathyroidism among different disease subgroups, and therefore, may reflect differences related to the underlying disease, rather than differences related to high iPTH. The same limitation applies to the multivariable logistic regression reported in Supplementary data, Table S8, and is underscored by the finding that the strongest predictor for hyperparathyroidism is the Bartter syndrome type I & II sub-class (OR: 3.20, 95% CI 1.68 - 6.08; $p < 0.001$). Other significant factors associated with hyperparathyroidism included higher serum magnesium (OR: 18.47 [2.67 - 127.93] per mmol/l, $p = 0.003$), lower serum calcium (OR: 0.03 [0.01 - 0.21] per mmol/l, $p < 0.001$), higher alkaline phosphatase-SDS (OR: 1.260 [1.11 - 1.431] per SDS, $p < 0.001$) and the use of NSAIDs (OR: 1.88 [1.08 - 3.28]; $p = 0.026$).

Hypophosphatemia (phosphate-SDS < -2) was observed in 22% of patients (Supplementary data, Table S9). The prevalence of hypophosphatemia was 19% in patients with Bartter syndrome type I & II, 32% in patients with Bartter syndrome type III, and 16% in patients with Gitelman syndrome. Patients with hypophosphatemia had higher serum calcium and lower TmP/GFR-SDS (Supplementary data, Table S10). Phosphate-SDS correlated with TRP ($r_s = 0.201$; $p = 0.004$) and with TmP/GFR-SDS ($r_s = 0.699$; $p < 0.001$) (Supplementary data, Table S11). To exclude an effect of age and/or the necessary modulation of serum phosphate to SDS score to be able to compare pediatric age groups and adults, we also evaluated serum phosphate only in adults. Hypophosphatemia was observed in 14.3% of adult patients (Supplementary data, Table S12). The prevalence of hypophosphatemia was 7% in patients with Bartter syndrome type I & II, 5% in patients with Bartter syndrome type III, and 15% in patients with Gitelman syndrome. However, patient numbers in the Bartter subtype groups were low.

We finally performed a sensitivity analysis, demonstrating that the overall results of Bartter patients were not significantly influenced by the inclusion of the Bartter syndrome NOS cohort (data not shown). Overall, patients included in the NOS subgroup tended to resemble more patients with type I & II Bartter syndrome, especially if they had elevated PTH.

B. Analysis of disease subgroups

When restricting the analysis to patients with Bartter syndrome type I & II, patients with hyperparathyroidism had lower serum calcium levels (2.42 versus 2.49 mmol/l; $p = 0.038$) and higher alkaline phosphatase-SDS (1.08 versus 0.27; $p = 0.010$) (Table 2). No differences were observed in serum creatinine, phosphate-SDS, 25OH vitamin D, TmP/GFR-SDS, urinary calcium/creatinine ratio, or the prevalence of nephrocalcinosis. No differences were observed in the prescription of NSAIDs or vitamin D supplements between patients with and without hyperparathyroidism.

In this cohort of patients with Bartter syndrome type I & II, a correlation was observed between iPTH and alkaline phosphatase-SDS ($r_s = 0.268$, $p = 0.015$) and an inverse correlation between iPTH and calcium ($r_s = -0.253$, $p = 0.009$) (Table 3; Figure 3). No correlation was observed between iPTH and phosphate-SDS, TmP/GFR-SDS, or urinary calcium/creatinine ratio. By multivariable logistic regression analysis, higher alkaline phosphatase-SDS (OR: 1.54 [1.09 – 2.19] per SDS, $p = 0.015$) was associated with high iPTH (Table 4). A trend was observed for lower serum calcium (OR: 0.025 [0.001 – 1.16] per SDS, $p = 0.059$), which did not reach statistical significance.

In the Gitelman cohort, we observed a significant correlation between iPTH and serum magnesium ($r_s = 0.191$, $p = 0.001$) and an inverse correlation between iPTH and 25OH vitamin D ($r_s = -0.207$, $p = 0.004$) (Supplementary data, Table S13).

In addition, as in the entire cohort, a significant correlation was observed between serum phosphate-SDS and TmP/GFR-SDS in patients with Bartter syndrome type III ($r_s = 0.744$; $p < 0.001$) and Gitelman syndrome ($r_s = 0.845$; $p < 0.001$) (Figure 4).

DISCUSSION

This cross-sectional study analyzed PTH and phosphate homeostasis in patients with Bartter and Gitelman syndrome. We observed high PTH (>7.0 pmol/L) in 56% of patients with type I & II disease. All disease subgroups, particularly patients with Bartter syndrome type III, had a tendency towards lower serum phosphate levels, and a significant proportion of patients had overt hypophosphatemia (phosphate-SDS < -2).

To date, hyperparathyroidism in these disorders has only been described in case reports and small series³⁻⁹. For example, Landau et al. observed high PTH levels in 10/14 patients with Bartter syndrome type II⁵. Additional small studies did not include genetic characterization^{3,6}, while others included only type I and II Bartter syndrome^{4,7-9}. Owing to the large number of patients that were recruited, we were able to analyze subgroups of patients, demonstrating that hyperparathyroidism developed primarily in patients with Bartter syndrome type I & II, and to a lesser degree, in patients with Bartter syndrome type III. A major confounding factor when analyzing PTH levels is kidney function. To prevent overestimating hyperparathyroidism, we excluded from the analysis all patients with elevated creatinine, using a very conservative threshold. In addition, we observed no correlation between PTH and creatinine levels. We did not observe age-dependent changes in PTH.

The aim of this study was to evaluate the prevalence of hyperparathyroidism and phosphate homeostasis abnormalities in patients with Bartter and Gitelman syndrome. To this end, we designed a cross-sectional study, favoring collection of data on a very large number of patients, over collecting longitudinal data on fewer patients that would have been more informative on pathophysiology, but would have limited our ability to estimate disease prevalence. However, some hypotheses on the pathophysiology and possible clinical implications can be proposed and discussed.

It has previously been hypothesized that hypercalciuria, a characteristic feature of patients with Bartter syndrome type I and type II, may contribute to increased PTH secretion, as a physiological response to maintain serum calcium within the normal range despite the excessive renal calcium loss^{4,5,7,19}. Indeed, in our whole cohort, a correlation between urinary calcium excretion and PTH was found. Calciuria is a crucial distinction between Bartter and Gitelman syndrome and our observational study cannot distinguish between a causative role of calciuria for hyperparathyroidism or a simple association because of other differences between Bartter and Gitelman syndrome. As expected, hypercalciuria was a consistent feature in patients with Bartter syndrome type I & II, but no correlation between urinary calcium excretion and PTH levels were observed in this subgroup. However, the cross-sectional design of the study and significant heterogeneity of patient treatments, do not allow excluding a possible role of hypercalciuria in promoting PTH secretion. Patients with

hyperparathyroidism in this subgroup had on average lower serum calcium levels, and, as expected, serum calcium was negatively correlated with PTH. These findings are concordant with the reports by Landau et al. and Bettinelli et al.^{3,5}. The cause of this lower serum calcium and whether it is sufficient to trigger PTH secretion in these patients remains undetermined.

In our study, high PTH levels in type I & II Bartter syndrome patients were not associated with lower serum phosphate or with higher urinary excretion of phosphate, as also observed by Landau et al.⁵. Taken together, these findings raise the possibility of renal hyporesponsiveness to PTH, which could contribute to PTH secretion. Yet, with renal hyporesponsiveness increased serum phosphate could be expected, which we did not observe.

Hyperparathyroidism in Bartter syndrome could also be secondary to prostaglandin secretion^{4,7}. Patients with type I & II disease produce large amounts of prostaglandins due to decreased intracellular NaCl concentration in the macula densa caused by impaired salt reabsorption via the NKCC2 pathway^{6,20,21}. Treatment of Bartter syndrome patients with NSAIDs reduces renal calcium excretion^{19,20,22,23}, and in one report, reduces PTH levels¹⁹. It is unclear in this latter study, if NSAIDs directly inhibit the parathyroid glands, or reduce PTH secretion indirectly, by inhibiting renal calcium excretion. In our study, we did not observe an association between NSAID prescription and PTH levels, nor between NSAID prescription and urinary calcium excretion.

It has also been hypothesized that higher PTH secretion in Bartter syndrome may be secondary to hyperaldosteronism⁵. Parathyroid glands express mineralocorticoid receptors allowing stimulation of PTH secretion by aldosterone²⁴. Accordingly, mild hyperparathyroidism has been reported in patients with primary hyperaldosteronism²⁵ and PTH levels in these patients decrease after adrenalectomy²⁶. This hypothesis seems unlikely in patients with salt-losing tubulopathies since patients with Gitelman syndrome, who typically have very high aldosterone levels, do not develop hyperparathyroidism. In addition, PTH levels in our cohort were not associated with treatment with aldosterone antagonists.

Finally, it has been suggested that hypokalemia could cause hyperparathyroidism and that potassium supplementation could normalize PTH²⁷. Here again, we did not observe an inverse correlation between serum potassium and PTH. Moreover, Bartter and Gitelman syndromes are both characterized by hypokalemia, whereas we observed hyperparathyroidism primarily in patients with Bartter syndrome.

Another noteworthy finding in our cohort was that 20% of patients with Gitelman syndrome had hypoparathyroidism. The calcium-sensing receptor signaling is known to be dependent on extracellular magnesium levels. Unlike calcium, PTH secretion is inhibited by supraphysiologically

high magnesium levels via stimulation of the calcium-sensing receptor, but also by low magnesium concentrations²⁸. This latter response is suggested to result from intracellular disinhibition of G-protein signaling downstream from the calcium-sensing receptor, precluding PTH release²⁹. Accordingly, decreased PTH secretion and decreased responsiveness of PTH to ionized calcium have been suggested in small cohorts of patients with Gitelman syndrome^{30,31,32}. In agreement with these data, we observed a significant correlation between PTH and serum magnesium in patients with Gitelman syndrome, suggesting that low serum magnesium might be, at least in part, responsible for hypoparathyroidism.

Hypophosphatemia was common in all disease subgroups, in particular in Bartter syndrome type III with a prevalence of 32%. The prevalence of hypophosphatemia in Bartter syndrome type III patients decreased in adulthood to 5%. In Gitelman syndrome, the prevalence of hypophosphatemia was not age-dependent. Low serum phosphate levels were associated with lower TmP/GFR in all subgroups, indicating that hypophosphatemia was most likely related to renal phosphate wasting. However, renal hypophosphatemia did not appear to be driven by hyperparathyroidism, since PTH did not correlate with phosphate-SDS or TmP/GFR-SDS. Furthermore, patients with Gitelman syndrome did not have high PTH levels, while they often had hypophosphatemia and renal phosphate wasting. The exact mechanism(s) of hypophosphatemia in these patients remains unclear. Notably, low serum phosphate levels have already been observed in anecdotal reports on Gitelman syndrome¹⁰⁻¹³, including a possible link with lower renal tubular phosphate reabsorption^{12,13}. Interestingly, it has been hypothesized that chronic hypokalemia can cause proximal tubular cell injury, resulting in a phosphate leak³³. However, serum potassium and serum phosphate were not correlated in our cohort. Furthermore, FGF23 could play a role. Interestingly, a recent study demonstrated that in the NCC knockout mouse model for Gitelman syndrome, FGF23 levels were increased³⁴. In contrast to our Gitelman cohort, these mice also showed increased PTH levels which, however, did not alter fractional phosphate excretion nor serum phosphate levels.

We also observed significantly higher serum calcium levels in patients with Bartter syndrome, compared to patients with Gitelman syndrome. Theoretically, since patients with Gitelman syndrome have hypocalciuria, they are expected to have higher serum calcium levels than patients with Bartter syndrome that have hypercalciuria. Furthermore, compared to healthy controls, patients with Gitelman syndrome have higher serum calcium levels¹³ and patients with Bartter syndrome tend to have lower serum calcium levels³. Vitamin D supplementation may have influenced our results. Although supplements were prescribed more frequently to patients with Bartter syndrome, patients treated with vitamin D did not have significantly higher serum calcium levels, compared to non-supplemented patients.

Whether in clinical practice hyperparathyroidism and hypophosphatemia require further diagnostic evaluation and treatments is unclear. In our cohort, high PTH was associated with higher alkaline phosphatase levels. This is likely due to increased bone turn over. Reduced bone mineral density has previously been observed in patients with Bartter syndrome^{35,36}. It may therefore be appropriate that Bartter syndrome patients undergo regular assessment of their bone density to detect incipient osteopenia or osteoporosis, particularly if they have high PTH. Since vitamin D deficiency stimulates PTH secretion, supplements could be prescribed in patients with low vitamin D levels, keeping in mind that they may worsen hypercalciuria in patients with Bartter syndrome. Of note, 25-OH vitamin D concentrations are usually normal in patients with Bartter syndrome²³, while high levels of 1,25-OH vitamin D have been reported^{23,36}. The use of calcimimetics in patients with hyperparathyroidism associated with Bartter syndrome has been described in five patients^{7,8}. In four of them, PTH levels normalized. However, all had high-normal or even high serum calcium levels. In theory, normocalcemic patients with Bartter syndrome could be at increased risk of developing symptomatic hypocalcemia if treated with calcimimetics, since they have metabolic alkalosis, which decreases the proportion of ionized calcium. It is unknown whether hypophosphatemia, especially if associated with bone changes, should be treated with phosphate supplementation.

The main limitations of our study relate to the cross-sectional design and the relatively large group of Bartter NOS patients in the analysis. The cross-sectional design allowed us to assemble data on a very large number of patients to estimate the prevalence of PTH and phosphate abnormalities in these diseases. However, longitudinal follow-up of patients would provide additional information about the course of PTH and other variables throughout the years, which could have helped identify possible causal effects. Future studies should collect data at the initial presentation, thus pre-treatment, and should also evaluate follow-up data of patients. An important opportunity to prospectively capture such data is by adding PTH and phosphate as variables to the ERKNet patient registry (ERKReg). The second main limitation is the relatively large group of patients with Bartter syndrome NOS, who lacked a genetic diagnosis and were diagnosed based on their phenotype. This subgroup was most likely composed of patients with mixed Bartter syndrome subtypes since their characteristics are intermediate between the type I & II and the type III subgroups. Sensitivity analyses performed removing the NOS subgroup showed that the inclusion of these patients did not modify our results.

Other limitations of our study are related to its retrospective, cross-sectional, and multicentric design. The number of variables that could be reliably captured in the database was therefore limited. For example, active vitamin D, ionized calcium, cyclic AMP, FGF-23, aldosterone levels, and data on bone metabolism were not available for most patients but could have been very instrumental in understanding the physiopathology of our findings. Treatments were decided by local

physicians according to local protocols. Laboratory values were measured in different laboratories, which may have used slightly different methods and reference values. These latter aspects were probably well compensated for by the large size of our cohort. Since the age range of our patients spanned from small children to late adulthood, some values had to be normalized for age using reference data from the literature. This may have introduced some distortions in our data.

In conclusion, this cross-sectional study demonstrates that PTH is frequently elevated in patients with Bartter syndrome, especially in those with type I & II disease. The cause of hyperparathyroidism in this population remains unclear. Hypophosphatemia was observed in a significant proportion of patients with Bartter and Gitelman syndrome and appeared to be primarily related to a PTH-independent renal phosphate leak. Further studies are needed to understand better the pathophysiology and the clinical relevance of these observations.

ACKNOWLEDGEMENTS

This project has been supported by the European Reference Network for Rare Kidney Diseases (ERKNet), which is partly co-funded by the European Union within the framework of the Third Health Programme 'ERN-2016-Framework Partnership Agreement 2017-2021'. This work is generated within the European Society for Pediatric Nephrology working group on inherited renal disorders. This work was supported by an Innovation Grant 19OI06 from the Dutch Kidney Foundation (to T.N.).

CONFLICT OF INTEREST STATEMENT

The authors have no conflicts of interest to declare.

DATA AVAILABILITY STATEMENT

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

1. Seyberth HW, Weber S, Komhoff M. Bartter's and Gitelman's syndrome. *Curr Opin Pediatr.* 2017;29(2):179-86.
2. Besouw MTP, Kleta R, Bockenhauer D. Bartter and Gitelman syndromes: Questions of class. *Pediatr Nephrol.* 2020;35(10):1815-24.
3. Bettinelli A, Vigano C, Provero MC, et al. Phosphate homeostasis in Bartter syndrome: a case-control study. *Pediatr Nephrol.* 2014;29(11):2133-8.
4. Gross I, Siedner-Weintraub Y, Simckes A, Gillis D. Antenatal Bartter syndrome presenting as hyperparathyroidism with hypercalcemia and hypercalciuria: a case report and review. *J Pediatr Endocrinol Metab.* 2015;28(7-8):943-6.
5. Landau D, Gurevich E, Sinai-Treiman L, Shalev H. Accentuated hyperparathyroidism in type II Bartter syndrome. *Pediatr Nephrol.* 2016;31(7):1085-90.
6. Leonhardt A, Timmermanns G, Roth B, Seyberth HW. Calcium homeostasis and hypercalciuria in hyperprostaglandin E syndrome. *J Pediatr.* 1992;120(4 Pt 1):546-54.
7. Li D, Tian L, Hou C, Kim CE, Hakonarson H, Levine MA. Association of Mutations in SLC12A1 Encoding the NKCC2 Cotransporter With Neonatal Primary Hyperparathyroidism. *J Clin Endocrinol Metab.* 2016;101(5):2196-200.
8. Wongsangsak S, Vidmar AP, Addala A, et al. A novel SLC12A1 gene mutation associated with hyperparathyroidism, hypercalcemia, nephrogenic diabetes insipidus, and nephrocalcinosis in four patients. *Bone.* 2017;97:121-5.
9. Zuo J, Guo W, Wang S, et al. Eight novel KCNJ1 variants and parathyroid hormone overaction or resistance in 5 probands with Bartter syndrome type 2. *Clin Chim Acta.* 2020;511:248-54.
10. Akhtar N, Hafeez F. A rare case of Gitelman's syndrome with hypophosphatemia. *J Coll Physicians Surg Pak.* 2009;19(4):257-9.
11. Ali A, Masood Q, Yaqub S, Kashif W. A case of Gitelman syndrome with severe hyponatraemia and hypophosphataemia. *Singapore Med J.* 2013;54(1):e18-20.
12. Katopodis K, Elisaf M, Siamopoulos KC. Hypophosphataemia in a patient with Gitelman's syndrome. *Nephrol Dial Transplant.* 1996;11(10):2090-2.
13. Vigano C, Amoruso C, Barretta F, et al. Renal phosphate handling in Gitelman syndrome--the results of a case-control study. *Pediatr Nephrol.* 2013;28(1):65-70.

14. Rego T, Fonseca F, Cerqueira R, Agapito A. Gitelman syndrome and primary hyperparathyroidism: a rare association. *BMJ Case Rep.* 2018;2018.
15. Wen YK. An unusual case of Gitelman's syndrome with hypercalcemia. *Ren Fail.* 2012;34(2):241-3.
16. Ardeshirpour L, Cole DE, Carpenter TO. Evaluation of bone and mineral disorders. *Pediatr Endocrinol Rev.* 2007;5 Suppl 1:584-98.
17. KDOQI Work Group. KDOQI Clinical Practice Guideline for Nutrition in Children with CKD: 2008 update. Executive summary. *Am J Kidney Dis.* 2009;53(3 Suppl 2):S11-104.
18. Zierk J, Arzideh F, Haeckel R, et al. Pediatric reference intervals for alkaline phosphatase. *Clin Chem Lab Med.* 2017;55(1):102-10.
19. Gasongo G, Greenbaum LA, Niel O, et al. Effect of nonsteroidal anti-inflammatory drugs in children with Bartter syndrome. *Pediatr Nephrol.* 2019;34(4):679-84.
20. Reinalter SC, Jeck N, Brochhausen C, et al. Role of cyclooxygenase-2 in hyperprostaglandin E syndrome/antenatal Bartter syndrome. *Kidney Int.* 2002;62(1):253-60.
21. Seyberth HW, Schlingmann KP. Bartter- and Gitelman-like syndromes: salt-losing tubulopathies with loop or DCT defects. *Pediatric Nephrology.* 2011;26(10):1789-802.
22. Reinalter SC, Grone HJ, Konrad M, Seyberth HW, Klaus G. Evaluation of long-term treatment with indomethacin in hereditary hypokalemic salt-losing tubulopathies. *J Pediatr.* 2001;139(3):398-406.
23. Restrepo de Rovetto C, Welch TR, Hug G, Clark KE, Bergstrom W. Hypercalciuria with Bartter syndrome: evidence for an abnormality of vitamin D metabolism. *J Pediatr.* 1989;115(3):397-404.
24. Maniero C, Fassina A, Guzzardo V, et al. Primary hyperparathyroidism with concurrent primary aldosteronism. *Hypertension.* 2011;58(3):341-6.
25. Tomaschitz A, Ritz E, Pieske B, et al. Aldosterone and parathyroid hormone interactions as mediators of metabolic and cardiovascular disease. *Metabolism.* 2014;63(1):20-31.
26. Lenzini L, Prisco S, Vanderriele PE, et al. PTH Modulation by Aldosterone and Angiotensin II is Blunted in Hyperaldosteronism and Rescued by Adrenalectomy. *J Clin Endocrinol Metab.* 2019;104(9):3726-34.
27. Bando Y, Miyakoshi H, Nagaoka T, Ohsawa K, Kobayashi K. [A case of pseudohypoparathyroidism (PHP) type II associated with Bartter's syndrome--restoration of phosphaturic response to

- parathyroid hormone (PTH) by treatment for hypopotassemia]. *Nihon Naibunpi Gakkai Zasshi*. 1992;68(7):676-87. (Article in Japanese; abstract in English)
28. Brown EM, Chen CJ. Calcium, magnesium and the control of PTH secretion. *Bone Miner*. 1989;5(3):249-57.
 29. Quitterer U, Hoffmann M, Freichel M, Lohse MJ. Paradoxical block of parathormone secretion is mediated by increased activity of G alpha subunits. *J Biol Chem*. 2001;276(9):6763-9.
 30. Bianchetti MG, Bettinelli A, Casez JP, et al. Evidence for disturbed regulation of calciotropic hormone metabolism in Gitelman syndrome. *J Clin Endocrinol Metab*. 1995;80(1):224-8.
 31. Nicolet-Barousse L, Blanchard A, Roux C, et al. Inactivation of the Na-Cl co-transporter (NCC) gene is associated with high BMD through both renal and bone mechanisms: analysis of patients with Gitelman syndrome and Ncc null mice. *J Bone Miner Res*. 2005;20(5):799-808.
 32. Blanchard A, Vallet M, Dubourg L, et al. Resistance to Insulin in Patients with Gitelman Syndrome and a Subtle Intermediate Phenotype in Heterozygous Carriers: A Cross-Sectional Study. *J Am Soc Nephrol*. 2019;30(8):1534-45.
 33. Bouchireb K, Boyer O, Mansour-Hendili L, et al. Fanconi syndrome and severe polyuria: an uncommon clinicobiological presentation of a Gitelman syndrome. *BMC Pediatr*. 2014;14:201.
 34. Pathare G, Anderegg M, Albano G, Lang F, Fuster DG. Elevated FGF23 Levels in Mice Lacking the Thiazide-Sensitive NaCl cotransporter (NCC). *Sci Rep*. 2018;8(1):3590.
 35. Rodriguez-Soriano J, Vallo A, Aguirre M. Bone mineral density and bone turnover in patients with Bartter syndrome. *Pediatr Nephrol*. 2005;20(8):1120-5.
 36. Shoemaker L, Welch TR, Bergstrom W, Abrams SA, Yergey AL, Vieira N. Calcium kinetics in the hyperprostaglandin E syndrome. *Pediatr Res*. 1993;33(1):92-6.

Table 1. Patient characteristics

	Bartter syndrome type I & II	Bartter syndrome type III	Bartter syndrome NOS	Gitelman syndrome	Total
Number of patients	107	88	90	304	589
<i>Characteristics</i>					
Male : Female ratio, <i>n</i> (%)	51 : 56 (48 : 52)	50 : 38 (57 : 43)	48 : 42 (53 : 47)	133 : 171 (44 : 56)	282 : 307 (48 : 52)
Age (years)	9.0 [3.5 – 16.0]	10.4 [3.3 – 17.2]	9.6 [3.6 – 16.3]	28.8 [16.4 – 43.9]	16.6 [8.2 – 33.5]
Nephrocalcinosis, <i>n</i> (%)	90/101 (89)	15/85 (18)	46/82 (56)	4/257 (2)	155/525 (30)
<i>Laboratory results</i>					
iPTH (pmol/L)	7.5 [4.8 – 11.2]	4.4 [3.0 – 6.7]	5.1 [2.9 – 8.3]	3.1 [2.3 – 4.0]	3.8 [2.5 – 6.6]
Creatinine (umol/L)	55 [37 – 76]	43 [31 – 62]	48 [34 – 66]	58 [47 – 71]	55 [41 – 70]
Sodium (mmol/L)	140 [138 – 142]	139 [137 – 141]	139 [137 – 140]	139 [138 – 141]	139 [138 – 141]
Potassium (mmol/L)	3.6 [3.3 – 3.8]	3.2 [2.8 – 3.6]	3.4 [3.0 – 3.9]	3.0 [2.7 – 3.4]	3.2 [2.8 – 3.7]
Chloride (mmol/L)	99 [97 – 101]	97 [92 – 100]	96 [93 – 101]	98 [95 – 100]	98 [95 – 100]
Bicarbonate (mmol/L)	26.2 [24.3 – 28.1]	29.0 [26.3 – 32.0]	28.4 [27.0 – 30.7]	29.0 [27.0 – 31.0]	28.3 [26.3 – 31.0]
Phosphate (mmol/L)	1.42 [1.21 – 1.63]	1.28 [1.09 – 1.44]	1.30 [1.12 – 1.53]	1.07 [0.92 – 1.25]	1.21 [0.99 – 1.42]
Phosphate-SDS	-0.67 [-1.76 – 0.43]	-1.34 [-2.31 – -0.15]	-1.32 [-2.32 – 0.05]	-0.77 [-1.51 – 0.29]	-0.91 [-1.82 – 0.18]
Calcium (mmol/L)	2.44 [2.37 – 2.53]	2.48 [2.40 – 2.58]	2.50 [2.38 – 2.60]	2.41 [2.31 – 2.50]	2.43 [2.34 – 2.53]
Magnesium (mmol/L)	0.89 [0.80 – 0.97]	0.77 [0.68 – 0.85]	0.81 [0.75 – 0.95]	0.63 [0.56 – 0.71]	0.71 [0.61 – 0.84]
Uric acid (μmol/l)	315 [246 – 405]	220 [162 – 319]	286 [190 – 409]	250 [190 – 321]	262 [190 – 350]
Alkaline phosphatase (IU/L)	227 [161 – 287]	212 [114 – 274]	230 [160 – 303]	66 [51 – 136]	148 [62 – 245]
Alkaline phosphatase-SDS	0.79 [-0.05 – 1.52]	0.21 [-0.50 – 0.99]	0.65 [-0.36 – 1.89]	-1.00 [-1.72 – 0.00]	-0.13 [-1.18 – 0.93]
Total protein (g/L)	73 [70 – 77]	73 [69 – 76]	75 [71 – 78]	73 [69 – 76]	73 [70 – 77]
Albumin (g/L)	46 [43 – 49]	45 [43 – 48]	46 [43 – 49]	46 [44 – 49]	46 [43 – 49]
25OH vitamin D (nmol/L)	39 [21 – 68]	40 [23 – 82]	43 [29 – 64]	61 [31 – 85]	51 [28 – 80]
Urinary calcium/creatinine (mmol/mmol)	1.55 [0.94 – 2.62]	0.33 [0.14 – 0.71]	0.55 [0.23 – 1.14]	0.08 [0.04 – 0.17]	0.21 [0.07 – 0.75]
TRP	0.90 [0.81 – 0.92]	0.90 [0.86 – 0.93]	0.88 [0.82 – 0.95]	0.91 [0.86 – 0.94]	0.90 [0.85 – 0.94]

TmP/GFR (mmol/l)	1.41 [1.10 – 1.48]	1.17 [1.02 – 1.33]	1.11 [0.84 – 1.35]	0.95 [0.82 – 1.10]	1.04 [0.84 – 1.23]
TmP/GFR-SDS	0.23 [-0.46 – 0.78]	-0.62 [-1.48 – 0.44]	-0.66 [-2.03 – 0.14]	-0.98 [-1.47 – -0.46]	-0.86 [-1.54 – -0.02]
<i>Treatment</i>					
NSAIDs, <i>n</i> (%)	69 (65)	52 (59)	55 (61)	26 (9)	202 (34)
Indomethacin, <i>n</i> (%)	53 (50)	48 (55)	54 (60)	26 (9)	181 (31)
Ibuprophen, <i>n</i> (%)	16 (15)	4 (5)	2 (2)	0 (0)	22 (4)
Other NSAIDs, <i>n</i> (%)	5 (5)	7 (8)	4 (4)	1 (0.3)	17 (3)
Potassium-sparing diuretics, <i>n</i> (%)	2 (2)	12 (14)	4 (4)	63 (21)	81 (14)
Amilorid, <i>n</i> (%)	2 (2)	12 (14)	4 (4)	58 (19)	76 (13)
Triamterene, <i>n</i> (%)	0 (0)	0 (0)	0 (0)	5 (2)	5 (1)
Aldosterone antagonists, <i>n</i> (%)	5 (5)	17 (19)	23 (26)	54 (18)	99 (17)
Eplerenone, <i>n</i> (%)	1 (1)	2 (2)	6 (7)	15 (5)	24 (4)
Spironolactone, <i>n</i> (%)	4 (4)	15 (17)	16 (18)	37 (12)	72 (12)
Canrenone, <i>n</i> (%)	0 (0)	0 (0)	1 (1)	2 (1)	3 (1)
ACE inhibitors/ARBs, <i>n</i> (%)	4 (4)	5 (6)	0 (0)	6 (2)	15 (3)
Hydrochlorothiazide, <i>n</i> (%)	3 (3)	0 (0)	6 (7)	0 (0)	9 (2)
Potassium supplements, <i>n</i> (%)	71 (66)	83 (94)	76 (84)	280 (92)	510 (87)
Sodium supplements, <i>n</i> (%)	28 (26)	44 (50)	21 (23)	43 (14)	136 (23)
Magnesium supplements, <i>n</i> (%)	11 (10)	21 (24)	18 (20)	226 (74)	276 (47)
Phosphate supplements, <i>n</i> (%)	0 (0)	2 (2)	3 (3)	1 (0.3)	6 (1)
Proton pump inhibitors, <i>n</i> (%)	20 (19)	24 (27)	14 (16)	39 (13)	97 (17)
Other gastric protectors, <i>n</i> (%)	12 (11)	10 (11)	8 (9)	11 (4)	41 (7)
Oral contraceptive, <i>n</i> (%)	1 (1)	1 (1)	1 (1)	20 (7)	23 (4)
Vitamin D supplements, <i>n</i> (%)	41 (38)	20 (23)	17 (19)	26 (9)	104 (18)

iPTH = intact parathyroid hormone; phosphate-SDS = age-related phosphate standard deviation score; alkaline phosphatase-SDS = age-related alkaline phosphatase standard deviation score; 25OH vitamin D = 25-hydroxy vitamin D; TRP = tubular reabsorption of phosphate; TmP/GFR = ratio of tubular maximum reabsorption of phosphate to GFR; TmP/GFR-SDS: age-related TmP/GFR standard deviation score; NSAIDs = nonsteroidal anti-inflammatory drugs; ACE inhibitors = angiotensin converting enzyme inhibitors; ARBs = angiotensin II receptor blockers

Categorical values are presented as number and percent given in parentheses. Continuous values are presented as median with interquartile range (IQR) given in square brackets

Nota bene: some patients are using more than one NSAID according to the provided data

Table 2. Characteristics of patients with Bartter syndrome type I & II with and without hyperparathyroidism

Variable	Hyperparathyroidism	No hyperparathyroidism	p-value
Number of patients	60 (56%)	47 (44%)	
<i>Characteristics</i>			
Age (years)	9.6 [3.8 – 16.3]	8.9 [2.7 – 15.5]	0.476
Sex (male)	27 (45)	24 (51)	0.563
Nephrocalcinosis, <i>n</i> (%)	49/57 (86)	41/44 (93)	0.340
<i>Laboratory results</i>			
iPTH (pmol/l)	10.7 [8.6 – 14.0]	4.5 [3.2 – 5.6]	NT
Creatinine (μmol/l)	54 [36 – 71]	55 [37 – 77]	0.863
Sodium (mmol/l)	140 [138 – 142]	140 [138 – 143]	0.838
Potassium (mmol/l)	3.6 [3.2 – 3.8]	3.6 [3.3 – 3.9]	0.538
Chloride (mmol/l)	99 [97 – 102]	99 [97 – 101]	0.390
Bicarbonate (mmol/l)	26.0 [24.0 – 28.0]	26.4 [24.8 – 28.8]	0.414
Phosphate-SDS	-0.66 [-1.56 – 0.47]	-0.82 [-1.94 – 0.21]	0.418
Calcium (mmol/l)	2.42 [2.34 – 2.51]	2.49 [2.39 – 2.60]	0.038
Magnesium (mmol/l)	0.90 [0.81 – 0.98]	0.89 [0.79 – 0.98]	0.651
Alkaline phosphatase-SDS	1.08 [0.17 – 1.89]	0.27 [-0.28 – 1.09]	0.010
Uric acid (μmol/l)	315 [240 – 403]	327 [262 – 421]	0.697
Total protein (g/l)	74 [72 – 78]	72 [68 – 75]	0.147
Albumin (g/l)	46 [43 – 48]	47 [43 – 50]	0.321
25OH vitamin D (nmol/l)	40 [24.5 – 62.9]	35 [15 – 94]	0.761
Urinary calcium/creatinine (mmol/mmol)	1.46 [0.91 – 2.42]	1.73 [1.02 – 2.76]	0.452
TRP	0.87 [0.79 – 0.93]	0.91 [0.81 – 0.93]	0.673
TmP/GFR-SDS	-0.13 [-1.26 – 0.67]	0.56 [-0.36 – 0.90]	0.481
<i>Treatment</i>			
Indomethacin or other NSAID, <i>n</i> (%)	37 (62)	32 (68)	0.545
Potassium-sparing diuretics, <i>n</i> (%)	2 (3)	0 (0)	0.503
Aldosterone antagonists, <i>n</i> (%)	2 (3)	3 (6)	0.652
ACE inhibitors/ARBs, <i>n</i> (%)	2 (3)	2 (4)	1.000
Hydrochlorothiazide, <i>n</i> (%)	2 (3)	1 (2)	1.000
Potassium supplements, <i>n</i> (%)	42 (70)	29 (62)	0.413
Sodium supplements, <i>n</i> (%)	17 (28)	11 (23)	0.660
Magnesium supplements, <i>n</i> (%)	9 (15)	2 (4)	0.108
Phosphate supplements, <i>n</i> (%)	0 (0)	0 (0)	N/A
Proton pump inhibitors, <i>n</i> (%)	10 (17)	10 (21)	0.621
Other gastric protectors, <i>n</i> (%)	5 (8)	7 (15)	0.360
Oral contraceptives, <i>n</i> (%)	1 (2)	0 (0)	1.000
Vitamin D supplements, <i>n</i> (%)	26 (43)	15 (32)	0.238

iPTH = intact parathyroid hormone; phosphate-SDS = age-related phosphate standard deviation score; alkaline phosphatase-SDS = age-related alkaline phosphatase standard deviation score; 25OH vitamin D = 25-hydroxy vitamin D; TRP = tubular reabsorption of phosphate; TmP/GFR-SDS: age-related ratio of tubular maximum reabsorption of phosphate to GFR standard deviation score; NSAID = nonsteroidal anti-

inflammatory drug; ACE inhibitors = angiotensin converting enzyme inhibitors; ARBs = angiotensin II receptor blockers; NT = not tested; N/A = not applicable
Categorical data is presented as number and percent and was analyzed by Fisher's exact test. Continuous data is presented as median and interquartile range and was analyzed by Mann-Whitney U test.

ORIGINAL UNEDITED MANUSCRIPT

Table 3. Correlation coefficients of iPTH in patients with Bartter syndrome type I & II

Variable	N	r_s	p-value
Sex (male)	107	-0.056	0.569
Age	107	0.116	0.236
Nephrocalcinosis	101	-0.064	0.526
Creatinine	107	0.171	0.078
Sodium	104	0.003	0.974
Potassium	107	-0.112	0.249
Chloride	98	-0.027	0.792
Bicarbonate	99	0.044	0.663
Phosphate-SDS	104	0.104	0.293
Calcium	105	-0.253	0.009
Magnesium	93	0.014	0.897
Uric acid	56	0.002	0.987
Alkaline phosphatase-SDS	82	0.268	0.015
Total protein	48	0.115	0.435
Albumin	75	-0.126	0.283
25OH vitamin D	60	-0.044	0.739
Urinary calcium/creatinine ratio	82	-0.083	0.459
TRP	17	-0.409	0.103
TmP/GFR-SDS	17	-0.085	0.747
Indomethacin or other NSAID	107	-0.095	0.331
Potassium-sparing diuretics	107	0.137	0.158
Aldosterone antagonists	107	-0.115	0.239
ACE inhibitors/ARBs	107	-0.049	0.619
Hydrochlorothiazide	107	-0.003	0.978
Potassium supplements	107	0.102	0.295
Sodium supplements	107	0.042	0.668
Magnesium supplements	107	0.175	0.072
Proton pump inhibitors	107	0.010	0.918
Other gastric protectors	107	-0.083	0.393
Oral contraceptives	107	0.108	0.266
Vitamin D supplements	107	0.102	0.294

id hormone; phosphate-SDS = age-related phosphate standard deviation score; alkaline phosphatase-SDS = age-related alkaline phosphatase standard deviation score; 25OH vitamin D = 25-hydroxy vitamin D; TRP = tubular reabsorption of phosphate; TmP/GFR-SDS: age-related ratio of tubular maximum reabsorption of phosphate to GFR standard deviation score; NSAID = nonsteroidal anti-inflammatory drug; ACE inhibitors = angiotensin converting enzyme inhibitors; ARBs = angiotensin II receptor blockers.

r_s = Spearman's rank correlation coefficient

i
P
T
H
=
i
n
t
a
c
t
p
a
r
a
t
h
y
r
o

ORIGINAL


Table 4. Multivariable logistic regression of patients with Bartter syndrome type I & II for hyperparathyroidism (PTH > 7.0 pmol/l)

Variable	Units	N	OR	95% C.I.		p-value
Calcium	mmol/l	80	0.03	0.001	– 1.16	0.059
Phosphate-SDS		80	1.29	0.93	– 1.81	0.128
Alkaline phosphatase-SDS		80	1.54	1.09	– 2.19	0.015
Indomethacin or other NSAID	Yes	80	1.12	0.38	– 3.34	0.833
Vitamin D supplements	Yes	80	1.34	0.49	– 3.66	0.566

Phosphate-SDS = age-related phosphate standard deviation score; alkaline phosphatase-SDS = age-related alkaline phosphatase standard deviation score; NSAID = nonsteroidal anti-inflammatory drug; OR = Odds ratio; 95% C.I. = 95% confidence interval

ORIGINAL UNEDITED MANUSCRIPT

Figure 1. Patient cohort




NOS = not otherwise specified

Patients were grouped for analysis according to the genetic confirmation of disease (subtype).

* Creatinine and PTH had to be provided.

51 patients with Bartter syndrome type I and 56 patients with Bartter syndrome type II

ORIGINAL UNEDITED MANUSCRIPT

Figure 2. Boxplots of iPTH, calcium, phosphate-SDS, and TmP/GFR-SDS according to disease subtype

iPTH = intact parathyroid hormone; phosphate-SDS = age-related phosphate standard deviation score; TmP/GFR-SDS: age-related ratio of tubular maximum reabsorption of phosphate to GFR standard deviation score;


Boxplot graphs represent the median and IQR; the upper and lower whiskers represent data points within 25th percentile minus 1.5x IQR and 75th percentile plus 1.5x IQR. Outliers are plotted individually. * p < 0.05; ** p < 0.01; *** p < 0.001.

ORIGINAL

A. iPTH in pmol/L; grey area represents normal range (2.0 – 7.0 pmol/l); statistically significant difference in iPTH level between patients with Bartter syndrome type I & II and patients with Bartter syndrome type III, Bartter syndrome NOS and Gitelman syndrome. **B.** Calcium in mmol/l, grey area represents normal range (2.20 – 2.60 mmol/l); statistically significant difference in calcium level between patients with Bartter syndrome type I & II and patients with Gitelman syndrome. **C.** Phosphate-SDS; grey area represents normal range (-2.0 – 2.0); statistically significant difference in age-adjusted phosphate level between patients with Bartter syndrome type III and patients with Bartter syndrome type I & II and Gitelman syndrome. **D.** TmP/GFR-SDS; grey area represents normal range (-2.0 – 2.0); statistically significant difference between Bartter syndrome type I & II and Bartter syndrome NOS and Gitelman syndrome.

ORIGINAL UNEDITED MANUSCRIPT

Figure 3. Scatterplots of calcium, urinary calcium/creatinine ratio, phosphate-SDS, TmP/GFR-SDS and alkaline phosphatase-SDS with iPTH of patients with Bartter syndrome type I & II




iPTH = intact parathyroid hormone; phosphate-SDS = age-related phosphate standard deviation score; alkaline phosphatase-SDS = age-related alkaline phosphatase standard deviation score; TmP/GFR-SDS: age-related ratio of tubular maximum reabsorption of phosphate to GFR standard deviation score;

A. Scatterplot of calcium with iPTH. **B.** Scatterplot of urinary calcium/creatinine ratio with iPTH. **C.** Scatterplot of phosphate-SDS with iPTH. **D.** Scatterplot of TmP/GFR-SDS with iPTH. **E.** Scatterplot of alkaline phosphatase-SDS with iPTH

ORIGINAL MANUSCRIPT

Figure 4. Scatterplots of TmP/GFR with age-adjusted phosphate



Phosphate-SDS = age-related phosphate standard deviation score; TmP/GFR-SDS: age-related ratio of tubular maximum reabsorption of phosphate to GFR standard deviation score;

A. Scatterplot of TmP/GFR with age-adjusted phosphate of all patients. **B.** Scatterplot of TmP/GFR with age-adjusted phosphate of patients with Bartter syndrome type III. **C.** Scatterplots of TmP/GFR with age-adjusted phosphate of patients with Gitelman syndrome

ORIGINAL