

What are the unmet needs and most relevant treatment outcomes according to patients with inflammatory bowel disease?

A qualitative patient preference study

Elise Schoefs^{1*}, Séverine Vermeire^{2,3}, Marc Ferrante^{2,3}, João Sabino^{2,3}, Tessy Lambrechts², Luisa Avedano⁴, Isabella Haaf⁴, Maria Stella De Rocchis⁴, Andrea Broggi⁴, Magdalena Sajak-Szczerba⁴, Roberto Saldaña⁵, Rosanne Janssens^{1†}, Isabelle Huys^{1†}

[†]These authors share last authorship

¹Department of Pharmaceutical and Pharmacological Sciences, KU Leuven, Leuven, Belgium

²Department of Gastroenterology and Hepatology, University Hospitals Leuven, KU Leuven, Leuven, Belgium

³Department of Chronic Diseases, Metabolism and Aging, KU Leuven, Leuven, Belgium

⁴European Federation of Crohn's & Ulcerative Colitis Associations (EFCCA), Brussels, Belgium

⁵European Federation of Crohn's & Ulcerative Colitis Associations (EFCCA), Madrid, Spain

© The Author(s) 2022. Published by Oxford University Press on behalf of European Crohn's and Colitis Organisation.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

*** Correspondence:**

Elise Schoefs

Herestraat 49 box 521,

3000 Leuven, Belgium

Tel: 0032 476 08 14 36

Mail: elise.schoefs@kuleuven.be

Contact details

Séverine Vermeire	-	severine.vermeire@uzleuven.be
Marc Ferrante	-	marc.ferrante@uzleuven.be
João Sabino	-	joao.sabino@uzleuven.be
Tessy Lambrechts	-	tessy.lambrechts@uzleuven.be
Luisa Avedano	-	luisa.avedano@efcca.org
Isabelle Haaf	-	bella.haaf@efcca.org
Maria Stella De Rocchis	-	mariastella.derocchis@efcca.org
Andrea Broggi	-	andrea.broggi@gmail.com
Magdalena Sajak-Szczerba	-	magdalena.sajak@efcca.org
Roberto Saldaña	-	roberto.saldana@efcca.org
Rosanne Janssens	-	rosanne.janssens@kuleuven.be
Isabelle Huys	-	isabelle.huys@kuleuven.be

A part of this work was presented as a poster presentation at DIA Europe 2022 on 30 March 2022.

Abstract

BACKGROUND AND AIMS: As more therapeutic options with their own characteristics become available for inflammatory bowel disease (IBD), drug development and individual treatment decision-making needs to be tailored towards patients' preferences and needs. This study aimed to understand patient preferences among IBD patients, and their most important treatment outcomes and unmet needs.

METHODS: This qualitative study consisted of 1) a scoping literature review, 2) two focus group discussions (FGDs) with IBD patients (n=11) using the nominal group technique, and 3) two expert panel discussions.

RESULTS: IBD patients discussed a multitude of unmet needs regarding their symptoms, side-effects, psychological and social issues for which they would welcome improved outcomes. Particularly, IBD patients elaborated on the uncertainties and fears they experienced regarding the possible need for surgery or an ostomy, the effectiveness and onset of action of their medication, and its long-term effects. Furthermore, participants extensively discussed the mental impact of IBD and their need for more psychological guidance, support, and improved information and communication with healthcare workers regarding their disease and emotional well-being. The following five characteristics were identified during the attribute grading as most important: prevent surgery, long-term clinical remission, improved quality of life (QoL), occurrence of urgency, and improved labor rate.

CONCLUSIONS: This study suggests that IBD drug development and treatment decision-making needs to improve IBD symptoms and adverse events that significantly impact IBD patients' QoL. Furthermore, this study underscores patients need for a shared decision-making process where their desired treatment outcomes and uncertainties are explicitly discussed and considered.

Keywords: Inflammatory bowel disease, patient-centric decision-making, qualitative patient preference study

Accepted Manuscript

1 Introduction

Inflammatory bowel disease (IBD) is a collective term for a range of clinical phenotypes caused by chronic, idiopathic, and remitting inflammation of the gastrointestinal tract. Crohn's disease (CD) and ulcerative colitis (UC) are the two most common forms^{1,2}. IBD is a lifelong disease affecting more than 10 million people worldwide, without any cure currently being available^{3,4}. Symptoms include abdominal pain, chronic and relapsing episodes of (bloody) diarrhea, urgency, nausea, vomiting, weight loss, anorexia, and fatigue, which negatively impact patients' health and quality of life⁵.

Over the past decades, a transition took place in the therapeutic landscape of IBD; where IBD patients were previously highly dependent on conventional therapies such as aminosalicylates, corticosteroids, and immunosuppressants, more advanced therapies are being developed and entering the market (e.g., biologicals and small molecules)⁶. IBD treatments in development and use today differ in terms of benefits, side-effects, route of administration, cost, mechanism and speed of action, treatment schedule, and uncertainties related to their treatment outcomes on the long and short-term. As more therapeutic options with their own characteristics come on the market and are in the pipeline, it is increasingly recognized that drug development and evaluation need to be tailored towards patients' preferences and needs. In addition, it is increasingly expected that IBD patients are involved in individual treatment decision-making, whereby shared decision-making (SDM) is considered the ideal model^{7,8}. This model encourages patients, in consultation with their clinician to 1) consider the available screening, treatment, or management options including their associated benefits, risks, and uncertainties, 2) communicate their preferences, and 3) express their desire for involvement in decision-making⁹. However, in order to facilitate SDM in the clinical encounter, a better understanding of factors that IBD patients find important is needed.

Patient preference studies, usually consisting of a qualitative and a quantitative phase, can help to identify which treatment characteristics are important to patients, how important, which trade-offs patients are willing to make between various characteristics, how preferences may vary according to individual patient characteristics, and how much uncertainty patients accept¹⁰. The qualitative phase of a patient preference study can generate information about patients' experiences and perspectives to gain in-depth and broad knowledge on the value of medical products and characteristics associated with them¹¹⁻¹³. In addition, qualitative research is recommended to inform the identification of attributes (treatment features) used in subsequent quantitative research^{14,15}.

A systematic literature review by Bewtra *et al.*¹⁶ showed that previous patient preference studies among IBD patients were mostly quantitative in nature, did not use a particular qualitative data collection method (such as individual interviews or group discussions), and/or resulted in limited evidence on the perceived value of treatment attributes according to patients. In addition, these studies were conducted before newer treatments such as biologicals and Janus Kinase (JAK) inhibitors were available, which have been described to affect patients' preferences¹⁶. Moreover, other published patient preference studies had a specific focus on preferences for surgery and medical management versus colectomy, had the goal to identify preferences for the administration of treatments, or were focused on a specific drug formulation¹⁷⁻²².

Given the lack of comprehensible and recent qualitative research on IBD patients' preferences that considers different (novel) treatments and their associated characteristics, the present qualitative patient preference study aimed to investigate which patient-relevant treatment characteristics of both treatments in development and use today IBD patients find important and why. More specifically, this study aimed to understand: 1) IBD patients' unmet needs regarding their disease and treatment, and 2) treatment outcomes considered important by IBD patients. Findings from this study were furthermore intended to support the identification and development of attributes and

levels for inclusion in a subsequent quantitative patient preference study to obtain more representative quantitative results in a larger sample of IBD patients. Furthermore, findings may inform the development of a patient decision aid to facilitate SDM in clinical practice.

2 Materials and methods

A qualitative study consisting out of three phases was designed following the general steps described in the PREFER framework²³. The three phases (Figure 1) consisted of 1) a scoping literature review, 2) two Focus Group Discussions (FGDs) with IBD patients (n=11) using the Nominal Group Technique (NGT), and 3) two expert panel discussions with patients (n=2), patient representatives (n=4), clinicians (n=3), and preference research experts (n=2). The methodology and results of the FGDs were reported according to the recommendations of Hollin *et al.*¹⁴ for reporting qualitative preference research, and the recommendations of Coast *et al.*¹⁵ and Bridges *et al.*²⁴ on collecting and analyzing methods for initial attribute development.

[Insert Figure 1 here]

2.1 Scoping literature review to develop the study documents for the focus group discussions

To develop the attribute grading exercise and other study documents for the FGDs, a scoping literature review was performed of 1) attributes and key results of previous published preference studies conducted among IBD patients, 2) favorable and unfavorable aspects of IBD treatments already being prescribed to patients as assessed by the European Public Assessment Reports (EPARs) of the European Medicines Agency (EMA) or as indicated on the products' leaflet, and 3) primary and secondary endpoints and adverse events of phase 3 IBD clinical trials in the European Union, to ensure also potential 'future' treatment outcomes and adverse events could be integrated in the

discussion. A detailed description of the databases, search queries, and in-and exclusion criteria can be found in Appendices 1-3.

2.2 Focus group discussions

2.2.1 Participant recruitment

Ethical approval was obtained from the Ethische Commissie Onderzoek UZ/KU Leuven in Belgium (S65034). Participants were recruited in April 2021 by an IBD-nurse at the University Hospitals of Leuven. Patients who were 18 years or older, diagnosed with IBD (CD or UC), and able to give informed consent were eligible to participate. Because individual patient characteristics could have an influence on participants' opinions and gradings, and the goal was to identify characteristics important across the IBD patient population, purposive sampling was used to reach heterogeneity among participants with regards to age, subtype of IBD, disease stage (active disease/remission), and treatment experience (including surgery)²⁵. Following recommendations of McMillan *et al.*²⁶, recruitment sought to include between five and seven IBD patients per FGD.

2.2.2 Completion of the online form

As preparation for the FGDs, participants were invited to read the information sheet (Appendix 4) and to complete the online form, which consisted of 1) the informed consent form (Appendix 5 Part A), 2) a short survey to assess patients' personal and disease-specific characteristics, and health literacy (Appendix 5 Part B), 3) open questions to identify treatment characteristics important to participants (Appendix 5 Part C), and 4) a grading exercise where participants needed to grade the treatment characteristics identified in the scoping literature review from 1 to 10, with 10 being the most important (Appendix 5 Part D). All participants completed the online form and gave their informed consent prior to the discussion. There was no drop-out due to the length of the survey.

2.2.3 Conduct of the focus group discussions using the Nominal Group Technique

FGDs using the NGT were chosen as preference exploration method in close dialogue with clinicians and patient representatives since there was interest in group dynamics and the subject of the discussion was not deemed to be too sensitive to discuss with others. The NGT is a consensus methodology where participants prioritize their thoughts and perspectives, both as group and individually, with the advantage of reaching consensus in a short period of time. As the NGT is structured in four-stages (idea, round-robin, clarification and finalization, grading and consensus), different techniques are combined including open questions to detect new characteristics not identified during literature review (bottom-up) and a grading exercise and discussion to question participants about the importance of characteristics identified in literature (top-down); the grading allows researchers to select and understand which treatment characteristics are most important and should be integrated in a subsequent preference survey²⁶.

During the FGDs, a predefined FGD guide (Appendix 6) was followed, taking into account the four steps of the NGT. Each discussion was executed by the same moderator (ES) and in the native language of the participants (Flemish). Considering the implications of the coronavirus (COVID-19) pandemic, the discussions were organized online. The discussions lasted around one hour and 45 minutes and were audio-recorded. The content of the online form and the FGD guide was provided with written feedback from a gastroenterologist and a member of a patient organization.

2.2.4 Analysis of the focus group discussions

2.2.4.1 Quantitative descriptive analysis

Participants' self-reported personal and disease-specific characteristics, and their scores on the attribute grading exercise were analyzed descriptively. Health Literacy was determined using the Chews' set of brief screening questions²⁷. To obtain a final rank of the treatment characteristics, the

average, standard deviation, minimal score, and maximal score were calculated for each characteristic by combining the grades of all participants.

2.2.4.2 Qualitative thematic analysis

The audio-recordings of all FGDs were transcribed ad verbatim in the original language by a member of the research team (ES) and pseudonymized. Subsequently, the transcripts were subjected to the framework analysis method (using NVivo software), a qualitative content analysis for text data where overarching themes are developed²⁸. Analysis started with familiarization through the conduct, transcription, and reading of the FGDs. Themes of the discussion guide informed the creation of deductive codes. Analytical notes were written in the margins of the transcripts and informed the creation of inductive codes. Based on the deductive and inductive codes a 'coding tree' with characteristics was developed. Regular meetings between the researchers that conducted the FGDs (ES, RS, and IH) were held to verify the characteristics were interpreted in the same manner. After consensus was reached, the coding tree was uploaded in NVivo, and applied to all transcripts, where literal quotations of transcripts relating to a particular theme were classified under the respective code. NVivo was used for charting (summarizing) the data per code in order to have an oversight of all results, making interpretation and analysis of the results both between participants and FGDs possible.

2.3 Expert panel discussions

After the conduct of the FGDs, expert panel discussions with patients, patient representatives, clinicians, and preference research experts were held to verify the outcomes of the FGDs and to provide further depth into the identified themes. Furthermore, insights were gathered regarding possible attributes and levels for inclusion in the subsequent quantitative patient preference study, taking into account the criteria and steps as described by Durosini *et al.*²⁹, Petrocchi *et al.*³⁰, and Janssens *et al.*³¹.

3 Results

3.1 Scoping literature review to identify characteristics potentially relevant to IBD patients

In total, 22 patient preference studies, 58 phase 3 IBD clinical trials, and 45 IBD treatments fulfilled the inclusion criteria. From the included publications, data was extracted and summarized (Appendices 1-3). Based on the results of the literature review, the attribute grading exercise was created, wherein 55 identified characteristics were grouped in three overarching categories namely, characteristics related to the treatment efficacy, characteristics related to the administration of the treatment, and characteristics related to symptoms of the disease or complications and adverse events of treatments (Appendix 5 Part D).

3.2 Focus group discussions to identify patient relevant characteristics

3.2.1 Participants' characteristics

First contact was established with 13 IBD patients of which 11 joined the FGDs, resulting in a response rate of 84%. The mean age was 37 years [age range: 27-56; median age: 38 years; interquartile range (IQR): 28.5-42 years] (Appendix 7). On average, there was a 10.5-year gap between diagnosis and participation in the FGD (age at diagnosis range: 16-35 years, median age at diagnosis; 26 years; IQR: 23.5-30 years). There was an imbalance in IBD subtype (73% diagnosed with UC), being the most outspoken in the second FGD. More than half of the participants (55%) had undergone surgery due to disease complications. Health literacy was adequate in all participants.

Of all participants, three patients were in remission with medical treatment and one participant was not taking any medication on advice of his treating physician. Participants were most frequently treated with biologicals only (n=7). Other than biologicals alone, participants were taking a combination of: aminosalicylates with JAK inhibitors (n=1), corticosteroids with immunosuppressants

and biologicals (n=1), and aminosaliclates with corticosteroids (n=1). Table 1 gives an overview of the treatment experience of the participants.

[Insert Table 1 here]

3.2.2 Unmet needs and treatment outcomes considered important by IBD patients

IBD patients discussed a multitude of treatment outcomes and unmet needs for which they would welcome improvements both in the context of drug development as well as in the context of decision-making in clinical practice.

3.2.2.1 Gastrointestinal symptoms

Gastrointestinal symptoms (e.g., abdominal pain and cramps, diarrhea, urgency, incontinence, nocturnal defecation, bowel movements, and constipation) had a significant effect on the participants' life and was seen as detrimental in three ways. First, some participants frequently experienced nocturnal defecation, having an incremental effect on their sleep and therefore causing reduced energy (as further described in 'Reduced energy'). Second, work, sports, or social activities could be hampered, limiting their freedom. Last, knowing that gastrointestinal symptoms could appear at any time had an important psychological impact, making participants worrisome about always having a toilet nearby. Therefore, an improvement in both the frequency and the severity of gastrointestinal problems was seen as the most important treatment outcome for all participants: *"Something that really comes at 1 actually."* #FG1_E.

3.2.2.2 Reduced energy

The theme of reduced energy caused by participants' therapy and illness was extensively discussed in both FGDs and was experienced by the majority of the participants. Reduced energy could hinder participants ability to normally participate in daily activities, limiting their quality of life. Therefore, participants wanted to avoid reduced energy caused by their therapy: *"For me fatigue is something I*

could not accept as a side-effect." #FG1_E. Nevertheless, reduced energy could not be avoided for every participant and therefore patients had to adapt their lifestyle around this problem. As a consequence, a couple of participants were not able to exercise, which was seen as another important limiting factor. Reduced energy and fatigue could also result from sleeping problems participants experienced due to their disease or nocturnal defecation, but also other causes of reduced energy were shared (e.g., following psychological problems patients were struggling with or weight fluctuations).

3.2.2.3 Changes in physical appearance

Patients highlighted that changes in physical appearance, caused both by their altered gut function or as a side-effect of their therapy, made them insecure about their body image. Especially participants treated with corticosteroids had experienced a spectrum of changes (namely moon face, developing facial hair, hair loss, and weight changes) in their physical appearance, which had a detrimental impact on their mental wellbeing. For these participants, the side-effects outranged the benefits of their treatment.

Among the several types of physical changes that emerged during the discussions, (excessive) weight fluctuations were seen as the most important, and the topic seemed to receive particular attention and discussion from female participants. Weight fluctuations caused both by participants' therapy or altered gut function were found detrimental because it made participants uncertain and had a negative effect on their energy levels. Therefore, all participants found it important that their weight was more or less under control. One participant even stated that weight gain was the most important treatment characteristic, and if a choice had to be made between different treatment options, the treatment in which weight gain might occur as a side-effect was to be dropped first.

3.2.2.4 Skin manifestations

Several participants previously experienced skin related issues as a side-effect of their treatment. In general, participants could cope with less severe skin manifestations. Nevertheless, the severity in which this side-effect occurred determined how much participants valued this treatment characteristic: *“If your whole body gets covered [with skin lesions], you start to look at it differently. So, I also think that it depends a little bit in which degree the side-effect occurs.”* #FG1_B.

3.2.2.5 Vision problems

One participant stated to have a deteriorated eyesight: *“I am suffering from tear duct inflammation. (...) Whether that's linked to Crohn's or my treatment with corticosteroids, I don't know.”* #FG1_B.

However, it remained largely unclear whether vision problems could also be an extra-intestinal manifestation associated with having a chronic autoimmune disease like IBD. As participants knew that a lot of medication could cause eye infections or other visual disturbances, suffering from (permanent) vision problems was seen as an important treatment characteristic.

3.2.2.6 Risk of surgery

Most of the participants saw surgery as their last resort and were resistant to it: *“I think surgery that is often the last step.”* #FG2_D. Despite this, more than half of the participants underwent surgery in the past because they had to. Participants, especially the persons diagnosed with CD, knew that repeatedly removing sections of their gut was not sustainable in the long run and were afraid that their bowels would become too short to take in enough nutrients. In addition, they looked at it as ‘a path without an end’. Nevertheless, some participants stated that their symptoms were more or less under control after they underwent surgery.

3.2.2.7 Risk of an ostomy

The need for an ostomy was extensively discussed during both focus groups, with some participants expressing their fear about ever going to need it. An ostomy was something they wanted to avoid at all costs because it would be an additional limitation and would hurt their self-image. Additionally,

some participants raised their concerns about needing additional care if they received an ostomy. However, others stated that they did not really think about ever going to need it. Nonetheless, one participant with CD had a permanent ostomy and was very pleased with it: *"I was 24 when I got my stoma and I have really resisted it. I already had every kind of medication (...). And then I first had a temporary stoma and even then, I felt that burden fell of my shoulders. That I did not have to run to the toilet, that I could go back to places without worrying (...). The only thing is that my disease is still not 100% under control."* #FG2_D. In addition, this participant mentioned that the ostomy enabled to relive a normal life without having the restrictions from the disease.

3.2.2.8 An effective and fast working medication

On several occasions, participants discussed the psychological burden of uncertainties regarding the effectiveness and onset of action of their medication. Participants found it frustrating to start with a new treatment, not knowing if the medication was ever going to work for them and for how long it was going to work before losing its efficacy. As a result, most participants had tried various medications until they found (or were still searching for) the one that worked for them. In addition, participants complained about the uncertainty regarding the onset of action of the new treatment. This had a negative impact in two ways: 1) because they did not know if the medication would eventually work for them, and whether they would need to switch to another medication, or whether they would need to give it more time to be effective, and 2) because during this bridging period, participants' symptoms stayed present. Therefore, one participant explicitly stated that a fast clinical response was the most important treatment characteristic.

3.2.2.9 Normal social interactions and activities

All participants experienced limitations on their social interactions and activities due to their disease and the unpredictable nature thereof. Making plans, such as going on a vacation, was seen as difficult by some participants. In addition, some participants raised that they frequently needed to cancel activities with their friends because they were not feeling well or were experiencing pain, and

were therefore forced to stay at home, resulting in isolation and loneliness. For some participants, this even resulted in deterioration of relationships with their friends or losing their job. Therefore, participants found it key that their treatment enabled them to have a normal social life where they were able to keep their social contacts, without being restricted by their (gastrointestinal) symptoms. Furthermore, some participants clearly addressed the lack of understanding of people in their surroundings for their situation and disease. Participants attributed this to the unawareness of IBD and all that it entails: *“Very often people say, 'it's all in your head', or 'you ate something bad', or... there is little understanding of the fact that sometimes you can sit on the toilet for more than half an hour.”* #FG2_A.

3.2.2.10 Mental and psychological support

In general, participants tried to maintain a positive mindset: *“No matter how sick I've been, I did always persevere and believe in it.”* #FG1_B. However, as many participants suffered greatly from their disease; the disease, to some extent, dictated what they could or could not do in their daily lives. As a result, this had a giant impact on the participants mental wellbeing and some patients were even struggling with mental disorders (e.g., loneliness, anxiety, depression). Therefore, multiple participants indicated that they needed professional mental and psychological help to cope with their disease: *“I have been at the point where I thought: 'I just can't get out of here, not mentally, not physically'. That I also sat with very dark thoughts.”* #FG2_D. Participants found that too little attention in the clinical encounter was given to the whole mental aspect of the disease and liked to have had more guidance. They experienced sometimes that clinicians did not have a full understanding of the impact of the disease on their daily lives. As a result, participants extensively discussed the lack of mental support some of them needed. Some participants stated that their clinician did advise them to contact a psychologist if they needed to. However, participants felt that once they left the hospital, they were completely on their own.

3.2.2.11 Information regarding the effect of the disease and treatments on a long and short term

Some participants expressed the lack of information they received on how their disease would progress and felt uncertain about the long-term effects of their medication, as these effects are relatively unknown. Participants found it very important that they would not struggle with other diseases later in their lives and some participants expressed their fear about the possible emergence of cancer. Despite this, participants still used these medication as they wanted to stay in remission. In addition, certain participants had tried so many different medications, they had no other options: *“Sometimes you are desperate, and you want to feel better. So, at that moment it was clear to me: I want to get better now, and we'll see what comes in the future.”* #FG2_D. Furthermore, participants would have liked to know more from the beginning on how the disease would affect their life in understandable, layman’s terms: *“You do get medical information about what Crohn's disease is, but you don't really get to know how to live with it.”* #FG2_A. In an ideal situation, most participants would have liked to be brought in contact with other patients who could share their experiences. Although participants knew that there was limited literature on the effect of different nutrition on their symptoms, they would like to receive more information on what they could or could not eat.

3.2.3 Attribute grading

In addition to the open discussion revealing patients’ unmet needs and treatment outcomes and goals, the attribute grading allowed to explore and trigger further discussion on which treatment and disease related characteristics matter most to patients. In general, most participants gave a very high score to all the 55 treatment characteristics on the attribute grading exercise (bottom-up) and several participants stated during the discussion of the list that they found all characteristics important. The top ten characteristics, based upon the attribute grading exercise, are shown in Table 2 (full results can be found in Appendix 8). To a large extent, these characteristics align with the qualitative results; gastrointestinal problems, characteristics related to quality of life, and surgery received a very high score (see 3.2.2). Characteristics found relatively less important were related to

the administration of the treatment, namely treatment frequency, place of treatment, and route of administration, which were also not spontaneously raised by patients during the open discussions. The attribute grading exercise also revealed that the characteristics included in the grading exercise did not include the level of detail and full range of aspects patients freely elaborated on in the open discussion (see 3.2.2); patients for example indicated and elaborated on the influence of IBD on their mental wellbeing, a characteristic not included in the attribute grading exercise.

While some participants stated that their reason for scoring an attribute was based on long-term reasoning, other participants only looked at the present. However, scoring could also be influenced by patients' previous treatment or disease experience. Particularly, whether they had experienced a certain symptom or side-effect could have determined their grading. In addition, some patients stated that when side-effects were temporary or could be resolved with other medication, they became manageable and therefore less 'important'.

[Insert Table 2 here]

3.3 Expert panel discussions to ensure the relevance and correct interpretation of the identified characteristics

Two expert panel discussions were held with i) patients (n=2) and patient representatives (n=4) to verify the importance, relevance, comprehensiveness, and understandability of the identified characteristics, ii) clinicians (n=3) to ensure plausibility from a clinical perspective, and iii) preference research experts (n=2) to give insights regarding possible attributes and levels compliant with the corresponding rules^{14,15}. During the expert panel discussions, the unmet needs and treatment outcomes important to IBD patients as identified during the FGDs were presented and the relevance and correct interpretation thereof was confirmed. In particular, in-depth discussions were held regarding the following characteristics: reduced energy, mental disorders that patients could struggle with, and the impact of IBD on patients'

social interactions and activities. As an outcome, the expert panel discussions led to more detail and insights into the identified themes.

4 Discussion

This qualitative study revealed in-depth insights into symptoms, side-effects, psychological and social issues that IBD patients find important. During the open discussions, patients highlighted gastrointestinal symptoms, reduced energy, changes in physical appearance, skin manifestations, and vision problems as being major issues reducing their quality of life. In addition, IBD patients elaborated on the uncertainties and fears they experienced regarding the possible need for surgery and in particular an ostomy. The effectiveness and onset of action of medication, and its long-term effects were further identified as major concerns. Lastly, the mental impact of IBD and the need for more psychological guidance, support, and improved communication with healthcare workers regarding patients' emotional well-being was found key.

Our findings show that IBD patients are confronted with both physical and psychological issues that require further support, which has been confirmed in several studies investigating the impact of IBD on patients' health related quality of life. For example, Spagnuolo *et al.*³² assessed the satisfaction with social roles and physical function in IBD patients by evaluating patient reported outcomes (PROs) and revealed that IBD patients had an impaired ability to perform normal daily activities compared to a control group. Likewise, Iaquina *et al.*³³ collected PRO data and concluded that IBD patients have increased fatigue and anxiety compared with healthy persons. The study from Dudley-Brown *et al.*³⁴ identified the following main themes corresponding to our study findings: uncertainty surrounding exacerbations, the desperate need to find successful treatment, and the feeling of being controlled by the disease.

A possible way to cope with patients' uncertainties is to improve the communication with healthcare workers; patients who are informed about the side-effects of medication and about the possible need for surgery or an ostomy may experience less uncertainty about their future disease progression. In addition, if patients are given the opportunity to ask for help and advice, symptoms and side-effects are likely better managed. Based on patients' individual needs, there are three possible efforts to support IBD patients. First, patients may find support in IBD associations where they can receive reliable support from other patients. As confirmed during our FGDs, participants were overall very happy to have participated in the discussion and share their experiences, feelings, and thoughts with other IBD patients as this took away some of their insecurities. Furthermore, Sewitch *et al.*³⁵ showed that improving social support can have a favorable impact on psychological distress, ultimately improving health outcomes. Second, staff could be trained in IBD centers to provide more psychological support in the IBD medical care pathway. Third, the results of this study can be used to develop patient decision aids (e.g., flyers, online tools ...) that can be integrated in IBD clinical practice to support SDM. During the discussions, patients discussed a multitude of treatment outcomes and unmet needs that should be discussed in clinical practice to enable an informed decision-making. For example, patients mentioned the need for more information regarding the effect of the disease and their treatments on a long and short term, which suggest an opportunity for the development of patient decision aids that can inform patients about these aspects. As a result, the communication between the patient and the health care provider can be facilitated and uncertainties can be addressed. Furthermore, by aligning individual medical treatment decisions with patients' needs and preferences, higher quality decisions, improved patients' compliance, and thereby improved patient's overall health can be achieved^{36,37}.

Comparing the disease and treatment characteristics found in this study to those identified in previous IBD patient preference studies (Appendix 1) reveals that some attributes overlap, while others were new. Important to note is that almost all previous studies had a specific scope and

involved both patients and gastroenterologists, making it not clear which attributes were raised as important by whom. In contrast, our study identified patient-relevant characteristics across different therapies, with an open scope, directly from IBD patients. Differences can further be explained by the fact that qualitative studies are highly context-specific in view of the research setting. As a result, different variables such as the selected patient sample, the time of the study, and the specific questions asked may explain a difference in the identified attributes, making comparison between the results of different qualitative patient preference studies difficult. Therefore, the results need to be interpreted considering the specific period and context the study took place, as well as in view of the type of participants that took part. For example, this study took place during the COVID-19 pandemic, wherein the social isolation participants experienced during the pandemic may have amplified the need and frequency patients expressed their wish for more mental support. Results should also be viewed in the specific drug therapy context and the influence of the individual drug therapy experience of participants on the identified symptoms and side-effects; the IBD treatment landscape is rapidly evolving, with new therapies that have entered the market or are emerging. These new treatments have different (long-term) side-effects and efficacy profiles from those currently available. This may explain why patients expressed uncertainties regarding the effectiveness and onset of action of their medication, and the long-term side-effects such as the increased risk of cancer, particularly regarding these treatments that have only being marketed and described for a relative short time.

4.1 Implications and further use

The characteristics identified in this study will be useful in informing the design of a subsequent quantitative online patient preference survey to obtain insights from a geographically dispersed and heterogeneous sample of IBD patients. Such a survey will enable to obtain quantitative insights regarding the relative importance of attributes, and whether patient characteristics (e.g., type of treatments received, subtype of IBD, disease state, and experience with symptoms) have a

statistically significant impact on patients' preferences. Furthermore, the results of this qualitative study can aid the development of patient decision aids used to facilitate treatment decision-making in clinical practice by revealing which aspects of treatments (e.g., long-term efficacy and safety of treatments) should be explicitly discussed in clinical practice and communicated to patients.

The identified characteristics important to IBD patients may further be useful to assist and optimize the identification of clinical trial endpoints and outcomes that matter most to IBD patients. Including such outcomes in clinical trials could enable an understanding of how treatments perform regarding patient-relevant outcomes. Obtaining evidence on the extent to which (new) drugs perform on relevant outcomes within and outside the clinical trial setting are two questions which are difficult to resolve at the time of regulatory benefit-risk assessment and Health Technology Assessment (HTA) evaluation. Such evidence could hence support regulators and HTA assessors in understanding whether (novel) IBD treatments address unmet needs and treatment outcomes that matter to patients, and thereby support initial and post-marketing benefit-risk and reimbursement decisions. Qualitative and quantitative preference results could also be considered to inform (joint) scientific advice discussions by HTA bodies and regulators on the relevance of the clinical trial endpoints proposed by industry applicants e.g., allowing to understand whether relevant measures of quality of life are included in clinical trials²³. In addition, resource allocation could be optimized to drugs that target the symptoms and side-effects found important to patients. This could result in a drug development process more tailored to patients' needs^{13,23,38,39}. Additionally, the concerns of IBD patients about uncertainties of long-term efficacy and safety can inform future real-world evidence studies.

4.2 Strengths and limitations

By performing discussions with patients as drug end-users to understand their experiences with both their illness as their treatment, this study demonstrates the value of performing a qualitative patient preference study. Furthermore, the use of focus group discussions with open questions allowed to obtain both broad and in-depth information on the treatment and disease related characteristics that IBD patients find most important. In addition, the plausibility, quality, and comprehensiveness of both the patient material and the study, as the interpretation of the results, was ensured by collaborating with gastroenterologist and members of patient organizations before and after the conduct of the focus group discussions.

Patients with different treatment experiences who may or may not have been in remission at the time of the discussions were included in the study. As a result, participant heterogeneity was introduced to avoid biases in the data; it was envisioned that these individual patient characteristics and experiences could have an influence on participants' opinions. As a consequence, interaction between patients with a different disease and treatment background emerged, ensuring the identified characteristics are found important for patients along the IBD spectrum. Likewise, an extensive literature review was performed to ensure patients also discussed potential 'future' treatment outcomes and side-effects of novel therapies, as well as favorable and unfavorable aspects associated with therapies already being prescribed to IBD patients, and thus not only treatment outcomes and side-effects participants already experienced themselves.

We did not differentiate the results according to drug therapy and IBD subtype as the goal was to provide an overarching view of important themes to IBD patients across therapies. Furthermore, it is worth noting that during the focus group discussions, it was not always possible to differentiate the raised characteristics between symptoms, side-effects of treatments, or extra-intestinal manifestations associated with having a chronic autoimmune disease like IBD. This further

underscores the complex etiology of IBD and the limited amount of published information on clinical evidence, resulting in uncertainty regarding the cause of problems that patients experience. Hence, more insights about the underlying mechanisms of IBD and the side-effects of different therapies would be useful in order to link adverse events to certain therapies, symptoms of the disease, or extraintestinal manifestations, which could increase treatment adherence and promote the treatment of IBD patients in general.

This study was performed during the COVID-19 pandemic, wherein the social distancing and hospital guidelines in Belgium did not allow face-to-face focus group discussions. Therefore, the discussions were organized online. This may have resulted in participants withholding to interact with others. Furthermore, it is likely that participants who were not comfortable with online discussions were less likely to participate (e.g., older participants). However, the median age of diagnosis of UC patients in this study was 28.3 and of CD patients 25.3, which is respectively 6.6 years and 3.7 years younger than the average age of diagnosis⁴⁰. Furthermore, as only two focus group discussions were performed and the discussions only contained participants living in Belgium, it cannot be determined if data saturation was reached and if the results can be extrapolated to patients outside Belgium. Therefore, we could speculate that the inclusion of additional patients from other countries could affect the derived themes.

Finally, it should be noted that the qualitative analysis of the focus group discussions was conducted by one researcher as opposed to multiple researchers, meaning that no cross-check was performed. Nevertheless, regular meetings between the researchers that conducted the focus group discussions minimized subjective interpretation of the data.

5 Conclusion

This study shows that IBD drug development and individual treatment decision-making does not only need to focus on controlling IBD patients' gastrointestinal symptoms, but also on other symptoms and adverse events that significantly impact IBD patients' quality of life. Furthermore, this study underscores the need for shared decision-making between healthcare providers and IBD patients where treatment outcomes and uncertainties about long-term efficacy and safety of medication are explicitly addressed and considered. Findings from this study may serve pharmaceutical industry, regulators, and HTA bodies/payers in understanding patient-relevant unmet needs and treatment outcomes to enable value-based and patient-centric decision-making that is aligned with the true patients' needs across the IBD drug life cycle.

Accepted Manuscript

6 Funding

This work received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

7 Acknowledgments

The authors are extremely grateful to the patients participating in the focus group discussions, for sharing their experiences and valuable contributions. Marc Ferrante and João Sabino are Senior Clinical Investigators of the FWO, Belgium.

8 Author Contributions

ES, RJ, and IHu were involved in the design of the study. ES designed the study materials. ES, TL, SV, MF, and JS were involved in the recruitment of the participants. ES, RJ, and IHu held the FGDs and analysed results. ES, RJ, IHu, SV, MF, JS, LA, IHa, SdR, AB, MS, and RS participated in meetings and reviewed the study materials. ES produced the first draft of the manuscript, which was subsequently revised and finalized with all authors. All the authors have read and approved the final manuscript.

9 Ethics approval and Consent to Participate

This study was reviewed and approved by the Ethische Commissie Onderzoek UZ/KU Leuven (Belgium; reference S65034). The patients/participants provided their informed consent to participate in this study.

10 Conflict of interest

SV reports research support from AbbVie, J&J, Pfizer, Galapagos, Takeda, and consultancy and/or speaker fees from AbbVie, AbolerIS Pharma, AgomAb, Alimentiv, Arena Pharmaceuticals, AstraZeneca, Avaxia, BMS, Boehringer Ingelheim, Celgene, CVasThera, Dr Falk Pharma, Ferring, Galapagos, Genentech-Roche, Gilead, GSK, Hospira, Imidomics, Janssen, J&J, Lilly, Materia Prima, MiroBio, Morphic, MrMHealth, Mundipharma, MSD, Pfizer, Prodigest, Progenity, Prometheus, Robarts Clinical Trials, Second Genome, Shire, Surrozen, Takeda, Theravance, Tillots Pharma AG, Zealand Pharma. MF reports research support from from Abbvie, Amgen, Biogen, Janssen, Pfizer, Takeda, and Viatrix, consultancy fees from Abbvie, Boehringer-Ingelheim, Celltrion, Janssen, Lilly, Medtronic, MSD, Pfizer, Sandoz, Takeda, Thermo Fisher, and speaker fees from Abbvie, Amgen, Biogen, Boehringer-Ingelheim, Falk, Ferring, Janssen, Lamepro, MSD, Pfizer, Sandoz, Takeda, Truvion Healthcare, Viatrix. JS reports consultancy fees from Janssen and Fresenius, and speaker fees from Abbvie, Takeda, Ferring. The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

11 Data availability statement

The datasets presented in this article are not publicly available because they contain information that could compromise participants' privacy and consent. Requests to access the datasets should be directed to elise.schoefs@kuleuven.be.

12 References

1. Pullen N., Gale JD. Inflammatory Bowel Disease. *Elsevier*. 2007. p. 613–42.
2. Damião AOMC., de Azevedo MFC., de Sousa Carlos A., Wada MY., Silva TVM., de Castro Feitosa F. Conventional therapy for moderate to severe inflammatory bowel disease: A systematic literature review. *World J Gastroenterol* 2019;**25**(9):1142–57. Doi: 10.3748/wjg.v25.i9.1142.
3. Lemmens B., de Hertogh G., Sagaert X. *Inflammatory Bowel Diseases*. Elsevier Inc.; 2014.
4. Guan Q. A Comprehensive Review and Update on the Pathogenesis of Inflammatory Bowel Disease. *J Immunol Res* 2019;**2019**. Doi: 10.1155/2019/7247238.
5. Badia X., Vilaseca J., Casellas F., Lo J. Influence of inflammatory bowel disease on different dimensions of quality of life. *Eur J Gastroenterol Hepatol* 2001:567–72.
6. Cohen NA., Rubin DT. New targets in inflammatory bowel disease therapy: 2021. *Curr Opin Gastroenterol* 2021:357–63. Doi: 10.1097/MOG.0000000000000740.
7. Barry MJ., Edgman-Levitan S. *Shared Decision Making-The Pinnacle of Patient-Centered Care*. 2012.
8. Charles C., Gafnv A., Whelan T. *SHARED DECISION-MAKING IN THE MEDICAL ENCOUNTER: WHAT DOES IT MEAN? (OR IT TAKES AT LEAST TWO TO TANGO)*. vol. 44. 1997.
9. Stiggelbout AM., van der Weijden T., de Wit MPT., Frosch D., Légaré F., Montori VM., et al. Shared decision making: Really putting patients at the centre of healthcare. *BMJ (Online)* 2012. Doi: 10.1136/bmj.e256.
10. FDA. Patient Preference Information – Voluntary Submission, Review in Premarket Approval Applications, Humanitarian Device Exemption Applications, and De Novo Requests, and Inclusion in Decision Summaries and Device Labeling. *Guidance for Industry* 2016.
11. van Overbeeke E., Whichello C., Janssens R., Veldwijk J., Cleemput I., Simoens S., et al. Factors and situations influencing the value of patient preference studies along the medical product lifecycle: a literature review. *Drug Discov Today* 2019. Doi: 10.1016/j.drudis.2018.09.015.
12. Whichello C., van Overbeeke E., Janssens R., Bywall KS., Russo S., Veldwijk J., et al. Factors and situations affecting the value of patient preference studies: Semi-structured interviews in

Europe and the US. *Front Pharmacol* 2019;**10**(September):1–11. Doi: 10.3389/fphar.2019.01009.

13. Janssens R., Huys I., van Overbeeke E., Whichello C., Harding S., Kübler J., et al. Opportunities and challenges for the inclusion of patient preferences in the medical product life cycle: A systematic review. *BMC Med Inform Decis Mak* 2019;**19**(1):1–16. Doi: 10.1186/s12911-019-0875-z.
14. Hollin IL., Craig BM., Coast J., Beusterien K., Vass C., DiSantostefano R., et al. Reporting Formative Qualitative Research to Support the Development of Quantitative Preference Study Protocols and Corresponding Survey Instruments: Guidelines for Authors and Reviewers. *Patient* 2020;**13**(1):121–36. Doi: 10.1007/s40271-019-00401-x.
15. Coast J., Al-Janabi H., Sutton EJ., Horrocks SA., Vosper AJ., Swancutt DR., et al. Using qualitative methods for attribute development for discrete choice experiments: Issues and recommendations. *Health Econ* 2012. Doi: 10.1002/hec.1739.
16. Bewtra M., Johnson FR. Assessing patient preferences for treatment options and process of care in inflammatory bowel disease: A critical review of quantitative data. *Patient* 2013;**6**(4):241–55. Doi: 10.1007/s40271-013-0031-2.
17. Byrne CM., Solomon MJ., Young JM., Selby W., Harrison JD. Patient preferences between surgical and medical treatment in Crohn's disease. *Dis Colon Rectum* 2007;**50**(5):586–97. Doi: 10.1007/s10350-006-0847-0.
18. Byrne CM., Tan KK., Young JM., Selby W., Solomon MJ. Patient and clinician preferences for surgical and medical treatment options in ulcerative colitis. *Colorectal Disease* 2014;**16**(4):285–92. Doi: 10.1111/codi.12538.
19. MacKenzie-Smith L., Marchi P., Thorne H., Timeus S., Young R., Le Calvé P. Patient Preference and Physician Perceptions of Patient Preference for Oral Pharmaceutical Formulations: Results from a Real-Life Survey. *Inflamm Intest Dis* 2018;**3**(1):43–51. Doi: 10.1159/000493346.
20. Almario C v., Keller MS., Chen M., Lasch K., Ursos L., Shklovskaya J., et al. Optimizing selection of biologics in inflammatory bowel disease: Development of an online patient decision aid using conjoint analysis. *American Journal of Gastroenterology* 2018;**113**(1):58–71. Doi: 10.1038/ajg.2017.470.
21. Kim ES., Kim KO., Jang BI., Lee CK., Kim HJ., Lee KM., et al. Factors contributing to the preference of Korean patients with Crohn's disease when selecting an anti-tumor necrosis factor agent (Choice study). *Gut Liver* 2016;**10**(3):391–8. Doi: 10.5009/gnl15126.
22. Vavricka SR., Bentele N., Scharl M., Rogler G., Zeitz J., Frei P., et al. Systematic assessment of factors influencing preferences of Crohn's disease patients in selecting an anti-tumor necrosis factor agent (CHOOSE TNF TRIAL). *Inflamm Bowel Dis* 2012;**18**(8):1523–30. Doi: 10.1002/ibd.21888.

23. The PREFER consortium. *PREFER recommendations: Why, when and how to assess and use patient preferences in medical product decision-making*. 2022.
24. Bridges JFP., Hauber AB., Marshall D., Lloyd A., Prosser LA., Regier DA., et al. Conjoint analysis applications in health - A checklist: A report of the ISPOR Good Research Practices for Conjoint Analysis Task Force. *Value in Health* 2011. Doi: 10.1016/j.jval.2010.11.013.
25. Wurtz CB. Quantitative Sampling. *Nautilus* 2016;**73**(4):131–5.
26. McMillan SS., King M., Tully MP. How to use the nominal group and Delphi techniques. *Int J Clin Pharm* 2016;**38**(3):655–62. Doi: 10.1007/s11096-016-0257-x.
27. Chew LD., Bradley KA., Boyko EJ. Brief questions to identify patients with inadequate health literacy. *Fam Med* 2004.
28. Gale NK., Heath G., Cameron E., Rashid S., Redwood S. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol* 2013. Doi: 10.1186/1471-2288-13-117.
29. Durosini I., Janssens R., Arnou R., Veldwijk J., Smith MY., Monzani D., et al. Patient Preferences for Lung Cancer Treatment: A Qualitative Study Protocol Among Advanced Lung Cancer Patients. *Front Public Health* 2021;**9**. Doi: 10.3389/fpubh.2021.622154.
30. Petrocchi S., Janssens R., Oliveri S., Arnou R., Durosini I., Guidi P., et al. What Matters Most to Lung Cancer Patients? A Qualitative Study in Italy and Belgium to Investigate Patient Preferences. *Front Pharmacol* 2021;**12**. Doi: 10.3389/fphar.2021.602112.
31. Janssens R., Lang T., Vallejo A., Galinsky J., Plate A., Morgan K., et al. Patient Preferences for Multiple Myeloma Treatments: A Multinational Qualitative Study. *Front Med (Lausanne)* 2021;**8**. Doi: 10.3389/fmed.2021.686165.
32. Spagnuolo R., Iaquinta FS., Mauro D., Pantano I., Dastoli S., Naty S., et al. Satisfaction in Social Roles and Physical Function in Immune-mediated Inflammatory Diseases: A Cross-Sectional Study. *Rev Recent Clin Trials* 2022;**17**. Doi: 10.2174/1574887117666220531162104.
33. Iaquinta FS., Grembiale RD., Mauro D., Pantano I., Naty S., Cosco C., et al. Fatigue and Associated Factors in an Immune-Mediated Inflammatory Disease Population: A Cross-Sectional Study. *J Clin Med* 2022;**11**(9). Doi: 10.3390/jcm11092455.
34. Dudley-Brown S. Living with ulcerative colitis. *Gastroenterology Nursing* 1996. Doi: 10.1097/00001610-199603000-00004.
35. Sewitch MJ., Abrahamowicz M., Bitton A., Daly D., Wild GE., Cohen A., et al. Psychological distress, social support, and disease activity in patients with inflammatory bowel disease. *American Journal of Gastroenterology* 2001;**96**(5):1470–9. Doi: 10.1016/S0002-9270(01)02363-2.
36. Coulter A., Härter M., Moumjid-Ferdjaoui N., Perestelo-Perez L., van der Weijden T. *European Experience with Shared Decision Making*. 2015.

37. Stiggelbout AM., Pieterse AH., de Haes JCJM. Shared decision making: Concepts, evidence, and practice. *Patient Educ Couns* 2015;**98**(10):1172–9. Doi: 10.1016/j.pec.2015.06.022.
38. van Overbeeke E., Janssens R., Whichello C., Bywall KS., Sharpe J., Nikolenko N., et al. Design, conduct, and use of patient preference studies in the medical product life cycle: A multi-method study. *Front Pharmacol* 2019;**10**. Doi: 10.3389/fphar.2019.01395.
39. van Overbeeke E., Forrester V., Simoens S., Huys I. Use of Patient Preferences in Health Technology Assessment: Perspectives of Canadian, Belgian and German HTA Representatives. *Patient* 2021;**14**(1):119–28. Doi: 10.1007/s40271-020-00449-0.
40. Crohn's & Colitis Foundation of America. The facts about Inflammatory Bowel Disease n.d.

Accepted Manuscript

13 Figures




Figure 1: Design of the qualitative study in three phases, where each phase informed the subsequent phase. FGD, Focus Group Discussion; IBD, Inflammatory Bowel Disease; NGT, Nominal Group Technique; EMA, European Medicines Agency.

14 Tables

Table 1: Treatment experience of participants (n=11)

Medicinal product classes*	Number of participants taking this class of medication at time of participation	Number of participants who have taken this class of medication (past and present)
Aminosalicylates	2 (18%)	6 (55%)
Corticosteroids	2 (18%)	11 (100%)
Immunosuppressants	1 (9%)	8 (73%)
Biologicals	8 (73%)	9 (82%)
Janus kinase inhibitors	1 (9%)	1 (9%)

* Taken alone or in combination with other medicines from the mentioned classes

Accepted Manuscript

Table 2: Top ten characteristics identified in the attribute grading exercise.

Rank	Characteristic	Mean score	Standard deviation	Maximal score	Minimal score
1	Prevent surgery	9,55	0,82	10	8
2	Long-term clinical remission	9,45	1,21	10	6
3	Improved quality of life	9,36	1,03	10	7
4	Occurrence of urgency	9,27	1,01	10	7
5	Improved labor rate	9,27	1,10	10	7
6	Occurrence of diarrhea	9,18	1,08	10	7
7	Occurrence of severe infections	9,18	1,25	10	6
8	Prevent hospitalization	9,09	1,45	10	6
9	Occurrence of joint pain	9,00	1,26	10	6
10	Prevent flare-ups	9,00	1,79	10	4