NANO EXPRESS

Magic Mathematical Relationships for Nanoclusters - Errata and Addendum

Forrest H Kaatz[1*] and Adhemar Bultheel[2]

[*]Correspondence:
fhkaatz@gmail.com
[1]Mesalands Community College,
911 South 10th Street,
Tucumcari, NM, 88401,
Full list of author information is
available at the end of the article

Abstract

We correct magic formulas for body centered cubic (bcc) structures. The logical rational for this is further corroborated by calculations of the radial distribution function (RDF) for several crystal structures. We add results for truncated cubes which may be found in nature.

Keywords: nanoclusters; topological indices; coordination; magic numbers

Introduction

We recently presented magic formulas for several crystal nanoclusters [1]. However, it is known to crystallographers that bcc structures have a bulk coordination of eight. The RDF determines nearest neighbor peaks from a central point and the integrated peak intensity reflects the corresponding coordination for those neighbors. We use an established method [2] to calculate the RDF for several crystals. Since ideal bcc cubes have coordination $cn = 1$, we provide results for truncated bcc and face centered cubic (fcc) clusters.

Main text

In reviewing the many magic formulas appearing in [1], it occurred to us that equation (1) below, which defines the adjacency matrix, depends on the crystal structure.

$$\mathbf{A}(i,j) = \begin{cases} 1 & \text{if } r_{ij} < r_c \text{ and } i \neq j \\ 0 & \text{otherwise} \end{cases} \tag{1}$$

Here r_{ij} is the Euclidean distance between atom i and atom j. While it is true that $r_c = 1.32{\cdot}r_{min}$ is necessary for the different bonds lengths in the dodecahedral structure, for the bcc structure this is not the case. We have calculated [2] the RDF for selected structures, and some of the nearest neighbors are tabulated below. The RDF has peak locations at neighbor sites, and the integrated intensity of the corresponding peak gives the coordination. We normalize the peaks in $R(r)$ by dividing by the first peak, thus the peak locations become dimensionless. As the table indicates, bcc structures have $r_c = 2/\sqrt{3} \cdot r_{min} \approx 1.15 \cdot r_{min}$, which means the adjacency matrix must be changed, and thus the magic formulas. Note that neighbor peaks are not the same as shells, which give rise to the 'magic numbers'. The dodecahedron is a complicated case, where the third neighbors appear at $r_2 = 1.31 \cdot r_{min}$. This case is challenging, and requires more analysis, which is in progress. The corrected bcc results are shown below. These results agree with those in van Hardeveld and Hartog [3] if one shifts the index by one,

i.e., we use the sequence 0,1,2... and they use 1,2,3... as their sequence. While perfect cubes may be of interest mathematically, they are not likely to appear in nature, due to single bonds at the corners. We have therefore generated truncated bcc and fcc cubes with the corners removed and their results are included here.

Table 1: Neighbor peaks in the normalized RDF for several structures

Structure	2nd Nearest Neighbor	3rd Nearest Neighbor	4th Nearest Neighbor
FCC	1.41	1.73	2.00
BCC	1.15	1.64	1.91
Hexagonal	1.41	1.73	1.91
Diamond	1.64	1.91	2.31
Simple Cubic	1.41	1.73	2.00
Tetrahedron	1.41	1.73	2.00
Decahedron	1.41	1.73	1.91
Icosahedron	1.41	1.69	1.91
Dodecahedron	1.17	1.31	1.37

Table 2: Magic formulas for the bcc cube



bcc cube $n = 3$

	bcc Cube
Atoms	$2n^3 + 3n^2 + 3n + 1,\ n \geq 1$
Bonds	$8n^3,\ n \geq 1$
$cn = 1$	$8,\ n \geq 1$
$cn = 2$	$12n - 12,\ n \geq 1$
$cn = 4$	$6n^2 - 12n + 6,\ n \geq 1$
$cn = 8$	$2n^3 - 3n^2 + 3n - 1,\ n \geq 1$

Table 3: Magic formulas for the bcc octahedron



bcc octahedron $n = 3$

bcc Octahedron	
Atoms	$\frac{8}{3}n^3 + 6n^2 + \frac{16}{3}n + 1,\ n \geq 1$
Bonds	$\frac{32}{3}n^3 + 12n^2 + \frac{28}{3}n,\ n \geq 1$
$cn = 4$	$4n^2 + 4n + 6,\ n \geq 1$
$cn = 6$	$12n - 12,\ n \geq 1$
$cn = 7$	$8n^2 - 16n + 8,\ n \geq 1$
$cn = 8$	$\frac{8}{3}n^3 - 6n^2 + \frac{16}{3}n - 1,\ n \geq 1$

Table 4: Magic formulas for the bcc truncated octahedron



bcc truncated octahedron $n = 2$

bcc Truncated Octahedron	
	n odd
Atoms	$8n^3 + \frac{9}{2}n^2 + \frac{5}{2},\ n \geq 1$ odd
Bonds	$32n^3 - 6n^2 + 6,\ n \geq 1$ odd
$cn = 4$	$9n^2 + 5, n \geq 1$ odd
$cn = 6$	$6n - 6,\ n \geq 1$ odd
$cn = 7$	$12n^2 - 12n, n \geq 1$ odd
$cn = 8$	$8n^3 - \frac{33}{2}n^2 + 6n + \frac{7}{2},\ n \geq 1$ odd
	n even
Atoms	$8n^3 + \frac{9}{2}n^2 + 3n + 1,\ n \geq 2$ even
Bonds	$32n^3 - 6n^2,\ n \geq 2$ even
$cn = 2$	6n+12, $n \geq 2$ even
$cn = 3$	$12n - 24,\ n \geq 2$ even
$cn = 4$	$9n^2 - 18n + 14, n \geq 2$ even
$cn = 6$	$6n, n \geq 2$ even
$cn = 7$	$12n^2 - 12n, n \geq 2$ even
$cn = 8$	$8n^3 - \frac{33}{2}n^2 + 9n - 1,\ n \geq 2$ even

Table 5:
Magic formulas for the bcc Rhombic Dodecahedron



bcc Rhombic Dodecahedron $n = 2$

bcc Rhombic Dodecahedron	
Atoms	$4n^3 + 6n^2 + 4n + 1,\ n \geq 1$
Bonds	$16n^3 + 12n^2 + 4n,\ n \geq 1$
$cn = 4$	$14,\ n\ \geq 1$
$cn = 5$	$24n - 24\ n\ \geq 1$
$cn = 6$	$12n^2 - 24n + 12,\ n\ \geq 1$
$cn = 8$	$4n^3 - 6n^2 + 4n - 1,\ n\ \geq 1$

Table 6: Magic formulas for the bcc cuboctahedron



bcc cuboctahedron $n = 2$

bcc Cuboctahedron	
Atoms	$\frac{5}{3}n^3 + 7n^2 + \frac{34}{3}n + 7,\ n \geq 1$ odd
	$\frac{5}{3}n^3 + 7n^2 + \frac{25}{3}n + 1,\ n \geq 2$ even
Bonds	$\frac{20}{3}n^3 + 19n^2 + \frac{64}{3}n + 9, n \geq 1$ odd
	$\frac{20}{3}n^3 + 19n^2 + \frac{46}{3}n,\ n \geq 2$ even
$cn = 2$	$12,\ n \geq 1$ odd; 0, n even
$cn = 3$	$12n - 12,\ n \geq 1$ odd; 0, n even
$cn = 4$	$4n^2 - 4n + 6,\ n \geq 1$ odd
	$4n^2 + 8n,\ n \geq 2$ even
$cn = 6$	$0,\ n \geq 1$ odd; 12, n even
$cn = 7$	$2n^2 + 4n + 2,\ n \geq 1$ odd
	$2n^2 + 4n - 16,\ n \geq 2$ even
$cn = 8$	$\frac{5}{3}n^3 + 1n^2 - \frac{2}{3}n - 1,\ n \geq 1$ odd
	$\frac{5}{3}n^3 + 1n^2 - \frac{11}{3}n + 5,\ n \geq 2$ even

Table 7: Magic formulas for the bcc Truncated Cube



truncated bcc cube $n = 3$

bcc Truncated Cube	
Atoms	$2n^3 + 3n^2 + 3n - 7,\ n \geq 2$
Bonds	$8n^3 - 8,\ n \geq 2$
$cn = 2$	$12n - 12,\ n \geq 2$
$cn = 4$	$6n^2 - 12n + 6,\ n \geq 2$
$cn = 7$	$8,\ n \geq 2$
$cn = 8$	$2n^3 - 3n^2 + 3n - 9,\ n \geq 2$

Table 8: Magic formulas for the fcc Truncated Cube



truncated bcc cube $n = 3$

fcc Truncated Cube	
Atoms	$4n^3 + 6n^2 + 3n - 7,\ n \geq 2$
Bonds	$24n^3 + 12n^2 - 24,\ n \geq 2$
$cn = 5$	$12n - 12,\ n \geq 2$
$cn = 7$	$24,\ n \geq 2$
$cn = 8$	$12n^2 - 12n - 18,\ n \geq 2$
$cn = 12$	$4n^3 - 6n^2 + 3n - 1,\ n \geq 2$

Table 9: Magic Topological Formulas for BCC and FCC Clusters

Index	Formula
	bcc Cube
Wiener	$\frac{76}{35}n^7+\frac{38}{5}n^6+\frac{74}{5}n^5+18n^4+\frac{206}{15}n^3+\frac{32}{5}n^2+\frac{136}{105}n$
Reverse Wiener	$\frac{64}{35}n^7+\frac{22}{5}n^6+\frac{31}{5}n^5+2n^4-\frac{26}{15}n^3-\frac{17}{5}n^2-\frac{136}{105}n$
HyperWiener	$\frac{47}{35}n^8+\frac{226}{35}n^7+\frac{469}{30}n^6+\frac{241}{10}n^5+\frac{119}{5}n^4+\frac{149}{10}n^3+\frac{1097}{210}n^2+\frac{19}{35}n$
Szeged	NA
	bcc Rhombic Dodecahedron
Wiener	$\frac{76}{7}n^7+38n^6+\frac{302}{5}n^5+56n^4+\frac{94}{3}n^3+10n^2+\frac{148}{105}n$
Reverse Wiener	$\frac{148}{7}n^7+58n^6+\frac{378}{5}n^5+48n^4+\frac{38}{3}n^3-2n^2-\frac{148}{105}n$
HyperWiener	$\frac{359}{42}n^8+\frac{832}{21}n^7+\frac{1217}{15}n^6+\frac{1454}{15}n^5+73n^4+\frac{103}{3}n^3+\frac{1957}{210}n^2+\frac{39}{35}n$
Szeged	$\frac{4637}{105}n^9+\frac{15655}{84}n^8+\frac{7661}{21}n^7+\frac{2615}{6}n^6+\frac{5194}{15}n^5+\frac{2245}{12}n^4+\frac{1412}{21}n^3+\frac{103}{7}n^2+\frac{32}{21}n$
	bcc Truncated Cube
Wiener	$\frac{76}{35}n^7+\frac{38}{5}n^6+\frac{74}{5}n^5-6n^4-\frac{394}{15}n^3-\frac{168}{5}n^2+\frac{4336}{105}n$
Reverse Wiener	$\frac{64}{35}n^7+\frac{22}{5}n^6+\frac{31}{5}n^5-6n^4-\frac{146}{15}n^3-\frac{57}{5}n^2+\frac{1544}{105}n$
HyperWiener	$\frac{47}{35}n^8+\frac{226}{35}n^7+\frac{469}{30}n^6+\frac{49}{10}n^5-\frac{121}{5}n^4-\frac{1313}{30}n^3+\frac{4457}{210}n^2+\frac{1933}{105}n$
Szeged	NA
	fcc Truncated Cube
Wiener	$\frac{956}{105}n^7+\frac{478}{15}n^6+\frac{1357}{30}n^5+\frac{110}{3}n^4+\frac{589}{30}n^3+\frac{97}{15}n^2+\frac{36}{35}n$
Reverse Wiener	$\frac{1564}{105}n^7+\frac{602}{15}n^6+\frac{1343}{30}n^5+\frac{70}{3}n^4+\frac{43}{15}n^3-\frac{59}{30}n^2-\frac{36}{355}n$
HyperWiener	$\frac{59}{10}n^8+\frac{2956}{105}n^7+\frac{1089}{20}n^6+\frac{701}{12}n^5+\frac{817}{20}n^4+\frac{1153}{60}n^3+\frac{53}{10}n^2+\frac{5}{7}n$
Szeged	$\frac{14822}{945}n^9+\frac{2099}{35}n^8+\frac{30781}{315}n^7+\frac{941}{10}n^6+\frac{1073}{18}n^5+\frac{251}{10}n^4+\frac{12629}{1890}n^3+\frac{29}{35}n^2+\frac{32}{105}n$

Conclusions

We have corrected magic formulas for bcc structures and added results from the RDF and for truncated bcc and fcc cubes.

Abbreviations

bcc: body centered cubic; fcc: face centered cubic; RDF: radial distribution function

Acknowledgments

We made use of the MATLAB file, Cluster Generator, which can be found in Mathworks File Exchange Central.

Funding

The authors did not receive external funding for this work.

Availability of Data and Materials

The dataset(s) supporting the conclusions of this article may be obtained from the corresponding author.

Author Contributions

FHK conceived of the project and analysis. AB wrote the code in MATLAB. Both authors contributed to writing the paper, and approved the final version of the manuscript.

ORCIDs
FHK ORCID: 0000-0001-7912-0627, AB ORCID: 0000-0001-9562-5297

Conflict of Interest Statement
The authors declare that they have no competing interests..

Author details
[1]Mesalands Community College, 911 South 10th Street, Tucumcari, NM, 88401,. [2]Dept. Computer Sci., KU Leuven, Celestijnenlaan 200A, 3001, Heverlee, Belgium.

References

1. Kaatz, F.H., Bultheel, A.: Magic mathematical relationships for nanoclusters. Nanoscale Res. Letters **14**(1), 150 (2019)
2. Juhas, P., Farrow, C.L., Yang, X., Knox, K.R., Billinge, S.J.L.: Complex modeling: a strategy and software program for combining multiple information sources to solve ill posed structure and nanostructure inverse problems. Acta Cryst. **A71**, 562–568 (2015)
3. van Hardeveld, R., Hartog, F.: The statistics of surface atoms and surface sites on metal clusters. Surf. Sci. **15**, 189–230 (1969)