

Manuscript

Intraspecific genetic variation is critical to robust toxicological predictions in *Daphnia*

René S. Shahmohamadloo^{*a,b}, Seth M. Rudman^{*a,b}, Catherine I. Clare^a, Judy A. Westrick^c, Luc De Meester^d, John M. Fryxell^b

^a School of Biological Sciences, Washington State University, Vancouver, Washington, 98686, United States

^b Department of Integrative Biology, University of Guelph, Guelph, Ontario, N1G 2W1, Canada

^c Department of Chemistry, Wayne State University, Detroit, Michigan, 48202, United States

^d Laboratory of Aquatic Ecology, Evolution, and Conservation, University of Leuven Charles Deberiotstraat 32, 3000 Leuven, Belgium

*These authors contributed equally to this work.

Address co-correspondence to:

René S. Shahmohamadloo, PhD

Liber Ero Postdoctoral Fellow, NSERC Postdoctoral Fellow

School of Biological Sciences | Washington State University

14204 NE Salmon Creek Avenue | Vancouver | Washington | 98686 | United States

Email: rene.shahmohamadloo@wsu.edu

Address co-correspondence to:

Seth M. Rudman, PhD

Assistant Professor

School of Biological Sciences | Washington State University

14204 NE Salmon Creek Avenue | Vancouver | Washington | 98686 | United States

Email: seth.rudman@wsu.edu

Author Contributions: R.S.S., S.M.R., L.D.M., and J.M.F. designed the research. R.S.S. and S.M.R. conducted the experiments. S.M.R., R.S.S., C.I.C., and J.A.W. analyzed the data. R.S.S. and S.M.R. wrote the paper. All authors read, amended, and approved the final manuscript.

Competing Interest Statement: The authors declare no competing interests.

Keywords: *Risk assessment, Genetic variation, Toxicology, Daphnia, Evolutionary toxicology*

Summary

Invertebrate toxicant exposure reveals substantial intrapopulation variation emphasizing the critical importance of accounting for intraspecies genetic variation in toxicity testing.

Abstract

Environmental risk assessment is a critical tool for protecting biodiversity and its effectiveness is predicated on predicting how natural populations respond to environmental stressors. Yet, routine toxicity testing typically examines only one genotype, which may render risk assessments inaccurate at the population scale. To determine the importance of intraspecific variation in the translation of toxicity testing to populations, we quantified the magnitude of genetic variation within 20 *Daphnia magna* clones derived from one lake using whole genome sequencing and phenotypic assays. We repeated these assays across two exposure levels of *Microcystis aeruginosa*, a cosmopolitan freshwater contaminant. We found considerable intraspecific genetic variation in survival, growth, and reproduction. Exposure to *Microcystis* amplified the degree of intraspecific variation. Simulations demonstrate assays employing one clone failed to produce an estimate within the 95% confidence interval over half of the time. These results illuminate the importance of incorporating intraspecific genetic variation, but not necessarily genome sequences, into toxicity testing to reliably predict how natural populations will respond to environmental stressors.

Main Text

Introduction

Reliable translation of toxicity testing into predictions of population-level consequences in nature is a cornerstone of environmental risk assessments^{1–6}. Such predictions are classically obtained by quantifying individual phenotypic responses under constant conditions in a laboratory⁷. Research has historically favored single-species toxicity testing that generates point estimates such as the lethal concentration at which a contaminant will kill 50% of a test population (LC50)⁸. For more than fifty years, regulatory agencies have relied on such lethality data to inform environmental risk assessments, mandated to safeguard biodiversity and associated ecosystem services by protecting rare or economically significant species⁶. Yet, reliably translating laboratory toxicity studies into population-level predictions requires building on traditional methods to incorporate aspects that are key to ecological realism⁸. Two particularly important, yet understudied aspects are the sublethal effects of toxicants on organismal life-history traits and intraspecific genetic variation in toxicological responses.

Genetic variation within species is recognized as a critical driver of both ecological and evolutionary processes^{9–11}. In ecological contexts, genetic variation begets variation in functional traits, which can have profound impacts on populations, communities, and ecosystems^{9–12}. Genetic variation within species is central to evolution, as the process of natural selection shaping genetic variation is what produces adaptation^{10,13}. Among the most visible and societally challenging examples of rapid adaptation are cases where populations adapt to agrochemicals¹⁴ or contaminants¹⁵. Despite an appreciation of the role of adaptation in resistance and some demonstrations of the amount of standing variation in tolerance^{4,5,16–18}, recognition of the importance of intraspecific genetic variation continues to be ignored by

regulators within the discipline of toxicology and, more broadly, the environmental risk assessment process.

Investigating the magnitude of intraspecific variation within a population represents a crucial and understudied element in toxicology. Little is known about the magnitude and ecological importance of intraspecific genetic variation in toxicological responses within natural populations. Choosing a genotype based on availability (*e.g.*, standard lab strain) may lead to inferences that are unrepresentative of imperiled populations. More broadly, conducting measurements on any single genotype is using a single point to estimate a distribution that likely has substantial variance^{19,20}. Detailed study of specific populations that includes both measures of the extent of genetic variation within focal populations and classical toxicity testing are needed to infer the potential for this variation to impact population-level responses under toxic insults²¹.

Daphnia magna (water fleas) are a powerful eco-genomic model system because they are simple to culture, have a fully-sequenced genome, and they play an important and well-documented ecological role in many freshwater ecosystems^{22,23}. With short generation times and considerable genetic diversity derived from diapausing eggs in sediments, *Daphnia* can undergo rapid adaptation in response to environmental variation^{22–26}. These eco-evolutionary insights have not translated to change in routine laboratory testing of *Daphnia*, which is a model system used in toxicology to test chemical compounds before they are produced, released or sold²³. Toxicologists previously advocated that toxicological studies should account for genetic variation in *Daphnia* based on discrepancies between several laboratory strains that were exposed to the same reference toxicants^{2,3,7,27}. Yet, determining the loss in accuracy associated with choosing a single clone from a natural genetically heterogeneous population for environmental risks assessments is crucial to demonstrating the overall importance of accounting for genetic variation.

Determining the magnitude of variation within a population is crucial both for predicting immediate toxicological responses and as an indicator of the potential for rapid adaptation. Doing so requires accounting for the scale and scope of genetic diversity within a population and conducting toxicity testing on a heterogeneous and realistic population of *Daphnia*. Genomic data can be useful in meeting these objectives as genome-wide data provide information on the scale and scope of genetic diversity being studied and can be translated to provide population-level insight. For investigations on the relationship between genetic diversity and toxicological responses whole genome sequencing data yields an ability to identify populations or clones and compare the extent to which they are genetically divergent, to determine whether the degree of genomic differentiation between populations is correlated with toxicological responses, and to test for relationships between genomic diversity and tolerance. Depending on the underlying genetic architecture of variation in toxin tolerance it may be possible to identify the genetic basis of resistance²⁸, though finding a ‘simple’ genetic basis may not be common²⁹. At a minimum, incorporating population genetic information can help standardized toxicity tests and the information generated also has the potential to better predict the risk of toxicant exposure on populations, and increase predictability on the associated community and ecosystem functions.

To determine the importance of intraspecific variation in toxicity projections key to the environmental risk assessment process, we sequenced the genomes of 20 *Daphnia magna* clones collected from a single Belgian lake and quantified the chronic effects of the

cyanobacterium *Microcystis* on their life-history traits (survival, growth, reproduction, broods). Harmful algal blooms (HABs) of *Microcystis* are a prominent and toxic ecological disturbance that strongly impact a wide-range of taxa^{30–35}. As a keystone zooplankton grazer, *Daphnia* shows substantial intraspecific diversity and rapid adaptation in response to HABs^{24,36,37}, including some evidence that particular genotypes can effectively feed on and reduce the abundance of toxic *Microcystis*^{38,39}. Determining whether intraspecific variation in toxin tolerance influences vital demographic rates and population dynamics by *Daphnia* clarifies the role of intraspecific genetic variation in toxicology, with implications for regulatory agencies tasked with maintaining biodiversity under toxic bloom conditions.

In this study, 30 individuals per genotype were divided and exposed to three treatments (i.e., common gardens): 100% green algae (*Chlorella vulgaris*, non-toxic diet), 2:1 chlorella:microcystis (a moderately toxic diet), and 1:1 chlorella:microcystis (a severely toxic diet). Common garden rearing keeps environmental conditions constant allowing for the estimation of the contribution of genetic variation to phenotypic variation. We chose 2:1 and 1:1 toxic treatments that resulted in microcystin concentrations of $3.3 \pm 0.001 \mu\text{g L}^{-1}$ and $5.1 \pm 0.001 \mu\text{g L}^{-1}$, respectively, which have been considered sublethal⁴⁰ to lethal^{41,42} in *Daphnia* laboratory assays and are common ratios in freshwater ecosystems with HABs³². We sought to address the following four questions: 1) How much intraspecific genomic and phenotypic variation occurs within a single population of *D. magna*? 2) Does exposure to toxic cyanobacteria reduce the magnitude of intraspecific phenotypic variation? 3) Is there an interaction between clonal growth and reproduction under favorable (green algae) and toxic (cyanobacteria) conditions? 4) Using data from these trials, how would sampling design influence the interpretation of reproductive output with and without contaminant exposure? We predicted that *D. magna* clones would exhibit considerable genetic and phenotypic intraspecific variation in toxicological responses, as *Daphnia* populations have been shown to be variable. Additionally, we predicted that phenotypic variation would be reduced when exposed to a toxic insult due to a decrease in trait values in high performing genotypes. Moreover, we predicted a trade-off between growth and reproduction under favorable and toxic conditions, with clones having the greatest reproductive output varying across cyanobacterial concentrations. Finally, we predicted that sampling a sufficient number of genotypes would be critical to producing an accurate estimate of reproduction in both favorable and toxic conditions.

Results

Intraspecific genomic and phenotypic variation

Genetic variation among *D. magna* clones was considerable in both phenotype and genotype (Figure 1). When reared on a *Chlorella*-only non-toxic diet, phenotypic variation among clones was significant for survival ($X^2 = 4.52$, $P = 0.033$), growth ($X^2 = 67.40$, $P < 0.0001$), and reproduction ($X^2 = 40.22$, $P < 0.0001$). The magnitude of genome-wide divergence between pairwise sets of clones varied (Figure 1a), with the most similar clones having $\sim 1.2\text{M}$ SNP differences and the most divergent $\sim 1.6\text{M}$.

We tested for an association between the degree of genomic divergence between clones and the magnitude of their phenotypic differentiation. We calculated a phenotypic distance between each pair of clones based on results of phenotypic assays for survival, growth, time to reproduction, and total reproduction when reared on non-toxic, moderately toxic, and severely toxic diets (Supporting Information, Figure S1). We found no significant association between the

degree of genomic divergence genome-wide and the magnitude of overall phenotypic divergence between clones (Figure 1) (mantel $r = 0.033$, $P = 0.781$). We also examined the relationship between genetic diversity found within clones (i.e., total heterozygosity) and clonal survival and reproduction (Supp Fig XX). We found no significant association between the amount of genetic diversity within clones and their ability to survive or reproduce across diet conditions.

Magnitude of phenotypic variation from toxic exposure

Exposure to cyanobacteria reduced the overall fitness of *D. magna* genotypes with increasing severity in the moderately toxic (2:1 *Chlorella* : *M. aeruginosa*) and severely toxic (1:1 *Chlorella* : *M. aeruginosa*) treatments (Figure 2). For example, mean survival across all clones for the 14-day trial declined from 94% in the non-toxic diet, to 86.5% in the moderately toxic diet, and to 53% in the severely toxic diet. We tested for differences in the magnitude of clonal variation with increasing exposure to *M. aeruginosa* for each of the three fitness-associated phenotypes (survival, growth, reproduction). Intraspecific variation in survival did not increase from the control to 2:1 ($F^* = 3.60$, $P = 0.066$) but did from 2:1 to 1:1 ($F^* = 22.59$, $P < 0.001$). Intraspecific variation in growth rate increased from the control to 2:1 ($F^* = 10.28$, $p = 0.003$) but decreased from 2:1 to 1:1 ($F^* = 11.82$, $P = 0.002$). Neonate production showed the most consistent pattern, with intraspecific variation between clones declining with increasing *M. aeruginosa* exposure (control to 2:1 ($F^* = 18.54$, $P = 0.0001$); 2:1 to 1:1 ($F^* = 7.44$, $P = 0.009$)).

Interactions under favorable and toxic conditions

We tested for interactions associated with performance across variations in diet toxicity. On average, survival declined with increasing toxicity by 8% in the moderately-toxic diet and 44% in the severely toxic diet relative to controls. However, we found evidence of an interaction between clonal identity and a *Microcystis* diet ($X^2 = 59.68$, $P = 0.014$). Some of these interactive effects were striking. Relative to rearing on the *Chlorella* diet, clones 5 and 15 showed an 80% decline in mean survival when reared on the severely toxic diet while clone 19 exhibited a 20% increase in survival. Similarly, we found a significant interaction associated with growth across treatments ($X^2 = 244.17$, $P < 0.0001$). Finally, neonate production showed a similar pattern, with a significant interaction between clonal identity and a *Microcystis* diet ($X^2 = 264.42$, $P < 0.0001$). Of particular interest was that certain genotypes performed better in the toxic treatments compared to the *Chlorella* diet (e.g., genotypes 6 and 13 across all life-history traits), while other genotypes performed better in the 1:1 versus 2:1 toxic treatments (e.g., genotypes 10, 14, 18, 19, 20). These findings indicate that relative performance of clones is dependent on the magnitude of toxicant exposure. reinforce the shifts in relative performance across clones associated with changing *M. aeruginosa* conditions.

Influence of genetic diversity on toxicological inferences

We used a bootstrapping approach to determine how variation in the number of clones in the experimental sampling design could influence the robustness of the data relative to including 20 clones. Our results demonstrate that sampling individuals from a larger number of genotypes is critical to generating a robust estimate of reproduction in all 3 rearing conditions (Figure 3a). To quantify the loss of robustness and accuracy that would result from assaying a reduced amount of genetic diversity we assessed how frequently samples at five reduced diversity levels would yield survival estimates within the 95% confidence interval (CI) obtained using all 20 clones

(Figure 3b). While including 15 clones always produced a survival estimate with the 95% CI assays using only a single clone failed to produce an estimate within the 95% CI over half the time in all three toxicity treatments. Overall, this reinforces the importance of accounting for genetic diversity in toxicity testing, as failing to do so means estimates of reproduction and survival are likely to be highly variable and inaccurate.

Discussion

The impact of intraspecific variation on toxicological inference and management

Genetic variation can meaningfully impact population dynamics⁴³, community composition⁴⁴, ecosystem function⁴⁵, and underlie rapid adaptation¹⁰. Incorporating natural genetic variation in toxicology, which directly informs environmental regulations and policy, is critical to increasing the reliability of benchtop toxicity assays to natural populations. Our data demonstrate that genetic variation, even within a population from a single locality, has a profound impact on toxicological responses in a widely used model system. For example, the moderately toxic diet used in this study could be interpreted as an LC50 (i.e., clone 4) or an LC0 (clone 18 and 19) depending on the clone selected for the assay. When reared on a severely toxic diet, mortality ranged from 0% to 90%. Toxicity studies that do not include intraspecific genetic variation necessarily cannot estimate this variation or provide context on whether the genotype or strain used in the assay is in any way representative of the species or genus. This can lead to considerable uncertainty when extrapolating results from laboratory to field, and have implications for the accuracy and efficacy of risk management strategies⁴⁶. The ramifications of this oversight are also pertinent in cases of pesticide resistance evolution, where natural selection acting on genetic variation can profoundly decrease the efficacy of agrochemicals⁴⁷.

The variation we observed came from a single population of *D. magna* and is doubtless an underestimate of the magnitude of variation that would be found across populations that experience dramatically different exposures to HABs. For example, if the species of focus spans a wide range of ecological conditions, particularly with regards to prior toxicant exposure that may have driven adaptation, there is likely to be substantially more intraspecific variation than we observed within the single population we studied here. Hence, we advocate for the inclusion of multiple genotypes in clonal test organisms or individuals from multiple populations in sexual organisms when conducting toxicity studies. Ideally, studies should survey across a sufficiently wide array of genetic diversity that is similar to populations that could be impacted by the toxicant to allow 'clone' or 'population origin' to be treated as a random effect in linear mixed models and hence generate reasonable estimates of the variation in key response variables across the populations of interest⁴⁸.

In practice, we recognize that such robust sampling designs may be challenging and would require shifting effort and focus in many toxicology studies. As a preliminary step, it must be acknowledged by toxicology regulatory agencies that LC50s derived from real, genetically heterogeneous populations will differ at local, regional, or global scale. Each geographic scale presents its own sources of variation and shapes the requirements in terms of the sampling required to achieve the stated objective of a meaningful safety limit recommendation based on an LC50. Therefore, testing multiple genotypes within a single population (local/regional scale) or multiple populations of genotypes (international scale) and reporting a CI of LC50s calculated across each would be a strong step forward. Doing so would provide both a more accurate and robust measure of the LC50, as it would be calculated from multiple genotypes or populations

and an indication of the uncertainty. The uncertainty of this estimate would not arise from experimental noise but from real differences in biology. Reporting LC50s would then include both the estimate and the variance around this estimate stemming from genetic variation. This would be useful in discussions about risk and the connection between laboratory assays and toxicological outcomes in natural settings, a key area of growth in translating benchtop experiments to wild populations.

The study here demonstrates that considerable variation is found in toxicological responses within a population. However, it is important to consider the overall scale and scope of variation in toxicological responses. Current methods often examine focal taxa across broad phylogenetic categories (e.g.,), which is important. But meaningful biological variation can be found across levels of organization. Future work examining variation across biological scales - within populations, across populations, between congeners, and across genera - is critical for parsing the levels of biodiversity most relevant for risk assessment and management action. Identifying the biological scales at which the largest fraction of functional variation is has been transformative in ecology ⁴⁹ and a similar approach could also lead to a major advance in toxicology.

Genomic variation, adaptation, and toxicology

The level of genomic variation observed across clones in our study demonstrates the large magnitude of genetic variation found within a single population of *Daphnia*. From a toxicological perspective this variation is important both as it provides phenotypic variation, as demonstrated here, and this variation is the raw material on which natural selection can act to drive adaptation. Previous work has suggested that rapid adaptation could be a critical mechanism of suppressing *Microcystis* blooms as field populations of *Daphnia* can rapidly adapt to tolerate *Microcystis* and can suppress the biomass of HABs that contain lethal concentrations of microcystins (3 to 6.5 $\mu\text{g L}^{-1}$) ³⁹. These data underscore the need for toxicological theory to integrate genetic diversity, and potentially rapid adaptation in routine testing dose-responses between chronically exposed ^{24,38,39,50} and unexposed ^{3,51} populations might well differ due to genomic variation.

The pace and ecological consequences of resistance evolution has been appreciated for decades ⁵², but a growing body of research is seeking to understand the molecular genetic underpinnings of variation in resistance to toxins ^{15,53,54}. We conducted whole genome sequencing of all 20 *D. magna* clones assayed in the study. These data provide a holistic view of the genetic diversity found within a population but do not provide the power required to discover the genetic basis of clonal differences in resistance to *M. aeruginosa*. This is underscored by the lack of associations both between phenotypic variation and genomic differentiation and genomic variation within clones and phenotypic responses. In short, it appears variation in responses to cyanobacteria are not the primary force shaping genetic variation and the raw amount of genetic variation itself is not a key determinant in performance when exposed to toxicants. These results largely fit with the prevailing population genetic theory for genome wide patterns ⁵⁵. The utility of genomic data in toxicology has been touted as potentially transformative - with a gene-level understanding of resistance presented as a major goal ⁵⁶. It is possible to conduct quantitative trait locus (QTL) mapping or genome wide association studies (GWAS) to find the underlying genomic basis of trait variation. However, generating the resolution needed to identify a causal genomic region, particularly specific genes, would require a herculean effort, a relatively simple genetic basis, and likely some

recombinational luck - efforts most likely to yield payoff only in model systems. Given the complex and environmentally variable genetic architecture of trait variation for many complex traits we advocate that toxicologists focus genetic studies on assessments of clonal, family, or population level variation. Explicitly including this information on origin in study designs and statistical models will yield powerful insights without requiring major changes in the methods most often employed in the discipline.

Conclusion

Studies of genetic variation have improved our understanding of ecological and evolutionary processes⁵³. Arguments in favor of incorporating genetic variation, and to an extent evolution, in the environmental risk assessment process are growing stronger as the recognition that meaningful levels of intraspecific genetic variation exist in key toxicological traits^{21,23,46,57,58}. Ignoring genetic variation can lead to inaccurate reporting of LC50 toxicity thresholds²³, which may obscure safety measures adhered to by companies invested in the commercial production of chemicals and other categories of stakeholders more concerned with toxicants threatening biodiversity. Here, we show that genotypic and phenotypic differences in *D. magna* tolerance to the aquatic contaminant *M. aeruginosa* were substantial across multiple life-history traits. Significant interactions between growth and reproduction were additionally observed among genotypes in green algae versus cyanobacteria treatments. These results suggest that genetic diversity can strongly influence population dynamics with and without toxicant exposure. Using simulations and bootstrapping approaches we further demonstrate that the common toxicological practice of using a single genotype to calculate toxicity tolerance leads to an estimate outside of the 95% confidence interval >50% of the time. We suggest that integrating intraspecific genetic variation into routine laboratory toxicity studies would improve the reliability and biological realism of LC50 calculations and improve both the environmental risk assessment process and our understanding of the transgenerational impacts of contaminants on natural populations.

Methods

Daphnia magna field collection and culturing

Twenty genotypes of *D. magna* were collected from 'Langerodevijver' (LRV; 50° 49' 42.08", 04° 38' 20.60"), a small lake (surface area = 140,000 m², max depth = 1 m) within the nature reserve of Doode Bemde, Vlaams-Brabant, Belgium⁵⁹. LRV has seasonal *Microcystis* HABs typical to many lakes globally and has a large resident population of *D. magna*. Parthenogenetic lines of each genotype were maintained for over five years in continuous cultures at 20 °C in UV-filtered dechlorinated municipal tap water containing 2 mg C L⁻¹ of the green alga *C. vulgaris* (strain CPCC 90; Canadian Phycological Culture Centre, Waterloo, ON, Canada). *C. vulgaris* was grown in COMBO medium⁶⁰.

Microcystis aeruginosa culturing

Following our previously described method⁶¹, *M. aeruginosa* (strain CPCC 300; Canadian Phycological Culture Centre, Waterloo, ON, Canada) was cultured in BG-11 media and kept in a growth chamber under axenic conditions with a fixed temperature of 21 ± 1 °C, cool-white fluorescent light of 600 ± 15 lx, with a photoperiod of 16:8 h light:dark. The culture was grown for a minimum of one month before preparation for the chronic study. *M. aeruginosa* CPCC 300

produces microcystins-LR (CAS: 101043-37-2, $C_{49}H_{74}N_{10}O_{12}$) and its desmethylated form [D-Asp³]-microcystin-LR (CAS: 120011-66-7, $C_{48}H_{72}N_{10}O_{12}$), which occur widely in freshwater ecosystems^{30,32} and are toxic to zooplankton.

To prepare *M. aeruginosa* for testing on *D. magna*, an aliquot of the stock was inoculated in 100% COMBO medium for two weeks prior to test initiation and cultured to a cell concentration of $1.25 \pm 0.02 \times 10^7$ cells mL⁻¹. This medium was chosen because it supports the growth of algae and cyanobacteria and is non-toxic to zooplankton⁶⁰.

Chronic toxicity study

A 14-d chronic toxicity study was performed in accordance with the test guideline of the Organization for Economic Cooperation and Development to assess the effects of contaminants on the survival, growth, and reproductive output of *D. magna*⁶². The OECD test guidelines are widely used in toxicity laboratory studies. Three treatments were included in this study: 100% chlorella (non-toxic diet), 2:1 chlorella:microcystis (moderately toxic diet), and 1:1 chlorella:microcystis (severely toxic diet). Following these ratios all *D. magna* were fed 2 mg C L⁻¹, corresponding to 3×10^6 cells total, which corroborates with previous literature exposing daphnids to dietary combinations of green algae and cyanobacteria^{37,40,63}.

Following the OECD test guidelines, *D. magna* juveniles born within 24 hr were collected from their respective genotype, and 10 juveniles were individually placed in 50-mL glass tubes containing 50 mL of UV-filtered water and its respective treatment. A total of 600 animals were used for this study (i.e., 10 replicates per genotype, 200 animals per treatment). Since this was a semi-static test, solutions were renewed 3 × wk on Mondays, Wednesdays, and Fridays by transferring *D. magna* from old to new glass tubes, followed by supplying each *D. magna* with 3×10^6 cells of food (e.g., the 2:1 treatment received 2×10^6 *C. vulgaris* cells and 1×10^6 *M. aeruginosa* cells, corresponding with 2 mg C L⁻¹). During this time, survival and reproductive output were recorded. Growth (mm) for each replicate was also measured on days 0, 7, and 14 to assess for interactions between genotype and treatment effects. The study was incubated under 400–800 lx cool-white fluorescent light at 20 ± 1 °C with a 16:8 light:dark cycle.

Genomics sequencing and analysis

Pools of individuals were collected from each clone and DNA was extracted using the QIAGEN MagAttract HMW DNA kit. Extractions were checked for DNA concentration using a Qubit 2.0 fluorometer and libraries for each clone were prepared using the KAPA Hyper Prep kit. Sequencing of these libraries was generated on a NovaSeq 6000 generating 150bp paired-end reads. Sequences were then trimmed using *trimmomatic/0.39* using a quality threshold of 20 and a min length of 50 and subsequently mapped to the *D. magna* reference genome (GCA_020631705.2) using *bwa/0.7.17* (default parameters). Duplicates were marked using *Picard tools/2.21.2* and then the bam files were input into *freebayes/1.2.0* to generate a VCF of all variant positions among all individuals on the 10 *D. magna* chromosomes (not scaffolds). *VcfTools* was then used to trim to a minimum quality score of 30. The total number of variant sites and the number of heterozygous sites for each clone was extracted using *VCFtools*. Genetic distance was calculated using *VCF2dis* (<https://github.com/BGI-shenzhen/VCF2Dis>). Genotype likelihood (beagle) files were generated using *angsd* (-GL 1 -out D23Genolike -doMaf 2 -SNP_pval 1e-6 -doMajorMinor 1 -doGlf 2), and *pcangsd* was used to generate principal component analysis.

Water sampling for analysis

Water samples were collected from each treatment at test initiation, solution changes, and termination to quantify the cell concentration, fluorescence, and standard water parameters and were quick-frozen at -80°C before analysis for microcystins and oligopeptides (Supporting Information, Dataset S1).

Chemicals and Reagents for water analysis

Water, acetonitrile, methanol, acetic acid, and formic acid were all Optima LC/MS grade solvents and purchased from Fisher Scientific (Tewksbury, MA, USA). Microcystin standards D-Asp³-RR, MC-RR, YR, HtyR, LR, D-Asp³-LR, HilR, WR, LA, LY, LW, and LF, as well as nodularin and cylindrospermopsin were purchased from Enzo Life Sciences (Farmingdale, NY, USA). Microcystin Dha⁷-LR was purchased from National Research Council Canada (Ottawa, ON, Canada). Microcystin Leu¹-LR was purchased from GreenWater Laboratories (Palatka, FL, USA). D-Asp³-E-Dhb⁷-RR was purchased from Sigma Aldrich. Anabaenopeptin A and B, oscillamide Y, oscillagin A, microginin 690 methyl ester, aeruginosamide B and C, cyanopeptolin 1040 MB and 1007, and micropeptin 1106 were purchased from MARBIONC (Wilmington, NC, USA). Anabaenopeptin F was synthetically created by the Stockill laboratory at Wayne State University (Detroit, MI, USA). ¹⁵N MC-LR was purchased from Cambridge Isotope Laboratories, INC. (Tewksbury, MA, USA).

Microcystin and cyanopeptide water analysis

Microcystin and cyanopeptide standards and samples were analyzed using a Thermo Scientific TSQ Altis™ triple quadrupole mass spectrometer (Waltham, MA, USA) with a TriPlus™ RSH EQuan 850 system. Both online concentration analysis and direct inject analysis were completed (Birbeck et al. 2019). Quantitative results shown use the direct inject analysis method. Detection limits for the online concentration method were between 0.5-5 ng L⁻¹, and the standard curve range was between 0.5-500 ng L⁻¹.

For direct injection analysis, 10 μL injections of standards and samples were loaded onto a Thermo Hypersil GOLD™ C18 2.1 x 50 mm, 1.9 μm column using a TriPlus RSH autosampler equipped with a 100 μL sampling syringe. The analytical separation gradient consisted of 0.1% formic acid in water (mobile phase A) and 0.1% formic acid in acetonitrile (mobile phase B). At a flow rate of 0.5 mL/min, the gradient started at 24% B and was held for 0.7 min. It was then ramped to 26% B from 0.7–1.7 min, from 26–50% B from 1.7–5.5 min. The column was washed at 98% B from 5.51–6.5 min at a flow rate 1.0 mL/min, then brought back down to original conditions to equilibrate. The total run time was 8.5 min. The column oven temperature was held at 35°C. The standard curve was prepped in LC-MS water and the range measured between 0.05-50 μg L⁻¹.

Mass spectrometry analysis was performed using electrospray ionization source (ESI). The MS source settings were as follows: spray voltage 3500 V, ion transfer tube temperature at 325 °C, vaporizer temperature at 350°C, sheath gas 35 (arbitrary units), auxiliary gas 10 (arbitrary units), sweep gas 0 (arbitrary units), collision gas at 1.5 mTorr, and Q1/Q3 resolution set to 0.7 Da.

Control samples and standards were performed every 20 samples during each run. Control standards included negative control (blank) and positive control (10 ppb standard) as well as duplicate and fortified samples. For each standard, a quantitative and qualitative ion transition were chosen for use with selected reaction monitoring (SRM).

Selective ion mode analysis

For selective ion mode (SIM) analysis, the following masses in Supporting Information, Dataset S1 were investigated because they are known to be produced by *M. aeruginosa* CPCC 300 (Racine et al. 2019). The gradient for the method consisted of 0.1% formic acid in water (A) and 0.1% formic acid in acetonitrile (B). At a flow rate of 0.35 mL min⁻¹, the gradient started at 0% B and was held for 1.0 min. It was then ramped to 60% B from 1.0-8.5 min. The column was washed at 98% B from 8.51–10.5 min at a flow rate 0.6 mL min⁻¹, then brought back down to original conditions to equilibrate. The total run time was 14 min. The injection volume was 10 μ L and the column oven temperature was held at 35 °C. Mass spectrometer source settings were the same as the quantitative method above. The SIM method was accomplished in Q3. The area under the m/z peak was calculated and reported.

Statistical analysis

For survival, growth, and neonate production

To test the magnitude of clonal variation, we used an LME (or GLM for binomial survival data) for each response with clone treated as a random effect and tested for significant differences in fit between models with and without clone included as random effect. Second, to determine whether the concentration of *Microcystis* influenced the magnitude of clonal variation we conducted Brown-Forsythe tests after generating means for each clone for each response variable. These Brown-Forsythe tests were done sequentially - meaning comparisons were between *Chlorella* and 2:1, then 2:1 and 1:1. Finally, to test for interactions between response variables across *Microcystis* concentrations we used linear models with concentration and clone as fixed effects (*glm* with binomial for survival, *lm* for growth, and *glm* with a gamma distribution for neonate production).

Association between genomic diversity, genomic divergence, and phenotypic distance

We tested for an association between the amount of genetic diversity found within clones and clonal survival and reproduction across each of the three diet conditions. To do so we regressed the total number of heterozygous sites and each response variable for each rearing condition. To test for associations between genomic diversity and phenotypic distance, we first generated distance matrices from mean values for all 20 clones. The genomic distance matrix was constructed from all polymorphic sites. The phenotypic distance matrix was constructed using neonate production, time to first brood, total broods, week one growth (day 0-7), week two growth (day 7-14), week one survival, and week two survival for each clone for each diet condition. A two-tailed Mantel test was then conducted to determine whether there was significant correlation between matrices that differed from 0.

Examining the effects of assays with reduced diversity

We sought to determine how the choice to conduct lower diversity assays might influence the estimate of the effects of the toxicant on reproduction and survival. For reproductive output we calculated means and standard errors for all 20 clones and for 15, 10, 5, 3, and 1 clonal diversity groups created by subsetting the full dataset 100 times at each diversity level. For survival, we calculated the 95% CI of survival probability using the means of all 20 clones. We then created subsamples containing 15, 10, 5, 3, and 1 clones and determined how frequently they would produce survival estimates within the 95% CI obtained from the full dataset.

Acknowledgements

This work was supported by the Canada First Research Excellence Fund through the Food from Thought Program (JMF), an NSERC Discovery Grant (JMF), the National Institute of General Medicine of the National Institute of Health (Award #1R35GM147264; SMR), a Banting Postdoctoral Fellowship (SMR), an NSERC Postdoctoral Fellowship (RSS), and a Liber Ero Postdoctoral Fellowship (RSS).

Supporting Information

Materials and Methods, Figures, Tables, References

References

1. Levin, S. A., Harwell, M. A., Kelly, J. R. & Kimball, K. D. Ecotoxicology: Problems and Approaches. in *Ecotoxicology: Problems and Approaches* (eds. Levin, S. A., Kelly, J. R., Harwell, M. A. & Kimball, K. D.) 3–7 (Springer New York, 1989).
2. Soares, A. M. V. M., Baird, D. J. & Calow, P. Interclonal variation in the performance of *Daphnia magna* Straus in chronic bioassays. *Environ. Toxicol. Chem.* **11**, 1477–1483 (1992).
3. Barata, C., Baird, D. J. & Soares, A. M. V. M. Determining genetic variability in the distribution of sensitivities to toxic stress among and within field populations of *Daphnia magna*. *Environ. Sci. Technol.* **36**, 3045–3049 (2002).
4. Clements, W. H. & Rohr, J. R. Community responses to contaminants: using basic ecological principles to predict ecotoxicological effects. *Environ. Toxicol. Chem.* **28**, 1789–1800 (2009).
5. Newman, M. C. *Fundamentals of ecotoxicology*. (CRC press, 2009).
6. Stark, J. D., Vargas, R. I. & Banks, J. E. Incorporating variability in point estimates in risk assessment: Bridging the gap between LC50 and population endpoints. *Environ. Toxicol. Chem.* **34**, 1683–1688 (2015).
7. Barata, C., Baird, D. J. & Markich, S. J. Influence of genetic and environmental factors on the tolerance of *Daphnia magna* Straus to essential and non-essential metals. *Aquat. Toxicol.* **42**, 115–137 (1998).
8. Relyea, R. & Hoverman, J. Assessing the ecology in ecotoxicology: a review and synthesis in freshwater systems. *Ecol. Lett.* **9**, 1157–1171 (2006).
9. Crutsinger, G. M. *et al.* Plant genotypic diversity predicts community structure and governs an ecosystem process. *Science* **313**, 966–968 (2006).
10. Barrett, R. D. H. & Schluter, D. Adaptation from standing genetic variation. *Trends Ecol. Evol.* **23**, 38–44 (2008).
11. Des Roches, S. *et al.* The ecological importance of intraspecific variation. *Nat Ecol Evol* **2**, 57–64 (2018).

12. Whitham, T. G., Allan, G. J., Cooper, H. F. & Shuster, S. M. Intraspecific Genetic Variation and Species Interactions Contribute to Community Evolution. *Annu. Rev. Ecol. Evol. Syst.* **51**, 587–612 (2020).
13. Rudman, S. M. *et al.* Direct observation of adaptive tracking on ecological time scales in *Drosophila*. *Science* **375**, eabj7484 (2022).
14. Weedall, G. D. *et al.* An Africa-wide genomic evolution of insecticide resistance in the malaria vector *Anopheles funestus* involves selective sweeps, copy number variations, gene conversion and transposons. *PLoS Genet.* **16**, e1008822 (2020).
15. Reid, N. M. *et al.* The genomic landscape of rapid repeated evolutionary adaptation to toxic pollution in wild fish. *Science* **354**, 1305–1308 (2016).
16. Lemaire, V. *et al.* Genotype × genotype interactions between the toxic cyanobacterium *Microcystis* and its grazer, the waterflea *Daphnia*. *Evol. Appl.* **5**, 168–182 (2012).
17. Hochmuth, J. D., De Meester, L., Pereira, C. M. S., Janssen, C. R. & De Schampelaere, K. A. C. Rapid Adaptation of a *Daphnia magna* Population to Metal Stress Is Associated with Heterozygote Excess. *Environ. Sci. Technol.* **49**, 9298–9307 (2015).
18. Jansen, M. *et al.* Experimental evolution reveals high insecticide tolerance in *Daphnia* inhabiting farmland ponds. *Evol. Appl.* **8**, 442–453 (2015).
19. Albert, C. H. *et al.* A multi-trait approach reveals the structure and the relative importance of intra- vs. interspecific variability in plant traits. *Funct. Ecol.* **24**, 1192–1201 (2010).
20. Lemmen, K. D., Butler, O. M., Koffel, T., Rudman, S. M. & Symons, C. C. Stoichiometric Traits Vary Widely Within Species: A Meta-Analysis of Common Garden Experiments. *Frontiers in Ecology and Evolution* **7**, (2019).
21. Bickham, J. W. The four cornerstones of Evolutionary Toxicology. *Ecotoxicology* **20**, 497–502 (2011).
22. Miner, B. E., De Meester, L., Pfrender, M. E., Lampert, W. & Hairston, N. G., Jr. Linking genes to communities and ecosystems: *Daphnia* as an ecogenomic model. *Proc. Biol. Sci.* **279**, 1873–1882 (2012).
23. Ebert, D. *Daphnia* as a versatile model system in ecology and evolution. *Evodevo* **13**, 16 (2022).
24. Hairston, N. G., Jr *et al.* Natural selection for grazer resistance to toxic cyanobacteria: evolution of phenotypic plasticity? *Evolution* **55**, 2203–2214 (2001).
25. Govaert, L., De Meester, L., Spaak, P. & Hairston, N. G. Eco-Evolutionary Dynamics in Freshwater Systems. in *Reference Module in Earth Systems and Environmental Sciences* (Elsevier, 2021).
26. Lynch, M., Wei, W., Ye, Z. & Pfrender, M. The Genome-wide Signature of Short-term Temporal Selection. *bioRxiv* 2023.04.28.538790 (2023) doi:10.1101/2023.04.28.538790.
27. Baird, D. J., Barber, I., Bradley, M., Soares, A. M. & Calow, P. A comparative study of genotype sensitivity to acute toxic stress using clones of *Daphnia magna* straus. *Ecotoxicol. Environ. Saf.* **21**, 257–265 (1991).
28. Rudman, S. M. *et al.* What genomic data can reveal about eco-evolutionary dynamics. *Nat Ecol Evol* **2**, 9–15 (2018).
29. Boyle, E. A., Li, Y. I. & Pritchard, J. K. An Expanded View of Complex Traits: From Polygenic to Omnigenic. *Cell* **169**, 1177–1186 (2017).
30. Harke, M. J. *et al.* A review of the global ecology, genomics, and biogeography of the toxic cyanobacterium, *Microcystis* spp. *Harmful Algae* **54**, 4–20 (2016).
31. Reid, A. J. *et al.* Emerging threats and persistent conservation challenges for freshwater biodiversity. *Biol. Rev. Camb. Philos. Soc.* **94**, 849–873 (2019).
32. Chorus, I. & Welker, M. *Toxic Cyanobacteria in Water: A Guide to Their Public Health Consequences, Monitoring and Management*. (CRC Press, 2021).

33. Shahmohamadloo, R. S. *et al.* Lake Erie fish safe to eat yet afflicted by algal hepatotoxins. *Sci. Total Environ.* **861**, 160474 (2023).
34. Shahmohamadloo, R. S. *et al.* Diseases and Disorders in Fish due to Harmful Algal Blooms. in *Climate Change on Diseases And Disorders Of Finfish In Cage Culture, 3rd Edition* (eds. Woo, P. T. K. & Subasinghe, R. P.) 387–429 (CAB International, 2023).
35. Shahmohamadloo, R. S. *et al.* Cyanotoxins accumulate in Lake St. Clair fish yet their fillets are safe to eat. *Sci. Total Environ.* **874**, 162381 (2023).
36. Hairston, N. G., Jr *et al.* Lake ecosystems: Rapid evolution revealed by dormant eggs. *Nature* **401**, 446–446 (1999).
37. Isanta-Navarro, J. *et al.* Reversed evolution of grazer resistance to cyanobacteria. *Nat. Commun.* **12**, 1945 (2021).
38. Sarnelle, O. & Wilson, A. E. Local adaptation of *Daphnia pulicariata* to toxic cyanobacteria. *Limnol. Oceanogr.* **50**, 1565–1570 (2005).
39. Chislock, M. F., Sarnelle, O., Jernigan, L. M. & Wilson, A. E. Do high concentrations of microcystin prevent *Daphnia* control of phytoplankton? *Water Res.* **47**, 1961–1970 (2013).
40. Shahmohamadloo, R. S., Poirier, D. G., Ortiz Almirall, X., Bhavsar, S. P. & Sibley, P. K. Assessing the toxicity of cell-bound microcystins on freshwater pelagic and benthic invertebrates. *Ecotoxicol. Environ. Saf.* **188**, 109945 (2020).
41. Ferrão-Filho, A. S., Azevedo, S. M. F. O. & DeMott, W. R. Effects of toxic and non-toxic cyanobacteria on the life history of tropical and temperate cladocerans. *Freshw. Biol.* **45**, 1–19 (2000).
42. Lürling, M. & van der Grinten, E. Life-history characteristics of *Daphnia* exposed to dissolved microcystin-LR and to the cyanobacterium *Microcystis aeruginosa* with and without microcystins. *Environ. Toxicol. Chem.* **22**, 1281–1287 (2003).
43. Yoshida, T., Jones, L. E., Ellner, S. P., Fussmann, G. F. & Hairston, N. G., Jr. Rapid evolution drives ecological dynamics in a predator-prey system. *Nature* **424**, 303–306 (2003).
44. Farkas, T. E., Mononen, T., Comeault, A. A., Hanski, I. & Nosil, P. Evolution of camouflage drives rapid ecological change in an insect community. *Curr. Biol.* **23**, 1835–1843 (2013).
45. Rudman, S. M. *et al.* Adaptive genetic variation mediates bottom-up and top-down control in an aquatic ecosystem. *Proc. Biol. Sci.* **282**, 20151234 (2015).
46. Coutellec, M.-A. & Barata, C. An introduction to evolutionary processes in ecotoxicology. *Ecotoxicology* **20**, 493–496 (2011).
47. Hawkins, N. J., Bass, C., Dixon, A. & Neve, P. The evolutionary origins of pesticide resistance. *Biol. Rev. Camb. Philos. Soc.* **94**, 135–155 (2019).
48. Bolker, B. M. *Ecological Models and Data in R*. (Princeton University Press, 2008).
49. McGill, B. J., Enquist, B. J., Weiher, E. & Westoby, M. Rebuilding community ecology from functional traits. *Trends Ecol. Evol.* **21**, 178–185 (2006).
50. Gustafsson, S. & Hansson, L.-A. Development of tolerance against toxic cyanobacteria in *Daphnia*. *Aquat. Ecol.* **38**, 37–44 (2004).
51. Forbes, V. E. Genetics and ecotoxicology—insights from the interface. in *Genetics and Ecotoxicology* 1–8 (CRC Press, 1998).
52. Mallet, J. The evolution of insecticide resistance: Have the insects won? *Trends Ecol. Evol.* **4**, 336–340 (1989).
53. Oziolor, E. M., Bickham, J. W. & Matson, C. W. Evolutionary toxicology in an omics world. *Evol. Appl.* **10**, 752–761 (2017).
54. Oziolor, E. M. *et al.* Adaptive introgression enables evolutionary rescue from extreme environmental pollution. *Science* **364**, 455–457 (2019).
55. Kimura, M. *The Neutral Theory of Molecular Evolution*. (Cambridge University Press, 1983).

56. Gaytán, B. D. & Vulpe, C. D. Functional toxicology: tools to advance the future of toxicity testing. *Front. Genet.* **5**, 110 (2014).
57. Breitholtz, M., Rudén, C., Hansson, S. O. & Bengtsson, B.-E. Ten challenges for improved ecotoxicological testing in environmental risk assessment. *Ecotoxicol. Environ. Saf.* **63**, 324–335 (2006).
58. Oziolor, E. M., DeSchampelaere, K., Lyon, D., Nacci, D. & Poynton, H. Evolutionary Toxicology-An Informational Tool for Chemical Regulation? *Environ. Toxicol. Chem.* **39**, 257–268 (2020).
59. Orsini, L., Spanier, K. I. & DE Meester, L. Genomic signature of natural and anthropogenic stress in wild populations of the waterflea *Daphnia magna*: validation in space, time and experimental evolution. *Mol. Ecol.* **21**, 2160–2175 (2012).
60. Kilham, S. S., Kreeger, D. A., Lynn, S. G., Goulden, C. E. & Herrera, L. COMBO: a defined freshwater culture medium for algae and zooplankton. *Hydrobiologia* **377**, 147–159 (1998).
61. Shahmohamadloo, R. S. *et al.* An efficient and affordable laboratory method to produce and sustain high concentrations of microcystins by *Microcystis aeruginosa*. *MethodsX* **6**, 2521–2535 (2019).
62. OECD. Test No. 211: *Daphnia magna* Reproduction Test, OECD Guidelines for the Testing of Chemicals, Section 2. in (OECD Publishing, 2012).
63. Rohrlack, T. *et al.* Ingestion of microcystins by *Daphnia*: Intestinal uptake and toxic effects. *Limnol. Oceanogr.* **50**, 440–448 (2005).

Figures

Figure 1. Multivariate a) genomic and b) phenotypic divergence between 20 *Daphnia magna* clones visualized using principal component analysis.




Figure 2. Intraspecific genetic variation in phenotypes of 20 *Daphnia magna* genotypes at the end of the 14-d chronic toxicity study. Phenotypes measured in parents were a) survival, b) growth from day 0 to 14, c) total number of neonates produced, and d) total number of broods produced. Cell concentration was standardized across three algal food treatments: non-toxic - “C”, moderately-toxic - “2:1 C:M”, and severely-toxic - “1:1 C:M”. Plots show means and standard errors for each clone when reared on each diet.




Figure 3. Panel A shows simulations of mean reproductive outputs for experimental designs with varying amounts of genetic diversity of *Daphnia magna* across three treatments of chlorella “C”, 2:1 chlorella:microcystis “2:1 C:M”, and 1:1 chlorella:microcystis “1:1 C:M”. Symbol and error bars represent the mean and standard deviation. A 95% confidence interval was calculated for survival using all 20 clones. Panel B shows simulations of how often estimates of survival fell within this 95% confidence interval when the data were subsetting to include a reduced number of genotypes.

