# Quadrature on the positive real line with quasi and pseudo orthogonal rational functions

Adhemar Bultheel
(joint work with P. González-Vera, E. Hendriksen, O. Njåstad)

Department of Computer Science
K.U.Leuven

SC2011
International Conference on Scientific Computing,
S. Margherita di Pula, Sardinia, Italy
October 10-14, 2011.

# Happy birthday Claude



# Happy birthday Claude



# Survey

- Quadrature on the positive real line and OP
- ORF (incl. OP /OLP)
- Quasi and Pseudo versions
- Quadrature with QORF and PORF
- Differences and similarities
- Numerical examples

# Gauss-type quadrature

- Consider integrals $I_\mu(f) = \int_0^\infty f(x) d\mu(x)$, $\mu > 0$.
- Gauss-type QF: inner product $\langle f, g \rangle_\mu = \int_0^\infty f(x) g(x) d\mu(x)$
- Construct OP $\{\varphi_n\}$: $\varphi_n \perp \mathcal{L}_{n-1} = \Pi_{n-1}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating polynomial for $f$ in these nodes $\Rightarrow$ weights $\{\lambda_{k,n} > 0\}_{k=1}^n$
- $$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n}).$$
- Equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} = \mathcal{L}_{2n-1} = \Pi_{2n-1}$

# Gauss-type quadrature

- Consider integrals $I_\mu(f) = \int_0^\infty f(x) d\mu(x)$, $\mu > 0$.
- Gauss-type QF: inner product $\langle f, g \rangle_\mu = \int_0^\infty f(x) g(x) d\mu(x)$
- Construct OP $\{\varphi_n\}$: $\varphi_n \perp \mathcal{L}_{n-1} = \Pi_{n-1}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating polynomial for $f$ in these nodes $\Rightarrow$ weights $\{\lambda_{k,n} > 0\}_{k=1}^n$
- $$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n}).$$
- Equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} = \mathcal{L}_{2n-1} = \Pi_{2n-1}$

# Gauss-type quadrature

- Consider integrals $I_\mu(f) = \int_0^\infty f(x) d\mu(x)$, $\mu > 0$.
- Gauss-type QF: inner product $\langle f, g \rangle_\mu = \int_0^\infty f(x) g(x) d\mu(x)$
- Construct OP $\{\varphi_n\}$: $\varphi_n \perp \mathcal{L}_{n-1} = \Pi_{n-1}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating polynomial for $f$ in these nodes $\Rightarrow$ weights $\{\lambda_{k,n} > 0\}_{k=1}^n$
- $$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n}).$$
- Equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} = \mathcal{L}_{2n-1} = \Pi_{2n-1}$

# Gauss-type quadrature

- Consider integrals $I_\mu(f) = \int_0^\infty f(x) d\mu(x)$, $\mu > 0$.
- Gauss-type QF: inner product $\langle f, g \rangle_\mu = \int_0^\infty f(x) g(x) d\mu(x)$
- Construct OP $\{\varphi_n\}$: $\varphi_n \perp \mathcal{L}_{n-1} = \Pi_{n-1}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating polynomial for $f$ in these nodes $\Rightarrow$ weights $\{\lambda_{k,n} > 0\}_{k=1}^n$
- $$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n}).$$
- Equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} = \mathcal{L}_{2n-1} = \Pi_{2n-1}$

# Gauss-type quadrature

- Consider integrals $I_\mu(f) = \int_0^\infty f(x) d\mu(x)$, $\mu > 0$.
- Gauss-type QF: inner product $\langle f, g \rangle_\mu = \int_0^\infty f(x) g(x) d\mu(x)$
- Construct OP $\{\varphi_n\}$: $\varphi_n \perp \mathcal{L}_{n-1} = \Pi_{n-1}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating polynomial for $f$ in these nodes $\Rightarrow$ weights $\{\lambda_{k,n} > 0\}_{k=1}^n$
- $$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n}).$$
- Equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} = \mathcal{L}_{2n-1} = \Pi_{2n-1}$

# Gauss-type quadrature

- Consider integrals $I_\mu(f) = \int_0^\infty f(x) d\mu(x)$, $\mu > 0$.
- Gauss-type QF: inner product $\langle f, g \rangle_\mu = \int_0^\infty f(x) g(x) d\mu(x)$
- Construct OP $\{\varphi_n\}$: $\varphi_n \perp \mathcal{L}_{n-1} = \Pi_{n-1}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating polynomial for $f$ in these nodes $\Rightarrow$ weights $\{\lambda_{k,n} > 0\}_{k=1}^n$
-

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n}).$$

- Equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} = \mathcal{L}_{2n-1} = \Pi_{2n-1}$

# Gauss-type quadrature

- Consider integrals $I_\mu(f) = \int_0^\infty f(x) d\mu(x)$, $\mu > 0$.
- Gauss-type QF: inner product $\langle f, g \rangle_\mu = \int_0^\infty f(x) g(x) d\mu(x)$
- Construct OP $\{\varphi_n\}$: $\varphi_n \perp \mathcal{L}_{n-1} = \Pi_{n-1}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating polynomial for $f$ in these nodes $\Rightarrow$ weights $\{\lambda_{k,n} > 0\}_{k=1}^n$
-
$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n}).$$
- Equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} = \mathcal{L}_{2n-1} = \Pi_{2n-1}$

# Orthogonal Rational Functions

- Introduce poles: define $\mathcal{L}_n = \{f_n/d_n : f_n \in \Pi_n\}$
- $d_n = r_1 r_2 \cdots r_n, \quad r_k(x) = \begin{cases} \zeta_k - x, & \text{if } -\infty < \zeta_k \leq 0 \\ 1, & \text{if } \zeta_k = -\infty \end{cases}$
- Orthogonalize $\varphi_n \perp \mathcal{L}_{n-1}$
- OP = all $\zeta$'s at $-\infty$
- OLP = all $\zeta$'s in $\{0, -\infty\}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n})$$

(equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} \neq^? \mathcal{L}_{2n-1}$)

# Orthogonal Rational Functions

- Introduce poles: define $\mathcal{L}_n = \{f_n/d_n : f_n \in \Pi_n\}$
- $d_n = r_1 r_2 \cdots r_n, \quad r_k(x) = \begin{cases} \zeta_k - x, & \text{if } -\infty < \zeta_k \leq 0 \\ 1, & \text{if } \zeta_k = -\infty \end{cases}$
- Orthogonalize $\varphi_n \perp \mathcal{L}_{n-1}$
- OP = all $\zeta$'s at $-\infty$
- OLP = all $\zeta$'s in $\{0, -\infty\}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n})$$

(equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} \neq^? \mathcal{L}_{2n-1}$)

# Orthogonal Rational Functions

- Introduce poles: define $\mathcal{L}_n = \{f_n/d_n : f_n \in \Pi_n\}$
- $d_n = r_1 r_2 \cdots r_n, \quad r_k(x) = \begin{cases} \zeta_k - x, & \text{if } -\infty < \zeta_k \leq 0 \\ 1, & \text{if } \zeta_k = -\infty \end{cases}$
- Orthogonalize $\varphi_n \perp \mathcal{L}_{n-1}$
- OP = all $\zeta$'s at $-\infty$
- OLP = all $\zeta$'s in $\{0, -\infty\}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n})$$

(equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} \neq^? \mathcal{L}_{2n-1}$)

# Orthogonal Rational Functions

- Introduce poles: define $\mathcal{L}_n = \{f_n/d_n : f_n \in \Pi_n\}$
- $d_n = r_1 r_2 \cdots r_n, \quad r_k(x) = \begin{cases} \zeta_k - x, & \text{if } -\infty < \zeta_k \leq 0 \\ 1, & \text{if } \zeta_k = -\infty \end{cases}$
- Orthogonalize $\varphi_n \perp \mathcal{L}_{n-1}$
- OP = all $\zeta$'s at $-\infty$
- OLP = all $\zeta$'s in $\{0, -\infty\}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n})$$

(equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} \neq^? \mathcal{L}_{2n-1}$)

# Orthogonal Rational Functions

- Introduce poles: define $\mathcal{L}_n = \{f_n/d_n : f_n \in \Pi_n\}$
- $d_n = r_1 r_2 \cdots r_n, \quad r_k(x) = \begin{cases} \zeta_k - x, & \text{if } -\infty < \zeta_k \leq 0 \\ 1, & \text{if } \zeta_k = -\infty \end{cases}$
- Orthogonalize $\varphi_n \perp \mathcal{L}_{n-1}$
- OP = all $\zeta$'s at $-\infty$
- OLP = all $\zeta$'s in $\{0, -\infty\}$
- Zeros $\varphi_n$: $\{x_{k,n}\}_{k=1}^n \subset \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n} f(x_{k,n})$$

(equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-1} \neq^? \mathcal{L}_{2n-1}$)

# QORF = Quasi ORF = ORF with a parameter

- Define

$$Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x), \quad \tau \in \mathbb{R} \cup \{\infty\}.$$

$$(Q_n(x,\infty) = \tfrac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x))$$

- $Q_n(x,\tau) \perp \mathcal{L}_{n-1}(\zeta_n) = \{\frac{p_{n-1}}{d_{n-1}} : p_{n-1}(\zeta_n) = 0\} \subset \mathcal{L}_{n-1}$
- Zeros $Q_n$: $\{x_{k,n}(\tau)\}_{k=1}^n \subset^? \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n}(\tau) f(x_{k,n}(\tau))$$

(equality for $f \in \mathcal{L}_{n-1} \cdot \mathcal{L}_{n-1} \neq^? \mathcal{L}_{2n-2}$)

- At most 1 node $x_{1,n}(\tau) < 0 \Rightarrow$ weight $\lambda_{1,n}(\tau) \not>^? 0$

# QORF = Quasi ORF = ORF with a parameter

- Define

$$Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x), \quad \tau \in \mathbb{R} \cup \{\infty\}.$$

$(Q_n(x,\infty) = \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x))$

- $Q_n(x,\tau) \perp \mathcal{L}_{n-1}(\zeta_n) = \{\frac{p_{n-1}}{d_{n-1}} : p_{n-1}(\zeta_n) = 0\} \subset \mathcal{L}_{n-1}$
- Zeros $Q_n$: $\{x_{k,n}(\tau)\}_{k=1}^n \subset^? \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n}(\tau) f(x_{k,n}(\tau))$$

(equality for $f \in \mathcal{L}_{n-1} \cdot \mathcal{L}_{n-1} \neq^? \mathcal{L}_{2n-2}$)

- At most 1 node $x_{1,n}(\tau) < 0 \Rightarrow$ weight $\lambda_{1,n}(\tau) \not>^? 0$

# QORF = Quasi ORF = ORF with a parameter

- Define

$$Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x), \quad \tau \in \mathbb{R} \cup \{\infty\}.$$

  $(Q_n(x,\infty) = \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x))$
- $Q_n(x,\tau) \perp \mathcal{L}_{n-1}(\zeta_n) = \{\frac{p_{n-1}}{d_{n-1}} : p_{n-1}(\zeta_n) = 0\} \subset \mathcal{L}_{n-1}$
- Zeros $Q_n$: $\{x_{k,n}(\tau)\}_{k=1}^n \subset^? \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n}(\tau) f(x_{k,n}(\tau))$$

  (equality for $f \in \mathcal{L}_{n-1} \cdot \mathcal{L}_{n-1} \neq^? \mathcal{L}_{2n-2}$)
- At most 1 node $x_{1,n}(\tau) < 0 \Rightarrow$ weight $\lambda_{1,n}(\tau) \not>^? 0$

# QORF = Quasi ORF = ORF with a parameter

- Define

$$Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x), \quad \tau \in \mathbb{R} \cup \{\infty\}.$$

  $(Q_n(x,\infty) = \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x))$
- $Q_n(x,\tau) \perp \mathcal{L}_{n-1}(\zeta_n) = \{\frac{p_{n-1}}{d_{n-1}} : p_{n-1}(\zeta_n) = 0\} \subset \mathcal{L}_{n-1}$
- Zeros $Q_n$: $\{x_{k,n}(\tau)\}_{k=1}^n \subset^? \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n}(\tau) f(x_{k,n}(\tau))$$

  (equality for $f \in \mathcal{L}_{n-1} \cdot \mathcal{L}_{n-1} \neq^? \mathcal{L}_{2n-2}$)
- At most 1 node $x_{1,n}(\tau) < 0 \Rightarrow$ weight $\lambda_{1,n}(\tau) \not>^? 0$

# QORF = Quasi ORF = ORF with a parameter

- Define

$$Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x), \quad \tau \in \mathbb{R} \cup \{\infty\}.$$

  $(Q_n(x,\infty) = \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x))$
- $Q_n(x,\tau) \perp \mathcal{L}_{n-1}(\zeta_n) = \{\frac{p_{n-1}}{d_{n-1}} : p_{n-1}(\zeta_n) = 0\} \subset \mathcal{L}_{n-1}$
- Zeros $Q_n$: $\{x_{k,n}(\tau)\}_{k=1}^n \subset^? \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n}(\tau) f(x_{k,n}(\tau))$$

  (equality for $f \in \mathcal{L}_{n-1} \cdot \mathcal{L}_{n-1} \neq^? \mathcal{L}_{2n-2}$)
- At most 1 node $x_{1,n}(\tau) < 0 \Rightarrow$ weight $\lambda_{1,n}(\tau) \not>^? 0$

# PORF = Pseudo ORF = ORF with a parameter

- Define

$$P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x), \quad \tau \in \mathbb{R} \cup \{\infty\}.$$

$(P_n(x,\infty) = \varphi_{n-1}(x))$

- $P_n(x,\tau) \perp \mathcal{L}_{n-2}$
- Zeros $P_n$: $\{x_{k,n}(\tau)\}_{k=1}^n \subset^? \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n}(\tau) f(x_{k,n}(\tau))$$

(equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-2} \neq^? \mathcal{L}_{2n-1}$)

- At most 1 node $x_{1,n}(\tau) < 0 \Rightarrow$ weight $\lambda_{1,n}(\tau) \not>^? 0$

# PORF = Pseudo ORF = ORF with a parameter

- Define

$$P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x), \quad \tau \in \mathbb{R} \cup \{\infty\}.$$

  $(P_n(x,\infty) = \varphi_{n-1}(x))$
- $P_n(x,\tau) \perp \mathcal{L}_{n-2}$
- Zeros $P_n$: $\{x_{k,n}(\tau)\}_{k=1}^n \subset^? \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n}(\tau) f(x_{k,n}(\tau))$$

  (equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-2} \neq^? \mathcal{L}_{2n-1}$)
- At most 1 node $x_{1,n}(\tau) < 0 \Rightarrow$ weight $\lambda_{1,n}(\tau) \not>^? 0$

# PORF = Pseudo ORF = ORF with a parameter

- Define

$$P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x), \quad \tau \in \mathbb{R} \cup \{\infty\}.$$

  $(P_n(x,\infty) = \varphi_{n-1}(x))$
- $P_n(x,\tau) \perp \mathcal{L}_{n-2}$
- Zeros $P_n$: $\{x_{k,n}(\tau)\}_{k=1}^n \subset^? \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n}(\tau) f(x_{k,n}(\tau))$$

  (equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-2} \neq^? \mathcal{L}_{2n-1}$)
- At most 1 node $x_{1,n}(\tau) < 0 \Rightarrow$ weight $\lambda_{1,n}(\tau) \not>^? 0$

# PORF = Pseudo ORF = ORF with a parameter

- Define

$$P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x), \quad \tau \in \mathbb{R} \cup \{\infty\}.$$

  $(P_n(x,\infty) = \varphi_{n-1}(x))$
- $P_n(x,\tau) \perp \mathcal{L}_{n-2}$
- Zeros $P_n$: $\{x_{k,n}(\tau)\}_{k=1}^n \subset^? \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n}(\tau) f(x_{k,n}(\tau))$$

  (equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-2} \neq^? \mathcal{L}_{2n-1}$)
- At most 1 node $x_{1,n}(\tau) < 0 \Rightarrow$ weight $\lambda_{1,n}(\tau) \not>^? 0$

# PORF = Pseudo ORF = ORF with a parameter

- Define

$$P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x), \quad \tau \in \mathbb{R} \cup \{\infty\}.$$

  $(P_n(x,\infty) = \varphi_{n-1}(x))$
- $P_n(x,\tau) \perp \mathcal{L}_{n-2}$
- Zeros $P_n$: $\{x_{k,n}(\tau)\}_{k=1}^n \subset^? \mathbb{R}_+ \Rightarrow$ nodes
- Interpolating RF for $f$ in zeros $\varphi_n \Rightarrow$

$$I_\mu(f) \approx I_{\mu_n}(f) = \sum_{k=1}^n \lambda_{k,n}(\tau) f(x_{k,n}(\tau))$$

  (equality for $f \in \mathcal{L}_n \cdot \mathcal{L}_{n-2} \neq^? \mathcal{L}_{2n-1}$)
- At most 1 node $x_{1,n}(\tau) < 0 \Rightarrow$ weight $\lambda_{1,n}(\tau) \not>^? 0$

# Zeros of QORF

**Theorem**

*The zeros of $Q_n(x,\tau)$ and $Q_{n-1}(x,\tau)$ interlace.*

- $Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x) = \frac{q_n(x,\tau)}{d_n(x)}$
  $q_n(x,\tau) = q_{0,n}(\tau) + q_{1,n}(\tau)x + \cdots + q_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $q_{0,n}(\tau) = f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}, \quad q_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_{n-1} \neq \infty$)
- $\mathcal{Z}\{Q_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau) \cdots x_{n,n}(\tau) = (-1)^n \frac{q_{0,n}(\tau)}{q_{n,n}(\tau)} = (-1)^n \frac{f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_{n-1} \neq \infty$)

# Zeros of QORF

**Theorem**

*The zeros of $Q_n(x,\tau)$ and $Q_{n-1}(x,\tau)$ interlace.*

- $Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x) = \frac{q_n(x,\tau)}{d_n(x)}$
  $q_n(x,\tau) = q_{0,n}(\tau) + q_{1,n}(\tau)x + \cdots + q_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $q_{0,n}(\tau) = f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}, \quad q_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_{n-1} \neq \infty$)
- $\mathcal{Z}\{Q_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau) \cdots x_{n,n}(\tau) = (-1)^n \frac{q_{0,n}(\tau)}{q_{n,n}(\tau)} = (-1)^n \frac{f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_{n-1} \neq \infty$)

# Zeros of QORF

**Theorem**

*The zeros of $Q_n(x,\tau)$ and $Q_{n-1}(x,\tau)$ interlace.*

- $Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x) = \frac{q_n(x,\tau)}{d_n(x)}$
  $q_n(x,\tau) = q_{0,n}(\tau) + q_{1,n}(\tau)x + \cdots + q_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $q_{0,n}(\tau) = f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}, \quad q_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_{n-1} \neq \infty$)
- $\mathcal{Z}\{Q_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau) \cdots x_{n,n}(\tau) = (-1)^n \frac{q_{0,n}(\tau)}{q_{n,n}(\tau)} = (-1)^n \frac{f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_{n-1} \neq \infty$)

# Zeros of QORF

**Theorem**

*The zeros of $Q_n(x,\tau)$ and $Q_{n-1}(x,\tau)$ interlace.*

- $Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x) = \frac{q_n(x,\tau)}{d_n(x)}$
  $q_n(x,\tau) = q_{0,n}(\tau) + q_{1,n}(\tau)x + \cdots + q_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $q_{0,n}(\tau) = f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}, \quad q_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_{n-1} \neq \infty$)
- $\mathcal{Z}\{Q_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau) \cdots x_{n,n}(\tau) = (-1)^n \frac{q_{0,n}(\tau)}{q_{n,n}(\tau)} = (-1)^n \frac{f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_{n-1} \neq \infty$)

# Zeros of QORF

**Theorem**

*The zeros of $Q_n(x,\tau)$ and $Q_{n-1}(x,\tau)$ interlace.*

- $Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x) = \frac{q_n(x,\tau)}{d_n(x)}$
  $q_n(x,\tau) = q_{0,n}(\tau) + q_{1,n}(\tau)x + \cdots + q_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $q_{0,n}(\tau) = f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}, \quad q_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_{n-1} \neq \infty$)
- $\mathcal{Z}\{Q_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau) \cdots x_{n,n}(\tau) = (-1)^n \frac{q_{0,n}(\tau)}{q_{n,n}(\tau)} = (-1)^n \frac{f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_{n-1} \neq \infty$)

# Zeros of QORF

**Theorem**

*The zeros of $Q_n(x,\tau)$ and $Q_{n-1}(x,\tau)$ interlace.*

- $Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x) = \frac{q_n(x,\tau)}{d_n(x)}$
  $q_n(x,\tau) = q_{0,n}(\tau) + q_{1,n}(\tau)x + \cdots + q_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $q_{0,n}(\tau) = f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}, \quad q_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_{n-1} \neq \infty$)
- $\mathcal{Z}\{Q_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau) \cdots x_{n,n}(\tau) = (-1)^n \frac{q_{0,n}(\tau)}{q_{n,n}(\tau)} = (-1)^n \frac{f_{0,n} - \tau\zeta_{n-1} f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_{n-1} \neq \infty$)

# Zeros of QORF

- $\Pi_1^n x_{k,n}(\tau) = (-1)^n \frac{f_{0,n} - \tau \zeta_{n-1} f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
- sign change at $\tau = \frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}$ and $\tau = -\frac{f_{n,n}}{f_{n-1,n-1}}$
  positive for $\tau = 0$ ($\Leftarrow Q_n(x,0) = \varphi_n(x)$)
  Hence all zeros are positive for $\tau \in (\frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}, -\frac{f_{n,n}}{f_{n-1,n-1}})$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_{n-1} \neq \infty$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_{n-1} = \infty$
  All zeros are positive for $\tau \in (\frac{f_{0,n}}{f_{0,n-1}}, +\infty)$
- For $\tau = \frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}$ or $\tau = \frac{f_{0,n}}{f_{0,n-1}}$ one zero at 0,
  i.e., a Radau-type QF.

# Zeros of QORF

- $\Pi_1^n x_{k,n}(\tau) = (-1)^n \frac{f_{0,n} - \tau \zeta_{n-1} f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
- sign change at $\tau = \frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}$ and $\tau = -\frac{f_{n,n}}{f_{n-1,n-1}}$
  positive for $\tau = 0$ ($\Leftarrow Q_n(x,0) = \varphi_n(x)$)
  Hence all zeros are positive for $\tau \in (\frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}, -\frac{f_{n,n}}{f_{n-1,n-1}})$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_{n-1} \neq \infty$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_{n-1} = \infty$
  All zeros are positive for $\tau \in (\frac{f_{0,n}}{f_{0,n-1}}, +\infty)$
- For $\tau = \frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}$ or $\tau = \frac{f_{0,n}}{f_{0,n-1}}$ one zero at 0,
  i.e., a Radau-type QF.

# Zeros of QORF

- $\Pi_1^n x_{k,n}(\tau) = (-1)^n \frac{f_{0,n} - \tau \zeta_{n-1} f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
- sign change at $\tau = \frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}$ and $\tau = -\frac{f_{n,n}}{f_{n-1,n-1}}$
  positive for $\tau = 0$ ($\Leftarrow Q_n(x,0) = \varphi_n(x)$)
  Hence all zeros are positive for $\tau \in (\frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}, -\frac{f_{n,n}}{f_{n-1,n-1}})$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_{n-1} \neq \infty$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_{n-1} = \infty$
  All zeros are positive for $\tau \in (\frac{f_{0,n}}{f_{0,n-1}}, +\infty)$
- For $\tau = \frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}$ or $\tau = \frac{f_{0,n}}{f_{0,n-1}}$ one zero at 0,
  i.e., a Radau-type QF.

# Zeros of QORF

- $\Pi_1^n x_{k,n}(\tau) = (-1)^n \frac{f_{0,n} - \tau \zeta_{n-1} f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
- sign change at $\tau = \frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}$ and $\tau = -\frac{f_{n,n}}{f_{n-1,n-1}}$
  positive for $\tau = 0$ ($\Leftarrow Q_n(x,0) = \varphi_n(x)$)
  Hence all zeros are positive for $\tau \in (\frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}, -\frac{f_{n,n}}{f_{n-1,n-1}})$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_{n-1} \neq \infty$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_{n-1} = \infty$
  All zeros are positive for $\tau \in (\frac{f_{0,n}}{f_{0,n-1}}, +\infty)$
- For $\tau = \frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}$ or $\tau = \frac{f_{0,n}}{f_{0,n-1}}$ one zero at 0,
  i.e., a Radau-type QF.

# Zeros of QORF

- $\Pi_1^n x_{k,n}(\tau) = (-1)^n \frac{f_{0,n} - \tau \zeta_{n-1} f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
- sign change at $\tau = \frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}$ and $\tau = -\frac{f_{n,n}}{f_{n-1,n-1}}$
  positive for $\tau = 0$ ($\Leftarrow Q_n(x,0) = \varphi_n(x)$)
  Hence all zeros are positive for $\tau \in (\frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}, -\frac{f_{n,n}}{f_{n-1,n-1}})$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_{n-1} \neq \infty$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_{n-1} = \infty$
  All zeros are positive for $\tau \in (\frac{f_{0,n}}{f_{0,n-1}}, +\infty)$
- For $\tau = \frac{f_{0,n}}{\zeta_{n-1} f_{0,n-1}}$ or $\tau = \frac{f_{0,n}}{f_{0,n-1}}$ one zero at 0,
  i.e., a Radau-type QF.

# Zeros of PORF (completely analogous)

**Theorem**

*The zeros of $P_n(x,\tau)$ and $P_{n-1}(x,\tau)$ interlace.*

- $P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x) = \frac{p_n(x,\tau)}{d_n(x)}$
  $p_n(x,\tau) = p_{0,n}(\tau) + p_{1,n}(\tau)x + \cdots + p_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $p_{0,n}(\tau) = f_{0,n} - \tau\zeta_n f_{0,n-1}, \quad p_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_n \neq \infty$)
- $\mathcal{Z}\{P_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau)\cdots x_{n,n}(\tau) = (-1)^n \frac{p_{0,n}(\tau)}{p_{n,n}(\tau)} = (-1)^n \frac{f_{0,n}-\tau\zeta_n f_{0,n-1}}{f_{n,n}+\tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_n \neq \infty$)

# Zeros of PORF (completely analogous)

## Theorem

*The zeros of $P_n(x,\tau)$ and $P_{n-1}(x,\tau)$ interlace.*

- $P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x) = \frac{p_n(x,\tau)}{d_n(x)}$
  $p_n(x,\tau) = p_{0,n}(\tau) + p_{1,n}(\tau)x + \cdots + p_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $p_{0,n}(\tau) = f_{0,n} - \tau\zeta_n f_{0,n-1}, \quad p_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_n \neq \infty$)
- $\mathcal{Z}\{P_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau) \cdots x_{n,n}(\tau) = (-1)^n \frac{p_{0,n}(\tau)}{p_{n,n}(\tau)} = (-1)^n \frac{f_{0,n} - \tau\zeta_n f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_n \neq \infty$)

# Zeros of PORF (completely analogous)

## Theorem

*The zeros of $P_n(x,\tau)$ and $P_{n-1}(x,\tau)$ interlace.*

- $P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x) = \frac{p_n(x,\tau)}{d_n(x)}$
  $p_n(x,\tau) = p_{0,n}(\tau) + p_{1,n}(\tau)x + \cdots + p_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $p_{0,n}(\tau) = f_{0,n} - \tau\zeta_n f_{0,n-1}, \quad p_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_n \neq \infty$)
- $\mathcal{Z}\{P_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau)\cdots x_{n,n}(\tau) = (-1)^n \frac{p_{0,n}(\tau)}{p_{n,n}(\tau)} = (-1)^n \frac{f_{0,n}-\tau\zeta_n f_{0,n-1}}{f_{n,n}+\tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_n \neq \infty$)

# Zeros of PORF (completely analogous)

**Theorem**

*The zeros of $P_n(x,\tau)$ and $P_{n-1}(x,\tau)$ interlace.*

- $P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x) = \frac{p_n(x,\tau)}{d_n(x)}$
  $p_n(x,\tau) = p_{0,n}(\tau) + p_{1,n}(\tau)x + \cdots + p_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $p_{0,n}(\tau) = f_{0,n} - \tau\zeta_n f_{0,n-1}, \quad p_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_n \neq \infty$)
- $\mathcal{Z}\{P_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau)\cdots x_{n,n}(\tau) = (-1)^n \frac{p_{0,n}(\tau)}{p_{n,n}(\tau)} = (-1)^n \frac{f_{0,n}-\tau\zeta_n f_{0,n-1}}{f_{n,n}+\tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_n \neq \infty$)

# Zeros of PORF (completely analogous)

## Theorem

*The zeros of $P_n(x,\tau)$ and $P_{n-1}(x,\tau)$ interlace.*

- $P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x) = \frac{p_n(x,\tau)}{d_n(x)}$
  $p_n(x,\tau) = p_{0,n}(\tau) + p_{1,n}(\tau)x + \cdots + p_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $p_{0,n}(\tau) = f_{0,n} - \tau\zeta_n f_{0,n-1}, \quad p_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_n \neq \infty$)
- $\mathcal{Z}\{P_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau)\cdots x_{n,n}(\tau) = (-1)^n \frac{p_{0,n}(\tau)}{p_{n,n}(\tau)} = (-1)^n \frac{f_{0,n}-\tau\zeta_n f_{0,n-1}}{f_{n,n}+\tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_n \neq \infty$)

# Zeros of PORF (completely analogous)

**Theorem**

*The zeros of $P_n(x,\tau)$ and $P_{n-1}(x,\tau)$ interlace.*

- $P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x) = \frac{p_n(x,\tau)}{d_n(x)}$
  $p_n(x,\tau) = p_{0,n}(\tau) + p_{1,n}(\tau)x + \cdots + p_{n,n}(\tau)x^n$
- $\varphi_k = \frac{f_k}{d_k}, \quad f_k(x) = f_{0,k} + f_{1,k}x + \cdots + f_{k,k}x^k$
- $p_{0,n}(\tau) = f_{0,n} - \tau\zeta_n f_{0,n-1}, \quad p_{n,n}(\tau) = f_{n,n} + \tau f_{n-1,n-1}$
  (if $\zeta_n \neq \infty$)
- $\mathcal{Z}\{P_n(x,\tau)\} = \{x_{1,n}(\tau), \ldots, x_{n,n}(\tau)\}$
- $x_{1,n}(\tau)\cdots x_{n,n}(\tau) = (-1)^n \frac{p_{0,n}(\tau)}{p_{n,n}(\tau)} = (-1)^n \frac{f_{0,n}-\tau\zeta_n f_{0,n-1}}{f_{n,n}+\tau f_{n-1,n-1}}$
  (proper sign normalization and $\zeta_n \neq \infty$)

# Zeros of PORF

- $\Pi_1^n x_{k,n}(\tau) = (-1)^n \frac{f_{0,n} - \tau\zeta_n f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
- sign change at $\tau = \frac{f_{0,n}}{\zeta_n f_{0,n-1}}$ and $\tau = -\frac{f_{n,n}}{f_{n-1,n-1}}$
  positive for $\tau = 0$ ($\Leftarrow P_n(x,0) = \varphi_n(x)$)
  Hence all zeros are positive for $\tau \in (\frac{f_{0,n}}{\zeta_n f_{0,n-1}}, -\frac{f_{n,n}}{f_{n-1,n-1}})$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_n \neq \infty$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_n = \infty$
  All zeros are positive for $\tau \in (\frac{f_{0,n}}{f_{0,n-1}}, +\infty)$
- For $\tau = \frac{f_{0,n}}{\zeta_n f_{0,n-1}}$ or $\tau = \frac{f_{0,n}}{f_{0,n-1}}$ one zero at 0,
  i.e., a Radau-type QF.

# Zeros of PORF

- $\Pi_1^n x_{k,n}(\tau) = (-1)^n \frac{f_{0,n} - \tau \zeta_n f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
- sign change at $\tau = \frac{f_{0,n}}{\zeta_n f_{0,n-1}}$ and $\tau = -\frac{f_{n,n}}{f_{n-1,n-1}}$
  positive for $\tau = 0$ ($\Leftarrow P_n(x,0) = \varphi_n(x)$)
  Hence all zeros are positive for $\tau \in (\frac{f_{0,n}}{\zeta_n f_{0,n-1}}, -\frac{f_{n,n}}{f_{n-1,n-1}})$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_n \neq \infty$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_n = \infty$
  All zeros are positive for $\tau \in (\frac{f_{0,n}}{f_{0,n-1}}, +\infty)$
- For $\tau = \frac{f_{0,n}}{\zeta_n f_{0,n-1}}$ or $\tau = \frac{f_{0,n}}{f_{0,n-1}}$ one zero at 0,
  i.e., a Radau-type QF.

# Zeros of PORF

- $\Pi_1^n x_{k,n}(\tau) = (-1)^n \frac{f_{0,n} - \tau\zeta_n f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
- sign change at $\tau = \frac{f_{0,n}}{\zeta_n f_{0,n-1}}$ and $\tau = -\frac{f_{n,n}}{f_{n-1,n-1}}$
  positive for $\tau = 0$ ($\Leftarrow P_n(x,0) = \varphi_n(x)$)
  Hence all zeros are positive for $\tau \in (\frac{f_{0,n}}{\zeta_n f_{0,n-1}}, -\frac{f_{n,n}}{f_{n-1,n-1}})$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_n \neq \infty$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_n = \infty$
  All zeros are positive for $\tau \in (\frac{f_{0,n}}{f_{0,n-1}}, +\infty)$
- For $\tau = \frac{f_{0,n}}{\zeta_n f_{0,n-1}}$ or $\tau = \frac{f_{0,n}}{f_{0,n-1}}$ one zero at 0,
  i.e., a Radau-type QF.

# Zeros of PORF

- $\Pi_1^n x_{k,n}(\tau) = (-1)^n \frac{f_{0,n} - \tau \zeta_n f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
- sign change at $\tau = \frac{f_{0,n}}{\zeta_n f_{0,n-1}}$ and $\tau = -\frac{f_{n,n}}{f_{n-1,n-1}}$
  positive for $\tau = 0$ ($\Leftarrow P_n(x,0) = \varphi_n(x)$)
  Hence all zeros are positive for $\tau \in (\frac{f_{0,n}}{\zeta_n f_{0,n-1}}, -\frac{f_{n,n}}{f_{n-1,n-1}})$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_n \neq \infty$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_n = \infty$
  All zeros are positive for $\tau \in (\frac{f_{0,n}}{f_{0,n-1}}, +\infty)$
- For $\tau = \frac{f_{0,n}}{\zeta_n f_{0,n-1}}$ or $\tau = \frac{f_{0,n}}{f_{0,n-1}}$ one zero at 0,
  i.e., a Radau-type QF.

# Zeros of PORF

- $\Pi_1^n x_{k,n}(\tau) = (-1)^n \frac{f_{0,n} - \tau\zeta_n f_{0,n-1}}{f_{n,n} + \tau f_{n-1,n-1}}$
- sign change at $\tau = \frac{f_{0,n}}{\zeta_n f_{0,n-1}}$ and $\tau = -\frac{f_{n,n}}{f_{n-1,n-1}}$
  positive for $\tau = 0$ ($\Leftarrow P_n(x,0) = \varphi_n(x)$)
  Hence all zeros are positive for $\tau \in (\frac{f_{0,n}}{\zeta_n f_{0,n-1}}, -\frac{f_{n,n}}{f_{n-1,n-1}})$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_n \neq \infty$
- Here is the result plotting $x_{k,n}(\tau)$, $\tau \in \mathbb{R}$ for $\zeta_n = \infty$
  All zeros are positive for $\tau \in (\frac{f_{0,n}}{f_{0,n-1}}, +\infty)$
- For $\tau = \frac{f_{0,n}}{\zeta_n f_{0,n-1}}$ or $\tau = \frac{f_{0,n}}{f_{0,n-1}}$ one zero at 0,
  i.e., a Radau-type QF.

# positive weights

**Theorem**

*If the zero* $x_{1,n}(\tau) \geq 0$ *or* $x_{1,n}(\tau) \neq \zeta_k$, $\forall k$
*then the weights of the QF* $\lambda_{k,n}(\tau) > 0$, $\forall k$ *QORF.*
*then the weights of the QF* $\lambda_{k,n}(\tau) > 0$, *if* $x_{k,n}(\tau) > 0$ *PORF.*

Skip arguments

- ORF of 2nd kind: $\sigma_n(x) = \int_0^\infty \frac{\varphi_n(t)-\varphi_n(x)}{t-x} dt.$
- QORF2: $T_n(x,\tau) = \sigma_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \sigma_{n-1}(x).$
- PORF2: $S_n(x,\tau) = \sigma_n(x) - \tau \sigma_{n-1}(x).$

# positive weights

**Theorem**

*If the zero* $x_{1,n}(\tau) \geq 0$ *or* $x_{1,n}(\tau) \neq \zeta_k$, $\forall k$
*then the weights of the QF* $\lambda_{k,n}(\tau) > 0$, $\forall k$ *QORF.*
*then the weights of the QF* $\lambda_{k,n}(\tau) > 0$, *if* $x_{k,n}(\tau) > 0$ *PORF.*

Skip arguments

- ORF of 2nd kind: $\sigma_n(x) = \int_0^\infty \frac{\varphi_n(t)-\varphi_n(x)}{t-x} dt.$
- QORF2: $T_n(x,\tau) = \sigma_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \sigma_{n-1}(x).$
- PORF2: $S_n(x,\tau) = \sigma_n(x) - \tau \sigma_{n-1}(x).$

# positive weights

**Theorem**

*If the zero* $x_{1,n}(\tau) \geq 0$ *or* $x_{1,n}(\tau) \neq \zeta_k$, $\forall k$
*then the weights of the QF* $\lambda_{k,n}(\tau) > 0$, $\forall k$ *QORF.*
*then the weights of the QF* $\lambda_{k,n}(\tau) > 0$, *if* $x_{k,n}(\tau) > 0$ *PORF.*

Skip arguments

- ORF of 2nd kind: $\sigma_n(x) = \int_0^\infty \frac{\varphi_n(t)-\varphi_n(x)}{t-x} dt.$
- QORF2: $T_n(x,\tau) = \sigma_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \sigma_{n-1}(x).$
- PORF2: $S_n(x,\tau) = \sigma_n(x) - \tau \sigma_{n-1}(x).$

# positive weights

**Theorem**

*If the zero* $x_{1,n}(\tau) \geq 0$ *or* $x_{1,n}(\tau) \neq \zeta_k$, $\forall k$
*then the weights of the QF* $\lambda_{k,n}(\tau) > 0$, $\forall k$ *QORF.*
*then the weights of the QF* $\lambda_{k,n}(\tau) > 0$, *if* $x_{k,n}(\tau) > 0$ *PORF.*

Skip arguments

- ORF of 2nd kind: $\sigma_n(x) = \int_0^\infty \frac{\varphi_n(t)-\varphi_n(x)}{t-x} dt.$
- QORF2: $T_n(x,\tau) = \sigma_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \sigma_{n-1}(x).$
- PORF2: $S_n(x,\tau) = \sigma_n(x) - \tau \sigma_{n-1}(x).$

# positive weights

Skip

- QORF2: $T_n(x,\tau) = \sigma_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \sigma_{n-1}(x)$
- Lagrange interpolant in $\mathcal{L}_{n-1}$: $\sum_k L_{k,n}(x,\tau) f(x_{k,n}(\tau))$
  $L_{k,n}(x,\tau) \in \mathcal{L}_{n-1}$, $L_{k,n}(x_{j,n}(\tau)) = \delta_{j,k}$
- QF exact in $\mathcal{L}_{n-1} \cdot \mathcal{L}_{n-1}$
- $I_\mu(L_{k,n}(\cdot,\tau)) = \sum_j I_\mu(L_{j,n}(\cdot,\tau)) L_{k,n}(x_{j,k}(\tau),\tau)$
  $= \lambda_{j,n}(\tau)\delta_{j,k} = \lambda_{k,n}(\tau)$
- $0 < I_\mu([L_{k,n}(\cdot,\tau)]^2) = \sum_j I_\mu(L_{j,n}(\cdot,\tau)) [L_{k,n}(x_{j,k}(\tau),\tau)]^2$
  $= \lambda_{j,n}(\tau)\delta_{j,k} = \lambda_{k,n}(\tau)$

# positive weights

Skip

- QORF2: $T_n(x,\tau) = \sigma_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \sigma_{n-1}(x)$
- Lagrange interpolant in $\mathcal{L}_{n-1}$: $\sum_k L_{k,n}(x,\tau) f(x_{k,n}(\tau))$
  $L_{k,n}(x,\tau) \in \mathcal{L}_{n-1}$, $L_{k,n}(x_{j,n}(\tau)) = \delta_{j,k}$
- QF exact in $\mathcal{L}_{n-1} \cdot \mathcal{L}_{n-1}$
- $I_\mu(L_{k,n}(\cdot,\tau)) = \sum_j I_\mu(L_{j,n}(\cdot,\tau)) L_{k,n}(x_{j,k}(\tau),\tau)$
  $= \lambda_{j,n}(\tau)\delta_{j,k} = \lambda_{k,n}(\tau)$
- $0 < I_\mu([L_{k,n}(\cdot,\tau)]^2) = \sum_j I_\mu(L_{j,n}(\cdot,\tau))[L_{k,n}(x_{j,k}(\tau),\tau)]^2$
  $= \lambda_{j,n}(\tau)\delta_{j,k} = \lambda_{k,n}(\tau)$

# positive weights

Skip

- QORF2: $T_n(x,\tau) = \sigma_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \sigma_{n-1}(x)$
- Lagrange interpolant in $\mathcal{L}_{n-1}$: $\sum_k L_{k,n}(x,\tau) f(x_{k,n}(\tau))$
  $L_{k,n}(x,\tau) \in \mathcal{L}_{n-1}$, $L_{k,n}(x_{j,n}(\tau)) = \delta_{j,k}$
- QF exact in $\mathcal{L}_{n-1} \cdot \mathcal{L}_{n-1}$
- $I_\mu(L_{k,n}(\cdot,\tau)) = \sum_j I_\mu(L_{j,n}(\cdot,\tau)) L_{k,n}(x_{j,k}(\tau),\tau)$
  $= \lambda_{j,n}(\tau)\delta_{j,k} = \lambda_{k,n}(\tau)$
- $0 < I_\mu([L_{k,n}(\cdot,\tau)]^2) = \sum_j I_\mu(L_{j,n}(\cdot,\tau))[L_{k,n}(x_{j,k}(\tau),\tau)]^2$
  $= \lambda_{j,n}(\tau)\delta_{j,k} = \lambda_{k,n}(\tau)$

# positive weights

Skip

- QORF2: $T_n(x,\tau) = \sigma_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \sigma_{n-1}(x)$
- Lagrange interpolant in $\mathcal{L}_{n-1}$: $\sum_k L_{k,n}(x,\tau) f(x_{k,n}(\tau))$
  $L_{k,n}(x,\tau) \in \mathcal{L}_{n-1}$, $L_{k,n}(x_{j,n}(\tau)) = \delta_{j,k}$
- QF exact in $\mathcal{L}_{n-1} \cdot \mathcal{L}_{n-1}$
- $I_\mu(L_{k,n}(\cdot,\tau)) = \sum_j I_\mu(L_{j,n}(\cdot,\tau)) L_{k,n}(x_{j,k}(\tau),\tau)$
  $= \lambda_{j,n}(\tau)\delta_{j,k} = \lambda_{k,n}(\tau)$
- $0 < I_\mu([L_{k,n}(\cdot,\tau)]^2) = \sum_j I_\mu(L_{j,n}(\cdot,\tau))[L_{k,n}(x_{j,k}(\tau),\tau)]^2$
  $= \lambda_{j,n}(\tau)\delta_{j,k} = \lambda_{k,n}(\tau)$

# positive weights

Skip

- QORF2: $T_n(x,\tau) = \sigma_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \sigma_{n-1}(x)$
- Lagrange interpolant in $\mathcal{L}_{n-1}$: $\sum_k L_{k,n}(x,\tau) f(x_{k,n}(\tau))$
  $L_{k,n}(x,\tau) \in \mathcal{L}_{n-1}$, $L_{k,n}(x_{j,n}(\tau)) = \delta_{j,k}$
- QF exact in $\mathcal{L}_{n-1} \cdot \mathcal{L}_{n-1}$
- $I_\mu(L_{k,n}(\cdot,\tau)) = \sum_j I_\mu(L_{j,n}(\cdot,\tau)) L_{k,n}(x_{j,k}(\tau),\tau)$
  $= \lambda_{j,n}(\tau)\delta_{j,k} = \lambda_{k,n}(\tau)$
- $0 < I_\mu([L_{k,n}(\cdot,\tau)]^2) = \sum_j I_\mu(L_{j,n}(\cdot,\tau))[L_{k,n}(x_{j,k}(\tau),\tau)]^2$
  $= \lambda_{j,n}(\tau)\delta_{j,k} = \lambda_{k,n}(\tau)$

# positive weights

Skip

- PORF2: $S_n(x,\tau) = \sigma_n(x) - \tau\sigma_{n-1}(x)$.

# positive weights

Skip

- PORF2: $S_n(x,\tau) = \sigma_n(x) - \tau\sigma_{n-1}(x)$.
- previous proof does not work: $[L_{k,n}]^2 \notin \mathcal{L}_n \cdot \mathcal{L}_{n-2}$

# positive weights

Skip

- PORF2: $S_n(x,\tau) = \sigma_n(x) - \tau\sigma_{n-1}(x)$.
- previous proof does not work: $[L_{k,n}]^2 \notin \mathcal{L}_n \cdot \mathcal{L}_{n-2}$
- weights $\lambda_{k,n}(\tau) = \left.\frac{S_n(x,\tau)}{P_n'(x,\tau)}\right|_{x=x_{k,n}(\tau)}$.

# positive weights


Skip

- PORF2: $S_n(x,\tau) = \sigma_n(x) - \tau\sigma_{n-1}(x)$.
- previous proof does not work: $[L_{k,n}]^2 \notin \mathcal{L}_n \cdot \mathcal{L}_{n-2}$
- weights $\lambda_{k,n}(\tau) = \left.\frac{S_n(x,\tau)}{P_n'(x,\tau)}\right|_{x=x_{k,n}(\tau)}$.
- $\mathcal{Z}(P_n(x,\tau))$ and $\mathcal{Z}(S_n(x,\tau))$ interlace

# positive weights


Skip

- PORF2: $S_n(x,\tau) = \sigma_n(x) - \tau\sigma_{n-1}(x)$.
- previous proof does not work: $[L_{k,n}]^2 \notin \mathcal{L}_n \cdot \mathcal{L}_{n-2}$
- weights $\lambda_{k,n}(\tau) = \left.\frac{S_n(x,\tau)}{P_n'(x,\tau)}\right|_{x=x_{k,n}(\tau)}$.
- $\mathcal{Z}(P_n(x,\tau))$ and $\mathcal{Z}(S_n(x,\tau))$ interlace
- $\operatorname{sgn}(P_n'(x,\tau)) = \operatorname{sgn}(S_n(x,\tau))$ for $x > x_{n,n}(\tau)$
  $\Rightarrow \operatorname{sgn}(P_n'(x_{k,n}(\tau),\tau)) = \operatorname{sgn}(S_n(x_{k,n}(\tau),\tau))$, $\forall k$



# positive weights

- This may not work if $\exists k$: $x_{1,n}(\tau) < \zeta_k < 0$



# Computation

- Recurrence

$$\varphi_n(x) \sim \frac{x^{\gamma_{n-1}} - c_n r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x) - d_{k-1} \frac{r_{n-2}(x)}{r_n(x)} \varphi_{n-2}(x)$$

$\gamma_n = 1$ if $\zeta_n = \infty$, $\gamma_n = 0$ otherwise

$$J_n = \operatorname{tridiag} \begin{pmatrix} d_1, d_2, \ldots, d_{n-1} \\ c_1, c_2, c_3, \ldots, c_{n-1}, c_n \\ d_1, d_2, \ldots, d_{n-1} \end{pmatrix}$$

$I_n^\gamma = \operatorname{diag}(\gamma_0, \gamma_1, \ldots, \gamma_{n-1}), \quad I_n^{1-\gamma} = I_n - I_n^\gamma$

$Z_n = \operatorname{diag}(\rho_0, \rho_1, \ldots, \rho_{n-1})$

$\rho_n = \zeta_n$ if $\zeta_n \neq \infty$, $\rho_n = 1$ otherwise

$A_n = J_n Z_n - I_n^{1-\gamma}, \quad B_n = J_n I_n^{1-\gamma} + I_n^\gamma.$

# Computation

## Theorem

*Consider the pencil* $(A_n, B_n)$ *as above.*
*Then the nodes* $x_{k,n}$ *of the rational Gauss QF are its eigenvalues. If* $\mathbf{e}_k = [e_{1,n}(k), \ldots, e_{n,n}(k)]^T$ *is the eigenvector corresponding to* $x_{k,n}(\tau = 0)$ *with* $\mathbf{e}_k^T \mathbf{e}_k = 1$*, then the corresponding weight in the rational Gauss QF is* $\lambda_{k,n}(\tau = 0) = e_{1,n}(k)^2$.

# Computation

## Theorem

*Consider the pencil* $(A_n, B_n)$ *as above.*
*Then the nodes* $x_{k,n}$ *of the rational Gauss QF are its eigenvalues. If* $\mathbf{e}_k = [e_{1,n}(k), \ldots, e_{n,n}(k)]^T$ *is the eigenvector corresponding to* $x_{k,n}(\tau = 0)$ *with* $\mathbf{e}_k^T \mathbf{e}_k = 1$*, then the corresponding weight in the rational Gauss QF is* $\lambda_{k,n}(\tau = 0) = e_{1,n}(k)^2$*.*

- Note that in the polynomial case all $\gamma_k = 1$, $I_n^\gamma = I_n$, $Z_n = I_n$, and thus $(A_n, B_n) = (J_n Z_n - I_n^{1-\gamma}, J_n I_n^{1-\gamma} + I_n^\gamma) = (J_n, I_n)$

# Computation QORF

- Use $Q_n(x,\tau) = \varphi_n(x) - \tau \frac{r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x)$ to get
  $Q_n(x,\tau) \sim \frac{x^{\gamma_n - 1} - c_n(\tau) r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x) - d_{n-1} \frac{r_{n-2}(x)}{r_n(x)} \varphi_{n-2}(x)$
  $c_n(\tau)$ is a simple modification of $c_n$ involving $\tau$.
  The same procedure as in the usual rational Gauss QF applies for nodes and weights.


# Computation PORF

- Use $P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x)$ to get
  $Q_n(x,\tau) \sim \frac{x^{\gamma_{n-1}} - c'_n(\tau) r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x) - d'_{n-1} \frac{r_{n-2}(x)}{r_n(x)} \varphi_{n-2}(x)$
  Now $c'_n$ and $d'_{n-1}$ are more complex modifications of $c_n$ and $d_{n-1}$ involving $\tau$ but also $\zeta_n$ and $\zeta_{n-1}$.
  Moreover the symmetry of the pencil is lost.
  The computation of the weights from the eigenvectors becomes a nonlinear procedure:
  (we drop $n$ and $\tau$ from the notation):
  $1/\lambda_k(\omega) = \sum_{i=0}^{n-2} \varphi_i^2(x_k) + \omega^{-1}\varphi_{n-1}^2(x_k)$, where
  $\sum_{k=1}^{n} \lambda_k(\omega) = 1$ (normalized measure: $\int d\mu = 1$).

# Computation PORF


- Use $P_n(x,\tau) = \varphi_n(x) - \tau\varphi_{n-1}(x)$ to get
  $Q_n(x,\tau) \sim \frac{x^{\gamma_{n-1}} - c'_n(\tau) r_{n-1}(x)}{r_n(x)} \varphi_{n-1}(x) - d'_{n-1} \frac{r_{n-2}(x)}{r_n(x)} \varphi_{n-2}(x)$
  Now $c'_n$ and $d'_{n-1}$ are more complex modifications of $c_n$ and $d_{n-1}$ involving $\tau$ but also $\zeta_n$ and $\zeta_{n-1}$.
  Moreover the symmetry of the pencil is lost.
  The computation of the weights from the eigenvectors becomes a nonlinear procedure:
  (we drop $n$ and $\tau$ from the notation):
  $1/\lambda_k(\omega) = \sum_{i=0}^{n-2} \varphi_i^2(x_k) + \omega^{-1}\varphi_{n-1}^2(x_k)$, where
  $\sum_{k=1}^{n} \lambda_k(\omega) = 1$ (normalized measure: $\int d\mu = 1$).
- It's a mess

# Example 1: Laguerre polynomials, PORF





$d\mu(x) = \exp(-x)dx,$
$n = 6,$
all $\zeta_k = \infty$

# Example 2, PORF





$$d\mu(x) = x^{10}\exp(-x)dx,$$
$$n = 6,$$
$$\zeta_k = -k,\ k = 1, 2, 3, 4, 5, 6$$

# Example 3, PORF




$$d\mu(x) = x^{10}\exp(-x^2)dx,$$
$$n = 6,$$
$$\zeta_k = \infty, -1, \infty, -3, \infty, -5$$

# Example 4, PORF




$$d\mu(x) = x^{10}\exp(-x^2)dx,$$
$$n = 6,$$
$$\zeta_k = -1, \infty, -3, \infty, -5, \infty$$

# The end


Thank you

# Reference



A. Bultheel, P. González-Vera, E. Hendriksen, and O. Njåstad. Quadratures associated with pseudo-orthogonal rational functions on the real half line with poles in $[-\infty, 0]$. Technical Report TW587, Department of Computer Science, K.U. Leuven, March 2011.

QORF $\tau(x_*) = \dfrac{f_n(x_*)}{r_{n-1}(x_*) f_{n-1}(x_*)} \quad (n = 4,\ \zeta_{n-1} \neq \infty)$





square $= \zeta_{n-1}$; circles $= \mathcal{Z}(\varphi_n)$; crosses $= \mathcal{Z}(\varphi_{n-1})$

note interlacing

Back


QORF $\tau(x_*) = \dfrac{f_n(x_*)}{r_{n-1}(x_*) f_{n-1}(x_*)}$ $\quad (n = 4,\ \zeta_{n-1} = \infty)$




circles $= \mathcal{Z}(\varphi_n)$; crosses $= \mathcal{Z}(\varphi_{n-1})$
note interlacing

Back

PORF $\tau(x_*) = \dfrac{f_n(x_*)}{r_n(x_*) f_{n-1}(x_*)} \quad (n = 4,\ \zeta_n \neq \infty)$





square $= \zeta_n$; circles $= \mathcal{Z}(\varphi_n)$; crosses $= \mathcal{Z}(\varphi_{n-1})$

note interlacing

Back

PORF $\tau(x_*) = \dfrac{f_n(x_*)}{r_n(x_*) f_{n-1}(x_*)} \quad (n = 4,\ \zeta_n = \infty)$




circles $= \mathcal{Z}(\varphi_n)$; crosses $= \mathcal{Z}(\varphi_{n-1})$
note interlacing

Back